

How often does a cubic hypersurface have a point?

Christopher Keyes (King's College London)

Show and Tell

15 March 2024

# Cubic hypersurfaces

A **cubic hypersurface**  $X_f \subset \mathbb{P}^n$  is cut out by a cubic form  $f$

$$X_f: f(x_0, \dots, x_n) = \sum_{0 \leq i \leq j \leq k \leq n} a_{ijk} x_i x_j x_k = 0.$$

# Cubic hypersurfaces

A **cubic hypersurface**  $X_f \subset \mathbb{P}^n$  is cut out by a cubic form  $f$

$$X_f: f(x_0, \dots, x_n) = \sum_{0 \leq i \leq j \leq k \leq n} a_{ijk} x_i x_j x_k = 0.$$

## Question

For **fixed**  $f$ , how many rational points are there on  $X_f$ ?

# Cubic hypersurfaces

A **cubic hypersurface**  $X_f \subset \mathbb{P}^n$  is cut out by a cubic form  $f$

$$X_f: f(x_0, \dots, x_n) = \sum_{0 \leq i \leq j \leq k \leq n} a_{ijk} x_i x_j x_k = 0.$$

## Question

For **fixed**  $f$ , how many rational points are there on  $X_f$ ?

## Question

As  $f$  **varies**, how often does  $X_f$  have a rational point?

# Counting points

## Question

*For fixed  $f$ , how many rational points are there on  $X_f$ ?*

Write  $P \in X_f(\mathbb{Q})$  as  $P = [x_0, \dots, x_n]$  with  $x_i \in \mathbb{Z}$  coprime.

Define **height** of a point

$$h(P) = \max_i \{|x_i|\}.$$

# Counting points

## Question

*For fixed  $f$ , how many rational points are there on  $X_f$ ?*

Write  $P \in X_f(\mathbb{Q})$  as  $P = [x_0, \dots, x_n]$  with  $x_i \in \mathbb{Z}$  coprime.

Define **height** of a point

$$h(P) = \max_i \{|x_i|\}.$$

Natural point-counting function

$$N_f(B) = \#\{P \in X_f(\mathbb{Q}) \mid h(P) \leq B\}.$$

# Heuristics

- A “random”  $\mathbf{x} \in \mathbb{P}^n(\mathbb{Q})$  with  $h(\mathbf{x}) \leq B$  has  $f(\mathbf{x}) = O_f(B^3)$ .
- Expect likelihood  $f(\mathbf{x}) = 0$  about  $\frac{1}{B^3}$ .

# Heuristics

- A “random”  $\mathbf{x} \in \mathbb{P}^n(\mathbb{Q})$  with  $h(\mathbf{x}) \leq B$  has  $f(\mathbf{x}) = O_f(B^3)$ .
- Expect likelihood  $f(\mathbf{x}) = 0$  about  $\frac{1}{B^3}$ .
- Heuristic:  $N_f(B) \sim c_f B^{n-3}$ .

Expectation:  $X_f(\mathbb{Q}) \neq \emptyset$  when  $n \geq 4$  and  $c_f > 0$ .



# Heuristics

- A “random”  $\mathbf{x} \in \mathbb{P}^n(\mathbb{Q})$  with  $h(\mathbf{x}) \leq B$  has  $f(\mathbf{x}) = O_f(B^3)$ .
- Expect likelihood  $f(\mathbf{x}) = 0$  about  $\frac{1}{B^3}$ .
- Heuristic:  $N_f(B) \sim c_f B^{n-3}$ .

Expectation:  $X_f(\mathbb{Q}) \neq \emptyset$  when  $n \geq 4$  and  $c_f > 0$ .

## Remark

When  $n = 2$ ,  $f = 0$  is an **elliptic curve**. In this case,

$$N_f(B) \sim c_f (\log B)^{\text{rk}(E)/2}.$$

# Circle method

Introduced by Hardy and Littlewood to count things.

$$N_f(B) = \sum_{\substack{\mathbf{x} \in \mathbb{Z}^{n+1} \\ h(\mathbf{x}) \leq B}} \int_0^1 \left( e^{2\pi i f(\mathbf{x})\alpha} d\alpha \right) = \int_0^1 \sum_{\substack{\mathbf{x} \in \mathbb{Z}^{n+1} \\ h(\mathbf{x}) \leq B}} e^{2\pi i f(\mathbf{x})\alpha} d\alpha$$

# Circle method

Introduced by Hardy and Littlewood to count things.

$$N_f(B) = \sum_{\substack{\mathbf{x} \in \mathbb{Z}^{n+1} \\ h(\mathbf{x}) \leq B}} \int_0^1 \left( e^{2\pi i f(\mathbf{x})\alpha} d\alpha \right) = \int_0^1 \sum_{\substack{\mathbf{x} \in \mathbb{Z}^{n+1} \\ h(\mathbf{x}) \leq B}} e^{2\pi i f(\mathbf{x})\alpha} d\alpha$$

Estimate sums when  $\alpha$  is

- (major arcs)  $\alpha$  well approximated by  $\frac{a}{q} \in \mathbb{Q}$ , otherwise
- (minor arcs) negligible contribution.

For major arc: count solutions **modulo**  $q$ .

# Some history

When  $n$  large enough<sup>1</sup>, circle method shows

$$N_f(B) \sim c_f B^{n-3}, \quad c_f > 0,$$

i.e.  $X_f$  always has a rational point.

---

<sup>1</sup>Recall  $n$  denotes the dimension of  $\mathbb{P}^n$ ; the number of variables is  $n + 1$

# Some history

When  $n$  large enough<sup>1</sup>, circle method shows

$$N_f(B) \sim c_f B^{n-3}, \quad c_f > 0,$$

i.e.  $X_f$  always has a rational point.

1950's Birch, Lewis, then Davenport:  $n \geq 15$

---

<sup>1</sup>Recall  $n$  denotes the dimension of  $\mathbb{P}^n$ ; the number of variables is  $n + 1$

# Some history

When  $n$  large enough<sup>1</sup>, circle method shows

$$N_f(B) \sim c_f B^{n-3}, \quad c_f > 0,$$

i.e.  $X_f$  always has a rational point.

1950's Birch, Lewis, then Davenport:  $n \geq 15$

1982 Heath-Brown:  $n \geq 9$  if  $X_f$  is nonsingular [HB83]

---

<sup>1</sup>Recall  $n$  denotes the dimension of  $\mathbb{P}^n$ ; the number of variables is  $n + 1$

# Some history

When  $n$  large enough<sup>1</sup>, circle method shows

$$N_f(B) \sim c_f B^{n-3}, \quad c_f > 0,$$

i.e.  $X_f$  always has a rational point.

1950's Birch, Lewis, then Davenport:  $n \geq 15$

1982 Heath-Brown:  $n \geq 9$  if  $X_f$  is nonsingular [HB83]

1988 Hooley:  $n \geq 8$  if  $X_f$  is everywhere locally soluble

---

<sup>1</sup>Recall  $n$  denotes the dimension of  $\mathbb{P}^n$ ; the number of variables is  $n + 1$

# Some history

When  $n$  large enough<sup>1</sup>, circle method shows

$$N_f(B) \sim c_f B^{n-3}, \quad c_f > 0,$$

i.e.  $X_f$  always has a rational point.

1950's Birch, Lewis, then Davenport:  $n \geq 15$

1982 Heath-Brown:  $n \geq 9$  if  $X_f$  is nonsingular [HB83]

1988 Hooley:  $n \geq 8$  if  $X_f$  is everywhere locally soluble

## Question

For  $n \leq 8$ , as  $f$  varies, how often does  $X_f$  have a rational point?

---

<sup>1</sup>Recall  $n$  denotes the dimension of  $\mathbb{P}^n$ ; the number of variables is  $n + 1$



# Back to cubic hypersurfaces

## Question

*How often does  $X_f$  have a rational point?*

$\mathbb{A}^{\binom{n+3}{3}} - 0$  is “moduli space” of cubic forms, each  $f$  is integer point.

# Back to cubic hypersurfaces

## Question

*How often does  $X_f$  have a rational point?*

$\mathbb{A}^{\binom{n+3}{3}} - 0$  is “moduli space” of cubic forms, each  $f$  is integer point.

Let  $h(f) = \|\mathbf{a}\|$  = Euclidean norm, define **natural density**

$$\rho_n = \lim_{B \rightarrow \infty} \frac{\#\{f \mid h(f) \leq B, X_f(\mathbb{Q}) \neq \emptyset\}}{\#\{f \mid h(f) \leq B\}}.$$

# Back to cubic hypersurfaces

## Question

*How often does  $X_f$  have a rational point?*

$\mathbb{A}^{\binom{n+3}{3}} - 0$  is “moduli space” of cubic forms, each  $f$  is integer point.

Let  $h(f) = \|\mathbf{a}\|$  = Euclidean norm, define **natural density**

$$\rho_n = \lim_{B \rightarrow \infty} \frac{\#\{f \mid h(f) \leq B, X_f(\mathbb{Q}) \neq \emptyset\}}{\#\{f \mid h(f) \leq B\}}.$$

## Remark

- Counting **primitive** forms gets same answer, i.e. using  $\mathbb{P}^{\binom{n+3}{3}-1}$

# Everywhere local solubility

A variety  $X/\mathbb{Q}$  is **everywhere locally soluble** (ELS) if

$$X(\mathbb{R}) \neq \emptyset \text{ and } X(\mathbb{Q}_p) \neq \emptyset \text{ for all } p.$$

# Everywhere local solubility

A variety  $X/\mathbb{Q}$  is **everywhere locally soluble** (ELS) if

$$X(\mathbb{R}) \neq \emptyset \text{ and } X(\mathbb{Q}_p) \neq \emptyset \text{ for all } p.$$

ELS is

- **necessary** for  $X_f$  to have  $\mathbb{Q}$ -point
- but not **sufficient**

# Everywhere local solubility

A variety  $X/\mathbb{Q}$  is **everywhere locally soluble** (ELS) if

$$X(\mathbb{R}) \neq \emptyset \text{ and } X(\mathbb{Q}_p) \neq \emptyset \text{ for all } p.$$

ELS is

- **necessary** for  $X_f$  to have  $\mathbb{Q}$ -point
- but not **sufficient**...or is it?

# Everywhere local solubility

A variety  $X/\mathbb{Q}$  is **everywhere locally soluble** (ELS) if

$$X(\mathbb{R}) \neq \emptyset \text{ and } X(\mathbb{Q}_p) \neq \emptyset \text{ for all } p.$$

ELS is

- **necessary** for  $X_f$  to have  $\mathbb{Q}$ -point
- but not **sufficient**...or is it?

Density of ELS cubic forms in  $\mathbb{Z}[x_0, \dots, x_n]$ :

$$\rho_n^{ELS} = \lim_{B \rightarrow \infty} \frac{\#\{f \mid h(f) \leq B, X_f \text{ ELS}\}}{\#\{f \mid h(f) \leq B\}}.$$

# (Lack of) obstructions

## Conjecture (Poonen–Voloch, 2004)

When  $n \geq 3$ ,  $\rho_n^{ELS} = \rho_n$ .

*i.e. local-global principle holds for 100% of cubic hypersurfaces.*



# (Lack of) obstructions

## Conjecture (Poonen–Voloch, 2004)

When  $n \geq 3$ ,  $\rho_n^{ELS} = \rho_n$ .

*i.e. local-global principle holds for 100% of cubic hypersurfaces.*

- Conj. of Colliot–Thélène: Brauer–Manin is only obstruction
- Brauer–Manin vacuous for cubics when  $n \geq 4$
- Brauer–Manin vacuous for 100% of cubic surfaces in  $\mathbb{P}^3$

# (Lack of) obstructions

## Conjecture (Poonen–Voloch, 2004)

When  $n \geq 3$ ,  $\rho_n^{ELS} = \rho_n$ .

*i.e. local-global principle holds for 100% of cubic hypersurfaces.*

- Conj. of Colliot–Thélène: Brauer–Manin is only obstruction
- Brauer–Manin vacuous for cubics when  $n \geq 4$
- Brauer–Manin vacuous for 100% of cubic surfaces in  $\mathbb{P}^3$

## Theorem (Browning–Le Boudec–Sawin, 2023)

When  $n \geq 4$ ,  $\rho_n^{ELS} = \rho_n$ .

# (Lack of) obstructions

## Conjecture (Poonen–Voloch, 2004)

When  $n \geq 3$ ,  $\rho_n^{ELS} = \rho_n$ .

*i.e. local-global principle holds for 100% of cubic hypersurfaces.*

- Conj. of Colliot–Thélène: Brauer–Manin is only obstruction
- Brauer–Manin vacuous for cubics when  $n \geq 4$
- Brauer–Manin vacuous for 100% of cubic surfaces in  $\mathbb{P}^3$

## Theorem (Browning–Le Boudec–Sawin, 2023)

When  $n \geq 4$ ,  $\rho_n^{ELS} = \rho_n$ . In fact *true for Fano deg.  $d$*   $(d, n) \neq (3, 3)$

# Computing $\rho^{ELS}$

Let  $\rho_n(p)$  = density of  $p$ -adic cubic forms  $f$  such that  $X_f(\mathbb{Q}_p) \neq \emptyset$ .

Theorem (Poonen–Voloch, 2004)

Let  $n \geq 2$ . We have

$$\rho_n^{ELS} = \prod_p \rho_n(p).$$

# Computing $\rho^{ELS}$

Let  $\rho_n(p)$  = density of  $p$ -adic cubic forms  $f$  such that  $X_f(\mathbb{Q}_p) \neq \emptyset$ .

Theorem (Poonen–Voloch, 2004)

Let  $n \geq 2$ . We have

$$\rho_n^{ELS} = \prod_p \rho_n(p).$$

Think

Local probabilities independent...even though infinitely many!

# Computing $\rho^{ELS}$

Let  $\rho_n(p)$  = density of  $p$ -adic cubic forms  $f$  such that  $X_f(\mathbb{Q}_p) \neq \emptyset$ .

Theorem (Poonen–Voloch, 2004)

Let  $n \geq 2$ . We have


$$\rho_n^{ELS} = \prod_p \rho_n(p).$$

Think

Local probabilities independent...even though infinitely many!

2016 Bright–Browning–Loughran: flexible generalization to families given by fibers of maps to affine/projective space.

# Varieties parameterized by fibers



# Main result

## Theorem (Beneish–K.)

Let  $4 \leq n \leq 8$ . There exist *explicit polynomials*  $g_n(t), h_n(t) \in \mathbb{Z}[t]$  describing  $\rho_n$  exactly as Euler product

$$\rho_n = \prod_p \rho_n(p) = \prod_p \left( 1 - \frac{g_n(p)}{h_n(p)} \right).$$



# Main result

## Theorem (Beneish–K.)

Let  $4 \leq n \leq 8$ . There exist *explicit polynomials*  $g_n(t), h_n(t) \in \mathbb{Z}[t]$  describing  $\rho_n$  exactly as Euler product

$$\rho_n = \prod_p \rho_n(p) = \prod_p \left( 1 - \frac{g_n(p)}{h_n(p)} \right).$$

## Remark

- We produce  $g_3, h_3$ , and conjectural formula for  $\rho_3$ .
- Recovers  $\rho_n(p) = 1$  for  $n \geq 9$ .

# Cubic 3-folds in $\mathbb{P}^4$

## Example

When  $n = 4$  we have

$$\begin{aligned} g_4(p) &= (p^{46} + 3p^{41} + p^{40} - p^{39} + p^{37} + p^{36} + p^{35} - 3p^{34} + 3p^{27} - p^{26} + p^{25} \\ &\quad + p^{19}) (p^2 + 1) (p + 1)^2 (p - 1)^4 \\ h_4(p) &= 9 (p^{19} - 1) (p^{17} - 1) (p^{10} + 1) (p^9 + 1) (p^9 - 1) (p^7 - 1) (p^5 + 1) \end{aligned}$$

Asymptotically,  $\frac{g_4(p)}{h_4(p)} \sim \frac{1}{9p^{22}}.$

Numerically,  $\rho_4 \approx 0.99999999497 = 1 - 5.022 \cdot 10^{-9}.$

# Asymptotics and numerics

## Theorem (Beneish–K.)

$$\rho_n = \prod_p \left( 1 - \frac{g_n(p)}{h_n(p)} \right)$$

$n$	$\rho_n \approx$	$g_n(p)/h_n(p) \sim$
3	0.999927 ( <i>conj.</i> )	$1/3p^{10}$
4	$1 - 5.022 \cdot 10^{-9}$	$1/9p^{22}$
5	$1 - 1.343 \cdot 10^{-15}$	$1/9p^{43}$
6	$1 - 3.502 \cdot 10^{-26}$	$1/9p^{78}$
7	$1 - 5.152 \cdot 10^{-42}$	$1/27p^{129}$
8	$1 - 6.222 \cdot 10^{-64}$	$1/27p^{201}$

# Asymptotics and numerics

## Theorem (Beneish–K.)

$$\rho_n = \prod_p \left( 1 - \frac{g_n(p)}{h_n(p)} \right)$$

$n$	$\rho_n \approx$	$g_n(p)/h_n(p) \sim$
3	0.999927 ( <i>conj.</i> )	$1/3p^{10}$
4	$1 - 5.022 \cdot 10^{-9}$	$1/9p^{22}$
5	$1 - 1.343 \cdot 10^{-15}$	$1/9p^{43}$
6	$1 - 3.502 \cdot 10^{-26}$	$1/9p^{78}$
7	$1 - 5.152 \cdot 10^{-42}$	$1/27p^{129}$
8	$1 - 6.222 \cdot 10^{-64}$	$1/27p^{201}$
9	1	0

# Roadmap

$\rho_n$  = density of cubic forms  $f \in \mathbb{Z}[x_0, \dots, x_n]$  with  $X_f(\mathbb{Q}) \neq \emptyset$ .

$$\textcircled{1} \quad \rho_n = \rho_n^{ELS} = \prod_p \rho_n(p) \quad [\text{BLBS23, PV04, BBL16}]$$

# Roadmap

$\rho_n$  = density of cubic forms  $f \in \mathbb{Z}[x_0, \dots, x_n]$  with  $X_f(\mathbb{Q}) \neq \emptyset$ .

- 1  $\rho_n = \rho_n^{ELS} = \prod_p \rho_n(p)$  [BLBS23, PV04, BBL16]
- 2 Study possible reductions  $\overline{X_f}$  modulo  $p$

# Roadmap

$\rho_n$  = density of cubic forms  $f \in \mathbb{Z}[x_0, \dots, x_n]$  with  $X_f(\mathbb{Q}) \neq \emptyset$ .

- 1  $\rho_n = \rho_n^{ELS} = \prod_p \rho_n(p)$  [BLBS23, PV04, BBL16]
- 2 Study possible reductions  $\overline{X_f}$  modulo  $p$
- 3 How often does  $\overline{P} \in \overline{X_f}$  lift to  $P \in X_f(\mathbb{Q}_p)$ ?

# Roadmap

$\rho_n$  = density of cubic forms  $f \in \mathbb{Z}[x_0, \dots, x_n]$  with  $X_f(\mathbb{Q}) \neq \emptyset$ .

- ①  $\rho_n = \rho_n^{ELS} = \prod_p \rho_n(p)$  [BLBS23, PV04, BBL16]
- ② Study possible reductions  $\overline{X_f}$  modulo  $p$
- ③ How often does  $\overline{P} \in \overline{X_f}$  lift to  $P \in X_f(\mathbb{Q}_p)$ ?

## Idea

Express lifting probabilities in terms of each other and recurse to get rational function  $\rho_n(p)$ .



Counting points  
○○○○○


Varying  $X_f$   
○○○○○

Results  
○○○○●○

Lifting probabilities  
○○○○○○○○○○○○○○

Final thoughts  
○○○○○

## Full picture



## Related results

Plane cubic curves [BCF16a, Bha14]

- $\rho_2^{ELS}$ ,  $\rho_2(p)$  computed by Bhargava–Cremona–Fisher
- Bhargava: pos. prop. fail Hasse principle (conj. 65%) [Bha14]

## Related results

### Plane cubic curves [BCF16a, Bha14]

- $\rho_2^{ELS}$ ,  $\rho_2(p)$  computed by Bhargava–Cremona–Fisher
- Bhargava: pos. prop. fail Hasse principle (conj. 65%) [Bha14]

### Quadric hypersurfaces [BCF<sup>+</sup>16b]

- BCF–Jones–Keating: explicit Euler product for density of quadratic forms with integral zero [BCF<sup>+</sup>16b]
- Hasse principle holds but archimedean place not trivial!
- 98.3% of quadric surfaces in  $\mathbb{P}^3$  soluble

# Related results

## Plane cubic curves [BCF16a, Bha14]

- $\rho_2^{ELS}$ ,  $\rho_2(p)$  computed by Bhargava–Cremona–Fisher
- Bhargava: pos. prop. fail Hasse principle (conj. 65%) [Bha14]

## Quadric hypersurfaces [BCF<sup>+</sup>16b]

- BCF–Jones–Keating: explicit Euler product for density of quadratic forms with integral zero [BCF<sup>+</sup>16b]
- Hasse principle holds but archimedean place not trivial!
- 98.3% of quadric surfaces in  $\mathbb{P}^3$  soluble

More: certain cubic surfaces [Bro17], (2,2)-curves in  $\mathbb{P}^1 \times \mathbb{P}^1$  [FHP21]

# Computing the local factors

## Goal

*Compute local probability  $\rho_n(p)$  that  $X_f$  has  $\mathbb{Q}_p$ -point.*

# Computing the local factors

## Goal

Compute local probability  $\rho_n(p)$  that  $X_f$  has  $\mathbb{Q}_p$ -point.

$\rho_n(p)$  is  $p$ -adic Haar measure with  $\mu_p \left( \mathbb{Z}_p^{\binom{n+3}{3}} \right) = 1$ :

$$\rho_n(p) = \mu_p \left( f \in \mathbb{Z}_p[x_0, \dots, x_n] \mid X_f(\mathbb{Q}_p) \neq \emptyset \right).$$

## Think

Reduce mod  $p$ , try to decide solubility with **Hensel's lemma**.

# Computing the local factors

## Goal

Compute local probability  $\rho_n(p)$  that  $X_f$  has  $\mathbb{Q}_p$ -point.

$\rho_n(p)$  is  $p$ -adic Haar measure with  $\mu_p \left( \mathbb{Z}_p^{\binom{n+3}{3}} \right) = 1$ :

$$\rho_n(p) = \mu_p \left( f \in \mathbb{Z}_p[x_0, \dots, x_n] \mid X_f(\mathbb{Q}_p) \neq \emptyset \right).$$


## Think

Reduce mod  $p$ , try to decide solubility with **Hensel's lemma**.


## Lemma (Hensel's Lemma)

If  $P_0 \in \overline{X}^{\text{sm}}(\mathbb{F}_p)$ , then **there exists lift**  $P \in X(\mathbb{Z}_p)$  with  $\overline{P} = P_0$ .

# Cubic hypersurfaces over $\mathbb{F}_p$



Def. over  
 $\mathbb{F}_p$   
 $\mathbb{F}_{p^2}$   
 $\mathbb{F}_{p^3}$





# When are there always $\mathbb{Q}_p$ -points?

## Proposition

If  $\overline{X_f}$  *not* a config. of conjugate hyperplanes, then  $X_f(\mathbb{Q}_p) \neq \emptyset$ .

## Proof for curves ( $n = 2$ ).

If geom. integral, use Hasse–Weil bounds on (normalization of)  $\overline{X_f}$

$$\#X_f(\mathbb{F}_p) \geq p + 1 - 2\sqrt{p} > 0.$$

All other possibilities contain line defined over  $\mathbb{F}_p$ .



# When are there always $\mathbb{Q}_p$ -points?

## Proposition

If  $\overline{X_f}$  *not* a config. of conjugate hyperplanes, then  $X_f(\mathbb{Q}_p) \neq \emptyset$ .

## Proof for curves ( $n = 2$ ).

If geom. integral, use Hasse–Weil bounds on (normalization of)  $\overline{X_f}$

$$\#X_f(\mathbb{F}_p) \geq p + 1 - 2\sqrt{p} > 0.$$

All other possibilities contain line defined over  $\mathbb{F}_p$ . □

For  $n \geq 3$ , use inductive argument reducing to *coordinate hypersurfaces*  $X_i = X_f \cap \{x_i = 0\} \subset \mathbb{P}^{n-1}$ .

# When are there always $\mathbb{Q}_p$ -points?

## Proposition

If  $\overline{X_f}$  *not* a config. of conjugate hyperplanes, then  $X_f(\mathbb{Q}_p) \neq \emptyset$ .

## Proof for curves ( $n = 2$ ).

If geom. integral, use Hasse–Weil bounds on (normalization of)  $\overline{X_f}$

$$\#X_f(\mathbb{F}_p) \geq p + 1 - 2\sqrt{p} > 0.$$

All other possibilities contain line defined over  $\mathbb{F}_p$ . □

For  $n \geq 3$ , use inductive argument reducing to *coordinate hypersurfaces*  $X_i = X_f \cap \{x_i = 0\} \subset \mathbb{P}^{n-1}$ .

## Warning

There exist irreducible deg.  $d > 3$  curves  $X \subset \mathbb{P}^2$  with  $X(\mathbb{Q}_p) = \emptyset$ .

# Some (non)examples

Observe 2 is not a cube in  $\mathbb{F}_7$ . Let  $u \in \mathbb{F}_{7^3}$  be root of  $x^3 - 2 = 0$ .

Example ( $n = 2$ ,  $p = 7$ )

$$f(x, y, z) = x^3 + 3x^2y + 3xy^2 - 6xyz + 3y^3 - 6y^2z + 4z^3 = 0$$

# Some (non)examples

Observe 2 is not a cube in  $\mathbb{F}_7$ . Let  $u \in \mathbb{F}_{7^3}$  be root of  $x^3 - 2 = 0$ .

Example ( $n = 2$ ,  $p = 7$ )

$$f(x, y, z) = x^3 + 3x^2y + 3xy^2 - 6xyz + 3y^3 - 6y^2z + 4z^3 = 0$$

Over  $\overline{\mathbb{F}_7}$  we have

$$\overline{f} = \prod_{\sigma \in \text{Gal}(\mathbb{F}_{7^3}/\mathbb{F}_7)} \sigma(x + (1 + u)y + u^2z).$$

$$\overline{X_f}(\mathbb{F}_p) = \emptyset \implies X_f(\mathbb{Z}_p) = \emptyset.$$

$\overline{X_f}$  is a triangle

# Some (non)examples

## Example

Suppose  $f = g_1(x_0, x_1, x_2) + pg_2(x_3, x_4, x_5) + p^2g_3(x_6, x_7, x_8)$   
for  $g_i = 0$  with **no nontrivial  $\mathbb{F}_p$ -solutions**.

If  $[x_0, \dots, x_8] \in X_f(\mathbb{Z}_p)$  then  $p \mid x_0, x_1, x_2$

# Some (non)examples

## Example

Suppose  $f = g_1(x_0, x_1, x_2) + pg_2(x_3, x_4, x_5) + p^2g_3(x_6, x_7, x_8)$   
for  $g_i = 0$  with **no nontrivial  $\mathbb{F}_p$ -solutions**.

If  $[x_0, \dots, x_8] \in X_f(\mathbb{Z}_p)$  then  $p \mid x_0, x_1, x_2$

$$\begin{aligned} \frac{1}{p} f(px_0, px_1, px_2, x_3, \dots, x_8) \\ = g_2(x_3, x_4, x_5) + pg_3(x_6, x_7, x_8) + p^2g_1(x_0, x_1, x_2) \end{aligned}$$

$$\implies X_f(\mathbb{Z}_p) = \emptyset.$$

# Some (non)examples

## Example

Suppose  $f = g_1(x_0, x_1, x_2) + pg_2(x_3, x_4, x_5) + p^2g_3(x_6, x_7, x_8)$   
for  $g_i = 0$  with **no nontrivial  $\mathbb{F}_p$ -solutions**.

If  $[x_0, \dots, x_8] \in X_f(\mathbb{Z}_p)$  then  $p \mid x_0, x_1, x_2$

$$\begin{aligned} \frac{1}{p} f(px_0, px_1, px_2, x_3, \dots, x_8) \\ = g_2(x_3, x_4, x_5) + pg_3(x_6, x_7, x_8) + p^2g_1(x_0, x_1, x_2) \end{aligned}$$

$$\implies X_f(\mathbb{Z}_p) = \emptyset.$$



## Remark

This generalizes, but it ignores **cross terms**...




# Configurations of conjugate lines

1 - triple line





2 - star




3 - triangle

# Configurations of conjugate hyperplanes

1-~~uple~~ (hyper)plane



3-triangle



2-star

# Strategy

## Goal

*Look modulo  $p$  and try to decide solubility*

$$\rho_n(p) = \xi_{n,0}\sigma_{n,0} + \xi_{n,1}\sigma_{n,1} + \xi_{n,2}\sigma_{n,2} + \xi_{n,3}\sigma_{n,3}$$

# Strategy

## Goal

*Look modulo  $p$  and try to decide solubility*

$$\rho_n(p) = \xi_{n,0}\sigma_{n,0} + \xi_{n,1}\sigma_{n,1} + \xi_{n,2}\sigma_{n,2} + \xi_{n,3}\sigma_{n,3}$$

- $\xi_{n,i}$  = prob.  $f$  has type  $i$

# Strategy

## Goal

*Look modulo  $p$  and try to decide solubility*

$$\rho_n(p) = \xi_{n,0}\sigma_{n,0} + \xi_{n,1}\sigma_{n,1} + \xi_{n,2}\sigma_{n,2} + \xi_{n,3}\sigma_{n,3}$$

- $\xi_{n,i}$  = prob.  $f$  has type  $i$
- $\sigma_{n,i}$  = prob.  $X_f(\mathbb{Q}_p) \neq \emptyset$  given  $f$  type  $i$

# Strategy

## Goal

*Look modulo  $p$  and try to decide solubility*

$$\rho_n(p) = \xi_{n,0}\sigma_{n,0} + \xi_{n,1}\sigma_{n,1} + \xi_{n,2}\sigma_{n,2} + \xi_{n,3}\sigma_{n,3}$$

- $\xi_{n,i}$  = prob.  $f$  has type  $i$
- $\sigma_{n,i}$  = prob.  $X_f(\mathbb{Q}_p) \neq \emptyset$  given  $f$  type  $i$
- $\xi_{n,0}$  = prob.  $f$  **not** config. of conj. hyperplanes

# Strategy

## Goal

*Look modulo  $p$  and try to decide solubility*

$$\rho_n(p) = \xi_{n,0} \cdot 1 + \xi_{n,1} \sigma_{n,1} + \xi_{n,2} \sigma_{n,2} + \xi_{n,3} \sigma_{n,3}$$

- $\xi_{n,i}$  = prob.  $f$  has type  $i$
- $\sigma_{n,i}$  = prob.  $X_f(\mathbb{Q}_p) \neq \emptyset$  given  $f$  type  $i$
- $\xi_{n,0}$  = prob.  $f$  not config. of conj. hyperplanes
- $\sigma_{n,0} = 1$

# Factorization probabilities

$$\xi_{n,0} = 1 - \frac{q^{3n-3} + 2q^{n+3} + 2q^{n+2} + 2q^{n+1} - 2q^2 - 2q - 3}{3(q^2 + q + 1) \left( q^{\binom{n+3}{3}} - 1 \right)}$$

$$\xi_{n,1} = \frac{q^{n+1} - 1}{q^{\binom{n+3}{3}} - 1}$$

$$\xi_{n,2} = \frac{(q^{2n+1} - q^{n+1} - q^n + 1)q}{3 \left( q^{\binom{n+3}{3}} - 1 \right)}$$

$$\xi_{n,3} = \frac{(q^{3n} - q^{2n} - q^{2n+1} - q^{2n-1} + q^{n+1} + q^{n-1} + q^n - 1)q^3}{3(q^2 + q + 1) \left( q^{\binom{n+3}{3}} - 1 \right)}$$

## Exercise

Convince yourself that probability of a polynomial factoring a certain way is given by a (uniform) rational function.



# Phases I, II, and III

Repeat this process three times:

- 1 Reduce mod  $p$ , lift any “easy” solutions with Hensel’s lemma

# Phases I, II, and III

Repeat this process three times:

- 1 Reduce mod  $p$ , lift any “easy” solutions with Hensel’s lemma
- 2 Reduction is type  $j = 1, 2, 3$  with explicit probability

# Phases I, II, and III

Repeat this process three times:

- 1 Reduce mod  $p$ , lift any “easy” solutions with Hensel’s lemma
- 2 Reduction is type  $j = 1, 2, 3$  with explicit probability
- 3 Introduce new lifting probability for each type

# Phases I, II, and III

Repeat this process three times:

- 1 Reduce mod  $p$ , lift any “easy” solutions with Hensel’s lemma
- 2 Reduction is type  $j = 1, 2, 3$  with explicit probability
- 3 Introduce new lifting probability for each type
- 4 Relate new lifting probabilities to others

Eventually this terminates: 64  $\mathbb{Q}(p)$ -linear relations in 64 unknowns

Counting points  
○○○○○


Varying  $X_f$   
○○○○○

Results  
○○○○○○

Lifting probabilities  
○○○○○○○○○○●●○

Final thoughts  
○○○○

## Setup



Counting points  
○○○○○


Varying  $X_f$   
○○○○○

Results  
○○○○○○

Lifting probabilities  
○○○○○○○○○○●○

Final thoughts  
○○○○

## Phase I



Counting points  
○○○○○


Varying  $X_f$   
○○○○○

Results  
○○○○○○

Lifting probabilities  
○○○○○○○○○○●●

Final thoughts  
○○○○

## Phase I



$\theta_{111}$   $\theta_{112}$   $\theta_{113}$   $\theta_{121}$   $\theta_{122}$   $\theta_{123}$   $\theta_{131}$   $\theta_{132}$   $\theta_{133}$   $\theta_{211}$   $\theta_{212}$   $\theta_{213}$   $\theta_{221}$   $\theta_{222}$   $\theta_{223}$   $\theta_{231}$   $\theta_{232}$   $\theta_{233}$   $\theta_{311}$   $\theta_{312}$   $\theta_{313}$   $\theta_{321}$   $\theta_{322}$   $\theta_{323}$   $\theta_{331}$   $\theta_{332}$   $\theta_{333}$

Counting points  
○○○○○


Varying  $X_f$   
○○○○○

Results  
○○○○○○

Lifting probabilities  
○○○○○○○○○○●●○

Final thoughts  
○○○○

## Phase II





Counting points  
○○○○○


Varying  $X_f$   
○○○○○

Results  
○○○○○○

Lifting probabilities  
○○○○○○○○○○●●○

Final thoughts  
○○○○

## Phase II



Counting points  
○○○○○


Varying  $X_f$   
○○○○○

Results  
○○○○○○

Lifting probabilities  
○○○○○○○○○○●●○

Final thoughts  
○○○○

### Phase III



Counting points  
○○○○○


Varying  $X_f$   
○○○○○

Results  
○○○○○○

Lifting probabilities  
○○○○○○○○○○●●○

Final thoughts  
○○○○○

When  $n = 8$



NO LIFT

# Some remarks

Practicalities:

- Solve with Sage symbolic solver
- Block variables ( $27 + 27 + 10$ ) to speed up

# Some remarks

## Practicalities:

- Solve with Sage symbolic solver
- Block variables  $(27 + 27 + 10)$  to speed up

In principle,  $\rho_n(p) = 1$  for  $n \geq 9$  can be seen “by hand”

- No “no lift” sink in the flow chart!
- Capture and describe **explicitly** Heath–Brown’s observation of necessary/sufficient conditions for  $f$  to have local solutions

# Density of cubics with a point

## Theorem (Beneish–K.)

Let  $n \geq 4$  (conjecturally  $n \geq 3$ ). Then  $\rho_n = 1$  when  $n \geq 9$  and

$$\rho_n = \prod_{p \text{ prime}} \left( 1 - \frac{g_n(p)}{h_n(p)} \right) \text{ when } n \leq 8$$

for *explicit polynomials*  $g_n(t), h_n(t) \in \mathbb{Z}[t]$ .

$n$	$\rho_n^{(ELS)} \approx$	$1 - \rho_n(p) \sim$
2	0.9726 [BCF16a]	$1/3p^3$
3	0.999927 (conj.)	$1/3p^{10}$
4	$1 - 5.022 \cdot 10^{-9}$	$1/9p^{22}$
5	$1 - 1.343 \cdot 10^{-15}$	$1/9p^{43}$
6	$1 - 3.502 \cdot 10^{-26}$	$1/9p^{78}$
7	$1 - 5.152 \cdot 10^{-42}$	$1/27p^{129}$
8	$1 - 6.222 \cdot 10^{-64}$	$1/27p^{201}$

## Further questions

Let  $\rho_{d,n}$  = density of degree  $d$  hypersurfaces in  $\mathbb{P}^n$  with  $\mathbb{Q}$ -point

How far can this approach go to compute  $\rho_{d,n}$ ?

## Further questions

Let  $\rho_{d,n}$  = density of degree  $d$  hypersurfaces in  $\mathbb{P}^n$  with  $\mathbb{Q}$ -point

How far can this approach go to compute  $\rho_{d,n}$ ?

- Lose uniformity in  $p$  in general
- Heath–Brown:  $\rho_{4,n}(p) = 1$  for  $n \geq 9126$ ,  
 $\rho_{5,n}(p) = 1$  known for  $n \geq 25$ ,  $p \geq 17$  [HB09]
- Can we predict asymptotics/numerics with less effort?



## Further questions

Let  $\rho_{d,n}$  = density of degree  $d$  hypersurfaces in  $\mathbb{P}^n$  with  $\mathbb{Q}$ -point

How far can this approach go to compute  $\rho_{d,n}$ ?

- Lose uniformity in  $p$  in general
- Heath–Brown:  $\rho_{4,n}(p) = 1$  for  $n \geq 9126$ ,  
 $\rho_{5,n}(p) = 1$  known for  $n \geq 25$ ,  $p \geq 17$  [HB09]
- Can we predict asymptotics/numerics with less effort?

Structure of  $\rho_{d,n}(p)$

- Always rational function? (for  $p \gg 0$ )
- Known for binary forms,  $\rho_{d,1}(p) = \rho_{d,1}(1/p)$  [BCFG22]

# Thank you I

Thank you for the invitation and for your attention!



M. J. Bright, T. D. Browning, and D. Loughran, Failures of weak approximation in families, *Compos. Math.* **152** (2016), no. 7, 1435–1475. MR 3530447



Manjul Bhargava, John Cremona, and Tom Fisher, The proportion of plane cubic curves over  $\mathbb{Q}$  that everywhere locally have a point, *Int. J. Number Theory* **12** (2016), no. 4, 1077–1092. MR 3484299



Manjul Bhargava, John E. Cremona, Tom Fisher, Nick G. Jones, and Jonathan P. Keating, What is the probability that a random integral quadratic form in  $n$  variables has an integral zero?, *Int. Math. Res. Not. IMRN* (2016), no. 12, 3828–3848. MR 3544620



Manjul Bhargava, John Cremona, Tom Fisher, and Stevan Gajović, The density of polynomials of degree  $nn$  over  $\mathbb{Z}_p$  having exactly  $r$  roots in  $\mathbb{q}_p$ , *Proceedings of the London Mathematical Society* **124** (2022), no. 5, 713–736 (English (US)).



Manjul Bhargava, A positive proportion of plane cubics fail the hasse principle, Preprint, available at <https://arxiv.org/pdf/1402.1131.pdf>, 2014.



Tim Browning, Pierre Le Boudec, and Will Sawin, The Hasse principle for random Fano hypersurfaces, *Ann. of Math. (2)* **197** (2023), no. 3, 1115–1203. MR 4564262



T. D. Browning, Many cubic surfaces contain rational points, *Mathematika* **63** (2017), no. 3, 818–839. MR 3731306

# Thank you II



Tom Fisher, Wei Ho, and Jennifer Park, [Everywhere local solubility for hypersurfaces in products of projective spaces](#), Res. Number Theory **7** (2021), no. 1, Paper No. 6, 27. MR 4199457



D. R. Heath-Brown, [Cubic forms in ten variables](#), Proc. London Math. Soc. (3) **47** (1983), no. 2, 225–257. MR 703978



———, [Zeros of p-adic forms](#), Proceedings of the London Mathematical Society **100** (2009), no. 2, 560–584.



Bjorn Poonen and José Felipe Voloch, [Random Diophantine equations](#), Arithmetic of higher-dimensional algebraic varieties (Palo Alto, CA, 2002), Progr. Math., vol. 226, Birkhäuser Boston, Boston, MA, 2004, With appendices by Jean-Louis Colliot-Thélène and Nicholas M. Katz, pp. 175–184. MR 2029869