

# Local solubility in families of superelliptic curves

Christopher Keyes (Emory University)

joint work with Lea Beneish (UC Berkeley)

<https://arxiv.org/abs/2111.04697>

CTNT

June 10, 2022

# Density

Consider the (hopefully) familiar statements:

- “50% of primes  $p$  split in  $\mathbb{Q}(i)$ .”

# Density

Consider the (hopefully) familiar statements:

- “50% of primes  $p$  split in  $\mathbb{Q}(i)$ .”
- “100% of polynomials are irreducible.”

# Density

Consider the (hopefully) familiar statements:

- “50% of primes  $p$  split in  $\mathbb{Q}(i)$ .”
- “100% of polynomials are irreducible.”
- “Most equations  $y^2 = f(x)$  **fail** to have a rational solution.”

# Density

Consider the (hopefully) familiar statements:

- “50% of primes  $p$  split in  $\mathbb{Q}(i)$ .”
- “100% of polynomials are irreducible.”
- “Most equations  $y^2 = f(x)$  **fail** to have a rational solution.”

## Definition

Given  $S \subseteq \mathbb{Z}^n$ , the **natural density** of  $S$  is the limit, if it exists:

$$\lim_{B \rightarrow \infty} \frac{\#(S \cap [-B, B]^n)}{(2B + 1)^n}.$$

# ((Everywhere) local) solubility

Let  $C/\mathbb{Q}$  be a curve and  $v$  a place of  $\mathbb{Q}$  (i.e.  $v = p$  or  $v = \infty$ ).

## Definition (soluble)

$C$  is **globally soluble** if  $C(\mathbb{Q})$  is nonempty.

$C$  is **locally soluble at  $v$**  if  $C(\mathbb{Q}_v)$  is nonempty.

$C$  is **everywhere locally soluble** if  $C(\mathbb{Q}_v) \neq \emptyset$  for all places  $v$ .

# ((Everywhere) local) solubility

Let  $C/\mathbb{Q}$  be a curve and  $v$  a place of  $\mathbb{Q}$  (i.e.  $v = p$  or  $v = \infty$ ).

## Definition (soluble)

$C$  is **globally soluble** if  $C(\mathbb{Q})$  is nonempty.

$C$  is **locally soluble at  $v$**  if  $C(\mathbb{Q}_v)$  is nonempty.

$C$  is **everywhere locally soluble** if  $C(\mathbb{Q}_v) \neq \emptyset$  for all places  $v$ .

## Question

*What proportion of curves over  $\mathbb{Q}$  (in some family) are globally soluble?*

# ((Everywhere) local) solubility

Let  $C/\mathbb{Q}$  be a curve and  $v$  a place of  $\mathbb{Q}$  (i.e.  $v = p$  or  $v = \infty$ ).

## Definition (soluble)

$C$  is **globally soluble** if  $C(\mathbb{Q})$  is nonempty.

$C$  is **locally soluble at  $v$**  if  $C(\mathbb{Q}_v)$  is nonempty.

$C$  is **everywhere locally soluble** if  $C(\mathbb{Q}_v) \neq \emptyset$  for all places  $v$ .

## Question

*What proportion of curves over  $\mathbb{Q}$  (in some family) are ~~globally soluble~~ **everywhere locally soluble**?*

Known for genus 1 curves [BCF21], plane cubics [BCF16], and some families of hypersurfaces [BBL16], [FHP21], [PV04], [Bro17].



# Superelliptic curves

## Definition

A **superelliptic curve**  $C/\mathbb{Q}$  is a smooth projective curve with a cyclic Galois cover of  $\mathbb{P}^1$  of degree  $m \geq 2$ .

More concretely: given by equation in weighted projective space

$$C: y^m = f(x, z)$$

where  $f$  is homogeneous of degree  $d$  divisible by  $m$ .

# Superelliptic curves

## Definition

A **superelliptic curve**  $C/\mathbb{Q}$  is a smooth projective curve with a cyclic Galois cover of  $\mathbb{P}^1$  of degree  $m \geq 2$ .

More concretely: given by equation in weighted projective space

$$C: y^m = f(x, z)$$

where  $f$  is homogeneous of degree  $d$  divisible by  $m$ .

## Warning

Some authors don't assume  $m \mid d$ , want  $f$  to be squarefree, or mean something else entirely by “superelliptic” (e.g. [Swa19]).

# Defining the proportion

For  $\mathbf{c} = (c_i)_{i=0}^d \in \mathbb{Z}^{d+1}$ , we associate a binary form

$$f(x, z) = \sum_{i=0}^d c_i x^i z^{d-i}$$

and a superelliptic curve  $C_f: y^m = f(x, z)$ .

# Defining the proportion

For  $\mathbf{c} = (c_i)_{i=0}^d \in \mathbb{Z}^{d+1}$ , we associate a binary form

$$f(x, z) = \sum_{i=0}^d c_i x^i z^{d-i}$$

and a superelliptic curve  $C_f: y^m = f(x, z)$ .

## Definition

We define

$$\rho_{m,d} = \lim_{B \rightarrow \infty} \frac{\#\{\mathbf{c} \in ([-B, B] \cap \mathbb{Z})^{d+1} \mid C_f \text{ E.L.S.}\}}{\#\{\mathbf{c} \in ([-B, B] \cap \mathbb{Z})^{d+1}\}},$$

the proportion of locally soluble superelliptic curves of this form.

# Main results

Fix  $(m, d) \neq (2, 2)$  such that  $m \mid d$ .

Theorem (Beneish–K. [BK21])

(A)  $0 < \rho_{m,d} < 1$ , and  $\rho_{m,d}$  is product of local densities,

$$\rho_{m,d} = \rho_{m,d}(\infty) \prod_p \rho_{m,d}(p).$$

$\rho_{m,d}(p)$  is (normalized) Haar measure of space of the  $\mathbb{Q}_p$ -soluble curves  $C_f: y^m = f(x, z)$ , with coefficients in  $\mathbb{Z}_p$ .

# Main results

Fix  $(m, d) \neq (2, 2)$  such that  $m$  is prime and  $m \mid d$ .

Theorem (Beneish–K. [BK21], continued)

(B) *We can find explicit (and sometimes good) bounds for  $\rho_{m,d}(p)$  and hence  $\rho_{m,d}$ . In particular,*

$$\liminf_{d \rightarrow \infty} \rho_{m,d} \geq \left(1 - \frac{1}{m^{m+1}}\right) \prod_{p \equiv 1(m)} \left(1 - \left(1 - \frac{p-1}{mp}\right)^{p+1}\right) \prod_{p \not\equiv 0,1(m)} \left(1 - \frac{1}{p^{2(p+1)}}\right).$$

# Main results

Fix  $(m, d) \neq (2, 2)$  such that  $m$  is prime and  $m \mid d$ .

Theorem (Beneish–K. [BK21], continued)

(B) *We can find explicit (and sometimes good) bounds for  $\rho_{m,d}(p)$  and hence  $\rho_{m,d}$ . In particular,*

$$\liminf_{d \rightarrow \infty} \rho_{m,d} \geq \left(1 - \frac{1}{m^{m+1}}\right) \prod_{p \equiv 1(m)} \left(1 - \left(1 - \frac{p-1}{mp}\right)^{p+1}\right) \prod_{p \not\equiv 0,1(m)} \left(1 - \frac{1}{p^{2(p+1)}}\right).$$

When  $m > 2$ , we have

$$0.83511 \leq \liminf_{d \rightarrow \infty} \rho_{m,d} \quad \text{and} \quad \limsup_{d \rightarrow \infty} \rho_{m,d} \leq 0.99804.$$

# Main results

Theorem (Beneish–K. [BK21], continued)

(C) *In the case  $(m, d) = (3, 6)$ , we compute  $\rho_{3,6} \approx 96.94\%$ .*



# Main results

## Theorem (Beneish–K. [BK21], continued)

(C) *In the case  $(m, d) = (3, 6)$ , we compute  $\rho_{3,6} \approx 96.94\%$ .*

*Moreover,  $\exists$  rational functions  $R_1(t)$  and  $R_2(t)$  such that*

$$\rho_{3,6}(p) = \begin{cases} R_1(p), & p \equiv 1 \pmod{3} \text{ and } p > 43 \\ R_2(p), & p \equiv 2 \pmod{3} \text{ and } p > 2. \end{cases}$$

*Asymptotically,*

$$1 - R_1(t) \sim \frac{2}{3}t^{-4},$$
$$1 - R_2(t) \sim \frac{53}{144}t^{-7}.$$

$$\rho = \begin{cases}
 \left( 1296p^{57} + 3888p^{56} + 9072p^{55} + 16848p^{54} + 27648p^{53} + 39744p^{52} + 53136p^{51} + 66483p^{50} + 80019p^{49} + 93141p^{48} \right. \\
 + 107469p^{47} + 120357p^{46} + 135567p^{45} + 148347p^{44} + 162918p^{43} + 176004p^{42} + 190278p^{41} + 203459p^{40} \\
 + 218272p^{39} + 232083p^{38} + 243639p^{37} + 255267p^{36} + 261719p^{35} + 264925p^{34} + 265302p^{33} + 261540p^{32} \\
 + 254790p^{31} + 250736p^{30} + 241384p^{29} + 226503p^{28} + 214137p^{27} + 195273p^{26} + 170793p^{25} + 151839p^{24} + 136215p^{23} \\
 + 118998p^{22} + 105228p^{21} + 94860p^{20} + 80471p^{19} + 67048p^{18} + 52623p^{17} + 40617p^{16} + 28773p^{15} + 19247p^{14} \\
 + 12109p^{13} + 7614p^{12} + 3420p^{11} + 756p^{10} - 2248p^9 - 4943p^8 - 6300p^7 - 6894p^6 - 5994p^5 - 2448p^4 - 648p^3 \\
 + 324p^2 + 1296p + 1296) / \left( 1296(p^{12} - p^{11} + p^9 - p^8 + p^6 - p^4 + p^3 - p + 1)(p^8 - p^6 + p^4 - p^2 + 1) \right. \\
 \times (p^6 + p^5 + p^4 + p^3 + p^2 + p + 1)(p^4 + p^3 + p^2 + p + 1)^3 (p^4 - p^3 + p^2 - p + 1)(p^2 + p + 1) \\
 \left. \times (p^2 + 1)p^{11} \right), & p \equiv 1 \pmod{3} \\
 \\
 \left( 144p^{57} + 432p^{56} + 1008p^{55} + 1872p^{54} + 3168p^{53} + 4608p^{52} + 6336p^{51} + 8011p^{50} + 9803p^{49} + 11357p^{48} \right. \\
 + 13061p^{47} + 14525p^{46} + 16295p^{45} + 17875p^{44} + 19654p^{43} + 21212p^{42} + 23030p^{41} + 24563p^{40} + 26320p^{39} \\
 + 27771p^{38} + 29711p^{37} + 30859p^{36} + 31135p^{35} + 31525p^{34} + 31510p^{33} + 29436p^{32} + 28502p^{31} + 28616p^{30} \\
 + 26856p^{29} + 25087p^{28} + 25057p^{27} + 23041p^{26} + 19921p^{25} + 18119p^{24} + 16287p^{23} + 13798p^{22} \\
 + 12140p^{21} + 10844p^{20} + 9191p^{19} + 7480p^{18} + 5839p^{17} + 4265p^{16} + 2909p^{15} + 1943p^{14} + 1109p^{13} \\
 + 590p^{12} + 604p^{11} + 372p^{10} - 144p^9 - 87p^8 - 84p^7 - 678p^6 - 618p^5 - 144p^4 - 168p^3 - 156p^2 \\
 + 144p + 144) / \left( 144(p^{12} - p^{11} + p^9 - p^8 + p^6 - p^4 + p^3 - p + 1)(p^8 - p^6 + p^4 - p^2 + 1) \right. \\
 \times (p^6 + p^5 + p^4 + p^3 + p^2 + p + 1)(p^4 + p^3 + p^2 + p + 1)^3 (p^4 - p^3 + p^2 - p + 1)(p^2 + p + 1) \\
 \left. \times (p^2 + 1)p^{11} \right), & p \equiv 2 \pmod{3}
 \end{cases}$$

# Local densities

Theorem (Beneish–K. [BK21])

(A)  $\rho_{m,d}$  exists and is given by the product of local densities,

$$\rho_{m,d} = \rho_{m,d}(\infty) \prod_p \rho_{m,d}(p) > 0.$$

$\rho_{m,d}(p)$  is (normalized) Haar measure of space of the  $\mathbb{Q}_p$ -soluble curves  $C_f: y^m = f(x, z)$ , with coefficients in  $\mathbb{Z}_p$ .

# Local densities

Theorem (Beneish–K. [BK21])

(A)  $\rho_{m,d}$  exists and is given by the product of local densities,

$$\rho_{m,d} = \rho_{m,d}(\infty) \prod_p \rho_{m,d}(p) > 0.$$

$\rho_{m,d}(p)$  is (normalized) Haar measure of space of the  $\mathbb{Q}_p$ -soluble curves  $C_f: y^m = f(x, z)$ , with coefficients in  $\mathbb{Z}_p$ .

## Idea

In *good situations*, imposing conditions at different primes looks independent...even if there's infinitely many.

# Local densities look independent

## Idea

In *good situations*, imposing conditions at different primes looks independent...*even if there's infinitely many*.

- Poonen–Stoll [[PS99a](#)] give criterion for when natural density is product of local densities.
- Apply to ELS in families of hyperelliptic curves [[PS99b](#)]; uses sieve of Ekedahl [[Eke91](#)].

# Local densities look independent


## Idea

In *good situations*, imposing conditions at different primes looks independent...*even if there's infinitely many*.

- Poonen–Stoll [[PS99a](#)] give criterion for when natural density is product of local densities.
- Apply to ELS in families of hyperelliptic curves [[PS99b](#)]; uses sieve of Ekedahl [[Eke91](#)].
- Bright–Browning–Loughran [[BBL16](#)] give *geometric criteria* when family comes from fibers of a morphism.


# Geometric picture

$$X: y^m = c_d x^d + \cdots + c_0 z^d \subset \mathbb{A}_{\mathbb{Q}}^{d+1} \times \mathbb{P}_{\mathbb{Q}} \left( 1 : \frac{d}{m} : 1 \right)$$



# Geometric picture

$$X: y^m = c_d x^d + \cdots + c_0 z^d \subset \mathbb{A}_{\mathbb{Q}}^{d+1} \times \mathbb{P}_{\mathbb{Q}} \left( 1 : \frac{d}{m} : 1 \right)$$



## Think

- A  $\mathbb{Q}$ -point  $(\mathbf{c}, [x : y : z])$  of  $X$  is the data of superelliptic curve  $C_f/\mathbb{Q}$  and a  $\mathbb{Q}$ -point  $[x : y : z] \in C_f(\mathbb{Q})$ .
- The fiber  $X_P$  of  $\pi$  over a point  $P \in \mathbb{A}^{d+1}(\mathbb{Q})$  is a superelliptic curve  $C_f/\mathbb{Q}$  whose coefficients are encoded in  $P$ .



# Computing local densities

## Question

*Once we know*

$$\rho_{m,d} = \rho(\infty) \prod_p \rho(p),$$

*how do we compute/estimate local densities  $\rho(p)$ ?*

# Computing local densities

## Question

*Once we know*

$$\rho_{m,d} = \rho(\infty) \prod_p \rho(p),$$

*how do we compute/estimate local densities  $\rho(p)$ ?*

Consider  $(m, d) = (3, 6)$  as an example,

$$C_f: y^3 = f(x, z) = c_6 x^6 + c_5 x^5 z + \cdots + c_1 x z^5 + c_0 z^6.$$

# Computing local densities

## Question

*Once we know*

$$\rho_{m,d} = \rho(\infty) \prod_p \rho(p),$$

*how do we compute/estimate local densities  $\rho(p)$ ?*

Consider  $(m, d) = (3, 6)$  as an example,

$$C_f: y^3 = f(x, z) = c_6 x^6 + c_5 x^5 z + \cdots + c_1 x z^5 + c_0 z^6.$$

(Easy mode)  $\rho_{3,6}(\infty) = 1$ , since *all*  $C_f/\mathbb{R}$  have real solutions.

# Computing local densities

$\rho_{3,6}(p)$  is (normalized) Haar measure of space of the  $\mathbb{Q}_p$ -soluble curves  $C_f: y^3 = f(x, z)$ , with coefficients in  $\mathbb{Z}_p$ .

## Think

Each possible reduction  $\bar{f}(x, z)$  occurs with equal probability.

Look mod  $p$  and check  $\mathbb{Q}_p$ -solubility with **Hensel's lemma!**

# Computing local densities

$\rho_{3,6}(p)$  is (normalized) Haar measure of space of the  $\mathbb{Q}_p$ -soluble curves  $C_f: y^3 = f(x, z)$ , with coefficients in  $\mathbb{Z}_p$ .

## Think

Each possible reduction  $\bar{f}(x, z)$  occurs with equal probability.

Look mod  $p$  and check  $\mathbb{Q}_p$ -solubility with **Hensel's lemma**!

Let  $\bar{C}_f: y^3 \equiv \bar{f}(x, z) \pmod{p}$  be the reduction.

- Smooth  $\mathbb{F}_p$ -points  $\bar{C}_f$  lift to  $\mathbb{Q}_p$ -solutions (Hensel),
- $\bar{C}_f(\mathbb{F}_p) = \emptyset$  means no  $\mathbb{Q}_p$ -solutions,
- If  $\bar{C}_f(\mathbb{F}_p)$  consists of non-smooth points, do more work.

# An extended example — bounds from geometry

If  $\overline{C}_f/\mathbb{F}_p$  smooth, irreducible, and  $p > 61$ ,

$$\#C(\mathbb{F}_p) \geq p + 1 - 8\sqrt{p} > 0,$$

so Hensel's lemma  $\implies \mathbb{C}_f$  has  $\mathbb{Q}_p$ -point!

# An extended example — bounds from geometry

If  $\overline{C}_f/\mathbb{F}_p$  smooth, irreducible, and  $p > 61$ ,

$$\#C(\mathbb{F}_p) \geq p + 1 - 8\sqrt{p} > 0,$$

so Hensel's lemma  $\implies \mathbb{C}_f$  has  $\mathbb{Q}_p$ -point!

- Refinements show  $\overline{C}_f^{\text{sm}}(\mathbb{F}_p) \neq \emptyset$  whenever  $\overline{C}_f/\mathbb{F}_p$  geom. irr. and  $p \geq 61$ .

# An extended example — bounds from geometry

If  $\overline{C}_f/\mathbb{F}_p$  smooth, irreducible, and  $p > 61$ ,

$$\#C(\mathbb{F}_p) \geq p + 1 - 8\sqrt{p} > 0,$$

so Hensel's lemma  $\implies \mathbb{C}_f$  has  $\mathbb{Q}_p$ -point!

- Refinements show  $\overline{C}_f^{\text{sm}}(\mathbb{F}_p) \neq \emptyset$  whenever  $\overline{C}_f/\mathbb{F}_p$  geom. irr. and  $p \geq 61$ .
- Irreducibility over  $\overline{\mathbb{F}}_p \iff \overline{f}(x, z) \not\equiv h(x, z)^3$  (when  $p \neq 3$ ).



# An extended example — bounds from geometry

If  $\overline{C}_f/\mathbb{F}_p$  smooth, irreducible, and  $p > 61$ ,

$$\#C(\mathbb{F}_p) \geq p + 1 - 8\sqrt{p} > 0,$$

so Hensel's lemma  $\implies \mathbb{C}_f$  has  $\mathbb{Q}_p$ -point!

- Refinements show  $\overline{C}_f^{\text{sm}}(\mathbb{F}_p) \neq \emptyset$  whenever  $\overline{C}_f/\mathbb{F}_p$  geom. irr. and  $p \geq 61$ .
- Irreducibility over  $\overline{\mathbb{F}}_p \iff \overline{f}(x, z) \not\equiv h(x, z)^3$  (when  $p \neq 3$ ).

$$\rho_{3,6}(p) \geq \frac{p^7 - p^3}{p^7} = 1 - \frac{1}{p^4} \text{ for all } p \geq 61.$$

# An extended example

- $\rho_{3,6}(p) \geq 1 - \frac{1}{p^7}$  when  $p \equiv 2 \pmod{3}$  and  $p > 2$

# An extended example

- $\rho_{3,6}(p) \geq 1 - \frac{1}{p^7}$  when  $p \equiv 2 \pmod{3}$  and  $p > 2$
- $\rho_{3,6}(p) \geq 1 - \frac{1}{p^4}$  when  $p \equiv 1 \pmod{3}$  and  $p > 43$

# An extended example

- $\rho_{3,6}(p) \geq 1 - \frac{1}{p^7}$  when  $p \equiv 2 \pmod{3}$  and  $p > 2$
- $\rho_{3,6}(p) \geq 1 - \frac{1}{p^4}$  when  $p \equiv 1 \pmod{3}$  and  $p > 43$
- Enumerate  $\bar{f}(x, z)$  and look for Hensel-liftable solutions:

$p$	$\rho_{3,6}(p) \geq$	$p$	$\rho_{3,6}(p) \geq$
2	$\frac{63}{64} \approx 0.98437$	19	$\frac{893660256}{893871739} \approx 0.99976$
3	$\frac{26}{27} \approx 0.96296$	31	$\frac{27512408250}{27512614111} \approx 0.99999$
7	$\frac{810658}{823543} \approx 0.98435$	37	$\frac{94931742132}{94931877133} \approx 0.999998$
13	$\frac{62655132}{62748517} \approx 0.99851$	43	$\frac{271818511748}{271818611107} \approx 0.9999996$

# An extended example

- $\rho_{3,6}(p) \geq 1 - \frac{1}{p^7}$  when  $p \equiv 2 \pmod{3}$  and  $p > 2$
- $\rho_{3,6}(p) \geq 1 - \frac{1}{p^4}$  when  $p \equiv 1 \pmod{3}$  and  $p > 43$
- Enumerate  $\bar{f}(x, z)$  and look for Hensel-liftable solutions:

$p$	$\rho_{3,6}(p) \geq$	$p$	$\rho_{3,6}(p) \geq$
2	$\frac{63}{64} \approx 0.98437$	19	$\frac{893660256}{893871739} \approx 0.99976$
3	$\frac{26}{27} \approx 0.96296$	31	$\frac{27512408250}{27512614111} \approx 0.99999$
7	$\frac{810658}{823543} \approx 0.98435$	37	$\frac{94931742132}{94931877133} \approx 0.999998$
13	$\frac{62655132}{62748517} \approx 0.99851$	43	$\frac{271818511748}{271818611107} \approx 0.9999996$

Taking products,

$$\rho_{3,6} = \prod_{p \text{ small}} \rho_{3,6}(p) \prod_{\substack{p \equiv 1 \pmod{3} \\ p > 43}} \left(1 - \frac{1}{p^4}\right) \prod_{\substack{p \equiv 2 \pmod{3} \\ p > 2}} \left(1 - \frac{1}{p^7}\right) \geq 0.93134.$$

# Directions from here

- 1 Varying  $(m, d)$ , similar approach produces

$$\rho_{m,d} \geq \prod_{p \text{ small}} \rho_{m,d}(p) \prod_{\substack{p \equiv 1 \pmod{m} \\ p \gg 0}} \left(1 - \frac{1}{p^{d(m-1)/m}}\right) \prod_{\substack{p \not\equiv 0,1 \pmod{m} \\ p \gg 0}} \left(1 - \frac{1}{p^{d+1}}\right).$$

# Directions from here

- 1 Varying  $(m, d)$ , similar approach produces

$$\rho_{m,d} \geq \prod_{p \text{ small}} \rho_{m,d}(p) \prod_{\substack{p \equiv 1 \pmod{m} \\ p \gg 0}} \left(1 - \frac{1}{p^{d(m-1)/m}}\right) \prod_{\substack{p \not\equiv 0,1 \pmod{m} \\ p \gg 0}} \left(1 - \frac{1}{p^{d+1}}\right).$$

- 2 Compute exact values for  $\rho_{3,6}(p)$ :
  - Determine how often factorization types occur modulo  $p$ ,
  - Determine lifting probabilities for each factorization type,
  - Solve equations relating these probabilities to others.

# Directions from here

- 1 Varying  $(m, d)$ , similar approach produces

$$\rho_{m,d} \geq \prod_{p \text{ small}} \rho_{m,d}(p) \prod_{\substack{p \equiv 1 \pmod{m} \\ p \gg 0}} \left(1 - \frac{1}{p^{d(m-1)/m}}\right) \prod_{\substack{p \not\equiv 0,1 \pmod{m} \\ p \gg 0}} \left(1 - \frac{1}{p^{d+1}}\right).$$

- 2 Compute exact values for  $\rho_{3,6}(p)$ :

- Determine how often factorization types occur modulo  $p$ ,
- Determine lifting probabilities for each factorization type,
- Solve equations relating these probabilities to others.

Treat small primes below with help from Magma

$$p = 2, 3, 7, 13, 19, 31, 37, 43.$$



## Getting exact answer

E.g. Here's what this looks like when  $f(x, z) \equiv px^6 \pmod{p^2}$ ,

$$\begin{aligned}\tau_9 &= \tau_9 a = \frac{1}{p} \tau_9 b \\ \tau_9 b &= \left(1 - \frac{1}{p}\right) + \frac{1}{p} \tau_9 c \\ \tau_9 c &= \Phi(p) + \frac{1}{p} \tau_9 d \\ \tau_9 d &= \left(1 - \frac{1}{p}\right) \left(\frac{p-1}{2p} + \frac{1}{p^2}\right) + \frac{1}{p} \tau_9 e \\ \tau_9 e &= \left(1 - \frac{1}{p}\right) + \frac{1}{p} \tau_9 f \\ \tau_9 f &= \frac{1}{p} \tau_9 g \\ \tau_9 g &= \left(1 - \frac{1}{p}\right) \alpha'' + \frac{1}{p} \tau_9 h \\ \tau_9 h &= \left(1 - \frac{1}{p}\right) \left(\frac{p-1}{2p} + \frac{\theta_2}{p}\right) + \frac{1}{p} \tau_9 i \\ \tau_9 i &= \left(1 - \frac{1}{p}\right) + \frac{1}{p} \tau_9 j \\ \tau_9 j &= \frac{1}{p} \tau_9 k \\ \tau_9 k &= \left(1 - \frac{1}{p}\right) + \frac{1}{p} \tau_9 \ell \\ \tau_9 \ell &= \Phi(p) + \left(1 - \Phi(p) - \frac{1}{p}\right) \beta + \frac{1}{p} \tau_9 m \\ \tau_9 m &= \left(1 - \frac{1}{p}\right) + \frac{1}{p} \tau_9 n \\ \tau_9 n &= \left(1 - \frac{1}{p}\right) + \frac{1}{p} \tau_9 o \\ \tau_9 o &= \Phi(p) + \frac{1}{p} \tau_9 p \\ \tau_9 p &= \sigma'_5\end{aligned}$$

[illegible]

# Final thoughts

If  $\rho_{m,d}$  is proportion of loc. sol.  $C_f: y^m = f(x, z)$  then

- (A)  $\rho_{m,d}$  given by product of local densities and  $0 < \rho_{m,d} < 1$ ;
- (B) For fixed prime  $m$ , bounding  $\rho_{m,d}(p)$  leads to explicit estimates, e.g.

$$0.83511 \leq \liminf_{d \rightarrow \infty} \rho_{m,d} \quad \text{and} \quad \limsup_{d \rightarrow \infty} \rho_{m,d} \leq 0.99804;$$

- (C) We compute  $\rho_{3,6}$  exactly,  $\approx 0.9694$ . Local densities  $\rho_{3,6}(p)$  given by rational functions.

# Further questions

What proportion of superelliptic curves  $C_f: y^m = f(x, z)$

- are *globally* soluble?
- satisfy/fail weak approximation?

Much is known for  $m = 2$  (see [BGW17]).

For  $m \nmid d$ , study density of  $y^m = f(x, z)$  with *primitive integral* solutions (see [Swa19] for  $m = 2$ ).

# Further questions

What proportion of superelliptic curves  $C_f: y^m = f(x, z)$

- are *globally* soluble?
- satisfy/fail weak approximation?

Much is known for  $m = 2$  (see [BGW17]).

For  $m \nmid d$ , study density of  $y^m = f(x, z)$  with *primitive integral* solutions (see [Swa19] for  $m = 2$ ).

Thank you!

Thank you to the organizers for hosting  
and thank **you** for your attention!

# References I



M. J. Bright, T. D. Browning, and D. Loughran, [Failures of weak approximation in families](#), Compos. Math. **152** (2016), no. 7, 1435–1475. MR 3530447



Manjul Bhargava, John Cremona, and Tom Fisher, [The proportion of plane cubic curves over  \$\mathbb{Q}\$  that everywhere locally have a point](#), Int. J. Number Theory **12** (2016), no. 4, 1077–1092. MR 3484299



———, [The proportion of genus one curves over  \$\mathbb{Q}\$  defined by a binary quartic that everywhere locally have a point](#), Int. J. Number Theory **17** (2021), no. 4, 903–923. MR 4262272



Manjul Bhargava, Benedict H. Gross, and Xiaoheng Wang, [A positive proportion of locally soluble hyperelliptic curves over  \$\mathbb{Q}\$  have no point over any odd degree extension](#), J. Amer. Math. Soc. **30** (2017), no. 2, 451–493, With an appendix by Tim Dokchitser and Vladimir Dokchitser. MR 3600041



Lea Beneish and Christopher Keyes, [On the proportion of locally soluble superelliptic curves](#), <https://arxiv.org/abs/2111.04697>, 2021.



T. D. Browning, [Many cubic surfaces contain rational points](#), Mathematika **63** (2017), no. 3, 818–839. MR 3731306



Torsten Ekedahl, [An infinite version of the Chinese remainder theorem](#), Comment. Math. Univ. St. Paul. **40** (1991), no. 1, 53–59. MR 1104780



Tom Fisher, Wei Ho, and Jennifer Park, [Everywhere local solubility for hypersurfaces in products of projective spaces](#), Res. Number Theory **7** (2021), no. 1, Paper No. 6, 27. MR 4199457



Bjorn Poonen and Michael Stoll, [The Cassels-Tate pairing on polarized abelian varieties](#), Ann. of Math. (2) **150** (1999), no. 3, 1109–1149. MR 1740984

# References II



———, A local-global principle for densities, Topics in number theory (University Park, PA, 1997), Math. Appl., vol. 467, Kluwer Acad. Publ., Dordrecht, 1999, pp. 241–244. MR 1691323



Bjorn Poonen and José Felipe Voloch, Random Diophantine equations, Arithmetic of higher-dimensional algebraic varieties (Palo Alto, CA, 2002), Progr. Math., vol. 226, Birkhäuser Boston, Boston, MA, 2004, With appendices by Jean-Louis Colliot-Thélène and Nicholas M. Katz, pp. 175–184. MR 2029869



Ashvin Swaminathan, Most integral odd-degree binary forms fail to properly represent a square, <https://arxiv.org/abs/1910.12409>, 2019.

# A geometric criterion

## Theorem (Bright–Browning–Loughran [BBL16])

Let  $\pi: X \rightarrow \mathbb{A}^n$  a dominant, quasiproj. morphism of  $\mathbb{Q}$ -varieties with geom. int. gen. fiber. Suppose

- (i) fibers above each codim. 1 point of  $\mathbb{A}^n$  are geom. integral,
- (ii)  $X(\mathbf{A}_{\mathbb{Q}}) \neq \emptyset$ ,
- (iii) For all  $B \geq 1$  we have  $B\pi(X(\mathbb{R})) \subseteq \pi(X(\mathbb{R}))$ .

Let  $\Psi' \subset \mathbb{R}^n$  be a bounded subset of positive measure lying in  $\pi(X(\mathbb{R}))$  whose boundary has measure zero. Then the limit

$$\lim_{B \rightarrow \infty} \frac{\#\{P \in \mathbb{Z}^n \cap B\Psi' \mid X_P(\mathbf{A}_{\mathbb{Q}}) \neq \emptyset\}}{\#\{P \in \mathbb{Z}^n \cap B\Psi'\}}$$

exists, is nonzero, and is equal to a product of local densities,

$$\prod_{p \nmid \infty} \mu_p(\{P \in \mathbb{Z}_p^n \mid X_P(\mathbb{Q}_p) \neq \emptyset\}).$$