

How often does a cubic hypersurface have a point?

Christopher Keyes
(King's College London)

Utrecht Geometry Center

9 April 2024

Cubic hypersurfaces

A **cubic hypersurface** $X_f \subset \mathbb{P}^n$ is cut out by a cubic form f

$$X_f: f(x_0, \dots, x_n) = \sum_{0 \leq i \leq j \leq k \leq n} a_{ijk} x_i x_j x_k = 0.$$

Cubic hypersurfaces

A **cubic hypersurface** $X_f \subset \mathbb{P}^n$ is cut out by a cubic form f

$$X_f: f(x_0, \dots, x_n) = \sum_{0 \leq i \leq j \leq k \leq n} a_{ijk} x_i x_j x_k = 0.$$

Question

For **fixed** f , how many rational points are there on X_f ?

Cubic hypersurfaces

A **cubic hypersurface** $X_f \subset \mathbb{P}^n$ is cut out by a cubic form f

$$X_f: f(x_0, \dots, x_n) = \sum_{0 \leq i \leq j \leq k \leq n} a_{ijk} x_i x_j x_k = 0.$$

Question

For **fixed** f , how many rational points are there on X_f ?

Question

As f **varies**, how often does X_f have a rational point?

Counting points

Question

For fixed f , how many rational points are there on X_f ?

Write $P \in X_f(\mathbb{Q})$ as $P = [x_0, \dots, x_n]$ with $x_i \in \mathbb{Z}$ coprime.

Define **height** of a point

$$h(P) = \max_i \{|x_i|\}.$$

Counting points

Question

For fixed f , how many rational points are there on X_f ?

Write $P \in X_f(\mathbb{Q})$ as $P = [x_0, \dots, x_n]$ with $x_i \in \mathbb{Z}$ coprime.

Define **height** of a point

$$h(P) = \max_i \{|x_i|\}.$$

Natural point-counting function

$$N_f(B) = \#\{P \in X_f(\mathbb{Q}) \mid h(P) \leq B\}.$$

Circle method

Introduced by Hardy and Littlewood to count things.

$$N_f(B) = \sum_{\substack{\mathbf{x} \in \mathbb{Z}^{n+1} \\ h(\mathbf{x}) \leq B}} \int_0^1 \left(e^{2\pi i f(\mathbf{x})\alpha} d\alpha \right) = \int_0^1 \sum_{\substack{\mathbf{x} \in \mathbb{Z}^{n+1} \\ h(\mathbf{x}) \leq B}} e^{2\pi i f(\mathbf{x})\alpha} d\alpha$$

Circle method

Introduced by Hardy and Littlewood to count things.

$$N_f(B) = \sum_{\substack{\mathbf{x} \in \mathbb{Z}^{n+1} \\ h(\mathbf{x}) \leq B}} \int_0^1 \left(e^{2\pi i f(\mathbf{x})\alpha} d\alpha \right) = \int_0^1 \sum_{\substack{\mathbf{x} \in \mathbb{Z}^{n+1} \\ h(\mathbf{x}) \leq B}} e^{2\pi i f(\mathbf{x})\alpha} d\alpha$$

Estimate sums when α is

- (major arcs) α well approximated by $\frac{a}{q} \in \mathbb{Q}$, otherwise
- (minor arcs) negligible contribution.

For major arc: count solutions **modulo** q .

Some history

When n large enough¹, circle method shows

$$N_f(B) \sim c_f B^{n-3}, \quad c_f > 0,$$

i.e. X_f always has a rational point.

¹Recall n denotes the dimension of \mathbb{P}^n ; the number of variables is $n + 1$

Some history

When n large enough¹, circle method shows

$$N_f(B) \sim c_f B^{n-3}, \quad c_f > 0,$$

i.e. X_f always has a rational point.

1950's Birch, Lewis, then Davenport: $n \geq 15$

¹Recall n denotes the dimension of \mathbb{P}^n ; the number of variables is $n + 1$

Some history

When n large enough¹, circle method shows

$$N_f(B) \sim c_f B^{n-3}, \quad c_f > 0,$$

i.e. X_f always has a rational point.

1950's Birch, Lewis, then Davenport: $n \geq 15$

1982 Heath-Brown: $n \geq 9$ if X_f is nonsingular [HB83]

¹Recall n denotes the dimension of \mathbb{P}^n ; the number of variables is $n + 1$

Some history

When n large enough¹, circle method shows

$$N_f(B) \sim c_f B^{n-3}, \quad c_f > 0,$$

i.e. X_f always has a rational point.

1950's Birch, Lewis, then Davenport: $n \geq 15$

1982 Heath-Brown: $n \geq 9$ if X_f is nonsingular [HB83]

1988 Hooley: $n \geq 8$ if X_f is everywhere locally soluble

¹Recall n denotes the dimension of \mathbb{P}^n ; the number of variables is $n + 1$

Some history

When n large enough¹, circle method shows

$$N_f(B) \sim c_f B^{n-3}, \quad c_f > 0,$$

i.e. X_f always has a rational point.

1950's Birch, Lewis, then Davenport: $n \geq 15$

1982 Heath-Brown: $n \geq 9$ if X_f is nonsingular [HB83]

1988 Hooley: $n \geq 8$ if X_f is everywhere locally soluble

Question

For $n \leq 8$, as f varies, how often does X_f have a rational point?

¹Recall n denotes the dimension of \mathbb{P}^n ; the number of variables is $n + 1$

Back to cubic hypersurfaces

Question

How often does X_f have a rational point?

$\mathbb{A}^{\binom{n+3}{3}} - 0$ is “moduli space” of cubic forms, each f is integer point.

Back to cubic hypersurfaces

Question

How often does X_f have a rational point?

$\mathbb{A}^{\binom{n+3}{3}} - 0$ is “moduli space” of cubic forms, each f is integer point.

Let $h(f) = \|\mathbf{a}\| = \left(\sum_{i,j,k} a_{ijk}^2 \right)^{1/2}$, define **natural density**

$$\rho_n = \lim_{B \rightarrow \infty} \frac{\#\{f \mid h(f) \leq B, X_f(\mathbb{Q}) \neq \emptyset\}}{\#\{f \mid h(f) \leq B\}}.$$

Back to cubic hypersurfaces

Question

How often does X_f have a rational point?

$\mathbb{A}^{\binom{n+3}{3}} - 0$ is “moduli space” of cubic forms, each f is integer point.

Let $h(f) = \|\mathbf{a}\| = \left(\sum_{i,j,k} a_{ijk}^2 \right)^{1/2}$, define **natural density**

$$\rho_n = \lim_{B \rightarrow \infty} \frac{\#\{f \mid h(f) \leq B, X_f(\mathbb{Q}) \neq \emptyset\}}{\#\{f \mid h(f) \leq B\}}.$$

Remark

Counting **primitive** forms gets same answer, i.e. using $\mathbb{P}^{\binom{n+3}{3}-1}$.

Everywhere local solubility

A variety X/\mathbb{Q} is **everywhere locally soluble** (ELS) if

$$X(\mathbb{R}) \neq \emptyset \text{ and } X(\mathbb{Q}_p) \neq \emptyset \text{ for all } p.$$

Everywhere local solubility

A variety X/\mathbb{Q} is **everywhere locally soluble** (ELS) if

$$X(\mathbb{R}) \neq \emptyset \text{ and } X(\mathbb{Q}_p) \neq \emptyset \text{ for all } p.$$

ELS is

- **necessary** for X_f to have \mathbb{Q} -point
- but not **sufficient**

Everywhere local solubility

A variety X/\mathbb{Q} is **everywhere locally soluble** (ELS) if

$$X(\mathbb{R}) \neq \emptyset \text{ and } X(\mathbb{Q}_p) \neq \emptyset \text{ for all } p.$$

ELS is

- **necessary** for X_f to have \mathbb{Q} -point
- but not **sufficient**...or is it?

Everywhere local solubility

A variety X/\mathbb{Q} is **everywhere locally soluble** (ELS) if

$$X(\mathbb{R}) \neq \emptyset \text{ and } X(\mathbb{Q}_p) \neq \emptyset \text{ for all } p.$$

ELS is

- **necessary** for X_f to have \mathbb{Q} -point
- but not **sufficient**...or is it?

Density of ELS cubic forms in $\mathbb{Z}[x_0, \dots, x_n]$:

$$\rho_n^{ELS} = \lim_{B \rightarrow \infty} \frac{\#\{f \mid h(f) \leq B, X_f \text{ ELS}\}}{\#\{f \mid h(f) \leq B\}}.$$

(Lack of) obstructions

Conjecture (Poonen–Voloch, 2004)

When $n \geq 3$, $\rho_n^{ELS} = \rho_n$.

i.e. local-global principle holds for 100% of cubic hypersurfaces.

(Lack of) obstructions

Conjecture (Poonen–Voloch, 2004)

When $n \geq 3$, $\rho_n^{ELS} = \rho_n$.

i.e. local-global principle holds for 100% of cubic hypersurfaces.

- Conj. of Colliot–Thélène: Brauer–Manin is only obstruction
- Brauer–Manin vacuous for cubics when $n \geq 4$
- Brauer–Manin vacuous for 100% of cubic surfaces in \mathbb{P}^3

(Lack of) obstructions

Conjecture (Poonen–Voloch, 2004)

When $n \geq 3$, $\rho_n^{ELS} = \rho_n$.

i.e. local-global principle holds for 100% of cubic hypersurfaces.

- Conj. of Colliot–Thélène: Brauer–Manin is only obstruction
- Brauer–Manin vacuous for cubics when $n \geq 4$
- Brauer–Manin vacuous for 100% of cubic surfaces in \mathbb{P}^3

Theorem (Browning–Le Boudec–Sawin, 2023)

When $n \geq 4$, $\rho_n^{ELS} = \rho_n$.

(Lack of) obstructions

Conjecture (Poonen–Voloch, 2004)

When $n \geq 3$, $\rho_n^{ELS} = \rho_n$.

i.e. local-global principle holds for 100% of cubic hypersurfaces.

- Conj. of Colliot–Thélène: Brauer–Manin is only obstruction
- Brauer–Manin vacuous for cubics when $n \geq 4$
- Brauer–Manin vacuous for 100% of cubic surfaces in \mathbb{P}^3

Theorem (Browning–Le Boudec–Sawin, 2023)

When $n \geq 4$, $\rho_n^{ELS} = \rho_n$. In fact *true for Fano deg. d* $(d, n) \neq (3, 3)$

Computing ρ^{ELS}

Let $\rho_n(p)$ = density of p -adic cubic forms f such that $X_f(\mathbb{Q}_p) \neq \emptyset$.

Theorem (Poonen–Voloch, 2004)

Let $n \geq 2$. We have

$$\rho_n^{ELS} = \prod_p \rho_n(p).$$

Computing ρ^{ELS}

Let $\rho_n(p)$ = density of p -adic cubic forms f such that $X_f(\mathbb{Q}_p) \neq \emptyset$.

Theorem (Poonen–Voloch, 2004)

Let $n \geq 2$. We have

$$\rho_n^{ELS} = \prod_p \rho_n(p).$$

Think

Local probabilities independent...even though infinitely many!

Computing ρ^{ELS}

Let $\rho_n(p)$ = density of p -adic cubic forms f such that $X_f(\mathbb{Q}_p) \neq \emptyset$.

Theorem (Poonen–Voloch, 2004)

Let $n \geq 2$. We have


$$\rho_n^{ELS} = \prod_p \rho_n(p).$$

Think

Local probabilities independent...even though infinitely many!

2016 Bright–Browning–Loughran: flexible generalization to families given by fibers of maps to affine/projective space.

Varieties parameterized by fibers



Main result

Theorem (Beneish–K.)

Let $4 \leq n \leq 8$. There exist *explicit polynomials* $g_n(t), h_n(t) \in \mathbb{Z}[t]$ describing ρ_n exactly as Euler product

$$\rho_n = \prod_p \rho_n(p) = \prod_p \left(1 - \frac{g_n(p)}{h_n(p)} \right).$$

Main result

Theorem (Beneish–K.)

Let $4 \leq n \leq 8$. There exist *explicit polynomials* $g_n(t), h_n(t) \in \mathbb{Z}[t]$ describing ρ_n exactly as Euler product

$$\rho_n = \prod_p \rho_n(p) = \prod_p \left(1 - \frac{g_n(p)}{h_n(p)} \right).$$

Remark

- We produce g_3, h_3 , and *conjectural* formula for ρ_3 .
- Recovers $\rho_n(p) = 1$ for $n \geq 9$.

Cubic 3-folds in \mathbb{P}^4

Example

When $n = 4$ we have

$$\begin{aligned} g_4(p) &= (p^{46} + 3p^{41} + p^{40} - p^{39} + p^{37} + p^{36} + p^{35} - 3p^{34} + 3p^{27} - p^{26} + p^{25} \\ &\quad + p^{19}) (p^2 + 1) (p + 1)^2 (p - 1)^4 \\ h_4(p) &= 9 (p^{19} - 1) (p^{17} - 1) (p^{10} + 1) (p^9 + 1) (p^9 - 1) (p^7 - 1) (p^5 + 1) \end{aligned}$$

Asymptotically, $\frac{g_4(p)}{h_4(p)} \sim \frac{1}{9p^{22}}.$

Numerically, $\rho_4 \approx 0.99999999497 = 1 - 5.022 \cdot 10^{-9}.$

Asymptotics and numerics

Theorem (Beneish–K.)

$$\rho_n = \prod_p \left(1 - \frac{g_n(p)}{h_n(p)} \right)$$

n	$\rho_n \approx$	$g_n(p)/h_n(p) \sim$
3	0.999927 (<i>conj.</i>)	$1/3p^{10}$
4	$1 - 5.022 \cdot 10^{-9}$	$1/9p^{22}$
5	$1 - 1.343 \cdot 10^{-15}$	$1/9p^{43}$
6	$1 - 3.502 \cdot 10^{-26}$	$1/9p^{78}$
7	$1 - 5.152 \cdot 10^{-42}$	$1/27p^{129}$
8	$1 - 6.222 \cdot 10^{-64}$	$1/27p^{201}$

Asymptotics and numerics

Theorem (Beneish–K.)

$$\rho_n = \prod_p \left(1 - \frac{g_n(p)}{h_n(p)} \right)$$

n	$\rho_n \approx$	$g_n(p)/h_n(p) \sim$
3	0.999927 (<i>conj.</i>)	$1/3p^{10}$
4	$1 - 5.022 \cdot 10^{-9}$	$1/9p^{22}$
5	$1 - 1.343 \cdot 10^{-15}$	$1/9p^{43}$
6	$1 - 3.502 \cdot 10^{-26}$	$1/9p^{78}$
7	$1 - 5.152 \cdot 10^{-42}$	$1/27p^{129}$
8	$1 - 6.222 \cdot 10^{-64}$	$1/27p^{201}$
9	1	0

Roadmap

ρ_n = density of cubic forms $f \in \mathbb{Z}[x_0, \dots, x_n]$ with $X_f(\mathbb{Q}) \neq \emptyset$.

$$\textcircled{1} \quad \rho_n = \rho_n^{ELS} = \prod_p \rho_n(p) \quad [\text{BLBS23, PV04, BBL16}]$$

Roadmap

ρ_n = density of cubic forms $f \in \mathbb{Z}[x_0, \dots, x_n]$ with $X_f(\mathbb{Q}) \neq \emptyset$.

- 1 $\rho_n = \rho_n^{ELS} = \prod_p \rho_n(p)$ [BLBS23, PV04, BBL16]
- 2 Study possible reductions $\overline{X_f}$ modulo p

Roadmap

ρ_n = density of cubic forms $f \in \mathbb{Z}[x_0, \dots, x_n]$ with $X_f(\mathbb{Q}) \neq \emptyset$.

- 1 $\rho_n = \rho_n^{ELS} = \prod_p \rho_n(p)$ [BLBS23, PV04, BBL16]
- 2 Study possible reductions $\overline{X_f}$ modulo p
- 3 How often does $\overline{P} \in \overline{X_f}$ lift to $P \in X_f(\mathbb{Q}_p)$?

Roadmap

ρ_n = density of cubic forms $f \in \mathbb{Z}[x_0, \dots, x_n]$ with $X_f(\mathbb{Q}) \neq \emptyset$.

- 1 $\rho_n = \rho_n^{ELS} = \prod_p \rho_n(p)$ [BLBS23, PV04, BBL16]
- 2 Study possible reductions $\overline{X_f}$ modulo p
- 3 How often does $\overline{P} \in \overline{X_f}$ lift to $P \in X_f(\mathbb{Q}_p)$?

Idea

Express lifting probabilities in terms of each other and recurse to get rational function $\rho_n(p)$.

Counting points
○○○○


Varying X_f
○○○○○

Results
○○○○●○

Lifting probabilities
○○○○○○○○○○○○○○

Final thoughts
○○○○

Full picture



Related results

Plane cubic curves [BCF16a, Bha14]

- ρ_2^{ELS} , $\rho_2(p)$ computed by Bhargava–Cremona–Fisher
- Bhargava: pos. prop. fail Hasse principle (conj. 65%)

Related results

Plane cubic curves [BCF16a, Bha14]

- ρ_2^{ELS} , $\rho_2(p)$ computed by Bhargava–Cremona–Fisher
- Bhargava: pos. prop. fail Hasse principle (conj. 65%)

Quadric hypersurfaces [BCF⁺16b]

- BCF–Jones–Keating: explicit Euler product for density of quadratic forms with integral zero
- Hasse principle holds but archimedean place not trivial!
- 98.3% of quadric surfaces in \mathbb{P}^3 soluble

Related results

Plane cubic curves [BCF16a, Bha14]

- ρ_2^{ELS} , $\rho_2(p)$ computed by Bhargava–Cremona–Fisher
- Bhargava: pos. prop. fail Hasse principle (conj. 65%)

Quadric hypersurfaces [BCF⁺16b]

- BCF–Jones–Keating: explicit Euler product for density of quadratic forms with integral zero
- Hasse principle holds but archimedean place not trivial!
- 98.3% of quadric surfaces in \mathbb{P}^3 soluble

More: certain cubic surfaces [Bro17], (2,2)-curves in $\mathbb{P}^1 \times \mathbb{P}^1$
[FHP21]

Computing the local factors

Goal

Compute local probability $\rho_n(p)$ that X_f has \mathbb{Q}_p -point.

Computing the local factors

Goal

Compute local probability $\rho_n(p)$ that X_f has \mathbb{Q}_p -point.

$\rho_n(p)$ is p -adic Haar measure with $\mu_p \left(\mathbb{Z}_p^{\binom{n+3}{3}} \right) = 1$:

$$\rho_n(p) = \mu_p \left(f \in \mathbb{Z}_p[x_0, \dots, x_n] \mid X_f(\mathbb{Q}_p) \neq \emptyset \right).$$

Think

Reduce mod p , try to decide solubility with **Hensel's lemma**.

Computing the local factors

Goal

Compute local probability $\rho_n(p)$ that X_f has \mathbb{Q}_p -point.

$\rho_n(p)$ is p -adic Haar measure with $\mu_p \left(\mathbb{Z}_p^{\binom{n+3}{3}} \right) = 1$:

$$\rho_n(p) = \mu_p \left(f \in \mathbb{Z}_p[x_0, \dots, x_n] \mid X_f(\mathbb{Q}_p) \neq \emptyset \right).$$


Think

Reduce mod p , try to decide solubility with **Hensel's lemma**.


Lemma (Hensel's Lemma)

If $P_0 \in \overline{X}^{\text{sm}}(\mathbb{F}_p)$, then **there exists lift** $P \in X(\mathbb{Z}_p)$ with $\overline{P} = P_0$.

Cubic hypersurfaces over \mathbb{F}_p



Def. over
 \mathbb{F}_p
 \mathbb{F}_{p^2}
 \mathbb{F}_{p^3}



When are there always \mathbb{Q}_p -points?

Proposition

If $\overline{X_f}$ *not* a config. of conjugate hyperplanes, then $X_f(\mathbb{Q}_p) \neq \emptyset$.

Proof for curves ($n = 2$).

If geom. integral, use Hasse–Weil bounds on (normalization of) $\overline{X_f}$

$$\#X_f(\mathbb{F}_p) \geq p + 1 - 2\sqrt{p} > 0.$$

All other possibilities contain line defined over \mathbb{F}_p .



When are there always \mathbb{Q}_p -points?

Proposition

If $\overline{X_f}$ *not* a config. of conjugate hyperplanes, then $X_f(\mathbb{Q}_p) \neq \emptyset$.

Proof for curves ($n = 2$).

If geom. integral, use Hasse–Weil bounds on (normalization of) $\overline{X_f}$

$$\#X_f(\mathbb{F}_p) \geq p + 1 - 2\sqrt{p} > 0.$$

All other possibilities contain line defined over \mathbb{F}_p . □

For $n \geq 3$, use inductive argument reducing to *coordinate hypersurfaces* $X_i = X_f \cap \{x_i = 0\} \subset \mathbb{P}^{n-1}$.

When are there always \mathbb{Q}_p -points?

Proposition

If $\overline{X_f}$ *not* a config. of conjugate hyperplanes, then $X_f(\mathbb{Q}_p) \neq \emptyset$.

Proof for curves ($n = 2$).

If geom. integral, use Hasse–Weil bounds on (normalization of) $\overline{X_f}$

$$\#X_f(\mathbb{F}_p) \geq p + 1 - 2\sqrt{p} > 0.$$

All other possibilities contain line defined over \mathbb{F}_p . □

For $n \geq 3$, use inductive argument reducing to *coordinate hypersurfaces* $X_i = X_f \cap \{x_i = 0\} \subset \mathbb{P}^{n-1}$.

Warning

There exist irreducible deg. $d > 3$ curves $X \subset \mathbb{P}^2$ with $X(\mathbb{Q}_p) = \emptyset$.

Some (non)examples

Example ($n = 2, p = 7$)

Consider the plane cubic curve over \mathbb{Z}_7 defined by

$$f(x, y, z) = x^3 + 3x^2y + 3xy^2 - 6xyz + 3y^3 - 6y^2z + 4z^3 = 0.$$

Some (non)examples

Example ($n = 2$, $p = 7$)

Consider the plane cubic curve over \mathbb{Z}_7 defined by

$$f(x, y, z) = x^3 + 3x^2y + 3xy^2 - 6xyz + 3y^3 - 6y^2z + 4z^3 = 0.$$

We have \bar{f} irreducible over \mathbb{F}_7 , but over \mathbb{F}_{7^3} ,

$$\bar{f} = \prod_{\sigma \in \text{Gal}(\mathbb{F}_{7^3}/\mathbb{F}_7)} \sigma(x + (1 + u)y + u^2z)$$

where $u \in \mathbb{F}_{7^3}$, $u^3 = 2$.

Some (non)examples

Example ($n = 2$, $p = 7$)

Consider the plane cubic curve over \mathbb{Z}_7 defined by

$$f(x, y, z) = x^3 + 3x^2y + 3xy^2 - 6xyz + 3y^3 - 6y^2z + 4z^3 = 0.$$

We have \bar{f} irreducible over \mathbb{F}_7 , but over \mathbb{F}_{7^3} ,

$$\bar{f} = \prod_{\sigma \in \text{Gal}(\mathbb{F}_{7^3}/\mathbb{F}_7)} \sigma(x + (1 + u)y + u^2z)$$

where $u \in \mathbb{F}_{7^3}$, $u^3 = 2$.

$\overline{X_f}$ is a **triangle**. $\overline{X_f}(\mathbb{F}_7) = \emptyset \implies X_f(\mathbb{Z}_7) = \emptyset$.

Some (non)examples

Example

Suppose $f = g_1(x_0, x_1, x_2) + pg_2(x_3, x_4, x_5) + p^2g_3(x_6, x_7, x_8)$
for $g_i = 0$ with **no nontrivial \mathbb{F}_p -solutions**.

If $[x_0, \dots, x_8] \in X_f(\mathbb{Z}_p)$ then $p \mid x_0, x_1, x_2$

Some (non)examples

Example

Suppose $f = g_1(x_0, x_1, x_2) + pg_2(x_3, x_4, x_5) + p^2g_3(x_6, x_7, x_8)$
for $g_i = 0$ with **no nontrivial \mathbb{F}_p -solutions**.

If $[x_0, \dots, x_8] \in X_f(\mathbb{Z}_p)$ then $p \mid x_0, x_1, x_2$

$$\frac{1}{p}f(px_0, px_1, px_2, x_3, \dots, x_8)$$

$$= g_2(x_3, x_4, x_5) + pg_3(x_6, x_7, x_8) + p^2g_1(x_0, x_1, x_2)$$

$$\implies X_f(\mathbb{Z}_p) = \emptyset.$$

Some (non)examples

Example

Suppose $f = g_1(x_0, x_1, x_2) + pg_2(x_3, x_4, x_5) + p^2g_3(x_6, x_7, x_8)$
for $g_i = 0$ with **no nontrivial \mathbb{F}_p -solutions**.

If $[x_0, \dots, x_8] \in X_f(\mathbb{Z}_p)$ then $p \mid x_0, x_1, x_2$

$$\begin{aligned} \frac{1}{p} f(px_0, px_1, px_2, x_3, \dots, x_8) \\ = g_2(x_3, x_4, x_5) + pg_3(x_6, x_7, x_8) + p^2g_1(x_0, x_1, x_2) \end{aligned}$$



$$\implies X_f(\mathbb{Z}_p) = \emptyset.$$

Remark


This generalizes, but it ignores **cross terms**...

Configurations of conjugate lines

1 - triple line





3 - triangle




2 - star

Configurations of conjugate hyperplanes

1-~~triple~~ (hyper)plane



3-triangle



2-star

Strategy

Goal

Look modulo p and try to decide solubility

$$\rho_n(p) = \xi_{n,0}\sigma_{n,0} + \xi_{n,1}\sigma_{n,1} + \xi_{n,2}\sigma_{n,2} + \xi_{n,3}\sigma_{n,3}$$

Strategy

Goal

Look modulo p and try to decide solubility

$$\rho_n(p) = \xi_{n,0}\sigma_{n,0} + \xi_{n,1}\sigma_{n,1} + \xi_{n,2}\sigma_{n,2} + \xi_{n,3}\sigma_{n,3}$$

- $\xi_{n,i}$ = prob. f has type i

Strategy

Goal

Look modulo p and try to decide solubility

$$\rho_n(p) = \xi_{n,0}\sigma_{n,0} + \xi_{n,1}\sigma_{n,1} + \xi_{n,2}\sigma_{n,2} + \xi_{n,3}\sigma_{n,3}$$

- $\xi_{n,i}$ = prob. f has type i
- $\sigma_{n,i}$ = prob. $X_f(\mathbb{Q}_p) \neq \emptyset$ given f type i

Strategy

Goal

Look modulo p and try to decide solubility

$$\rho_n(p) = \xi_{n,0}\sigma_{n,0} + \xi_{n,1}\sigma_{n,1} + \xi_{n,2}\sigma_{n,2} + \xi_{n,3}\sigma_{n,3}$$

- $\xi_{n,i}$ = prob. f has type i
- $\sigma_{n,i}$ = prob. $X_f(\mathbb{Q}_p) \neq \emptyset$ given f type i
- $\xi_{n,0}$ = prob. f **not** config. of conj. hyperplanes

Strategy

Goal

Look modulo p and try to decide solubility

$$\rho_n(p) = \xi_{n,0} \cdot 1 + \xi_{n,1} \sigma_{n,1} + \xi_{n,2} \sigma_{n,2} + \xi_{n,3} \sigma_{n,3}$$

- $\xi_{n,i}$ = prob. f has type i
- $\sigma_{n,i}$ = prob. $X_f(\mathbb{Q}_p) \neq \emptyset$ given f type i
- $\xi_{n,0}$ = prob. f not config. of conj. hyperplanes
- $\sigma_{n,0} = 1$

Factorization probabilities

$$\xi_{n,0} = 1 - \frac{p^{3n-3} + 2p^{n+3} + 2p^{n+2} + 2p^{n+1} - 2p^2 - 2p - 3}{3(p^2 + p + 1) \left(p^{\binom{n+3}{3}} - 1 \right)}$$

$$\xi_{n,1} = \frac{p^{n+1} - 1}{p^{\binom{n+3}{3}} - 1}$$

$$\xi_{n,2} = \frac{(p^{2n+1} - p^{n+1} - p^n + 1)p}{3 \left(p^{\binom{n+3}{3}} - 1 \right)}$$

$$\xi_{n,3} = \frac{(p^{3n} - p^{2n} - p^{2n+1} - p^{2n-1} + p^{n+1} + p^{n-1} + p^n - 1)p^3}{3(p^2 + p + 1) \left(p^{\binom{n+3}{3}} - 1 \right)}$$

Exercise

Convince yourself that probability of a polynomial factoring a certain way is given by a (uniform) rational function.

Phases I, II, and III

Repeat this process three times:

- 1 Reduce mod p , lift any “easy” solutions with Hensel’s lemma

Phases I, II, and III

Repeat this process three times:

- 1 Reduce mod p , lift any “easy” solutions with Hensel’s lemma
- 2 Reduction is type $j = 1, 2, 3$ with explicit probability

Phases I, II, and III

Repeat this process three times:

- 1 Reduce mod p , lift any “easy” solutions with Hensel’s lemma
- 2 Reduction is type $j = 1, 2, 3$ with explicit probability
- 3 Introduce new lifting probability for each type

Phases I, II, and III

Repeat this process three times:

- 1 Reduce mod p , lift any “easy” solutions with Hensel’s lemma
- 2 Reduction is type $j = 1, 2, 3$ with explicit probability
- 3 Introduce new lifting probability for each type
- 4 Relate new lifting probabilities to others

Eventually this terminates: 64 $\mathbb{Q}(p)$ -linear relations in 64 unknowns

Counting points
○○○○


Varying X_f
○○○○○

Results
○○○○○○

Lifting probabilities
○○○○○○○○○○●○

Final thoughts
○○○○

Setup



Counting points
○○○○


Varying X_f
○○○○○

Results
○○○○○○

Lifting probabilities
○○○○○○○○○○●●○

Final thoughts
○○○○

Phase I



Counting points
○○○○


Varying X_f
○○○○○

Results
○○○○○○

Lifting probabilities
○○○○○○○○○○●●

Final thoughts
○○○○

Phase I



θ_{111} θ_{112} θ_{113} θ_{121} θ_{122} θ_{123} θ_{131} θ_{132} θ_{133} θ_{211} θ_{212} θ_{213} θ_{221} θ_{222} θ_{223} θ_{231} θ_{232} θ_{233} θ_{311} θ_{312} θ_{313} θ_{321} θ_{322} θ_{323} θ_{331} θ_{332} θ_{333}

Counting points
○○○○


Varying X_f
○○○○○

Results
○○○○○○

Lifting probabilities
○○○○○○○○○○●●○

Final thoughts
○○○○

Phase II



Counting points
○○○○


Varying X_f
○○○○○

Results
○○○○○○

Lifting probabilities
○○○○○○○○○○●●○

Final thoughts
○○○○

Phase II



Counting points
○○○○


Varying X_f
○○○○○

Results
○○○○○○

Lifting probabilities
○○○○○○○○○○○○●●

Final thoughts
○○○○

Phase III



Counting points
oooo


Varying X_f
ooooo

Results
oooooo

Lifting probabilities
oooooooooooo●●o

Final thoughts
oooo

When $n = 8$



NO LIFT

Some remarks

Practicalities:

- Solve with Sage symbolic solver
- Block variables ($27 + 27 + 10$) to speed up

Some remarks

Practicalities:

- Solve with Sage symbolic solver
- Block variables $(27 + 27 + 10)$ to speed up

In principle, $\rho_n(p) = 1$ for $n \geq 9$ can be seen “by hand”

- No “no lift” sink in the flow chart!
- Capture and describe **explicitly** Heath–Brown’s observation of necessary/sufficient conditions for f to have local solutions

Density of cubics with a point

Theorem (Beneish–K.)

Let $n \geq 4$ (conjecturally $n \geq 3$). Then $\rho_n = 1$ when $n \geq 9$ and

$$\rho_n = \prod_{p \text{ prime}} \left(1 - \frac{g_n(p)}{h_n(p)} \right) \text{ when } n \leq 8$$

for *explicit polynomials* $g_n(t), h_n(t) \in \mathbb{Z}[t]$.

n	$\rho_n^{(ELS)} \approx$	$1 - \rho_n(p) \sim$
2	0.9726 [BCF16a]	$1/3p^3$
3	0.999927 (conj.)	$1/3p^{10}$
4	$1 - 5.022 \cdot 10^{-9}$	$1/9p^{22}$
5	$1 - 1.343 \cdot 10^{-15}$	$1/9p^{43}$
6	$1 - 3.502 \cdot 10^{-26}$	$1/9p^{78}$
7	$1 - 5.152 \cdot 10^{-42}$	$1/27p^{129}$
8	$1 - 6.222 \cdot 10^{-64}$	$1/27p^{201}$

Further questions

Let $\rho_{d,n}(p)$ = density of deg. d hypersurfaces in \mathbb{P}^n with \mathbb{Q}_p -point

Is $\rho_{d,n}(p)$ always rational function in p ?

Further questions

Let $\rho_{d,n}(p)$ = density of deg. d hypersurfaces in \mathbb{P}^n with \mathbb{Q}_p -point

Is $\rho_{d,n}(p)$ always rational function in p ?

- Known for **binary forms**, $\rho_{d,1}(p) = \rho_{d,1}(1/p)$ [BCFG22]

Further questions

Let $\rho_{d,n}(p)$ = density of deg. d hypersurfaces in \mathbb{P}^n with \mathbb{Q}_p -point

Is $\rho_{d,n}(p)$ always rational function in p ?

- Known for **binary forms**, $\rho_{d,1}(p) = \rho_{d,1}(1/p)$ [BCFG22]
- **Lose uniformity** in p in general:
E.g. $\rho_{4,17}(2) \neq 1$, but $\rho_{4,17}(p) = 1$ for $p \gg 0$
- Heath–Brown: $\rho_{4,n}(p) = 1$ for $n \geq 9126$,
 $\rho_{5,n}(p) = 1$ known for $n \geq 25$, $p \geq 17$ [HB09]

Further questions

Let $\rho_{d,n}(p)$ = density of deg. d hypersurfaces in \mathbb{P}^n with \mathbb{Q}_p -point

Is $\rho_{d,n}(p)$ always rational function in p ?

- Known for **binary forms**, $\rho_{d,1}(p) = \rho_{d,1}(1/p)$ [BCFG22]
- **Lose uniformity** in p in general:
E.g. $\rho_{4,17}(2) \neq 1$, but $\rho_{4,17}(p) = 1$ for $p \gg 0$
- Heath–Brown: $\rho_{4,n}(p) = 1$ for $n \geq 9126$,
 $\rho_{5,n}(p) = 1$ known for $n \geq 25$, $p \geq 17$ [HB09]
- Can we **predict** asymptotics/numerics with less effort?

Further questions

Let $\rho_{d,n}(p)$ = density of deg. d hypersurfaces in \mathbb{P}^n with \mathbb{Q}_p -point

Is $\rho_{d,n}(p)$ always rational function in p ?

- Known for **binary forms**, $\rho_{d,1}(p) = \rho_{d,1}(1/p)$ [BCFG22]
- **Lose uniformity** in p in general:
E.g. $\rho_{4,17}(2) \neq 1$, but $\rho_{4,17}(p) = 1$ for $p \gg 0$
- Heath–Brown: $\rho_{4,n}(p) = 1$ for $n \geq 9126$,
 $\rho_{5,n}(p) = 1$ known for $n \geq 25$, $p \geq 17$ [HB09]
- Can we **predict** asymptotics/numerics with less effort?

Thank you I

Thank you for the invitation and for your attention!



M. J. Bright, T. D. Browning, and D. Loughran, Failures of weak approximation in families, *Compos. Math.* **152** (2016), no. 7, 1435–1475. MR 3530447



Manjul Bhargava, John Cremona, and Tom Fisher, The proportion of plane cubic curves over \mathbb{Q} that everywhere locally have a point, *Int. J. Number Theory* **12** (2016), no. 4, 1077–1092. MR 3484299



Manjul Bhargava, John E. Cremona, Tom Fisher, Nick G. Jones, and Jonathan P. Keating, What is the probability that a random integral quadratic form in n variables has an integral zero?, *Int. Math. Res. Not. IMRN* (2016), no. 12, 3828–3848. MR 3544620



Manjul Bhargava, John Cremona, Tom Fisher, and Stevan Gajović, The density of polynomials of degree nn over \mathbb{Z}_p having exactly r roots in \mathbb{q}_p , *Proceedings of the London Mathematical Society* **124** (2022), no. 5, 713–736 (English (US)).



Manjul Bhargava, A positive proportion of plane cubics fail the hasse principle, Preprint, available at <https://arxiv.org/pdf/1402.1131.pdf>, 2014.



Tim Browning, Pierre Le Boudec, and Will Sawin, The Hasse principle for random Fano hypersurfaces, *Ann. of Math. (2)* **197** (2023), no. 3, 1115–1203. MR 4564262



T. D. Browning, Many cubic surfaces contain rational points, *Mathematika* **63** (2017), no. 3, 818–839. MR 3731306

Thank you II



Tom Fisher, Wei Ho, and Jennifer Park, [Everywhere local solubility for hypersurfaces in products of projective spaces](#), Res. Number Theory **7** (2021), no. 1, Paper No. 6, 27. MR 4199457



D. R. Heath-Brown, [Cubic forms in ten variables](#), Proc. London Math. Soc. (3) **47** (1983), no. 2, 225–257. MR 703978



———, [Zeros of p-adic forms](#), Proceedings of the London Mathematical Society **100** (2009), no. 2, 560–584.



Bjorn Poonen and José Felipe Voloch, [Random Diophantine equations](#), Arithmetic of higher-dimensional algebraic varieties (Palo Alto, CA, 2002), Progr. Math., vol. 226, Birkhäuser Boston, Boston, MA, 2004, With appendices by Jean-Louis Colliot-Thélène and Nicholas M. Katz, pp. 175–184. MR 2029869