

KERODON REMIX PART II: COCARTESIAN FIBRATIONS, TRANSPORT, AND THE YONEDA EMBEDDING

CRIS NEGRON

ABSTRACT. These are notes on ∞ -categories which are (mostly) adapted from Lurie’s digital text Kerodon [4]. The main distinctions are the length of the document, the order of presentation, and selective omissions. We also deviate from [4] in that we focus on derived categories and dg categories as our primary examples of interest. A distinction from the related text [2] would be the complete avoidance of model structures, though this approach is already adopted in [4].

Following Part I, which presented the basic foundations for studies of ∞ -categories, we discuss herein cartesian and cocartesian fibrations, transport functors (i.e. Grothendieck straightening and unstraightening), and limits and colimits. Specific topics include Hom functions and Yoneda embedding, calculations of limits and colimits in categories of spaces and cochains, and stable ∞ -categories. While we follow the analytic, rather than synthetic, approach, we’ve attempted to communicate these topics in a manner which is consumable to the working mathematician.

CONTENTS

1. Introduction to cartesian and cocartesian fibrations	3
1.1. Definitions	3
1.2. Imaginings: Cartesian fibrations as lax moduli	4
1.3. Discussion: Classifying functors etc.	5
2. Cartesian and cocartesian fibrations	5
2.1. Cartesian morphisms via overcategories	5
2.2. Cartesian morphisms via mapping spaces	6
2.3. Proof of Proposition 2.3	7
2.4. Uniqueness for q -cocartesian lifts	7
2.5. Exponentiating cocartesian fibrations	8
3. Maps between fibrations	10
4. Directional fibrations and Kan complexes	10
4.1. Exponentials for directional fibrations	10
4.2. Directional fibrations and Kan complexes	11
5. Deviation into $(\infty, 2)$ -categories	11
5.1. $(\infty, 2)$ -categories	11
5.2. The pith of an $(\infty, 2)$ -category	12
5.3. $(\infty, 2)$ -category via simplicial categories	13
5.4. The $(\infty, 2)$ -category of ∞ -categories	13

Date: May 17, 2024.

The author was supported by NSF Grant No. DMS-2149817, NSF CAREER Grant No. DMS-2239698, and Simons Collaboration Grant No. 999367.

5.5. Interior fibrations	14
5.6. Undercategories and overcategories and pointed ∞ -categories	16
5.7. Mapping categories in the homotopy coherent nerve	17
6. Transport functors	19
6.1. Preliminary discussion	19
6.2. Categories with objects and the universal cocartesian fibration	19
6.3. A remark on notation	20
6.4. Fibers of the universal map $\mathcal{P}\mathcal{C}at_\infty \rightarrow \mathcal{C}at_\infty$	20
6.5. Covariant transport: classifying cocartesian fibrations	21
6.6. Transport for cartesian fibrations	23
6.7. Transport along edges	23
6.8. Classification of left and right fibrations	24
7. Homotopy transport representations	25
7.1. Homotopy transport representations	25
7.2. The $h\mathcal{K}an$ -enriched category of ∞ -categories	28
7.3. Enriched homotopy transport	28
7.4. Transport functors induce homotopy transport	30
7.5. Homotopy transport and enriched Hom functors	30
8. Naturality of transport: straightening and unstraightening	32
8.1. Marked simplicial sets	32
8.2. The ∞ -category of cocartesian fibrations over a base	34
8.3. Simplicial enrichment for simplicial functors	35
8.4. Simplicial functors as functor categories	36
8.5. Non-enriched Straightening and unstraightening	37
8.6. Enriched unstraightening	38
8.7. The straightening and unstraightening equivalences	38
8.8. A remark on uniqueness	39
8.9. Recovering transport via straightening	39
8.10. Straightening and unstraightening for left fibrations	40
8.11. Some naturality over the base	41
8.12. Straightening for cartesian fibrations	41
9. Initial and terminal objects	42
9.1. Initial and terminal basics	42
9.2. Aside: trivial fibrations via the fibers	43
9.3. Initial objects and undercategories	45
9.4. Initial and terminal objects in simplicial nerves	46
9.5. Zero objects in pointed spaces	47
9.6. Zero objects in derived categories	47
9.7. Initial objects and weak contractibility	48
9.8. Equivalences of fibrations via initial objects	50
10. Representable and corepresentable functors	51
10.1. Left fibrations with initial transport	52
10.2. Corepresentable functors for simplicial categories	53
10.3. Aside: Dold-Kan at the ∞ -level	54
10.4. Corepresentable functors for dg categories	56
10.5. Representable functors for simplicial and dg categories	57
11. Twisted arrows and bifunctorial Homs	58
11.1. The twisted arrows category	58

11.2. Fibers of the twisted arrows fibration	60
11.3. Recognition for Hom functors	62
12. Hom functors for dg categories	62
References	62

1. INTRODUCTION TO CARTESIAN AND COCARTESIAN FIBRATIONS

1.1. Definitions.

Definition 1.1. Consider a map of simplicial sets $q : X \rightarrow S$. We call a 1-simplex $\alpha : x \rightarrow y$ in X a q -cartesian morphism if any lifting problem

$$\begin{array}{ccc} \Lambda_n^n & \xrightarrow{\bar{\sigma}} & X \\ \downarrow & \nearrow & \downarrow q \\ \Delta^n & \longrightarrow & S \end{array} \quad (1) \quad \boxed{\text{eq:121}}$$

with $n \geq 2$ and $\bar{\sigma}|_{\Delta^{\{n-1,n\}}} = \alpha$ admits a solution. We say $\alpha : x \rightarrow y$ is q -cocartesian if any lifting problem

$$\begin{array}{ccc} \Lambda_0^n & \xrightarrow{\bar{\tau}} & X \\ \downarrow & \nearrow & \downarrow q \\ \Delta^n & \longrightarrow & S \end{array} \quad (2) \quad \boxed{\text{eq:128}}$$

with $n \geq 2$ and $\bar{\tau}|_{\Delta^{\{0,1\}}} = \alpha$ admits a solution.

Though at some specific moments we will consider a case where X and S are not ∞ -categories, we are primarily invested in the ∞ -categorical setting.

Definition 1.2. We call a map of ∞ -categories $q : \mathcal{C} \rightarrow \mathcal{D}$ a cartesian fibration if q is an inner fibration and, for any map $\bar{\alpha} : \bar{x} \rightarrow \bar{y}$ in \mathcal{D} and y in \mathcal{C} with $q(y) = \bar{y}$, there is a q -cartesian map $\alpha : x \rightarrow y$ in \mathcal{C} with $q(\alpha) = \bar{\alpha}$.

Similarly, we call q a cocartesian fibration if it is an inner fibration and, for any map $\bar{\beta} : \bar{x} \rightarrow \bar{y}$ in \mathcal{D} and x with $q(x) = \bar{x}$, there is a q -cocartesian map $\beta : x \rightarrow y$ with $q(\beta) = \bar{\beta}$.

The following is obvious. Recall our definitions of right and left fibrations from Definition I-4.23.

Proposition 1.3. *If $q : \mathcal{C} \rightarrow \mathcal{D}$ is a right fibration (resp. left fibration) then q is a cartesian (resp. cocartesian).*

Proof. In this case any lifting problem of the form (1), or (2) respectively, admits a solution simply by the definition. \square

Obviously when $q : \mathcal{C} \rightarrow \mathcal{D}$ is a Kan fibration it is both cartesian and cocartesian.


Example 1.4. Consider a diagram $p : K \rightarrow \mathcal{C}$, with K some simplicial set. Then we have the overcategory $\mathcal{C}_{/p}$ and the undercategory $\mathcal{C}_{p/}$. The two forgetful functors

$$\mathcal{C}_{/p} \rightarrow \mathcal{C} \quad \text{and} \quad \mathcal{C}_{p/} \rightarrow \mathcal{C}$$

are, respectively, a right and left fibration Proposition I-4.25. Hence these maps are respectively a cartesian and cocartesian fibration.

In the case where K is a point $x : * \rightarrow \mathcal{C}$ we recall that the fibers of the fibration $\mathcal{C}_{/x} \rightarrow \mathcal{C}$ and $\mathcal{C}_{x/} \rightarrow \mathcal{C}$ are the right and left pinched mapping spaces $\mathrm{Hom}_{\mathcal{C}}^R(w, x)$ and $\mathrm{Hom}_{\mathcal{C}}^L(x, y)$.

Example 1.5 ([4, 01T8]). Consider an ∞ -category $q : \mathcal{C} \rightarrow *$. A morphism $\alpha : x \rightarrow y$ is q -cartesian if and only if α is an isomorphism. To see this consider fillings of the horns



One similarly finds that a morphism is q -cocartesian if and only if it is an isomorphism.

Via the existence of identity morphisms the structure map $q : \mathcal{C} \rightarrow *$ is always a cartesian and cocartesian fibration. Note that this map is not a left or right fibration unless \mathcal{C} is a Kan fibration.

1.2. Imaginings: Cartesian fibrations as lax moduli. Give a cartesian fibration $q : \mathcal{C} \rightarrow \mathcal{D}$ one might think of \mathcal{C} as a lax moduli of “stuff” varying over the objects in \mathcal{D} . The cartesian lifts of morphisms in \mathcal{D} provide transition functions between these fibers, i.e. the stuff we are parametrizing, over \mathcal{D} . In the case of the cartesian fibration $\mathcal{C}_{/x} \rightarrow \mathcal{C}$ the category $\mathcal{C}_{/x}$ is, in an obvious sense, the “moduli of maps to x ”. Let us leave the latter point about lifting maps for now, and try to make some comment on the moduli point.

Let us just consider how one classically constructs a moduli space. Here we consider the base space $\mathcal{D} = \mathrm{Sch}_k$ of schemes over k , which we can endow with some Grothendieck topology if we like, though we don’t care at the moment. Then a pre-stack is a choice of a functor of plain categories $q : \mathbb{M} \rightarrow \mathrm{Sch}_k$ which makes \mathbb{M} into a category fibered in groupoids over Sch_k [5, 003S]. One simply compares definitions to see that

$$\left\{ \begin{array}{l} \mathbb{M} \text{ is fibered in} \\ \text{groupoids over } \mathrm{Sch}_k \end{array} \right\} \Leftrightarrow \left\{ \begin{array}{l} q \text{ is a cartesian fibration} \\ \text{in which all maps in } \mathbb{M} \\ \text{are } q\text{-cartesian} \end{array} \right\}.$$

In this familiar setting one can now “invert” this functor q to produce an associated 2-functor

$$q^\vee : (\mathrm{Sch}_k)^{\mathrm{op}} \rightarrow \mathrm{Groupoids} \subseteq \mathrm{Cat}, \quad Y \mapsto \mathbb{M}_Y.$$

One establishes this functor via an abuse of the axiom of choice.

To elaborate a bit more, for any map of schemes $\alpha : X \rightarrow Y$ we take a lift $\alpha^*y \rightarrow y$ in \mathbb{M} . This lift is unique up to unique isomorphism, and via unique filling defines a functor between the fibers

$$\alpha^* : \mathbb{M}_Y \rightarrow \mathbb{M}_X, \quad y \mapsto \alpha^*y.$$

On morphisms $\xi : y_1 \rightarrow y_2$ in the fiber \mathbb{M}_Y , we note that the cartesian property for maps in \mathbb{M} asserts the existence of a unique map $\alpha^*\xi : \alpha^*y_1 \rightarrow \alpha^*y_2$ completing

the diagram

$$\begin{array}{ccc} \alpha^* y_1 & \longrightarrow & y_1 \\ \exists! \downarrow & & \downarrow \xi \\ \alpha^* y_2 & \longrightarrow & y_2, \end{array}$$

where we note that uniqueness comes from filling the appropriate 3-simplex in \mathbb{M} . Hence α^* is well-defined on morphisms via the assignment $\xi \mapsto \alpha^* \xi$.

We note that this inversion of $q : \mathbb{M} \rightarrow \text{Sch}_k$ into a functor $q^\vee : (\text{Sch}_k)^{\text{op}} \rightarrow \text{Cat}$ does not require all maps in \mathbb{M} to be cartesian. This is simply a consequence of q being a cartesian fibration between plain categories.

In the general ∞ -context, we again have this inversion property for (co)cartesian fibrations. Here a cartesian fibration $q : \mathcal{C} \rightarrow \mathcal{D}$ will specify, and be specified by, a functor

$$q^\vee : \mathcal{D}^{\text{op}} \rightarrow \text{Cat}_\infty$$

whose values over objects y in \mathcal{D} are the fibers \mathcal{C}_y . The functors between fibers $\alpha^* : \mathcal{C}_y \rightarrow \mathcal{C}_x$ are what we've referred to as *transport* along α (following Kerodon [4]).

While such fibrations play an essentially non-existent role in plain category theory, from the perspective of the working mathematician, they play an extraordinarily important role in the development of ∞ -category theory. The main point is that they tame choices in the ∞ -categorical setting. While in the plain category setting we can simply make a choice, and if that choice is not unique we can simply say it's unique up to a unique isomorphism, and then if I make two of the same types of choices then any ambiguities will vanish due to sufficient uniqueness, etc. etc., such a *laissez faire* attitude will lead to immediate intractable problems in the ∞ -context. So one generally bundles all choices of a certain “type” into a cartesian or cocartesian fibrations, and manipulates these bundles in order to make global movements between choices of different types.

1.3. Discussion: Classifying functors etc.

2. CARTESIAN AND COCARTESIAN FIBRATIONS

2.1. Cartesian morphisms via overcategories.

prop:232

Proposition 2.1 ([4, 01TF]). *Let $q : \mathcal{C} \rightarrow \mathcal{D}$ be a map between ∞ -categories. A morphism $\alpha : x \rightarrow y$ in \mathcal{C} is q -cartesian if and only if the natural map*

$$\mathcal{C}_{/\alpha} \rightarrow \mathcal{C}_{/y} \times_{\mathcal{D}_{/q(y)}} \mathcal{D}_{/q(\alpha)}$$

is a trivial Kan fibration. Similarly, α is q -cocartesian if and only if the map

$$\mathcal{C}_{/\alpha} \rightarrow \mathcal{C}_{x/} \times_{\mathcal{D}_{q(x)/}} \mathcal{D}_{q(\alpha)/}$$

is a trivial Kan fibration.

For the proof we employ a specific deconstruction of the relevant horn inclusions.

lem:joyal3.3

Lemma 2.2 ([1, Lemma 3.3]). *For non-negative integers p and q , and $n = p+q+1$, the inclusions*

$$(\Lambda_0^p \star \Delta^q) \coprod_{\Lambda_q^p \star \partial \Delta^q} (\Delta^p \star \partial \Delta^q) \rightarrow \Delta^p \star \Delta^q \cong \Delta^n$$

and

$$(\partial\Delta^p \star \Delta^q) \coprod_{\partial\Delta^p \star \Lambda_q^q} (\partial\Delta^p \star \Lambda_q^q) \rightarrow \Delta^p \star \Delta^q \cong \Delta^n$$

are identified with the inclusions of the extremal horns $\Lambda_0^n \rightarrow \Delta^n$ and $\Lambda_n^n \rightarrow \Delta^n$ respectively.

One can see the text [1], or [4, 018N] for the details. We now proceed with the proof of Proposition 2.1.

Proof of Proposition 2.1. We address the cartesian situation, the cocartesian one being similar. Let $F : \mathcal{C}/\alpha \rightarrow \mathcal{C}/y \times_{\mathcal{D}/q(y)} \mathcal{D}/q(\alpha)$ denote the map under consideration. A solution to a lifting problems of the form

$$\begin{array}{ccc} \partial\Delta^m & \xrightarrow{\quad} & \mathcal{C}/\alpha \\ \downarrow & \nearrow & \downarrow F \\ \Delta^m & \xrightarrow{\quad} & \mathcal{C}/y \times_{\mathcal{D}/q(y)} \mathcal{D}/q(\alpha), \end{array} \quad (3) \quad \text{eq:262}$$

with $m \geq 0$, admit a solution if and only if the equivalent lifting problem

$$\begin{array}{ccc} (\partial\Delta^m \star \Delta^1) \cup (\Delta^m \star \Lambda_1^1) & \xrightarrow{\quad} & \mathcal{C} \\ \downarrow & \nearrow & \downarrow q \\ \Delta^m \star \Delta^1 & \xrightarrow{\quad} & \mathcal{D} \end{array}$$

obtained by way of adjunction Lemma I-4.22 admits a solution. Via direct inspection the final edge $\Delta^1 \cong \emptyset \star \Delta^1 \rightarrow \mathcal{C}$ in the latter diagram is α , so that this diagram is identified, via Lemma 2.2, with a diagram of the form

$$\begin{array}{ccc} \Lambda_n^n & \xrightarrow{\quad} & \mathcal{C} \\ \downarrow & \nearrow & \downarrow q \\ \Delta^n & \xrightarrow{\quad} & \mathcal{D} \end{array} \quad (4) \quad \text{eq:276}$$

in which $n \geq 2$ the edge $\Delta^{\{n-1, n\}} \rightarrow \mathcal{C}$ is α . It follows that all lifting problems of the form (3) admit a solution if and only if all lifting problems of the form (4) admit a solution, i.e. that the map F is a trivial Kan fibration if and only if the map α is q -cartesian. \square

2.2. Cartesian morphisms via mapping spaces.

prop:cocart_maps

Proposition 2.3 ([4, 01TL]). *Consider an inner fibration $q : \mathcal{C} \rightarrow \mathcal{D}$, and a morphism $\alpha : x_1 \rightarrow x_2$ in \mathcal{C} with image $\bar{\alpha} : \bar{x}_1 \rightarrow \bar{x}_2$ in \mathcal{D} . The morphism α is q -cartesian if and only if for each third object x_0 in \mathcal{C} , with corresponding triples $x : \{0, 1, 2\} \rightarrow \mathcal{C}$ and $\bar{x} : \{0, 1, 2\} \rightarrow \mathcal{D}$, the diagram*

$$\begin{array}{ccc} \text{Fun}(\Delta^2, \mathcal{C})_x \times_{\text{Hom}_{\mathcal{C}}(x_1, x_2)} \{\alpha\} & \longrightarrow & \text{Hom}_{\mathcal{C}}(x_1, x_2) \\ q \downarrow & & \downarrow q \\ \text{Fun}(\Delta^2, \mathcal{D})_{\bar{x}} \times_{\text{Hom}_{\mathcal{D}}(\bar{x}_1, \bar{x}_2)} \{\bar{\alpha}\} & \longrightarrow & \text{Hom}_{\mathcal{D}}(\bar{x}_1, \bar{x}_2) \end{array}$$

is a homotopy pullback diagram of Kan complexes.

We cover the proof of Proposition ?? in Section 2.3 below. Let us record now a number of examples.

sect:cocart_maps_proof

2.3. Proof of Proposition 2.3.

2.4. Uniqueness for q -cocartesian lifts.

prop:cocart_uniqueness

Proposition 2.4 ([4, 01VK]). *Let $q : X \rightarrow S$ be an inner fibration of simplicial sets, and let Y be the full simplicial subset in $\text{Fun}(\Delta^1, X)$ whose vertices are q -cocartesian edges in X . Let Z be the full simplicial set in $\text{Fun}(\{0\}, X) \times_{\text{Fun}(\{0\}, S)} \text{Fun}(\Delta^1, S)$ whose edges lie in the image of the composition*

$$Y \rightarrow \text{Fun}(\Delta^1, X) \rightarrow \text{Fun}(\{0\}, X) \times_{\text{Fun}(\{0\}, S)} \text{Fun}(\Delta^1, S).$$

Then the induced map $Y \rightarrow Z$ is a trivial Kan fibration. The analogous statement holds when we replace Y with the full simplicial subset of q -cartesian edges in $\text{Fun}(\Delta^1, X)$ as well.

Said informally, Proposition 2.4 tells us that, if a cocartesian solution to the diagram

$$\begin{array}{ccc} \{0\} & \longrightarrow & X \\ \downarrow & & \downarrow q \\ \Delta^1 & \xrightarrow{\alpha} & S \end{array}$$

exists, then that solution is unique. We note that in the case where q itself is cocartesian, the simplicial subset Z is all of $\text{Fun}(\{0\}, X) \times_{\text{Fun}(\{0\}, S)} \text{Fun}(\Delta^1, S)$.

The proof of Proposition 2.4 employs a decomposition of a certain inclusion of simplicial sets which we record here.

lem:simpl_328

Lemma 2.5 ([4, 00TH]). *At any positive integer n , the inclusion*

$$(\Delta^1 \times \partial\Delta^n) \coprod_{\{0\} \times \partial\Delta^n} (\{0\} \times \Delta^n) \rightarrow \Delta^1 \times \Delta^n$$

decomposes into a sequence of inclusions

$$(\Delta^1 \times \partial\Delta^n) \cup (\{0\} \times \Delta^n) = X(0) \rightarrow \cdots \rightarrow X(n) \rightarrow X(n+1) = \Delta^1 \times \Delta^n$$

in which each $X(i+1)$ fits into a pushout diagram

$$\begin{array}{ccc} \Lambda_{n-i}^{n+1} & \longrightarrow & X(i) \\ \downarrow & & \downarrow \\ \Delta^{n+1} & \longrightarrow & X(i+1) \end{array}$$

and the sequence

$$\Delta^{\{0,1\}} \rightarrow \Delta^{n+1} \rightarrow X(n+1) = \Delta^1 \times \Delta^{n+1}$$

is an isomorphism onto the edge $\Delta^1 \times \{0\}$ in $\Delta^1 \times \Delta^{n+1}$.

To be clear, our filtration is obtained by applying the opposite to the specific sequence from [4, 00TH].

Idea of proof. Consider the simplices $\sigma_i : \Delta^{n+1} \rightarrow \Delta^1 \times \Delta^n$ defined by taking $\sigma_i(j) = (0, j)$ if $j \leq n-i$ and $(1, j-1)$ if $j > n-i$. We define sequentially $X(i+1) = X(i) \cup \text{im}(\sigma_i)$. We refer the reader to [4] for the specific details. \square

We now can prove our uniqueness result for cocartesian lefts.

Proof of Proposition 2.4. We deal with the case of cocartesian situation, the cartesian case following by taking opposites. We consider a lifting problem

$$\begin{array}{ccc} \partial\Delta^n & \longrightarrow & Y \\ \downarrow & \nearrow & \downarrow \\ \Delta^n & \longrightarrow & Z. \end{array} \quad (5) \quad \boxed{\text{eq:361}}$$

In the case $n = 0$, this problem admits a solution by the definition of Z . In the case $n \geq 0$, solving this problem is equivalent to solving a lifting problem of the form

$$\begin{array}{ccc} (\Delta^1 \times \partial\Delta^n) \cup (\{0\} \times \delta^{q_n}) & \longrightarrow & X \\ \text{incl} \downarrow & \nearrow & \downarrow \\ \Delta^1 \times \Delta^n & \longrightarrow & S \end{array} \quad (6) \quad \boxed{\text{eq:368}}$$

in which all of the constituent maps $\Delta^1 \times \{i\} \rightarrow X$ are q -cocartesian. In particular, the map $\Delta^1 \times \{0\} \rightarrow X$ is q -cocartesian. We decompose the map incl into a sequence of inclusion $X(i) \rightarrow X(i+1)$ as in Lemma 2.5, and produce sequential solutions to the problems

$$\begin{array}{ccc} X(i) & \xrightarrow{\sigma_i} & X \\ \downarrow & \nearrow \sigma_{i+1} & \downarrow q \\ X(i+1) & \longrightarrow & S \end{array}$$

for each $i < n$ since the inclusion $X(i) \rightarrow X(i+1)$ is inner anodyne in this case. For the final inclusion at $i = n$, we have the extended diagram

$$\begin{array}{ccccc} \Lambda_0^{n+1} & \longrightarrow & X(n) & \xrightarrow{\sigma_n} & X \\ \downarrow & & \downarrow & \nearrow & \downarrow q \\ \Delta^{n+1} & \longrightarrow & X(n+1) = \Delta^1 \times \Delta^n & \longrightarrow & S \end{array}$$

and can solve the external problem since the initial edge $\Lambda_0^n \rightarrow X$ has q -cocartesian image in X , and can therefore solve the internal lifting problem since the left-most square is a pushout diagram. We therefore obtain a solution to our original problems (5) and (6). \square

2.5. Exponentiating cocartesian fibrations.

prop:397

Proposition 2.6 ([4, 01VG]). *If $q : X \rightarrow S$ is a cocartesian fibration, then for any simplicial set K the map $q_* : \text{Fun}(K, X) \rightarrow \text{Fun}(K, S)$ is a cocartesian fibration. An edge $\xi : \Delta^1 \rightarrow \text{Fun}(K, X)$ is q_* -cocartesian if and only if, at each v in K , the composite $v^*\xi : \Delta^1 \rightarrow X$ is q -cocartesian in X .*

The proof follows by a hands on analysis of certain lifting problems which we won't reproduce here. The reader can see the [4, 01VG & 01VM] for the details.

Let us recall that, for any inner fibration $q : X \rightarrow S$ and fixed map $\xi : A \rightarrow S$ the simplicial set $\text{Fun}_S(A, X)$ is obtained as the fiber

$$\begin{array}{ccc} \text{Fun}_S(A, X) & \longrightarrow & \text{Fun}(A, X) \\ \downarrow & & \downarrow q_* \\ * & \xrightarrow{\xi} & \text{Fun}(A, S) \end{array}$$

Since the map q_* is an inner fibration (Corollary I-4.8) we understand that $\text{Fun}_S(A, X)$ is an ∞ -category. Of course, in the more restrictive case in which q is a cocartesian fibration, we have just seen that q_* is furthermore cocartesian.

thm:simp_lift

Theorem 2.7. *Let K be any simplicial set and $q : X \rightarrow S$ be a cocartesian fibration. Any lifting problem*

$$\begin{array}{ccc} \{0\} \times K & \longrightarrow & X \\ \downarrow & \nearrow & \downarrow \\ \Delta^1 \times K & \longrightarrow & S \end{array}$$

admits a solution $\Delta^1 \times K \rightarrow X$ for which, at each vertex v in K , the composite

$$\Delta^1 \cong \Delta^1 \times \{v\} \rightarrow \Delta^1 \times K \rightarrow X$$

is a q -cocartesian edge in X . Furthermore, the full ∞ -subcategory in $\text{Fun}_S(\Delta^1 \times K, X)$ spanned by such solutions is a contractible Kan complex.

Proof. By Proposition 2.6, the map $q_* : \text{Fun}(K, X) \rightarrow \text{Fun}(K, S)$ is a cocartesian fibration and solutions to the above lifting problem are identified with q_* -cocartesian solutions $\tilde{\xi} : \Delta^1 \rightarrow \text{Fun}(K, X)$ to the associated lifting problem

$$\begin{array}{ccc} \{0\} & \longrightarrow & \text{Fun}(K, X) \\ \downarrow & \nearrow & \downarrow q_* \\ \Delta^1 & \longrightarrow & \text{Fun}(K, S). \end{array}$$

Existence and uniqueness of such solutions now follow by Proposition 2.4. □

In the event that $q : X \rightarrow S$ is a left fibration, all morphisms in X are cocartesian. So we see that there is a unique solution to the above lifting problem.

cor:left_lift

Corollary 2.8. *Let K be any simplicial set and $q : X \rightarrow S$ be a left fibration. Any lifting problem*

$$\begin{array}{ccc} \{0\} \times K & \longrightarrow & X \\ \downarrow & \nearrow & \downarrow \\ \Delta^1 \times K & \longrightarrow & S \end{array}$$

admits a solution $\Delta^1 \times K \rightarrow X$, and the collection of all such solutions $\text{Fun}_S(\Delta^1 \times K, X)$ is a contractible Kan complex.

3. MAPS BETWEEN FIBRATIONS

prop:318

Proposition 3.1 ([4, 023R]). *Suppose that*

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{F} & \mathcal{D} \\ & \searrow q & \swarrow p \\ & S & \end{array}$$

(7)

eq:320

is a map of inner fibrations, and that F is an equivalence. Then a morphism α is \mathcal{C} is q -cocartesian if and only if $F\alpha$ is p -cocartesian.

Corollary 3.2. *Suppose we have a diagram (7) in which p and q are cocartesian fibrations and F is an equivalence of inner fibrations. Then F is an equivalence of cocartesian fibrations.*

4. DIRECTIONAL FIBRATIONS AND KAN COMPLEXES

4.1. Exponentials for directional fibrations.

Definition 4.1. A map of simplicial sets $A \rightarrow B$ is called left anodyne (resp. right anodyne) if any lifting problem

$$\begin{array}{ccc} A & \longrightarrow & X \\ \downarrow & \nearrow \text{dotted} & \downarrow f \\ B & \longrightarrow & S \end{array}$$

in which f is a left (resp. right) fibration admits a solution.

One can show that the class of left anodyne maps is the saturated class generated by the horn inclusions $\Lambda_i^n \rightarrow \Delta^n$, where $0 \leq i < n$ [4, 0151]. One similarly characterizes right anodyne maps.

lem:328

Lemma 4.2 ([4, kerodon]). *Let $i : A \rightarrow B$ and $j : K \rightarrow L$ be monomorphisms of simplicial sets. If one of i or j is left (resp. right) anodyne, then the induced map*

$$(B \times K) \coprod_{A \times K} (A \times L) \rightarrow B \times L$$

is left (resp. right) anodyne.

We refer the reader to Kerodon [4] for the proof.

prop:direct_tech

Proposition 4.3. *Let $f : X \rightarrow S$ be a map of simplicial sets, and $j : K \rightarrow L$ be a monomorphism of simplicial sets. Consider the induced map on the functor complexes*

$$\rho : \text{Fun}(L, X) \rightarrow \text{Fun}(K, X) \times_{\text{Fun}(K, S)} \text{Fun}(L, S).$$

- (1) *If f is a left fibration, then ρ is a left fibration.*
- (2) *If f is a right fibration, then ρ is a right fibration.*
- (3) *If f is a left fibration and j is left anodyne, then ρ is a trivial Kan fibration.*
- (4) *If f is a right fibration and j is right anodyne, then ρ is a trivial Kan fibration.*

Proof. Solving a lifting problem of the form

$$\begin{array}{ccc} A & \xrightarrow{\quad} & \text{Fun}(L, X) \\ \downarrow & \nearrow \text{dotted} & \downarrow f \\ B & \xrightarrow{\quad} & \text{Fun}(K, X) \times_{\text{Fun}(K, S)} \text{Fun}(L, S) \end{array}$$

is equivalent to solving the corresponding lifting problem

$$\begin{array}{ccc} (B \times K) \amalg_{(A \times K)} (A \times L) & \xrightarrow{\quad} & X \\ \downarrow & \nearrow \text{dotted} & \downarrow \\ B \times L & \xrightarrow{\quad} & S. \end{array}$$

So all of the claims follow from a consideration of Lemma 4.2. \square

4.2. Directional fibrations and Kan complexes.

Proposition 4.4. *A cocartesian (resp. cartesian) fibration $f : X \rightarrow S$ is a left (resp. right) fibration if and only if all of the fibers X_s , at arbitrary $s : * \rightarrow S$, are Kan complexes.*

5. DEVIATION INTO $(\infty, 2)$ -CATEGORIES

5.1. $(\infty, 2)$ -categories.

Definition 5.1. Let X be a simplicial set. A 2-simplex $\tau : \Delta^2 \rightarrow X$ is called thin if any horn for any $n > 2$, index $0 < i < n$, and inner horn

$$\bar{\sigma} : \Lambda_i^n \rightarrow X \quad \text{with} \quad \bar{\sigma}|_{\Delta^{\{i-1, i, i+1\}}} = \tau,$$

the lifting problem

$$\begin{array}{ccc} \Lambda_i^n & \xrightarrow{\bar{\sigma}} & X \\ \downarrow & \nearrow \text{dotted} & \downarrow \\ \Delta^n & \xrightarrow{\quad} & * \end{array}$$

admits a solution.

One sees immediately that every 2-simplex in an ∞ -category is thin, for example.

Recall our notation $s_i : [n] \rightarrow [n-1]$ for the weakly increasing surjection with $s_i(i) = s_i(i+1) = i$, for $0 \leq i \leq n-1$, and the corresponding degeneracies $s_i^* : \Delta^n \rightarrow \Delta^{n-1}$. We call an n -simplex $\sigma : \Delta^n \rightarrow X$ in a simplicial set *left degenerate* if σ factors through the degeneracy $s_0^* : \Delta^n \rightarrow \Delta^{n-1}$, and *right degenerate* if σ factors through the degeneracy $s_{n-1}^* : \Delta^n \rightarrow \Delta^{n-1}$.

def:infy2

Definition 5.2 ([4, 01W9, 01Y5]). A simplicial set X is called an $(\infty, 2)$ -category if the following hold:

- (a) Any horn $\Lambda_1^2 \rightarrow X$ admits an extension to a thin 2-simplex.
- (b) Every degenerate 2-simplex in X is thin.
- (c.l) For $n > 2$, any horn $\bar{\sigma} : \Lambda_0^n \rightarrow X$ in which the 2-simplex $\bar{\sigma}|_{\Delta^{\{0,1,n\}}}$ is left degenerate admits an extension to an n -simplex in X .
- (c.r) For $n > 2$, any horn $\bar{\sigma}' : \Lambda_n^n \rightarrow X$ in which the 2-simplex $\bar{\sigma}'|_{\Delta^{\{0,n-1,n\}}}$ is right degenerate admits an extension to an n -simplex in X .

A functor, or map, between $(\infty, 2)$ -categories is a map of simplicial sets $F : X \rightarrow Y$ which preserves thin 2-simplices.

Remark 5.3. Having introduced this notion, let us recall that the term ∞ -category is used interchangeably with the term $(\infty, 1)$ -category.

If we consider an ∞ -category \mathcal{C} , then in any horn $\Lambda_0^n \rightarrow \mathcal{C}$ as in (c.l) the initial edge $\Delta^{\{0,1\}} \rightarrow \mathcal{C}$ is degenerate, and hence an isomorphism in \mathcal{C} . Hence we have the proposed completion to an n -simplex $\Delta^n \rightarrow \mathcal{C}$, by Proposition I-4.33. Similarly any horn $\Lambda_n^n \rightarrow \mathcal{C}$ as in (c.r) completes to an n -simplex as well. So we observe the following.

Lemma 5.4. *Any ∞ -category is an $(\infty, 2)$ -category. Furthermore, an $(\infty, 2)$ -category X is an ∞ -category if and only if every 2-simplex in X is thin.*

Recall that each simplex Δ^n is an ∞ -category, and hence an $(\infty, 2)$ -category.

Example 5.5. Since any degenerate 2-simplex in an $(\infty, 2)$ -category is thin, any map of simplicial sets $* = \Delta^0 \rightarrow X$ is a map of $(\infty, 2)$ -categories. Similarly, any map of simplicial sets $\Delta^1 \rightarrow X$ is a map of $(\infty, 2)$ -categories.

One has the following practical check for maps between $(\infty, 2)$ -categories.

prop:infy2_check

Proposition 5.6 ([4, 01YC]). *Let X and Y be $(\infty, 2)$ -categories, and $F : X \rightarrow Y$ be a map of simplicial sets. Then F is a functor, i.e. preserves thin 2-simplices, if and only if any horn $\Lambda_1^2 \rightarrow X$ can be completed to a thin simplex with thin image in Y .*

Idea of proof. The result is a consequence of stability of thin simplices under various conditions. Namely one establishes an inner-exchange property for thin simplices, which we recall below, and a 4-of-5 property which one can find at [4, 01XX]. \square

5.2. The pith of an $(\infty, 2)$ -category.

Definition 5.7. Given an $(\infty, 2)$ -category X , the pith in X is the simplicial subset $X^{\text{Pith}} \subseteq X$ whose simplices $\Delta^n \rightarrow X^{\text{Pith}}$ consist of all simplices $\sigma : \Delta^n \rightarrow X$ in which each restriction along a 2-simplex

$$\Delta^2 \rightarrow \Delta^n \xrightarrow{\sigma} X$$

is thin.

Since functors between $(\infty, 2)$ -categories preserve thin simplices, by definition, we see that any map $F : \mathcal{C} \rightarrow X$ from an ∞ -category to an $(\infty, 2)$ -category factors through the pith.

lem:360

Lemma 5.8 ([4, 01XL], Inner-exchange property). *Consider a 3-simplex $\sigma : \Delta^3 \rightarrow X$ in an $(\infty, 2)$ -category, and suppose that the associated 2-simplices $\sigma|_{\Delta^{\{1,2,3\}}}$ and $\sigma|_{\Delta^{\{0,1,2\}}}$ are thin. Then the 2-simplex $\sigma|_{\Delta^{\{0,2,3\}}}$ is thin if and only if the 2-simplex $\sigma|_{\Delta^{\{0,1,3\}}}$ is thin.*

The proof employs certain facts about interior fibrations (see below), and is omitted. From Lemma 5.8 the proof of the following is immediate.

Proposition 5.9. *For any $(\infty, 2)$ -category X , the subcomplex X^{Pith} is an ∞ -category.*

Proof. For any completion $\Delta^3 \rightarrow X$ of an inner horn $\Lambda_i^3 \rightarrow X$ in which all of the associated face $\Delta^2 \rightarrow \Lambda_i^3 \rightarrow X$ are thin, the final face $\Delta^{[3] \setminus \{i\}} \rightarrow X$ is also thin, by Lemma 5.8. This shows that the pith is stable under the completion of inner horns $\Lambda_i^3 \rightarrow X^{\text{Pith}}$. Stability under completion of all inner horns $\Lambda_i^n \rightarrow X^{\text{Pith}}$ with $n > 3$ is immediate, since the horn Λ_i^n already contains all 2-faces in Δ^n in this case. Taken together with condition (a) of Definition 5.2, we see that any lifting problem

$$\begin{array}{ccc} \Lambda_i^n & \longrightarrow & X^{\text{Pith}} \\ \downarrow & & \downarrow \\ \Delta^n & \longrightarrow & * \end{array}$$

with $0 < i < n$ admits a solution, as required. \square

5.3. $(\infty, 2)$ -category via simplicial categories. Recall that one can associate to any simplicial category \underline{S} its associated homotopy coherent nerve $N^{\text{hc}}(\underline{S})$ Section I-2.7. The n -simplices in $N^{\text{hc}}(\underline{S})$ are simplicial functors from the path category $\text{Path}[n]$. For $S = N^{\text{hc}}(\underline{S})$ we have in low dimension

$$S[0] = \{ \text{objects in } \underline{S} \}$$

$$S[1] = \{ \text{pairs of object } (x_0, x_1) \text{ along with a map } f \in \underline{\text{Hom}}_{\underline{S}}(x_0, x_1)[0] \}$$

$$S[2] = \left\{ \begin{array}{l} \text{triples of objects } (x_0, x_1, x_2), \text{ maps } f_{ij} : x_i \rightarrow x_j \text{ for each } i < j, \text{ and} \\ \text{a 1-simplex } h : \Delta^1 \rightarrow \underline{\text{Hom}}_{\underline{S}}(x_0, x_2) \text{ with } h|_0 = f_{02}, h|_1 = f_{12}f_{01} \end{array} \right\}$$

Lemma I-2.16. We take the following theorem for granted.

thm:hc_infty2

Theorem 5.10 ([4, 01YM]). *Let \underline{S} be a simplicial category in which, at each pair of objects x and y in \underline{S} , the mapping complex $\underline{\text{Hom}}_{\underline{S}}(x, y)$ is an ∞ -category. Then the homotopy coherent nerve $N^{\text{hc}}(\underline{S})$ is an $(\infty, 2)$ -category.*

What we are most interested in here is the $(\infty, 2)$ -category of ∞ -categories. Recall that for any ∞ -categories \mathcal{C} and \mathcal{D} the simplicial set of functors $\text{Fun}(\mathcal{C}, \mathcal{D})$, whose simplicial are as expected

$$\text{Fun}(\mathcal{C}, \mathcal{D})[n] = \text{Hom}_{\text{sSet}}(\Delta^n \times \mathcal{C}, \mathcal{D}),$$

form another ∞ -category Corollar I-4.8. With these morphisms we obtain the simplicial category $\underline{\text{Cat}}_{\infty}$ of ∞ -categories and their functor categories. We note that $\underline{\text{Cat}}_{\infty}$ is a full simplicial subcategory in the ambient category sSet of simplicial sets.

5.4. The $(\infty, 2)$ -category of ∞ -categories.

Theorem 5.11. *The homotopy coherent nerve*

$$\text{Cat}_{\infty} := N^{\text{hc}}(\underline{\text{Cat}}_{\infty})$$

is an $(\infty, 2)$ -category.

According to the above analysis the 0-simplices in Cat_{∞} are ∞ -categories, the 1-simplices are functors between ∞ -categories, and 2-simplices are triples of functors

and a natural transformation

$$\begin{array}{ccc}
 & \mathcal{C}_2 & \\
 F_{12} \nearrow & & \searrow F_{01} \\
 \mathcal{C}_1 & & \mathcal{C}_0 \\
 & \xrightarrow{\zeta} & \\
 & \mathcal{C}_2 & \\
 & \uparrow F_{02} & \\
 & \mathcal{C}_0 &
 \end{array}$$

Definition 5.12. The $(\infty, 2)$ -category Cat_∞ is called the $(\infty, 2)$ -category of ∞ -categories.

Remark 5.13. The $(\infty, 2)$ -category Cat_∞ is in our universe of “large” sets, which is strictly larger than our universe of “normal sized” set in which all other ∞ -categories are assumed to live.

We recall our ∞ -category \mathcal{Cat}_∞ of ∞ -categories, which we obtain by restricting the morphisms $\text{Fun}(\mathcal{C}, \mathcal{D})$ to the associated Kan complex $\text{Fun}(\mathcal{C}, \mathcal{D})^{\text{Kan}}$ then applying the simplicial nerve. The inclusions of ∞ -categories

$$\text{Fun}(\mathcal{C}, \mathcal{D})^{\text{Kan}} \rightarrow \text{Fun}(\mathcal{C}, \mathcal{D})$$

imply an inclusion of $(\infty, 2)$ -categories $\mathcal{Cat}_\infty \rightarrow \text{Cat}_\infty$, and hence an inclusion into the pith

$$\mathcal{Cat}_\infty \rightarrow (\text{Cat}_\infty)^{\text{Pith}}. \quad (8) \quad \boxed{\text{eq:444}}$$

By a general result one finds that this inclusion is an equality.

Proposition 5.14 ([4, 01YT]). *The inclusion (8) is an equality, $\mathcal{Cat}_\infty = (\text{Cat}_\infty)^{\text{Pith}}$.*

5.5. Interior fibrations.

def:interior

Definition 5.15. A map of simplicial sets $q : X \rightarrow S$ is called an interior fibration if the following hold:

- (a) At each 0-simplex x in X , the identity $\text{id}_x : x \rightarrow x$ is both q -cartesian and cocartesian.
- (b) For any lifting problem

$$\begin{array}{ccc}
 \Lambda_i^n & \longrightarrow & X \\
 \downarrow & \nearrow & \downarrow \\
 \Delta^n & \xrightarrow{\sigma} & S
 \end{array} \quad (9) \quad \boxed{\text{eq:400}}$$

in which $0 < i < n$ and $\sigma|_{\Delta^{\{i-1, i, i\}}}$ is thin in S , (9) admits a solution.

It is clear that if $f : S' \rightarrow S$ is a map of simplicial sets which preserves thin 2-simplices, and the diagram

$$\begin{array}{ccc}
 X' & \longrightarrow & X \\
 q' \downarrow & & \downarrow q \\
 S' & \xrightarrow{f} & S
 \end{array}$$

is a pullback diagram of simplicial sets in which q is an interior fibration, then the map $q' : X' \rightarrow S'$ is an interior fibration as well.

One also observes the following.

lem:495

Lemma 5.16. *If S is an $(\infty, 2)$ -category, and $q : X \rightarrow S$ is an interior fibration, then X is also an $(\infty, 2)$ -category and q is a functor between $(\infty, 2)$ -categories.*

Proof. One sees via the lifting property for q that any 2-simplex $\Delta^2 \rightarrow X$ which has thin image in S is thin in X . From this we see that any horn $\Lambda_1^2 \rightarrow X$ can be completed to a thin 2-simplex in X . One obtains this thin completion by lifting a thin completion $\Delta^2 \rightarrow S$. We are left to prove that any appropriate degenerate horn $\Lambda_0^n \rightarrow X$ or $\Lambda_n^n \rightarrow X$, at $n > 2$, completes to an n -simplex. However this follows from the fact that identity maps in X are both q -cartesian and cocartesian, and the fact that the corresponding horns in S admit completions. We now see that X is an $(\infty, 2)$ -category. One sees that q is a functor, i.e. preserves thin 2-simplices, by applying Proposition 5.6. \square

As we see in the above proof, given an interior fibration $q : X \rightarrow S$ over an $(\infty, 2)$ -category, one can detect thin simplices in X by considering their images in S along q .

lem:505

Lemma 5.17. *If $q : X \rightarrow S$ is an interior fibration then a 2-simplex in X is thin if and only if its image in S is thin.*

cor:interior_pullback

Corollary 5.18. *Consider a pullback diagram*

$$\begin{array}{ccc} Z & \xrightarrow{p_2} & X \\ p_1 \downarrow & & \downarrow q \\ Y & \xrightarrow{F} & S \end{array}$$

in which q is an interior fibration and F is a map between $(\infty, 2)$ -categories. Then Z is an $(\infty, 2)$ -category, p_1 is an interior fibration, and p_2 is a map of $(\infty, 2)$ -categories.

Proof. The fact that Z is an $(\infty, 2)$ -category and p_1 is a map of $(\infty, 2)$ -categories follows by Lemma 5.16. As for p_1 , we consider a thin 2-simplex $\Delta^2 \rightarrow Z$, and note that its image in Y is thin. Hence its image in S is thin, and so its image in X is thin by Lemma 5.17. It follows that p_2 is a map of $(\infty, 2)$ -categories, by definition. \square

We are especially interested in the fiberings of interior fibrations over ∞ -categories.

lem:502

Lemma 5.19. *Let \mathcal{C} be an ∞ -category. A map of simplicial sets $q : X \rightarrow \mathcal{C}$ is an interior fibration if and only if it is an inner fibration.*

Proof. If q is an interior fibration then it is an inner fibration since all 2-simplices in \mathcal{C} are thin. Conversely, if q is an inner fibration then X is an ∞ -category and q is therefore an inner fibration between ∞ -categories. Condition (a) of Definition 5.15 now follows from the fact that the identity in an ∞ -category is an isomorphism, and an application of Proposition I-4.33. \square

One combines Lemma 5.19 with the above discussion of fiber products to obtain the following corollary.

Corollary 5.20. *Consider an interior fibration $q : X \rightarrow S$ over an $(\infty, 2)$ -category S .*

- (a) *For any ∞ -category \mathcal{C} , and any functor of $(\infty, 2)$ -categories $\mathcal{C} \rightarrow S$, the fiber product $X \times_S \mathcal{C}$ is an ∞ -category. Furthermore, the projection $X \times_S \mathcal{C} \rightarrow \mathcal{C}$ is an inner fibration.*

(b) At each point $s : * \rightarrow S$ the fiber X_s is an ∞ -category.

cor:interior_pith

Corollary 5.21. *If $q : X \rightarrow S$ is an interior fibration over an $(\infty, 2)$ -category S then the diagram*

$$\begin{array}{ccc} X^{\text{Pith}} & \longrightarrow & X \\ \downarrow & & \downarrow \\ S^{\text{Pith}} & \longrightarrow & S \end{array}$$

is a pullback diagram, and the map $X^{\text{Pith}} \rightarrow S^{\text{Pith}}$ is an interior fibration.

Proof. In this case the pullback $X \times_S S^{\text{Pith}}$ is an ∞ -category and the projection to X is a map of $(\infty, 2)$ -categories. So the identification

$$X^{\text{Pith}} = X \times_S S^{\text{Pith}}$$

follows via an application of the universal property for the pullback and the universal property for the pith. \square

5.6. Undercategories and overcategories and pointed ∞ -categories. In the $(\infty, 2)$ -setting we can define overcategories and undercategories exactly as in the ∞ -setting. Namely, for a map of simplicial sets $p : K \rightarrow X$ the overcategory $X_{p/}$ is the simplicial set with n -simplices provided by the join

$$X_{p/}[n] := \text{Hom}_{\text{sSet}}(K \star \Delta^n, X)_p,$$

and similarly for the undercategory

$$X_{/p}[n] := \text{Hom}_{\text{sSet}}(\Delta^n \star K, X)_p$$

Section I-4.6. In the case in which X is an ∞ -category, we saw that the forgetful functors

$$\begin{array}{ccc} X_{p/} & & X_{/p} \\ & \searrow & \swarrow \\ & X & \end{array}$$

obtained by restricting along the inclusions $\Delta^n \rightarrow \Delta^n \star K$ and $\Delta^n \rightarrow K \star \Delta^n$ are directional fibrations, and in particular isofibrations. We have a similar result in the 2-categorical context.

prop:504

Proposition 5.22 ([4, 01WU]). *Let X be an $(\infty, 2)$ -category and $p : K \rightarrow X$ be a map of simplicial sets. The forgetful maps*

$$X_{p/} \rightarrow X \quad \text{and} \quad X_{/p} \rightarrow X$$

are both interior fibrations.

At this point we'll begin to leave many of the details unaccounted for. In particular, we direct the reader to the original text [4] for the details on Proposition 5.22. In any case, we record some corollaries.

Corollary 5.23. *For an $(\infty, 2)$ -category X and a diagram $p : K \rightarrow X$, the simplicial sets $X_{p/}$ and $X_{/p}$ are $(\infty, 2)$ -category and the forgetful maps are both functors between $(\infty, 2)$ -categories.*

cor:601

Corollary 5.24. *Let $p : K \rightarrow X$ be a map from a simplicial set into an $(\infty, 2)$ -category. At any point $x : * \rightarrow X$ the fibers $(X_{p/})_x$ and $(X_{/p})_x$ are both ∞ -categories.*

We apply this corollary in the case where the diagram p is a point $x : * \rightarrow X$ to obtain mapping categories for any $(\infty, 2)$ -category X .

Definition 5.25. For any $(\infty, 2)$ -category X , and objects $x, y : * \rightarrow X$, the left pinched mapping ∞ -category is the fiber

$$\mathrm{Hom}_X^L(x, y) := (X_{x/}) \times_X \{y\}.$$

Similarly, the right pinched mapping ∞ -category is the fiber

$$\mathrm{Hom}_X^R(x, y) = \{x\} \times_X (X_{/y}).$$

As with any interior fibration, we can restrict the forgetful functor to the piths to obtain inner fibrations of ∞ -categories

$$(X_{p/})^{\mathrm{Pith}} \rightarrow X^{\mathrm{Pith}} \quad \text{and} \quad (X_{/p})^{\mathrm{Pith}} \rightarrow X^{\mathrm{Pith}}.$$

In this particular instance one can observe a stronger characterization of these functors.

`prop:over_cartesian`

Proposition 5.26 ([4, 01YE]). For X and $p : K \rightarrow X$ as above, the restrictions of the forgetful functors

$$(X_{p/})^{\mathrm{Pith}} \rightarrow X^{\mathrm{Pith}} \quad \text{and} \quad (X_{/p})^{\mathrm{Pith}} \rightarrow X^{\mathrm{Pith}}.$$

are, respectively, a cocartesian fibration and a cartesian fibration.

One might view this result in analogy with the ∞ -setting, where the forgetful functors were observed to be right and left fibrations Corollary I-4.27.

5.7. Mapping categories in the homotopy coherent nerve. Let \underline{S} be a simplicial category whose morphism complexes are weak Kan complexes, and let S be the homotopy coherent nerve, $S = \mathrm{N}^{\mathrm{hc}}(\underline{S})$. We recall that S is an $(\infty, 2)$ -category in this case. By an abuse of notation take

$$\underline{\mathrm{Hom}}_S(x, y) = \underline{\mathrm{Hom}}_{\underline{S}}(x, y)$$

for any given pair of objects in S . We construct a map of simplicial sets

$$\theta : \underline{\mathrm{Hom}}_S(x, y) \rightarrow \mathrm{Hom}_S^L(x, y)$$

[4, 01LD] which is subsequently found to be an equivalence of ∞ -categories.

To begin, for any simplicial set K we consider the simplicial category $E(K)$ with objects x_- and x_+ and morphisms

$$\mathrm{Hom}_{E(K)}(x_-, x_-) = \mathrm{Hom}_{E(K)}(x_+, x_+) = * \quad \text{and} \quad \mathrm{Hom}_{E(K)}(x_-, x_+) = K.$$

We consider the $(n+1)$ -simplex $\{-1\} \star \Delta^n \cong \Delta^{n+1}$ and the simplicial path category $\mathrm{Path}(\{-1\} \star \Delta^n)$ whose morphisms are given by the nerves

$$\mathrm{Hom}_{\mathrm{Path}(\{-1\} \star \Delta^n)}(l, m) = \mathrm{N}(\mathrm{Subsets}_{l, m}^{\mathrm{op}})$$

where $\mathrm{Subsets}_{l, m}$ is the partially ordered set of subsets $S \subseteq [n]$ with $\min S = l$ and $\max S = m$, ordered by inclusion.

At each integer n we have a simplicial functor

$$\theta_n^* : \mathrm{Path}(\{-1\} \star \Delta^n) \rightarrow E\Delta^n$$

which is define on objects by taking $\theta_n^*(-1) = x_-$, and $\theta_n^*(i) = x_+$ for all $i \geq 0$, and defined on morphisms by the simplicial map

$$\theta_n^* : \mathrm{Hom}_{\mathrm{Path}(\{-1\} \star \Delta^n)}(l, m) = \mathrm{N}(\mathrm{Subsets}_{l, m}^{\mathrm{op}}) \rightarrow \mathrm{Hom}_E(x_-, x_+) = \Delta^n = \mathrm{N}([n])$$

associated to the functor $\text{Subsets}_{l,m}^{\text{op}} \rightarrow [n]$ which sends each subset $S = \{l < s_1 < \dots < s_r < m\}$ to s_1 and each inclusion $S' \supseteq S$ to the inequality $s'_1 \leq s_1$.

For objects x and y in S , n -simplices in $\underline{\text{Hom}}_S(x, y)$ are identified with simplicial functors $\text{Fun}_{\text{sCat}}(E\Delta^n, \underline{S})$ in the fiber over (x, y) in $\text{Fun}(E\emptyset, \underline{S})$. Each such functor now defines an $(n+1)$ -simplex in S via a consideration of the identification

$$S[n+1] = \text{Fun}_{\text{sCat}}(\text{Path}(\{-1\} \star \Delta^n), \underline{S})$$

and restricting along θ_n^* . One sees, by the definition of θ_n^* that this associated $(n+1)$ -simplex has initial vertex x and all other vertices y , and restricts trivially to $\Delta^n \subseteq \{-1\} \star \Delta^n$. So we obtain a map of sets

$$\begin{aligned} \theta_n : \underline{\text{Hom}}_S(x, y)[n] &\rightarrow (S_{x/}) \times_S \{y\} = \text{Hom}_S^{\text{L}}(x, y)[n], \\ (f : E\Delta^n \rightarrow \underline{S}) &\mapsto (f\theta_n^* : \text{Path}\{-1\} \star \Delta^n \rightarrow \underline{S}). \end{aligned}$$

One observes directly that any increasing function $t : [n] \rightarrow [n']$ produces a commutative diagram

$$\begin{array}{ccc} \text{Path}(\{-1\} \star \Delta^n) & \xrightarrow{\theta_n} & E\Delta^n \\ t_* \downarrow & & \downarrow t_* \\ \text{Path}(\{-1\} \star \Delta^{n'}) & \xrightarrow{\theta_{n'}} & E\Delta^{n'}, \end{array}$$

from which we see that the θ_n assemble into a map of simplicial sets, or a map of ∞ -categories,

$$\theta : \underline{\text{Hom}}_S(x, y) \rightarrow \text{Hom}_S^{\text{L}}(x, y).$$

thm:pinched_simplicial

Theorem 5.27 ([4, 01LG]). *Let \underline{S} be a simplicial category whose morphism complexes are ∞ -categories. Take $S = \text{N}^{\text{hc}}(\underline{S})$. For any objects $x, y : * \rightarrow S$ there is a natural equivalence of ∞ -categories*

$$\theta : \underline{\text{Hom}}_S(x, y) \rightarrow \text{Hom}_S^{\text{L}}(x, y).$$

We do not cover the details, and refer the reader to the text [4].

cor:simplicial_pullback

Corollary 5.28. *Take \underline{S} and S as above. For any pair of points $x, y : * \rightarrow S$ there is a categorical pullback diagram*

$$\begin{array}{ccc} \underline{\text{Hom}}_S(x, y) & \xrightarrow{\theta} & (S_{x/})^{\text{Pith}} \\ \downarrow & & \downarrow \\ * & \xrightarrow{y} & S^{\text{Pith}}. \end{array} \tag{10} \quad \text{eq:704}$$

Proof. By Corollary 5.18 and Proposition 5.22 the projection map $\text{Hom}_S^{\text{L}}(x, y) \rightarrow S_{x/}$ has image in the $\text{Pith}(S_{x/})^{\text{Pith}}$. Applying this fact in conjunction with Corollary 5.21, we observe a pullback diagram of ∞ -categories

$$\begin{array}{ccc} \text{Hom}_S^{\text{L}}(x, y) & \xrightarrow{\theta} & (S_{x/})^{\text{Pith}} \\ \downarrow & & \downarrow \\ * & \xrightarrow{y} & S^{\text{Pith}}. \end{array}$$

in which the right-hand map is an inner fibration. This diagram is additionally a categorical pullback square by Corollary I-5.22 and Proposition 5.26. Since $\theta :$

$\underline{\mathrm{Hom}}_S(x, y) \rightarrow \mathrm{Hom}_S^L(x, y)$ is an equivalence of ∞ -categories it follows that the corresponding diagram (10) is a categorical pullback square as well (see Proposition I-5.23). \square

sect:transport

6. TRANSPORT FUNCTORS

6.1. Preliminary discussion. In analogy with the plain categorical setting, we claim that cocartesian fibrations $q : \mathcal{E} \rightarrow \mathcal{C}$ over a given ∞ -category are “the same thing” as functors into the ∞ -category of ∞ -categories $F : \mathcal{C} \rightarrow \mathcal{Cat}_\infty$. In our imaginations, the functor F should evaluate as the fibers $F(x) \cong \mathcal{E}_x$ and the image of a given map $\alpha : x \rightarrow y$ should be some kind of pushforward functor $\alpha_* : \mathcal{E}_x \rightarrow \mathcal{E}_y$ which “moves along” cartesian lifts $\tilde{\alpha} : \tilde{x} \rightarrow \tilde{y}$, so that $\alpha_*(\tilde{x}) \cong \tilde{y}$.

Of course one can not simply construct the desired functor $F : \mathcal{C} \rightarrow \mathcal{Cat}_\infty$ by hand. We (or rather, Lurie) instead proceed(s) by establishing a universal cocartesian fibration over ∞ -categories

$$U : \mathcal{Z} \rightarrow \mathcal{Cat}_\infty.$$

It is then shown that each cocartesian fibration is realized as a (categorical) pullback along U ,

$$\begin{array}{ccc} \mathcal{E} & \longrightarrow & \mathcal{Z} \\ q \downarrow & & \downarrow \\ \mathcal{C} & \xrightarrow{F} & \mathcal{Cat}_\infty, \end{array}$$

and furthermore that the space of such pullback diagrams assembles into a contractible space. We refer to this uniquely determined functor F as the covariant transport functor along q , or as the functor which classifies q . One obtains a completely similar analysis of cartesian fibrations and classification via an applications of the opposite involution.

In this section we outline the above construction. Unlike at other points in this text we are not especially concerned with (all of) the technical details, and seek only to provide a coherent narrative which explains clearly what’s going on and how this stuff works.

We begin with a detour into $(\infty, 2)$ -categories. We then construct the universal cocartesian fibration \mathcal{Z} via a certain “category of objects”, and explain how each fiber $\mathcal{Z}_{\mathcal{E}}$ over a given ∞ -category $\mathcal{E} : * \rightarrow \mathcal{Cat}_\infty$ reproduces \mathcal{E} itself, up to equivalence. We define the space $\mathcal{T}(q)$ of classifying diagrams and recall the contractibility of this space from [4]. The section concludes with a description of the pushforward functors α_* appearing the transport F .

6.2. Categories with objects and the universal cocartesian fibration. From the $(\infty, 2)$ -category \mathcal{Cat}_∞ we can produce the simplicial of pointed ∞ -categories

$$(\mathcal{Cat}_\infty)_* := (\mathcal{Cat}_\infty)_{\Delta^0/}.$$

By Proposition 5.22 the simplicial set $(\mathcal{Cat}_\infty)_*$ is an $(\infty, 2)$ -category and the forgetful functor $(\mathcal{Cat}_\infty)_* \rightarrow \mathcal{Cat}_\infty$ is a interior fibration.

Definition 6.1. The ∞ -category of ∞ -categories with a distinguished object is the pith of the $(\infty, 2)$ -category of pointed ∞ -categories,

$$\mathcal{P}\mathcal{Cat}_\infty := ((\mathcal{Cat}_\infty)_*)^{\mathrm{Pith}}.$$

Remark 6.2. The \mathcal{P} suffix stands for “pointed”, though we heed the warning from [4, 020W] and do not label this ∞ -category as such.

Remark 6.3. There is a comparison functor $\mathcal{P}\mathcal{C}at_\infty \rightarrow (\mathcal{C}at_\infty)_*/$ which is, apparently, bijective on objects. However this map is not bijective on 1-morphisms so that it is not an isomorphism [4, 020Z].

Via an application of Corollary 5.21 we see that the forgetful functor restricts to provide a pullback diagram

$$\begin{array}{ccc} \mathcal{P}\mathcal{C}at_\infty & \longrightarrow & (\mathcal{C}at_\infty)_*/ \\ \downarrow & & \downarrow \\ \mathcal{C}at_\infty & \longrightarrow & \mathcal{C}at_\infty. \end{array}$$

The forgetful functor $\mathcal{P}\mathcal{C}at_\infty \rightarrow \mathcal{C}at_\infty$ is furthermore seen to be a cocartesian fibration in this case, via an application of Proposition 5.26. We record this result.

Proposition 6.4 ([4, 0213]). *The forgetful functor $\mathcal{P}\mathcal{C}at_\infty \rightarrow \mathcal{C}at_\infty$ is a cocartesian fibration.*

We call the above forgetful functor the *universal cocartesian fibration*, for reasons which will be apparent shortly.

Definition 6.5. We let $\text{univ} : \mathcal{P}\mathcal{C}at_\infty \rightarrow \mathcal{C}at_\infty$ denote the cocartesian fibration induced by the forgetful functor $(\mathcal{C}at_\infty)_* \rightarrow \mathcal{C}at_\infty$, as considered above.

6.3. A remark on notation. Our $(\infty, 2)$ -category $\mathcal{C}at_\infty$ is the $(\infty, 2)$ -category denoted by a bold \mathcal{QC} in [4, 020K]. Our $(\mathcal{C}at_\infty)_*/$ is the $(\infty, 2)$ -category denoted by a bold $\mathcal{QC}_{\text{Obj}}$ in [4, 0210]. The associated piths, which we’ve denoted $\mathcal{C}at_\infty$ and $\mathcal{P}\mathcal{C}at_\infty$ respectively, are the non-bolded ∞ -categories \mathcal{QC} and $\mathcal{QC}_{\text{Obj}}$ in [4].

6.4. Fibers of the universal map $\mathcal{P}\mathcal{C}at_\infty \rightarrow \mathcal{C}at_\infty$. In considering the universal cocartesian fibration $\mathcal{P}\mathcal{C}at_\infty \rightarrow \mathcal{C}at_\infty$, any point $e : \Delta^0 \rightarrow \mathcal{C}at_\infty$ corresponds to an ∞ -category $\mathcal{E} = e(0)$ and we have the pullback

$$\mathcal{P}\mathcal{C}at_\infty \times_{\mathcal{C}at_\infty} \{e\}$$

which is some other ∞ -category. Now, objects in this fiber are simply maps of ∞ -categories $* \rightarrow \mathcal{E}$, and hence are identified with objects in \mathcal{E} . Similarly, 1-simplices in the fiber are identified 2-simplices in the ∞ -category of ∞ -categories

$$\begin{array}{ccc} & * & \\ x \swarrow & & \searrow y \\ \mathcal{E} & \xrightarrow{=} & \mathcal{E}. \end{array}$$

These are, by definition, natural transformations $\alpha \in \text{Hom}_{\text{sSet}}(\Delta^1, \mathcal{E})$ with $\alpha|_0 = x$ and $\alpha|_1 = y$, i.e. 1-simplices $\alpha : x \rightarrow y$. So we observe an identification of 1-skeleta

$$\mathcal{E}[\leq 1] = \mathcal{P}\mathcal{C}at_\infty \times_{\mathcal{C}at_\infty} \{e\}[\leq 1].$$

As an application of Theorem 5.27 and Corollary 5.28, we see that this direct identification of simplices in low-dimension expands to an equivalence of ∞ -categories which calculates the fiber.

prop:univ_fibs

Proposition 6.6. *For any ∞ -category \mathcal{E} , which we can understand as a point $\mathcal{E} : * \rightarrow \mathcal{Cat}_\infty$, we have a categorical pullback square*

$$\begin{array}{ccc} \mathcal{E} & \longrightarrow & \mathcal{P}\mathcal{Cat}_\infty \\ \downarrow & & \downarrow \\ * & \xrightarrow[\mathcal{E}]{} & \mathcal{Cat}_\infty, \end{array}$$

and a corresponding equivalence between ∞ -categories $\theta : \mathcal{E} \xrightarrow{\sim} \mathcal{P}\mathcal{Cat}_\infty \times_{\mathcal{Cat}_\infty} \{\mathcal{E}\}$.

6.5. Covariant transport: classifying cocartesian fibrations.

Definition 6.7. Let $q : \mathcal{E} \rightarrow \mathcal{C}$ be a cocartesian fibration of ∞ -categories. We say a functor $F : \mathcal{C} \rightarrow \mathcal{Cat}_\infty$ classifies the cocartesian fibration q if the functors q and F fit into a categorical pullback diagram

$$\begin{array}{ccc} \mathcal{E} & \xrightarrow{\tilde{F}} & \mathcal{P}\mathcal{Cat}_\infty \\ q \downarrow & & \downarrow \text{univ} \\ \mathcal{C} & \xrightarrow{F} & \mathcal{Cat}_\infty. \end{array} \tag{11} \quad \boxed{\text{eq:672}}$$

In this case we say the above diagram witnesses F as a (covariant) transport functor along q .

Remark 6.8. The term “classifies” is used with some frequency in the works [2, 3]. However, [4] seems to prefer the term “transport representation” for a functor F as above. We will usually just refer to F as a, or the, transport functor for q .

We have an alternate characterization of transport functors via the fiber product $\mathcal{C} \times_{\mathcal{Cat}_\infty} \mathcal{P}\mathcal{Cat}_\infty$.

Lemma 6.9. *Let $q : \mathcal{E} \rightarrow \mathcal{C}$ be a cocartesian fibration. A diagram (14) witnesses $F : \mathcal{C} \rightarrow \mathcal{Cat}_\infty$ as transport along q if and only if the induced map to the fiber product*

$$\mathcal{E} \rightarrow \mathcal{C} \times_{\mathcal{Cat}_\infty} \mathcal{P}\mathcal{Cat}_\infty$$

is an equivalence of cocartesian fibrations over \mathcal{C} .

Proof. The fact that the map to the fiber is an equivalence follows by Proposition I-5.23. It follows by Proposition ?? and the diagram

$$\begin{array}{ccc} \mathcal{E} & \xrightarrow{\sim} & \mathcal{C} \times_{\mathcal{Cat}_\infty} \mathcal{P}\mathcal{Cat}_\infty \\ & \searrow q & \swarrow \\ & \mathcal{C} & \end{array}$$

that the equivalence in question is an equivalence of cocartesian fibrations. □

The fiber product considered above is often denoted

$$\int_{\mathcal{C}} F := \mathcal{C} \times_{\mathcal{Cat}_\infty} \mathcal{P}\mathcal{Cat}_\infty,$$

so that any functors $F : \mathcal{C} \rightarrow \mathcal{Cat}_\infty$ determines an associated cocartesian fibration $\int_{\mathcal{C}} F \rightarrow \mathcal{C}$. We find that F is a transport functor for $q : \mathcal{E} \rightarrow \mathcal{C}$ if there is an equivalence

$$\mathcal{E} \xrightarrow{\sim} \int_{\mathcal{C}} F \text{ in the overcategory } \mathbf{sSet}/_{\mathcal{C}}.$$

We also notes that transport functors are stable under restriction.

1em:687

Lemma 6.10. *Suppose we have a pullback square*

$$\begin{array}{ccc} \mathcal{E}' & \xrightarrow{g} & \mathcal{E} \\ q' \downarrow & & \downarrow q \\ \mathcal{C}' & \xrightarrow{f} & \mathcal{C} \end{array} \quad (12) \quad \text{eq:689}$$

in which q and q' are cocartesian fibrations between ∞ -categories, and consider a diagram of the form (14) which witnesses a functor $F : \mathcal{C} \rightarrow \mathcal{Cat}_\infty$ as transport along q . Then for $\tilde{F}' = \tilde{F}g$ and $F' = Ff$, the diagram

$$\begin{array}{ccc} \mathcal{E}' & \xrightarrow{\tilde{F}'} & \mathcal{P}\mathcal{Cat}_\infty \\ q' \downarrow & & \downarrow \\ \mathcal{C}' & \xrightarrow{F'} & \mathcal{Cat}_\infty. \end{array}$$

witnesses F' as a transport functor along q' .

Proof. Since q is a cocartesian fibration, and in particular an isofibration, the diagram (12) is a categorical pullback square Corollary I-5.22. The result now follows from the fact that categorical pullback squares are stable under composition [4, 033J]. \square

Given a cocartesian fibration we can now consider the simplicial subset in the functor category

$$\mathbf{Fun}(\mathcal{C}, \mathcal{Cat}_\infty) \times_{\mathbf{Fun}(\mathcal{E}, \mathcal{Cat}_\infty)} \mathbf{Fun}(\mathcal{E}, \mathcal{P}\mathcal{Cat}_\infty) \quad (13) \quad \text{eq:710}$$

which consists of diagrams witnessing transport for a given cocartesian fibration $q : \mathcal{E} \rightarrow \mathcal{C}$.

Definition 6.11. For a given cocartesian fibration $q : \mathcal{E} \rightarrow \mathcal{C}$, we let $\mathcal{T}(q)$ denote the simplicial subset in the fiber product (13) whose simplices correspond to diagrams

$$\begin{array}{ccc} \Delta^n \times \mathcal{E} & \xrightarrow{\tilde{F}} & \mathcal{P}\mathcal{Cat}_\infty \\ \Delta^n \times q \downarrow & & \downarrow \\ \Delta^n \times \mathcal{C} & \xrightarrow{F} & \mathcal{Cat}_\infty. \end{array}$$

which witness F as a covariant transport functor along $\Delta^n \times q$.

Stability of such diagrams under restriction (Lemma 6.10) assures us that $\mathcal{T}(q)$ is in fact a simplicial subset in the given fiber product.

thm:transport

Theorem 6.12 (Universality theorem [4, 02SC]). *For any cocartesian fibration $q : \mathcal{E} \rightarrow \mathcal{C}$, the terminal map $\mathcal{T}(q) \rightarrow *$ is a trivial Kan fibration.*

This result says that any cocartesian fibration q admits a covariant transport functor $F : \mathcal{C} \rightarrow \mathcal{Cat}_\infty$, and that this functor is uniquely determined up to a contractible space of choices.

6.6. Transport for cartesian fibrations. Given any cartesian fibration $p : \mathcal{E} \rightarrow \mathcal{C}$, we have the associated cocartesian fibration $p^{\text{op}} : \mathcal{E}^{\text{op}} \rightarrow \mathcal{C}^{\text{op}}$. So our analysis of classifying functors for cocartesian fibrations dualizes in the obvious ways to provide an analysis of classifying functors for cartesian fibrations.

Definition 6.13. Let $p : \mathcal{E} \rightarrow \mathcal{C}$ be a cartesian fibration of ∞ -categories. We say a functor $F : \mathcal{C}^{\text{op}} \rightarrow \mathcal{Cat}_\infty$ classifies the cartesian fibration q if the functors p^{op} and F fit into a categorical pullback diagram

$$\begin{array}{ccc} \mathcal{E}^{\text{op}} & \xrightarrow{\tilde{F}} & \mathcal{P}\mathcal{Cat}_\infty \\ q \downarrow & & \downarrow \text{univ} \\ \mathcal{C}^{\text{op}} & \xrightarrow{F} & \mathcal{Cat}_\infty. \end{array} \quad (14) \quad \boxed{\text{eq:672}}$$

In this case we say the above diagram witnesses F as a (contravariant) transport functor along p .

Given a cartesian fibration $p : \mathcal{E} \rightarrow \mathcal{C}$, we define the space of transport functors with witness in the obvious way $\mathcal{T}(p) := \mathcal{T}(p^{\text{op}})$. Theorem 6.12 implies contractibility of this space immediately.

thm:uniq_transp

Theorem 6.14 (Contravariant universality). *For any cartesian fibration $p : \mathcal{E} \rightarrow \mathcal{C}$, the terminal map $\mathcal{T}(p) \rightarrow *$ from the space of transport functors along p is a trivial Kan fibration.*

Again this establishes both the existence and uniqueness of contravariant transport.

6.7. Transport along edges. Let $q : \mathcal{E} \rightarrow \Delta^1$ be a cocartesian fibration, and $f : \Delta^1 \rightarrow \mathcal{Cat}_\infty$ be a transport functor for q . The functor f specifies a pair of ∞ -categories and a functor $F' : \mathcal{E}'_0 \rightarrow \mathcal{E}'_1$. We then have partial diagrams

$$\begin{array}{ccc} \mathcal{E}_i & & \mathcal{E}'_i \\ & \searrow & \swarrow \theta \\ & \{i\} \times_{\Delta^1} (\int_{\Delta^1} f), & \end{array}$$

where as before $\int_{\Delta^1} f$ is the fiber product $\Delta^1 \times_{\mathcal{Cat}_\infty} \mathcal{P}\mathcal{Cat}_\infty$, θ is the equivalence of Section ??, and $\mathcal{E}_i \rightarrow \int_{\Delta^1} f$ is the equivalence witnessing f as a transport functor. We choose equivalences $\xi_i : \mathcal{E}_i \rightarrow \mathcal{E}'_i$ which complete a 2-simplex

$$\begin{array}{ccc} \mathcal{E}_i & \xrightarrow{\xi_i} & \mathcal{E}'_i \\ & \searrow & \swarrow \theta \\ & \{i\} \times_{\Delta^1} (\int_{\Delta^1} f), & \end{array}$$

in \mathcal{Cat}_∞ . By pulling back along the ξ_i , the defining functor $F' : \mathcal{E}_0 \rightarrow \mathcal{E}_1$ is identified with a functor between the \mathcal{E}_i . The following determines this functor via implicit criterion.

prop:1_transport

Proposition 6.15. Fix $q : \mathcal{E} \rightarrow \Delta^1$ and $f : \Delta^1 \rightarrow \mathcal{Cat}_\infty$ as above. There is a functor $F : \mathcal{E}_1 \rightarrow \mathcal{E}_2$ and a natural transformation

$$\tilde{F} : \Delta^1 \times \mathcal{E}_0 \rightarrow \mathcal{E}$$

from the inclusion $\mathcal{E}_1 \rightarrow \mathcal{E}$ to the composite of F with the inclusion $\mathcal{E}_1 \rightarrow \mathcal{E}$ for which, at each object $s : * \rightarrow \mathcal{E}_0$, the corresponding map

$$\tilde{F}_s : \Delta^1 = \Delta^1 \times \{s\} \rightarrow \mathcal{E}$$

is q -cocartesian. This functor is unique up to equivalence, and fits into a diagram

$$\begin{array}{ccc} \mathcal{E}_0 & \xrightarrow{F} & \mathcal{E}_1 \\ \xi_0 \downarrow & \searrow & \downarrow \xi_1 \\ \mathcal{E}'_0 & \xrightarrow{F'} & \mathcal{E}'_1 \end{array}$$

in the ∞ -category \mathcal{Cat}_∞ , where the ξ_i are as above.

Proof. Combine [4, 01VS] and [4, 027K]. □

Now consider the global situation, where we have some cocartesian fibration $q : \mathcal{E} \rightarrow \mathcal{C}$ and classifying functor $\mathcal{C} \rightarrow \mathcal{Cat}_\infty$. At any morphism $\alpha : x \rightarrow y$ in \mathcal{C} , transport restricts to a functor $F' : \mathcal{E}'_x \rightarrow \mathcal{E}'_y$. According to Proposition 6.15, this functor is uniquely determined by a functorial choice of q -cocartesian maps between the fiber categories \mathcal{E}_x and \mathcal{E}_y . There is, in particular, a unique functor $F : \mathcal{E}_x \rightarrow \mathcal{E}_y$ and a transformation which fits into a diagram

$$\begin{array}{ccc} \Delta^1 \times \mathcal{E}_x & \longrightarrow & \mathcal{E} \\ \downarrow & & \downarrow \\ \Delta^1 & \xrightarrow{\alpha} & \mathcal{C} \end{array}$$

and sends each edge $\Delta^1 \times \{s\}$ in $\Delta^1 \times \mathcal{E}_x$ to a q -cartesian morphism in \mathcal{E} which lies over α . This functor fits into a diagram

$$\begin{array}{ccc} \mathcal{E}_x & \xrightarrow{F} & \mathcal{E}_y \\ \sim \downarrow & \searrow & \downarrow \sim \\ \mathcal{E}'_x & \xrightarrow{F'} & \mathcal{E}'_y \end{array}$$

6.8. Classification of left and right fibrations. We have the simplicial subcategory $\underline{\mathbf{Kan}} \rightarrow \underline{\mathbf{Cat}}_\infty$ and subsequent simplicial subset $\mathcal{Kan} \subseteq \mathbf{Cat}_\infty$. This simplicial subset is the full $(\infty, 2)$ -subcategory whose objects are precisely those ∞ -categories which are Kan complexes, and so the inclusion preserves thin 2-simplices. We now have the full $(\infty, 2)$ -subcategory $\mathcal{Kan}_{*/} \rightarrow (\mathbf{Cat}_\infty)_{*/}$ of pointed Kan complexes and the pullback diagram

$$\begin{array}{ccc} \mathcal{Kan}_{*/} & \longrightarrow & (\mathbf{Cat}_\infty)_{*/} \\ \downarrow & & \downarrow \\ \mathcal{Kan} & \longrightarrow & \mathbf{Cat}_\infty \end{array}$$

which restricts to a pullback diagram into the piths

$$\begin{array}{ccc} \mathcal{K}an_{*/} & \longrightarrow & (\mathcal{C}at_{\infty})_{*/} \\ \downarrow & & \downarrow \\ \mathcal{K}an & \longrightarrow & \mathcal{C}at_{\infty}. \end{array}$$

We recall that the map $\mathcal{K}an_{*/} \rightarrow \mathcal{K}an$ is a left fibration, by Corollary I-??.

`prop:kan_transp`

Proposition 6.16. *A cocartesian fibration $q : \mathcal{E} \rightarrow \mathcal{C}$ is a left fibration if and only if the corresponding transport functor $F : \mathcal{C} \rightarrow \mathcal{C}at_{\infty}$ has image in $\mathcal{K}an$. Similarly, a cartesian fibration $p : \mathcal{E} \rightarrow \mathcal{C}$ is a right fibration if and only if the corresponding transport functor $G : \mathcal{C}^{\text{op}} \rightarrow \mathcal{C}at_{\infty}$ has image in $\mathcal{K}an$.*

Proof. By Proposition ??, a cocartesian (resp. cartesian) fibration $\mathcal{E} \rightarrow \mathcal{C}$ is a left (resp. right) fibration if and only if its fibers over object $\text{sin } \mathcal{C}$ are Kan complexes. So the result follows by the calculation of the fibers of the pullback fibration $\int_{\mathcal{C}} F \rightarrow \mathcal{C}$ provided in Proposition 6.6. \square

This proposition tells us that any left fibration $q : \mathcal{E} \rightarrow \mathcal{C}$ fits into a categorical pullback square

$$\begin{array}{ccc} \mathcal{E} & \longrightarrow & \mathcal{K}an_{*/} \\ q \downarrow & & \downarrow \\ \mathcal{C} & \xrightarrow{F} & \mathcal{K}an \end{array}$$

for some functor F , and that F , considered as a functor into $\mathcal{C}at_{\infty}$, is the transport functor for q . In this way left fibrations are classified by maps into the ∞ -category of Kan complexes. One obtains similar statements for right fibrations by applying the opposite functor.

`sect:htf`

7. HOMOTOPY TRANSPORT REPRESENTATIONS

7.1. Homotopy transport representations. Consider a cocartesian fibration $q : X \rightarrow S$ and an edge $\alpha : s \rightarrow t$ in S . We then have the fibers X_s and X_t over these respective points, both of which are ∞ -categories.

We consider the diagram

$$\begin{array}{ccc} \{0\} \times X_s & \longrightarrow & X \\ \downarrow & & \downarrow q \\ \Delta^1 \times X_s & \longrightarrow & S \end{array}$$

where the top arrow is the inclusion and the bottom arrow is the composite of the projection with α ,

$$\Delta^1 \times X_s \rightarrow \Delta^1 \times \{s\} \xrightarrow{ev_{\alpha}} S.$$

By Theorem 2.7 the above diagram is split by a transformation

$$\xi_{\alpha} : \Delta^1 \times X_s \rightarrow X$$

which has $\alpha_1|_{\{0\} \times X_s}$ equal to the inclusion and has

$$\xi_{\alpha}|_{\Delta^1 \times \{s'\}} : \Delta^1 \times \{s'\} \rightarrow X$$

a q -cocartesian morphism over α . In particular, the restriction at 1 produces a functor

$$\alpha_! := \xi_\alpha|_{\{1\} \times X_s} : X_s \rightarrow X_t.$$

Furthermore, this transformation ξ_α is uniquely determined up to a contractible space of choices, so that $\alpha_!$ is similarly uniquely determined up to a contractible space as well.

Definition 7.1. Given a cocartesian fibration $q : X \rightarrow S$ and any edge $\alpha : s \rightarrow t$ in the base, we let $\alpha_! : X_s \rightarrow X_t$ denote the uniquely determined functor which comes equipped with a cocartesian transformation ξ_α over α , as above. We call $\alpha_! : X_s \rightarrow X_t$ the homotopy transport functor over α .

prop:1297

Proposition 7.2. Let $X \rightarrow S$ be a cocartesian fibration. Suppose that we have a 2-simplex $A : \Delta^2 \rightarrow S$ and take $\alpha_{ij} = A|_{\Delta^{\{i,j\}}} : s_i \rightarrow s_j$. Then there is an isomorphism

$$(\alpha_{02})_! \cong (\alpha_{12})_!(\alpha_{01})_!$$

in $\text{Fun}(X_{s_0}, X_{s_2})$.

Proof. Take $X_i = X_{s_i}$. Consider a diagram $\tilde{A} : \Delta^1 \times \Delta^1 \rightarrow S$ which appears as

$$\begin{array}{ccc} s_2 & \xrightarrow{id} & s_2 \\ \alpha_{02} \uparrow & \nearrow \alpha_{02} & \uparrow \alpha_{12} \\ s_0 & \xrightarrow{\alpha_{01}} & s_1 \end{array},$$

which we might obtain by expanding A for example. Then we have a lifting problem

$$\begin{array}{ccc} \{0\} \times \Delta^1 \times X_0 & \xrightarrow{\xi_{\alpha_{02}}} & X \\ \downarrow & \nearrow \text{dotted} & \downarrow \\ \Delta^1 \times \Delta^1 \times X_0 & \xrightarrow{\tilde{A} \times \{s_0\}} & S \end{array}$$

and can take $\xi_A : \Delta^1 \times \Delta^1 \times X_0 \rightarrow X$ to be the unique cocartesian solution. Restricting ξ_A to $\Delta^1 \times \{0\} \times X_0$ provides a cocartesian lift of α_{01} and so is identified with $\xi_{\alpha_{01}}$, and similarly restricting ξ_A to the edge $\Delta^1 \times \{1\} \times X_0$ is identified with $id_{\alpha_{02}}$. By the 2-of-3 property for q -cocartesian maps, we also see that the restriction of ξ_A to the diagonal $\Delta^1 \times X_0$ is a cocartesian lift of α_{02} and so is identified with $\xi_{\alpha_{02}}$ as well.

We have only the edge $\{1\} \times \Delta^1 \times X_0 \rightarrow X$ to be identified. By the 2-of-3 property again we see that F_A provides a cocartesian solution to the diagram

$$\begin{array}{ccccc} \{1\} \times \{0\} \times X_0 & \xrightarrow{\xi_A|_{\dots}} & \{0\} \times X_1 & \xrightarrow{\quad} & X \\ \downarrow & \searrow (\alpha_{01})_! & \downarrow & \downarrow & \downarrow \\ \{1\} \times \Delta^1 \times X_0 & \xrightarrow{id \times (\alpha_{01})_!} & \Delta^1 \times X_1 & \longrightarrow & S \end{array}$$

However we have the alternate cocartesian lift

$$\begin{array}{ccccc} \{1\} \times \{0\} \times X_0 & \xrightarrow{\quad (\alpha_{01})_! \quad} & \{0\} \times X_1 & \longrightarrow & X \\ \downarrow & & \downarrow & \nearrow \xi_{\alpha_{12}} & \downarrow \\ \{1\} \times \Delta^1 \times X_0 & \xrightarrow{\quad id \times (\alpha_{01})_! \quad} & \Delta^1 \times X_1 & \longrightarrow & S \end{array}$$

so that there is a unique isomorphism

$$\xi_A|_{\{1\} \times \Delta^1 \times X_0} \cong \xi_{\alpha_{12}}(id \times (\alpha_{01})_!).$$

So we restrict further to $\{1\} \times \{1\} \times X_0$ to get

$$(\alpha_{02})_! = \xi_{\alpha_{02}}|_{\{1\}} \cong \xi_A|_{\{1\} \times \{1\}} \cong \xi_{\alpha_{12}}(id \times (\alpha_{01})_!)|_{\{1\}} = (\alpha_{12})_!(\alpha_{01})_!.$$

□

Corollary 7.3. *Let $q : \mathcal{E} \rightarrow \mathcal{C}$ be a cocartesian fibration over an ∞ -category \mathcal{C} . The functors $\alpha_! : X_s \rightarrow X_t$ assemble into a functor into the homotopy category of ∞ -categories $\bar{q}_! : \mathbf{h}\mathcal{C} \rightarrow \mathbf{h}\mathcal{Cat}_\infty$.*

Definition 7.4. Let $q : \mathcal{E} \rightarrow \mathcal{C}$ be a cocartesian fibration over an ∞ -category \mathcal{C} . The homotopy transport representation for q is the functor on homotopy categories

$$\bar{q}_! : \mathbf{h}\mathcal{C} \rightarrow \mathbf{h}\mathcal{Cat}_\infty$$

is the functor whose value at each object $x : * \rightarrow \mathcal{C}$ is the fiber $\bar{q}_!(x) = \mathcal{E}_x$, and whose value at any morphism $\alpha : x \rightarrow y$ in \mathcal{C} is the associated homotopy transport functor $\alpha_! : \mathcal{E}_x \rightarrow \mathcal{E}_y$.

More generally, we call any functor $F : \mathbf{h}\mathcal{C} \rightarrow \mathbf{h}\mathcal{Cat}_\infty$ which comes equipped with a natural isomorphism $\zeta : F \xrightarrow{\sim} \bar{q}_!$ a homotopy transport representation for q .

One observes that the homotopy transport representation is natural in diagrams of cocartesian fibrations.

lem:1366

Lemma 7.5. *Let*

$$\begin{array}{ccc} Y & \xrightarrow{F} & X \\ \downarrow & & \downarrow \\ T & \xrightarrow{f} & S \end{array}$$

be a diagram of cocartesian fibrations, i.e. a diagram in which F preserves cocartesian maps. Then for any edge $\alpha : t \rightarrow t'$ in T the homotopy transport functors fit into a diagram

$$\begin{array}{ccc} Y_t & \xrightarrow{\alpha_!} & Y_{t'} \\ F \downarrow & & \downarrow F \\ X_{f(t)} & \xrightarrow{f(\alpha)_!} & X_{f(t')} \end{array}$$

in \mathcal{Cat}_∞ .

Proof. Follows from the fact that both $F\xi_\alpha$ and $\xi_{f(\alpha)}f$ provide cocartesian lifts for the diagram

$$\begin{array}{ccccc} \{0\} \times Y_t & \longrightarrow & \{0\} \times X_{f(t)} & \longrightarrow & X \\ \downarrow & & \downarrow & & \downarrow \\ \Delta^1 \times Y_t & \longrightarrow & \Delta^1 \times X_{f(t)} & \longrightarrow & S. \end{array}$$

□

7.2. The $\mathbf{h}\mathcal{K}an$ -enriched category of ∞ -categories. Let \underline{A} be a simplicial category whose morphism complexes are all Kan complexes. Via the functor to the homotopy category $\pi : \mathbf{Kan} \rightarrow \mathbf{h}\mathcal{K}an$ we obtain a new category $\pi \underline{A}$ which is enriched in $\mathbf{h}\mathcal{K}an$. (Here we note that the usual product of Kan complexes endows $\mathbf{h}\mathcal{K}an$ with a unique symmetric monoidal structure under which the projection $\pi : \mathbf{Kan} \rightarrow \mathbf{h}\mathcal{K}an$ is symmetric monoidal.) We compare this Kan enriched category to the Kan enriched category $\pi N^{\mathbf{hc}}(\underline{A})$ obtained via the mapping spaces in the homotopy coherent nerve and their associated composition functions of Section I-8.

prop:pi_hcnerv

Proposition 7.6 ([4, 02LN]). *Let \underline{A} be a simplicial category whose morphism complexes are Kan complexes, and let $\mathcal{A} = N^{\mathbf{hc}}(\underline{A})$ denote its associated ∞ -category. Then the natural equivalences*

$$\underline{\mathrm{Hom}}_{\underline{A}}(x, y) \xrightarrow{\sim} \mathrm{Hom}_{\mathcal{A}}^L(x, y) \xrightarrow{\sim} \mathrm{Hom}_{\mathcal{A}}(x, y)$$

supplied by Theorems 5.27 and I-9.3 define an equivalence of $\mathbf{h}\mathcal{K}an$ -enriched categories $\pi \underline{A} \rightarrow \pi \mathcal{A}$.

sect:ehf

7.3. Enriched homotopy transport. Given a simplicial set S and vertices $s, t : * \rightarrow S$ we take $\mathrm{Hom}_S(s, t) = \mathrm{Fun}(\Delta^1, S) \times_{S \times S} \{(s, t)\}$. For any cocartesian fibration $q : X \rightarrow S$ we consider the evaluation map

$$\Delta^1 \times \mathrm{Hom}_S(s, t) \times X_s \rightarrow \Delta^1 \times \mathrm{Fun}(\Delta^1, S) \times \{s\} \xrightarrow{ev} S$$

and the diagram

$$\begin{array}{ccc} \{0\} \times \mathrm{Hom}_S(s, t) \times X_s & \longrightarrow & X \\ \downarrow & & \downarrow \\ \Delta^1 \times \mathrm{Hom}_S(s, t) \times X_s & \longrightarrow & S. \end{array}$$

The top map is the composite of the projection to X_s with the inclusion $X_s \rightarrow X$. By Theorem 2.7 there is a unique functor

$$\xi : \Delta^1 \times \mathrm{Hom}_S(s, t) \times X_s \rightarrow X$$

which splits the above diagram and sends each edge $\Delta^1 \times \{(\alpha, x)\}$ to a q -cocartesian morphism in X . The uniqueness claim of Theorem 2.7 tells us that, and each α in $\mathrm{Hom}_S(s, t)$, ξ restricts to the transformation ξ_α appearing in the definition of the homotopy transport functor α_t . So the map

$$\xi|_{\{1\}} : \mathrm{Hom}_S(s, t) \times X_s \rightarrow X_t$$

provides a parameterized family of morphisms whose fibers are the homotopy transport functors α_t .

def:pht

Definition 7.7. Given a cocartesian fibration $q : X \rightarrow S$, we call the functor $\xi|_{\{1\}} : \text{Hom}_S(s, t) \times X_s \rightarrow X_t$ constructed above the parametrized homotopy transport functor for q .

Note that we can view $\xi|_{\{1\}}$ as a functor

$$q_{s,t} : \text{Hom}_S(s, t) \rightarrow \text{Fun}(X_s, X_t)$$

via adjunction. In the case that S is an ∞ -category, we note that $\tilde{q}_{s,t}$ is a functor between ∞ -categories.

Consider now a cocartesian fibration $\mathcal{E} \rightarrow \mathcal{C}$ over an ∞ -category \mathcal{C} . Recall our notation $\pi\mathcal{C}$ for the $\text{h}\mathcal{K}an$ -enriched category with By similar arguments to those employed in our analysis of the homotopy transport functors $\alpha_! : X_s \rightarrow X_t$, one sees that these maps assemble into a functor of $\text{h}\mathcal{K}an$ -enriched categories

$$\pi\mathcal{C} \rightarrow \pi\text{Cat}_\infty^+$$

which lifts the homotopy transport functor $\tilde{q}_!$ of Section 7.1.

Definition 7.8. Given a cocartesian fibration $q : \mathcal{E} \rightarrow \mathcal{C}$ over an ∞ -category \mathcal{C} , we let

$$q_! : \pi\mathcal{C} \rightarrow \pi\mathcal{C}at_\infty$$

denote the $\text{h}\mathcal{K}an$ -enriched functor whose value at any object $x : * \rightarrow \mathcal{C}$ is the fiber \mathcal{E}_x , and whose values on morphisms

$$q_! : \text{Hom}_{\mathcal{C}}(x, y) \rightarrow \text{Fun}(\mathcal{E}_x, \mathcal{E}_y)^{\text{Kan}} \cong \text{Hom}_{\mathcal{C}at_\infty}(\mathcal{E}_x, \mathcal{E}_y)$$

are the functors induced by parametrized homotopy transport. We call $q_!$ the enriched homotopy transport representation associated to q .

We generally, we call any enriched functor $F : \pi\mathcal{C} \rightarrow \pi\mathcal{C}at_\infty$ which comes equipped with a natural isomorphism $\zeta : F \xrightarrow{\sim} q_!$ a homotopy transport representation for q .

Note that here we've employed the natural identification $\pi\text{Cat}_\infty^+ \cong \mathcal{C}at_\infty$ provided by Proposition 7.6 here when we replace the functor categories with the Hom-spaces for $\mathcal{C}at_\infty$.

One again sees that the enriched homotopy transport representation is natural

Lemma 7.9. *Consider a diagram of ∞ -categories*

$$\begin{array}{ccc} \mathcal{E} & \xrightarrow{G} & \mathcal{K} \\ q \downarrow & & \downarrow p \\ \mathcal{C} & \xrightarrow{F} & \mathcal{D} \end{array} \quad (15) \quad \boxed{\text{eq:1464}}$$

in which p and q are cocartesian fibrations and G preserves cocartesian maps. Suppose also that \mathcal{C} and \mathcal{D} are ∞ -categories. Then the maps $G|_{\mathcal{E}_x} : \mathcal{E}_x \rightarrow \mathcal{K}_{F(x)}$ define a natural transformation between the enriched homotopy transport representations

$$G_! : q_! \rightarrow p_!F.$$

The proof is the same as that of Lemma 7.5. We note that when the diagram (15) is a categorical pullback diagram then the maps $G|_{\mathcal{E}_x} : \mathcal{E}_x \rightarrow \mathcal{K}_{F(x)}$ are isomorphisms in $\text{h}\mathcal{K}an$, so that $\tilde{G}_!$ is a natural isomorphism of $\text{h}\mathcal{K}an$ -enriched functors.

Lemma 7.10. *If a diagram (15) is a pullback diagram of cocartesian fibrations, then the composite functor*

$$\pi\mathcal{C} \xrightarrow{F} \pi\mathcal{D} \xrightarrow{p_!} \pi\mathcal{Cat}_\infty$$

is an enriched homotopy transport functor for q .

7.4. Transport functors induce homotopy transport. We have the following fundamental result concerning homotopy transport.

Theorem 7.11 ([4, 02S5]). *Consider the universal cocartesian fibration $U : \mathcal{P}\mathcal{Cat}_\infty \rightarrow \mathcal{Cat}_\infty$. The equivalences*

$$\theta : \mathcal{C} \rightarrow (\mathcal{P}\mathcal{Cat}_\infty)_{\mathcal{C}}$$

from Corollary 5.28 define a natural isomorphism $id_{\pi\mathcal{Cat}_\infty} \xrightarrow{\sim} U_!$. This isomorphism realizes the identity functor $id : \pi\mathcal{Cat}_\infty \rightarrow \pi\mathcal{Cat}_\infty$ as enriched homotopy transport for the universal fibration.

The proof proceeds by a more general analysis of homotopy transport for cocartesian fibrations of the form

$$(N^{\text{hc}}(\underline{A})_{x/})^{\text{Pith}} \rightarrow N^{\text{hc}}(\underline{A})^{\text{Pith}},$$

where \underline{A} a simplicial category which is enriched in ∞ -categories [4, 02RZ]. We omit the details and refer the reader instead to the text [4].

Now, given an arbitrary cocartesian fibration $q : \mathcal{E} \rightarrow \mathcal{C}$ we have a categorical pullback diagram

$$\begin{array}{ccc} \mathcal{E} & \longrightarrow & \mathcal{P}\mathcal{Cat}_\infty \\ q \downarrow & & \downarrow \\ \mathcal{C} & \xrightarrow{F} & \mathcal{Cat}_\infty \end{array} \quad (16) \quad \text{eq:1525}$$

which identifies a given functor F as the transport functor for q . Since homotopy transport is preserved under categorical pullback, we conclude that transport functors always induce enriched homotopy transport at the level of the enriched homotopy category.

Corollary 7.12. *For any cocartesian fibration $q : \mathcal{E} \rightarrow \mathcal{C}$, and classifying functor $F : \mathcal{C} \rightarrow \mathcal{Cat}_\infty$, the induced map on enriched homotopy categories*

$$\pi F : \pi\mathcal{C} \rightarrow \pi\mathcal{Cat}_\infty$$

is an enriched homotopy transport functor for q . More specifically, the isomorphisms $\mathcal{E}_x \rightarrow F(x)$ in $\mathbf{h}\mathcal{K}\mathbf{an}$ provided by the diagram (16) and Theorem 5.27 define an isomorphism of enriched functors $\pi F \xrightarrow{\sim} q_!$.

7.5. Homotopy transport and enriched Hom functors. For an ∞ -category \mathcal{C} and an object $x : * \rightarrow \mathcal{C}$, we recall the oriented fiber product $\{x\} \tilde{\times}_{\mathcal{C}} \mathcal{C}$, which is the explicitly the isofibration

$$\{x\} \times_{\text{Fun}(\{0\}, \mathcal{C})} \text{Fun}(\Delta^1, \mathcal{C}) \rightarrow \text{Fun}(\{1\}, \mathcal{C}) = \mathcal{C}.$$

We have the equivalence of isofibrations

$$\begin{array}{ccc} \mathcal{C}_{x/} & \xrightarrow{\sim} & \{x\} \tilde{\times}_{\mathcal{C}} \mathcal{C} \\ & \searrow & \swarrow \\ & \mathcal{C} & \end{array}$$

of Theorem I-9.14, from which we conclude that the projection $\{x\} \times_{\mathcal{C}} \mathcal{C} \rightarrow \mathcal{C}$ is in fact a left fibration.

We assess homotopy transport $\mathcal{C}_{x/} \rightarrow \mathcal{C}$ by considering this equivalent fibration. We note that the fibers of $\{x\} \times_{\mathcal{C}} \mathcal{C}$ over \mathcal{C} are simply the mapping spaces $\text{Hom}_{\mathcal{C}}(x, y)$.

Proposition 7.13. *Let \mathcal{C} be an ∞ -category and $x_0 : * \rightarrow \mathcal{C}$ be any object. The composition functions*

$$\circ : \text{Hom}_{\mathcal{C}}(x_1, x_2) \times \text{Hom}_{\mathcal{C}}(x, x_1) \rightarrow \text{Hom}_{\mathcal{C}}(x, x_2)$$

from Section I-8.1 are parametrized homotopy transport for the left fibration $\{x\} \times_{\mathcal{C}} \mathcal{C} \rightarrow \mathcal{C}$.

Proof. Take $x_0 = x$ and $\vec{x} = (x_0, x_1, x_2)$. A morphism $\Delta^1 \times \text{Hom}(x_1, x_2) \times \text{Hom}(x_0, x_1) \rightarrow \{x_0\} \times_{\mathcal{C}} \mathcal{C}$ is equivalent to a choice of a morphism

$$\Delta^1 \times \Delta^1 \times \text{Hom}(x_1, x_2) \times \text{Hom}(x_0, x_1) \rightarrow \mathcal{C}$$

whose restriction to $\{0\}$ in the first argument is of constant value x_0 . Let $h : \Delta^1 \times \Delta^1 \rightarrow \Delta^2$ be the map which sends $(0, j)$ to 0 and $(1, j)$ to $j + 1$ and let

$$\omega : \text{Hom}(x_1, x_2) \times \text{Hom}(x_0, x_1) \rightarrow \text{Fun}(\Delta^2, \mathcal{C})_{\vec{x}}$$

be any section of the trivial Kan fibration

$$\text{Fun}(\Delta^2, \mathcal{C})_{\vec{x}} \rightarrow \text{Fun}(\Delta_1^2, \mathcal{C})_{\vec{x}} = \text{Hom}(x_1, x_2) \times \text{Hom}(x_0, x_1).$$

We consider the composite

$$\begin{aligned} \Delta^1 \times \Delta^1 \times \text{Hom}(x_1, x_2) \times \text{Hom}(x_0, x_1) &\xrightarrow{h \times id} \Delta^2 \times \text{Hom}(x_1, x_2) \times \text{Hom}(x_0, x_1) \quad (17) \\ &\xrightarrow{id \times \omega} \Delta^2 \times \text{Fun}(\Delta^2, \mathcal{C})_{\vec{x}} \xrightarrow{ev} \mathcal{C}. \end{aligned} \quad \boxed{\text{eq:1582}}$$

One sees directly that this composite is of constant value x_0 when restricted to $\{0\}$ in the first argument, and the restriction to $\{1\}$ in the first argument yields the map

$$\Delta^1 \times \text{Hom}(x_1, x_2) \times \text{Hom}(x_0, x_1) \rightarrow \Delta^1 \times \text{Hom}(x_1, x_2) \xrightarrow{ev} \mathcal{C}$$

since ω is a section of the aforementioned fibration. This implies commutativity of the diagram

$$\begin{array}{ccc} \{0\} \times \text{Hom}(x_1, x_2) \times \text{Hom}(x_0, x_1) & \longrightarrow & \{x\} \times_{\mathcal{C}} \mathcal{C} \\ \downarrow & \nearrow \xi & \downarrow \\ \Delta^1 \times \text{Hom}(x_1, x_2) \times \text{Hom}(x_0, x_1) & \longrightarrow & \mathcal{C}, \end{array}$$

where ξ is adjoint to the composite (17). So therefore realize the restriction

$$\xi|_{\{1\}} : \Delta^1 \times \text{Hom}(x_1, x_2) \times \text{Hom}(x_0, x_1) \rightarrow \text{Hom}(x_0, x_2)$$

as enriched transport for the given fibration, which one checks directly is simply the composition function for $\text{Hom}_{\mathcal{C}}$, i.e. the uniquely determined composite

$$\text{Hom}(x_1, x_2) \times \text{Hom}(x_0, x_1) \rightarrow \text{Fun}(\Delta^2, \mathcal{C})_{\vec{x}} \rightarrow \text{Fun}(\Delta^{\{0,1\}}, \mathcal{C})_{(x_0, x_2)} = \text{Hom}(x_0, x_2)$$

in $\mathcal{H}an$. \square

We now find that the Hom-functor

$$\mathrm{Hom}_{\mathcal{C}}(x, -) : \pi\mathcal{C} \rightarrow \pi\mathcal{K}an$$

is the enriched homotopy transport representation for the oriented fiber product $\{x\} \tilde{\times}_{\mathcal{C}} \mathcal{C} \rightarrow \mathcal{C}$, and hence also for the fibration $\mathcal{C}_{x/} \rightarrow \mathcal{C}$.

Corollary 7.14. *Let \mathcal{C} be any ∞ -category and $x : * \rightarrow \mathcal{C}$ be any object. The Hom-functor*

$$\mathrm{Hom}_{\mathcal{C}}(x, -) : \pi\mathcal{C} \rightarrow \pi\mathcal{K}an$$

is an enriched homotopy transport functor for the left fibration $\mathcal{C}_{x/} \rightarrow \mathcal{C}$.

8. NATURALITY OF TRANSPORT: STRAIGHTENING AND UNSTRAIGHTENING

In this section we explain how the assignment which sends a functor $F : \mathcal{C} \rightarrow \mathcal{Cat}_{\infty}$ to the corresponding cocartesian fibration

$$\int_{\mathcal{C}} F = \mathcal{C} \times_{\mathcal{Cat}_{\infty}} \mathcal{P}\mathcal{Cat}_{\infty} \rightarrow \mathcal{C}$$

extends to an equivalence of ∞ -categories

$$\mathrm{Un} : \mathrm{Fun}(\mathcal{C}, \mathcal{Cat}_{\infty}) \xrightarrow{\sim} \mathrm{Cocart}(\mathcal{C}),$$

where $\mathrm{Cocart}(\mathcal{C})$ is an ∞ -category of cocartesian fibrations over \mathcal{C} . Similarly, we have an equivalence for cartesian fibrations

$$\mathrm{Un} : \mathrm{Fun}(\mathcal{C}^{\mathrm{op}}, \mathcal{Cat}_{\infty}) \xrightarrow{\sim} \mathrm{Cart}(\mathcal{C}),$$

which one obtains simply by applying the opposite involution. These functors are referred to as the unstraightening functors [2].

Our investigation centers not the unstraightening functor per-se, but its discrete inverse, the aptly named straightening functor. In our discussion we generalize our base to a simplicial set S rather than an ∞ -category.

8.1. Marked simplicial sets. A marked simplicial set is a pair (K, W) consisting of a simplicial set K and a choice of 1-simplices $W \subseteq K[1]$ which contains all degenerate 1-simplices. A map between marked simplicial sets $f : (K, W) \rightarrow (K', W')$ is a map of simplicial sets which sends W into W' . In this way we obtain the category sSet^+ of marked simplicial sets.

The forgetful functor $\mathrm{sSet}^+ \rightarrow \mathrm{sSet}$ has a right adjoint $-^{\#} : \mathrm{sSet} \rightarrow \mathrm{sSet}^+$, which sends a simplicial set K to the simplicial set K with all 1-simplices marked. The category sSet^+ also admits products, with $(K, W) \times (K', W') = (K \times K', W \times W')$, and the right adjoint $-^{\#}$ is seen to be symmetric monoidal. Via this monoidal functor we obtain an action of sSet on sSet^+ .

As a notational point, from this point on we often omit markings from our notation, and simply say K is a marked simplicial set to indicate that K is a simplicial set equipped with some specified marking (K, W) .

Lemma 8.1. *The sSet -module category sSet^+ admits inner-Homs $\underline{\mathrm{Hom}}_{\mathrm{sSet}^+}(K, K')$. The n -simplices in the underlying simplicial set maps*

$$\underline{\mathrm{Hom}}_{\mathrm{sSet}^+}(K, K')[n] := \mathrm{Hom}_{\mathrm{sSet}^+}((\Delta^n)^{\#} \times K, K').$$

Note that this simplicial set is a simplicial subset in the usual inner-Homs for simplicial sets.

lem:1074

Lemma 8.2. *Let (K, W) and (K', W') be marked simplicial sets and suppose that the marked vertices in K' are stable under compositions, i.e. that for any simplex*

$$s : \Delta^2 \rightarrow K'$$

in which edges $s|_{\Delta^{\{i,i+1\}}}$ are marked, the edge $s|_{\Delta^{\{0,2\}}}$ is marked as well. A map of unmarked simplicial sets $F : \Delta^n \times K \rightarrow K'$ is an n -simplex in $\underline{\text{Hom}}_{\text{sSet}^+}(K, K')$ if and only if the following hold:

- (a) *The restrict to each vertex $F|_{\Delta^{\{i\}}} : K \rightarrow K'$ is a map of marked simplicial sets.*
- (b) *At each vertex $x : * \rightarrow K$, and each $0 \leq i < j \leq n$, the edge $F|_{\Delta^{\{i,j\}} \times \{x\}} \rightarrow K$ is marked in K' .*

Proof. The marked edges in $(\Delta^n)^\# \times K$ are all pairs (α_{ij}, w) where $\alpha_{ij} : [1] \rightarrow [n]$ is the unique increasing map with image $\{i, j\}$ and $w : x \rightarrow y$ is any marked edge in K . Since all degenerate 1-simplices are marked, it is clear that any marked map F must satisfy (a) and (b). So let us suppose now that F satisfies (a) and (b), and consider a marked edge (α_{ij}, w) with $i \leq j$. Such an edge appears as the $\{0, 3\}$ edge in a 2-simplex

$$t : \Delta^2 \rightarrow \Delta^n \times K \quad \text{with} \quad t|_{\Delta^{\{0,1\}}} = (\alpha_{i,j}, x) \quad \text{and} \quad t|_{\Delta^{\{1,2\}}} = (\alpha_{i,j}, w).$$

and $Ft : \Delta^2 \rightarrow K'$ is now a 2-simplex in K' with both edges $Ft|_{\Delta^{\{i,i+1\}}}$ marked. (In the case $i = j$ the simplex $Ft|_{\Delta^{\{i,j\}}}$ is marked simply because it is degenerate, otherwise this follows by (a).) It follows by stability under composition that $Ft|_{\Delta^{\{0,2\}}} = F(\alpha_{ij}, w)$ is marked as well. So F preserves markings. \square

The following is an alternate phrasing of this lemma

lem:1098

Lemma 8.3. *The mapping complex $\underline{\text{Hom}}_{\text{sSet}^+}(K, K')$ is a subcomplex in the complex of simplicial maps $\text{Fun}(K, K')$. An n -simplex $F : \Delta^n \rightarrow \text{Fun}(K, K')$ lies in $\underline{\text{Hom}}_{\text{sSet}^+}(K, K')$ if and only if each functor $F_i : \Delta^{\{i\}} \rightarrow \text{Fun}(K, K')$ preserved markings and, at each $x : * \rightarrow K'$ and $i < j$, the image*

$$\Delta^1 \cong \Delta^{\{i,j\}} \xrightarrow{F_{ij}} \text{Fun}(K, K') \xrightarrow{x^*} \text{Fun}(*, K') = K'$$

is a marked map in K' .

We are also interested in the nature of 2-simplices in $\underline{\text{Hom}}_{\text{sSet}^+}(K, K')$. The following is obtained from Lemma 8.3, essentially immediately.

cor:1108

Corollary 8.4. *Suppose that K and K' are marked simplicial sets, and that the markings in K' satisfy the 2-of-3 property. A 2-simplex $\Delta^2 \rightarrow \text{Fun}(K, K')$ lies in $\underline{\text{Hom}}_{\text{sSet}^+}(K, K')$ if and only if each constituent map $\Delta^{\{i\}} \rightarrow \text{Fun}(K, K')$ is marked, and 2 of the 3 edges $\Delta^{\{i < j\}} \rightarrow \text{Fun}(K, K')$ is marked.*

Given a simplicial set S we let sSet_S^+ denote the simplicial category whose objects are morphisms of marked simplicial sets $p : K \rightarrow S^\#$, whose morphism complexes $\underline{\text{Hom}}_S(K, K')$ are the fiber products

$$\begin{array}{ccc} \underline{\text{Hom}}_S(K, K') & \longrightarrow & \underline{\text{Hom}}_{\text{sSet}^+}(K, K') \\ \downarrow & & \downarrow (p')_* \\ * & \xrightarrow{p} & \underline{\text{Hom}}_{\text{sSet}^+}(K, S^\#) = \text{Fun}(K, S) \end{array}$$

prop:cocart_kan

Proposition 8.5. *Let $q : X \rightarrow S$ be a cocartesian fibration, and consider the associated map of marked simplicial sets $X^q \rightarrow S^\#$ where X^q is X paired with the collection of all q -cocartesian maps. Then for any marked simplicial set K over $S^\#$, the mapping complex $\underline{\mathrm{Hom}}_S(K, X^q)$ is a Kan complex.*

Proof. By 14.8 the map $\mathrm{Fun}(K, X) \rightarrow \mathrm{Fun}(K, S)$ is an inner fibration, so that the fiber $\mathrm{Fun}_S(K, X)$ over the diagram p is an ∞ -category. We have now that $\underline{\mathrm{Hom}}_S(K, X^q)$ is a subcomplex in the ∞ -category $\mathrm{Fun}_S(K, X)$ whose n -simplices are precisely those n -simplices in $\mathrm{Fun}_S(K, X)$ which satisfy the conditions specified in Lemma 8.2 (a) and (b), since q -cocartesian maps in X are stable under composition. This stability under composition in X also implies that any completion $\Delta^2 \times K \rightarrow X$ of an inner horn $\Lambda_1^2 \times K \rightarrow X$ which lies in $\underline{\mathrm{Hom}}_S(K, X^q)$ also lies in $\underline{\mathrm{Hom}}_S(K, X^q)$. Since the subcomplex $\underline{\mathrm{Hom}}_S(K, X^q)$ in $\mathrm{Fun}(K, X)$ is characterized by a restriction on the 1-simplices, it follows that all higher dimensional horns in $\underline{\mathrm{Hom}}_S(K, X^q)$ complete to simplices in $\underline{\mathrm{Hom}}_S(K, X^q)$. So $\underline{\mathrm{Hom}}_S(K, X^q)$ is an ∞ -subcategory in $\mathrm{Fun}_S(K, X)$.

Now, for any 1-simplex $\zeta : \Delta^1 \rightarrow \underline{\mathrm{Hom}}_S(K, X^q)$ and vertex $x : * \rightarrow K$ with image s in S , the composite

$$\Delta^1 \rightarrow \underline{\mathrm{Hom}}_S(K, X^q) \xrightarrow{x^*} X$$

has q -cocartesian image in the ∞ -category X_s , and in particular is an isomorphism in X_s . This implies that ζ is an isomorphism in the ambient category $\mathrm{Fun}_S(K, X)$, by Proposition 1-6.8. From the 2 of 3 property for q -cocartesian morphisms, and Corollary 8.4, it follows that any inverse $\zeta^{-1} : \Delta^1 \rightarrow \mathrm{Fun}_S(K, X)$ to ζ is also in $\underline{\mathrm{Hom}}_S(K, X^q)$. So we see that every morphism in $\underline{\mathrm{Hom}}_S(K, X^q)$ is an isomorphism, and so this complex is a Kan complex. \square

8.2. The ∞ -category of cocartesian fibrations over a base. We consider each cocartesian fibration $q : X \rightarrow S$ as a morphism in sSet^+ by applying the q -cocartesian marking X^q on X and the maximal marking $S^\#$ on S , and we have a the full simplicial subcategory $\underline{\mathrm{Cocart}}(S) \subseteq \mathrm{sSet}_S^+$ of cocartesian fibrations over S . By Proposition 8.5 this simplicial subcategory is enriched in Kan complexes, so that the homotopy coherent nerve is an ∞ -category.

Definition 8.6. Given a simplicial set S , the ∞ -category of cocartesian fibrations over S is the homotopy coherent nerve

$$\mathrm{Cocart}(S) := \mathrm{N}^{\mathrm{hc}}(\underline{\mathrm{Cocart}}(S))$$

Example 8.7. When $S = *$ a cocartesian fibration over $*$ is an ∞ -category \mathcal{C} . The associated cocartesian marking marks equivalences in \mathcal{C} . Since all functors between ∞ -categories preserve equivalences we have

$$\underline{\mathrm{Hom}}_*(\mathcal{C}, \mathcal{D}) = \mathrm{Fun}(\mathcal{C}, \mathcal{D}) \quad \text{and} \quad \underline{\mathrm{Hom}}_{\mathrm{Cocart}(*)}(\mathcal{C}, \mathcal{D}) = \mathrm{Fun}(\mathcal{C}, \mathcal{D})^{\mathrm{Kan}}.$$

Hence $\mathrm{Cocart}(*) = \mathcal{Cat}_\infty$.

Remark 8.8. Of course, we have an $(\infty, 2)$ -category of cocartesian fibrations, which we obtain by applying the homotopy coherent nerve to the simplicial category $\underline{\mathrm{Cocart}}(S)'$. We have no intentions of using this $(\infty, 2)$ -category in this work, and so disregard it.

To state things clearly, objects in $\text{Cocart}(S)$ are cocartesian fibrations $X \rightarrow S$ and morphisms are functors $F : X \rightarrow Y$ which preserve cocartesian maps and fit into a diagram over the base

$$\begin{array}{ccc} X & \xrightarrow{F} & Y \\ & \searrow & \swarrow \\ & S & \end{array} .$$

A 2-simplex $\Delta^2 \rightarrow \text{Cocart}(S)$ is a pair of partial diagrams of cocartesian fibrations

$$\begin{array}{ccc} X_0 & \xrightarrow{F_{01}} X_1 & \xrightarrow{F_{12}} X_2 \\ & \searrow q_0 & \downarrow q_1 \swarrow q_2 \\ & S & \end{array} \quad \begin{array}{ccc} X_0 & \xrightarrow{F_{02}} & X_2 \\ & \searrow q_0 & \swarrow q_2 \\ & S & \end{array}$$

and a map

$$\begin{array}{ccc} \Delta^1 \times X_0 & \xrightarrow{F} & X_2 \\ & \searrow q_0 p_2 & \swarrow q_2 \\ & S & \end{array}$$

which satisfies $F|_0 = F_{12}F_{01}$ and $F|_1 = F_{02}$, and for which $F|_{\Delta^1 \times \{x\}}$ is an isomorphism in the ∞ -category $X_2 \times_S \{s\}$.

8.3. Simplicial enrichment for simplicial functors. For any simplicial set K , and any simplicial functor $F : \underline{A} \rightarrow \underline{\text{sSet}}^+$ we define $K \times F : \underline{A} \rightarrow \underline{\text{sSet}}^+$ to be the composite

$$\underline{A} \xrightarrow{F} \underline{\text{sSet}}^+ \xrightarrow{K \times -} \underline{\text{sSet}}^+.$$

One sees that the action of K is in fact a simplicial functor via the symmetry $K \times \Delta^n \times - \cong \Delta^n \times K \times -$.

For two functors $F, F' : \underline{A} \rightarrow \underline{\text{sSet}}^+$ we let $\text{Nat}(F, F')$ denote the simplicial set with n -simplices

$$\text{Nat}(F, F')[n] := \{\text{Natural transformations } \Delta^n \times F \rightarrow F'\}.$$

The composition operation

$$\text{Nat}(F', F'') \times \text{Nat}(F, F') \rightarrow \text{Nat}(F, F'')$$

takes transformations $\Delta^n \times F' \rightarrow F''$ and $\Delta^n \times F \rightarrow F'$ to the transformation

$$\Delta^n \times F \xrightarrow{\delta \times 1} (\Delta \times \Delta) \times F = \Delta^n \times (\Delta^n \times F) \rightarrow \Delta^n \times F' \rightarrow F''.$$

Definition 8.9. For any simplicial category \underline{A} , we let $\text{Fun}(\underline{A}, \underline{\text{sSet}}^+)$ denote the simplicial category of simplicial functors, with morphism complexes $\text{Nat}(F, F')$.

We have the evaluation functor

$$ev : \text{Fun}(\underline{A}, \underline{\text{sSet}}^+) \times \underline{A} \rightarrow \underline{\text{sSet}}^+.$$

This functor sends a pair (F, a) of a functor and an object in \underline{A} to $F(a)$, and on morphisms the map of simplicial sets

$$\text{Nat}(F, F') \times \underline{\text{Hom}}_{\underline{A}}(a, a') \rightarrow \underline{\text{Hom}}_{\underline{\text{sSet}}}^+(Fa, F'a')$$

sends a pair

$$(\Delta^n \times F \rightarrow F', \sigma : \Delta^n \rightarrow \underline{\text{Hom}}_{\underline{A}}(a, a'))$$

to the composite

$$\Delta^n \times Fa \xrightarrow{\delta \times 1} \Delta^n \times \Delta^n \times Fa \xrightarrow{1 \times F \sigma} \Delta^n \times Fa' \rightarrow F'a'.$$

Lemma 8.10. *There is a natural isomorphism*

$$\begin{aligned} \text{Hom}_{\text{sCat}}(\underline{A}', \text{Fun}(\underline{A}, \underline{\text{sSet}}^+)) &\xrightarrow{\sim} \text{Hom}_{\text{sCat}}(\underline{A}' \times \underline{A}, \underline{\text{sSet}}^+) \\ G &\mapsto \text{ev}(G \times \text{id}_{\underline{A}}). \end{aligned}$$

Proof. The inverse sends a functor $\Theta : A' \times A \rightarrow \underline{\text{sSet}}^+$ to the functor

$$\theta : \underline{A}' \rightarrow \text{Fun}(\underline{A}, \underline{\text{sSet}}^+), \quad \theta(a) = \Theta(a, -)$$

$$\theta_{ab} : \underline{\text{Hom}}_{\underline{A}'}(a, b) \rightarrow \text{Nat}(\Theta(a, -), \Theta(b, -)),$$

where θ_{ab} sends an n -simplex σ to the transformation which evaluates at each x in \underline{A} as

$$\Theta(\sigma, x) : \Delta^n \times \Theta(a, x) \rightarrow \Theta(b, x).$$

□

8.4. Simplicial functors as functor categories. As the category of simplicial categories is cocomplete [4, 00K3], one sees that the path category construction admits a unique extension from the class of simplices to the entire category of simplicial sets.

Lemma 8.11 ([4, 00L4]). *The association $\Delta^n \mapsto \text{Path } \Delta^n$ extends to a functor $\text{Path} : \text{sSet} \rightarrow \text{sCat}$ which provides a left adjoint to the homotopy coherent nerve,*

$$\text{Hom}_{\text{sSet}}(-, \text{N}^{\text{hc}}-) \cong \text{Hom}_{\text{sCat}}(\text{Path } -, -).$$

The product of the unit map $S \rightarrow \text{N}^{\text{hc}} \text{Path } S$, and commutativity of the homotopy coherent nerve with products, provide natural maps

$$\text{Path}(S' \times S) \rightarrow \text{Path}(S') \times \text{Path}(S)$$

from which the path category functor becomes op-lax monoidal. Via this op-lax structure we obtain a map

$$\begin{aligned} \text{Hom}_{\text{sCat}}(\text{Path } \Delta^n, \text{Fun}(\text{Path } S, \underline{\text{sSet}}^+)) &\xrightarrow{\sim} \text{Hom}_{\text{sCat}}(\text{Path } \Delta^n \times \text{Path } S, \underline{\text{sSet}}^+) \\ &\rightarrow \text{Hom}_{\text{sCat}}(\text{Path}(\Delta^n \times S), \underline{\text{sSet}}^+) \cong \text{Hom}_{\text{sSet}}(\Delta^n \times S, \text{N}^{\text{hc}} \underline{\text{sSet}}^+). \end{aligned}$$

Taking these maps collectively across various n provides a map of simplicial sets

$$\text{N}^{\text{hc}} \text{Fun}(\text{Path } S, \underline{\text{sSet}}^+) \rightarrow \text{Fun}(S, \text{N}^{\text{hc}} \underline{\text{sSet}}^+).$$

We restrict to consider those functors which land in the non-full simplicial subcategory of ∞ -categories, with Kanified morphism complexes, and consider the resultant map

$$\text{comp} : \text{N}^{\text{hc}} \text{Fun}(\text{Path } S, \underline{\text{Cat}}_{\infty}^+) \rightarrow \text{Fun}(S, \mathcal{C}at_{\infty}). \quad (18)$$

eq:comp_NN

prop:nerv_to_nerv

Proposition 8.12. *The simplicial category $\text{Fun}(\text{Path } S, \underline{\text{Cat}}_{\infty}^+)$ is enriched in Kan complexes, and the comparison functor (18) is an equivalence of ∞ -categories.*

We outline how this result occurs, according to the logic of [2]. So the proof is not so much a proof as an “authentication ticket” which the reader might verify for themselves.

Proof. The categories $\underline{\mathbf{sSet}}^+$ and $\mathbf{Fun}(\mathbf{Path} S, \underline{\mathbf{sSet}}^+)$ admit combinatorial simplicial model structures under which the subcategories of fibrant-cofibrant objects are precisely the subcategories

$$\underline{\mathbf{Cat}}_\infty^+ \subseteq \underline{\mathbf{sSet}}^+ \quad \text{and} \quad \mathbf{Fun}(\mathbf{Path} S, \underline{\mathbf{Cat}}_\infty^+) \subseteq \mathbf{Fun}(\mathbf{Path} S, \underline{\mathbf{sSet}}^+)$$

[2, Proposition 3.1.3.7, Corollary 3.1.4.4, Proposition A.3.3.2, Remark A.3.3.4]. It follows that the simplicial category $\mathbf{Fun}(\mathbf{Path} S, \underline{\mathbf{Cat}}_\infty^+)$ is enriched in Kan complexes [2, Remark A.3.1.8], and also that $\underline{\mathbf{sSet}}^+$ provides a $\mathbf{Path} S$ -chunk [2, Definition A.3.4.9] of itself [2, Example A.3.4.4]. We now see from [2, Proposition 4.2.4.4] that the comparison functor \square

Remark 8.13. The specific claim of Proposition 8.12 seems not to appear explicitly in [2], though it may be implicit. We found this particular claim in [?, Remark 3.7].

8.5. Non-enriched Straightening and unstraightening. Let $p : K \rightarrow S$ be a cocartesian fibration from a marked simplicial set (K, W) , and consider the pushout S_p of the following diagram

$$\begin{array}{ccc} K & \xrightarrow{i} & \{*\} \star K \\ p \downarrow & & \downarrow \\ S & \xrightarrow{i_p} & S_p. \end{array}$$

We now have the functor of plain categories

$$\mathbf{St}[0] : \underline{\mathbf{sSet}}_{/S}^+[0] \rightarrow \mathbf{Fun}(\mathbf{Path} S, \underline{\mathbf{sSet}}_\infty^+)[0]$$

which sends each object $q : K \rightarrow S$ to the representable functor $\underline{\mathbf{Hom}}_{\mathbf{Path} S_p}(*, -)$, where each value $\underline{\mathbf{Hom}}_{\mathbf{Path} S_p}(*, s)$ is equipped with a natural marking W_s . (Here we are writing s for $i_p(s)$ by an abuse of notation.) The marked edges are all those which appear as follows:

- Take a marked edge $\alpha : x' \rightarrow y'$ in K , and consider the extension to a 2-simplex

$$\mathbf{Ext}(\alpha) := \{*\} \star \alpha : \Delta^2 \rightarrow \{*\} \star K$$

which then has image $\mathbf{Ext}'(\alpha) : \Delta^2 \rightarrow S_p$. This 2-simplex appears as

$$\mathbf{Ext}'(\alpha) : \begin{array}{ccc} & px & \\ \nearrow & & \searrow p\alpha \\ * & \xrightarrow{\quad} & py. \end{array}$$

which then determines a 1-simplex $\mathbf{E}(\alpha) : \Delta^1 \rightarrow \underline{\mathbf{Hom}}_{\mathbf{Path} S_p}(*, py)$.

- Take any 1-simplex $B : \Delta^1 \rightarrow \underline{\mathbf{Hom}}_{\mathbf{Path} S}(py, s)$.
- Consider the composite $B \mathbf{E}(\alpha) : \Delta^1 \rightarrow \underline{\mathbf{Hom}}_{\mathbf{Path} S_p}(*, s)$.

The marked edges in W_s in $\underline{\mathbf{Hom}}_{\mathbf{Path} S_p}(*, s)$ are precisely those edges which appear as above.

Definition 8.14. The functor $\mathbf{St}[0] : \underline{\mathbf{sSet}}_{/S}^+[0] \rightarrow \mathbf{Fun}(\mathbf{Path} S, \underline{\mathbf{sSet}}_\infty^+)[0]$ constructed above is called the non-enriched straightening functor.

It can be shown that the straightening functor is cocontinuous [2, Proposition 3.2.1.4] and hence admits a right adjoint.

Proposition 8.15 ([2, Corollary 3.2.1.5]). *The functor $\text{St}[0]$ admits a right adjoint*

$$\text{Un}[0] : \text{Fun}(\text{Path } S, \underline{\text{sSet}}_\infty^+)[0] \rightarrow \underline{\text{sSet}}_S^+[0].$$

8.6. Enriched unstraightening. At each simplicial set L and map $p : K \rightarrow S$ from a marked simplicial set, we have the new map

$$Lp : L^\# \times K \rightarrow K \rightarrow S$$

and hence a new object in $\underline{\text{sSet}}_S^+$. This product construction endows $\underline{\text{sSet}}_S^+$ with a module category structure over $\underline{\text{sSet}}_S^+$ whose inner-Homs recover the simplicial mapping complexes for the enhancement $\underline{\text{sSet}}_S^+$.

We have the natural map of simplicial sets

$$\text{St}[0](L^\# \times K) \rightarrow L^\# \times \text{St}[0](K)$$

[2, Corollary 3.2.1.15] which endows the straightening functor with a op-lax module category structure. The op-lax structure on $\text{St}[0]$ endow the adjoint $\text{Un}[0]$ with a lax module category structure, and this lax structure provides a canonical enrichment on the unstraightening functor to the simplicial setting. Specifically, we have the natural maps

$$\begin{aligned} \text{Hom}_{\text{sSet}}(L, \underline{\text{Hom}}(F, G)) &= \text{Hom}_{\text{Fun}}(L^\# \times F, G) \rightarrow \text{Hom}_{\underline{\text{sSet}}_S^+}(\text{Un}[0](L^\# \times F) \\ &\rightarrow \text{Hom}_{\underline{\text{sSet}}_S^+}(L^\# \times \text{Un}[0]F, \text{Un}[0]G) = \text{Hom}_{\text{sSet}}(L, \underline{\text{Hom}}(\text{Un}[0]F, \text{Un}[0]G)) \end{aligned}$$

across all simplicial sets L . Via Yoneda this lifts the application of the unstraightening functor on morphism sets to the simplicial level,

$$\text{Un} : \underline{\text{Hom}}_{\text{Fun}(\text{Path } S, \underline{\text{sSet}}^+)}(F, G) \rightarrow \underline{\text{Hom}}_{\underline{\text{sSet}}_S^+}(\text{Un}[0]F, \text{Un}[0]G),$$

and so enriches the non-simplicial unstraightening functor.

Definition 8.16. The unstraightening functor

$$\underline{\text{Un}} : \text{Fun}(\text{Path } S, \underline{\text{sSet}}^+) \rightarrow \underline{\text{sSet}}_S^+$$

is the simplicial enrichment of the non-enriched unstraightening functor, as constructed above.

8.7. The straightening and unstraightening equivalences.

thm:1382

Theorem 8.17. *The unstraightening functor restricts to an equivalence of simplicial categories*

$$\underline{\text{Un}} : \text{Fun}(\text{Path } S, \underline{\text{Cat}}_\infty^+) \rightarrow \underline{\text{Cocart}}(S).$$

Proof. Follows by [2, Lemma 3.2.4.1] and [2, Theorem 3.2.0.1]. \square

In the statement of Theorem 8.17 by an equivalence of simplicial categories we mean an equivalence specifically in the sense of [2, Definition A.3.2.1]. Equivalently, we are saying that the induced functor on homotopy categories is an equivalence and that the maps on morphism complexes are equivalences of Kan complexes. We refer to this latter property as fully faithfulness.

Lemma 8.18. *If $\Theta : \underline{A} \rightarrow \underline{B}$ is an equivalence of Kan-enriched simplicial categories, then the induced functor on homotopy coherent nerves*

$$\text{N}^{\text{hc}} \Theta : \text{N}^{\text{hc}} \underline{A} \rightarrow \text{N}^{\text{hc}} \underline{B}$$

is an equivalence of ∞ -categories.

Proof. Since the induced functor on homotopy categories is an equivalence, $N^{\text{hc}} \Theta$ is essentially surjective. Fully faithfulness follows from fully faithfulness of Θ and the calculation of the (left pinched) mapping spaces in the homotopy coherent nerve via the mapping complexes in the original categories (see Theorem 5.27). \square

We now conclude that unstraightening induces an equivalence on the associated ∞ -categories

$$N^{\text{hc}} \underline{\text{Un}} : \text{Fun}(\text{Path } S, \underline{\text{Cat}}_{\infty}^+) \rightarrow N^{\text{hc}} \underline{\text{Cocart}}(S) = \text{Cocart}(S)$$

is an equivalence of ∞ -categories. Finally, we compose with the inverse comparison equivalence to obtain an equivalence of ∞ -categories

$$\text{Fun}(S, \mathcal{C}at_{\infty}) \xrightarrow{\sim} \text{Cocart}(S).$$

We record this finding.

thm:unstrt_equiv

Theorem 8.19. *The functor*

$$\text{Un} := \text{comp}^{-1} N^{\text{hc}} \text{Un} : \text{Fun}(S, \mathcal{C}at_{\infty}) \xrightarrow{\sim} \text{Cocart}(S).$$

is an equivalence of ∞ -categories.

Definition 8.20. The unstraightening equivalence is the equivalence of Theorem 8.19. The straightening equivalence is the inverse functor

$$\text{St} : \text{Cocart}(S) \xrightarrow{\sim} \text{Fun}(S, \mathcal{C}at_{\infty}).$$

8.8. A remark on uniqueness.

8.9. Recovering transport via straightening. For any map of simplicial sets $f : S \rightarrow S'$, we have the enriched pullback functor

$$f^* : \underline{\text{Cocart}}(S') \rightarrow \underline{\text{Cocart}}(S), \quad (K \rightarrow S') \mapsto (K \times_{S'} S \rightarrow S).$$

lem:1434

Lemma 8.21 ([?, Observation 2.13]). *For any map of simplicial sets $f : S \rightarrow S'$, there is a commutative diagram at the level of homotopy categories*

$$\begin{array}{ccc} \text{hCocart}(S) & \xleftarrow{\text{Un}} & \text{hFun}(S, \mathcal{C}at_{\infty}) \\ f^* \uparrow & & \uparrow f^* \\ \text{hCocart}(S') & \xleftarrow{\text{Un}} & \text{hFun}(S', \mathcal{C}at_{\infty}) \end{array}$$

Proof. The functor $f^* : \text{Fun}(\text{Path } S', \underline{\text{sSet}}^+) \rightarrow \text{Fun}(\text{Path } S, \underline{\text{sSet}}^+)$ has a left adjoint $f_!$ and we have a natural isomorphism $\text{St}_f[0] \cong f_! \text{St}[0]$ [2, Proposition 3.2.1.4], where the functor St_f is as in [2]. This implies and identification of non-enriched functors $\text{Un}_f[0] \cong \text{Un}[0]f^*$. But, by construction, $\text{St}_f[0] = \text{St}[0]f$, where

$$f : \underline{\text{sSet}}_{/S}^+ \rightarrow \underline{\text{sSet}}_{/S'}^+$$

just composes maps over S with f . Now one can see directly that the pullback functor on marked simplicial sets over S' is right adjoint to this composition functor f , so that we have $\text{Un}_f[0] = f^* \text{Un}[0]$. So we have finally $f^* \text{Un}[0] \cong \text{Un}[0]f^*$, and this identification of non-enriched functors implies a corresponding identification of the ∞ -functors at the level of homotopy categories. \square

Remark 8.22. We expect that all of the identifications employed in the proof are compatible with (op)-lax module category structures, so that the identification $f^* \mathrm{Un}[0] \cong \mathrm{Un}[0] f^*$ enriches to an identification at the simplicial level, and hence at the ∞ -level $f^* \mathrm{Un} \cong \mathrm{Un} f^*$. This homotopy-level identification suffices for our purposes however.

We can now consider the universal cocartesian fibration, i.e. the cocartesian fibration over $\mathcal{C}at_\infty$ which is associated to the identity functor

$$\mathcal{Z} := \mathrm{Un}(id_{\mathcal{C}at_\infty}).$$

By the materials of Section 6 we understand that there is an equivalence of fibrations

$$\begin{array}{ccc} \mathcal{Z} & \xrightarrow{\sim} & \mathcal{P}\mathcal{C}at_\infty \\ & \searrow & \swarrow \\ & \mathcal{C}at_\infty & . \end{array}$$

Now the diagram of Lemma 8.21 implies the existence of an equivalence of cocartesian fibrations

$$\mathrm{Un}(F) \cong S \times_{\mathcal{C}at_\infty} \mathcal{Z} =: \int_S F$$

at any functor F in $\mathrm{Fun}(S, \mathcal{C}at_\infty)$.

`prop:str_t_transport`

Proposition 8.23 ([2, §3.3.2]). *For any cocartesian fibration $q : X \rightarrow S$ there is an equivalence of cocartesian fibrations*

$$\begin{array}{ccc} X & \xrightarrow{\sim} & \int_S \mathrm{St}(q) \\ & \searrow q & \swarrow \\ & S & . \end{array}$$

In particular, when the base S is an ∞ -category the straightening functor $\mathrm{St}(q) : S \rightarrow \mathcal{C}_\infty$ is a transport functor for $q : X \rightarrow S$.

Proof. We have $q \cong \mathrm{Un} \mathrm{St}(q) \cong \int_S \mathrm{St}(q)$. □

We can consider now the ∞ -category of left fibrations.

Definition 8.24. Let $\mathrm{LFib}(S)$ denote the full ∞ -subcategory in $\mathrm{Cocart}(S)$ whose objects are precisely the left fibrations $q : X \rightarrow S$.

8.10. Straightening and unstraightening for left fibrations. Since the straightening functor $\mathrm{St}(q)$ produces a transport functor for any cocartesian fibration $q : X \rightarrow S$, and the unstraightening functor pulls back along the universal fibrations, Proposition ?? and the fibration calculation of Proposition 6.6 now imply that unstraightening and straightening restrict to equivalences between left fibrations over S and functors into the ∞ -subcategory of spaces $\mathcal{K}an \subseteq \mathcal{C}at_\infty$.

`cor:str_t_kan`

Corollary 8.25. *The straightening and unstraightening equivalences restrict to inverse equivalences*

$$\mathrm{St} : \mathrm{LFib}(S) \xrightarrow{\sim} \mathrm{Fun}(S, \mathcal{K}an) \quad \text{and} \quad \mathrm{Un} : \mathrm{Fun}(S, \mathcal{K}an) \xrightarrow{\sim} \mathrm{LFib}(S).$$

8.11. Some naturality over the base.

`prop:small_nat`**Proposition 8.26.** *Any diagram of cocartesian fibrations*

$$\begin{array}{ccc} Y & \longrightarrow & X \\ p \downarrow & & \downarrow q \\ T & \xrightarrow{\xi} & S \end{array}$$

determines a uniquely associated transformation $\mathrm{St}(p) \rightarrow \mathrm{St}(q)\xi$ in the ∞ -category $\mathrm{Fun}(T, \mathcal{C}at_\infty)$.

In the case where the bases are ∞ -categories, this transformation is visualized as a diagram

$$\begin{array}{ccc} T & \xrightarrow{\quad} & S \\ \mathrm{St}(p) \searrow & & \swarrow \mathrm{St}(q) \\ & \mathcal{C}at_\infty & \end{array}$$

in the $(\infty, 2)$ -category of (big) ∞ -categories.

Construction. Take $F = \mathrm{St}(q)$, $G = \mathrm{St}(p)$. The composite $F\xi$ is a transport functor for the fibration $q' : X \times_S T \rightarrow T$, which then determines an isomorphism $F\xi \cong \mathrm{St}(q')$ which is unique up to a contractible space of choices, by Theorem 6.14 (or rather its generalized form [4, 02SC]). Additionally the given diagram specifies, and is specified by, a morphism

$$\begin{array}{ccc} Y & \xrightarrow{l} & X \times_S T \\ p \searrow & & \swarrow q' \\ & T & \end{array}$$

in $\mathrm{Cocart}(T)$. Via straightening this morphism determines a map

$$\mathrm{St}(l) : G \rightarrow \mathrm{St}(q') \cong F\xi.$$

□

8.12. Straightening for cartesian fibrations. As with the cocartesian case, one can show that the full simplicial subcategory $\underline{\mathrm{Cart}}(S) \subseteq \underline{\mathrm{sSet}}_S$ consisting of cartesian fibrations over S is enriched in Kan complexes. We can therefore consider the ∞ -category

$$\mathrm{Cart}(S) = \mathrm{N}^{\mathrm{hc}} \underline{\mathrm{Cart}}(S).$$

An application of the opposite functor provides an isomorphism of ∞ -categories $\mathrm{Cart}(S) \cong \mathrm{Cocart}(S^{\mathrm{op}})$. We therefore obtain the following results for cartesian fibrations:

There are mutually inverse equivalences $\mathrm{St} : \mathrm{Cart}(S) \xrightarrow{\sim} \mathrm{Fun}(S^{\mathrm{op}}, \mathcal{C}at_\infty)$ and $\mathrm{Un} : \mathrm{Fun}(S, \mathcal{C}at_\infty) \xrightarrow{\sim} \mathrm{Cart}(S)$ (Theorem 8.19). These equivalences restrict to equivalences

$$\mathrm{St} : \mathrm{RFib}(S) \xrightarrow{\sim} \mathrm{Fun}(S^{\mathrm{op}}, \mathcal{K}an) \quad \text{and} \quad \mathrm{Un} : \mathrm{Fun}(S^{\mathrm{op}}, \mathcal{K}an) \xrightarrow{\sim} \mathrm{RFib}(S)$$

(Corollary 8.25). The value $\mathrm{St}(q) : S^{\mathrm{op}} \rightarrow \mathcal{C}at_{\infty}$ at any cartesian fibration $q : X \rightarrow S$ is a transport functor for q (Proposition 8.23). Finally, any diagram of cocartesian fibrations

$$\begin{array}{ccc} Y & \longrightarrow & X \\ p \downarrow & & \downarrow q \\ T & \xrightarrow{\xi} & S \end{array}$$

determines a uniquely associated transformation between the transport functors $\mathrm{St}(p) \rightarrow \mathrm{St}(q)\xi$ (Proposition 8.26).

9. INITIAL AND TERMINAL OBJECTS

Before beginning with our study in earnest, with the introduction of Hom functors and the Yoneda embedding for ∞ -categories, we discuss the notions of initial and terminal objects in an ∞ -category.

9.1. Initial and terminal basics.

Definition 9.1. Let \mathcal{C} be an ∞ -category. An object x in \mathcal{C} is called initial if, for each object z in \mathcal{C} , the mapping space $\mathrm{Hom}_{\mathcal{C}}(x, z)$ is contractible. An object z in \mathcal{C} is called terminal if, for each object x in \mathcal{C} , the space $\mathrm{Hom}_{\mathcal{C}}(x, z)$ is contractible.

One sees that an object x is initial (resp. terminal) in \mathcal{C} if and only if x is terminal (resp. initial) in the opposite category $\mathcal{C}^{\mathrm{op}}$. So we can freely translate between results for initial versus terminal objects. Note also that we can replace the mapping space $\mathrm{Hom}_{\mathcal{C}}(x, y)$ with either the left or right pinched spaces when evaluating initial-ness or terminal-ness of objects.

lem:init_unique

Lemma 9.2. Let \mathcal{C} be an ∞ -category, and let $\mathcal{C}_{\mathrm{Init}}$ and $\mathcal{C}_{\mathrm{Term}}$ denote the full ∞ -subcategories whose objects are the initial and terminal objects in \mathcal{C} , respectively. Then each of the categories $\mathcal{C}_{\mathrm{Init}}$ and $\mathcal{C}_{\mathrm{Term}}$ is either empty or a contractible Kan complex.

This is to say, the initial (or terminal) object in an ∞ -category \mathcal{C} is unique, provided any such object exists.

Proof. We only consider the case of $\mathcal{C}_{\mathrm{Init}}$. Let us suppose that this subcategory is nonempty. Via contractibility of the mapping spaces we conclude that the functor $\mathcal{C}_{\mathrm{Init}} \rightarrow *$ is fully faithful and essentially surjective, and hence an equivalence of ∞ -categories. So $\mathcal{C}_{\mathrm{Init}}$ is a contractible Kan complex. \square

Lemma 9.3. If x is initial (resp. terminal) in \mathcal{C} , then another object x' is initial (resp. terminal) in \mathcal{C} if and only if x' is isomorphic to x .

Proof. For any isomorphism $\alpha : x \rightarrow x'$ the induced maps

$$\alpha^* : \mathrm{Hom}_{\mathcal{C}}(x, y) \rightarrow \mathrm{Hom}_{\mathcal{C}}(x', y) \quad \text{and} \quad \alpha_* : \mathrm{Hom}_{\mathcal{C}}(y, x) \rightarrow \mathrm{Hom}_{\mathcal{C}}(y, x')$$

are isomorphisms in $\mathbf{h}\mathcal{K}an$, at all y in \mathcal{C} . So contractibility of the left-hand spaces implies contractibility of the right-hand spaces. \square

One also sees that equivalences of ∞ -categories preserve initial and terminal objects.

lem:equiv_initial

Lemma 9.4. *If $F : \mathcal{C} \rightarrow \mathcal{D}$ is an equivalence between ∞ -categories, and x is initial (resp. terminal) in \mathcal{C} , then $F(x)$ is initial (resp. terminal) in \mathcal{D}*

Proof. Suppose that x is initial in \mathcal{C} . First note that any isomorphism $\beta : y \rightarrow y'$ in \mathcal{D} induces isomorphisms

$$\beta_* : \mathrm{Hom}_{\mathcal{D}}(z, y) \xrightarrow{\sim} \mathrm{Hom}_{\mathcal{D}}(z, y')$$

in the homotopy category of Kan complexes. So an object z in \mathcal{D} is initial if and only if the relevant mapping spaces are contractible at a dense collection of objects in \mathcal{D} . (By a dense collection we mean a collection which contains a representative for every isoclass in \mathcal{D} .) Since any equivalence is both fully faithful and essentially surjective, we have that the mapping spaces $\mathrm{Hom}_{\mathcal{D}}(Fx, y)$ are contractible at all y in the image of \mathcal{C} , and hence at all y in \mathcal{D} . So $F(x)$ is initial in \mathcal{D} . The case where x is terminal is proved similarly. \square

Warning 9.5. Initial and terminal objects are not well-behaved under fibering. Consider for example the cone $C = \{x^2 + y^2 = z : x, y, z \in \mathbb{R}\}$ and its projection onto the z -axis line $R_z \cong \mathbb{R}$. The projection $\mathrm{Sing}(C) \rightarrow \mathrm{Sing}(R_z)$ is a Kan fibration and the objects $\vec{1} = (1, 1, 1)$ and 1 are both initial and terminal in $\mathrm{Sing}(C)$ and $\mathrm{Sing}(R_z)$ respectively, since these spaces are contractible. However, $\vec{1}$ is not initial or terminal in the fiber $\mathrm{Sing}(C)_1 = \mathrm{Sing}(S^1)$. In fact, this fiber admits no such objects.

9.2. Aside: trivial fibrations via the fibers. For the analysis that follows, it is convenient to have a characterization of trivial Kan fibrations which can be checked on the fibers.

prop:triv_fibs

Proposition 9.6. *A map of simplicial sets $f : \mathcal{C} \rightarrow S$ is a trivial Kan fibration if and only if f is a left (or right) fibration and, at each point $s : * \rightarrow S$, the fiber \mathcal{C}_s is a contractible Kan complex.*

Sketch proof. If f is a trivial Kan fibration then it is both a left and right fibration, and all of its fibers are contractible. As for the other direction, assume now that f is a left fibration and that all of its fibers are contractible. (The case of a right fibrations is then obtained by taking opposites.)

We must show that each lifting problem of the form

$$\begin{array}{ccc} \partial\Delta^n & \longrightarrow & \mathcal{C} \\ \downarrow & \nearrow & \downarrow \\ \Delta^n & \longrightarrow & S \end{array}$$

admits a solution. In the case that $n = 0$, such a solution exists since the fibers \mathcal{C}_s are all non-empty. So we assume $n > 0$. By replacing S with Δ^n , and \mathcal{C} with $\Delta^n \times_S \mathcal{C}$, we may assume also that both S and \mathcal{C} are ∞ -categories. For fun we can finally replace this lifting problem with the related lifting problem

$$\begin{array}{ccc} \{0\} \times \partial\Delta^n & \xrightarrow{\bar{\sigma}} & \mathcal{C} \\ \downarrow & \nearrow & \downarrow \\ \Delta^1 \times \Delta^n & \xrightarrow{\sigma} & S, \end{array} \tag{19}$$

eq:1722

which is obtained by restricting along the projections

$$p : \Delta^1 \times \Delta^n \rightarrow \Delta^n, \quad p(0, i) = i, \quad p(1, i) = n,$$

and which recovers our original problem after restricting to $\{0\} \times \Delta^n$. It suffices to solve this second problem.

By [4, 0153] the class of left anodyne maps is stable under the cartesian action of \mathbf{sSet} on itself, so that the inclusion $\{0\} \times \partial\Delta^n \rightarrow \Delta^1 \times \partial\Delta^n$ is left anodyne. Since the map f is left anodyne, it follows that the lifting problem (19) extends to a problem

$$\begin{array}{ccc} \Delta^1 \times \partial\Delta^n & \xrightarrow{\bar{\sigma}} & \mathcal{C} \\ \downarrow & \nearrow & \downarrow \\ \Delta^1 \times \Delta^n & \xrightarrow{\sigma} & S, \end{array}$$

Since the fibers of f are trivial Kan fibrations, the above problem extends further to a problem of the form

$$\begin{array}{ccc} Y(0) & \xrightarrow{\bar{\sigma}} & \mathcal{C} \\ \downarrow & \nearrow & \downarrow \\ \Delta^1 \times \Delta^n & \xrightarrow{\sigma} & S, \end{array}$$

where $Y(0)$ is the pushout

$$Y(0) = (\Delta^1 \times \partial\Delta^n) \coprod_{\{1\} \times \partial\Delta^n} (\{1\} \times \Delta^n).$$

By [4, Proof of 00TH] the inclusion $Y(0) \rightarrow \Delta^1 \times \Delta^n$ can be factored into a sequence $Y(0) \rightarrow Y(1) \rightarrow \cdots \rightarrow Y(n+1) = \Delta^1 \times \Delta^n$ with each $Y(i+1)$ fitting into a pushout diagram

$$\begin{array}{ccc} \Delta_{i+1}^{n+1} & \longrightarrow & Y(i) \\ \downarrow & & \downarrow \\ \Delta^{n+1} & \longrightarrow & Y(i+1). \end{array}$$

Furthermore, this sequence can be constructed so that at $Y(n) = \Delta^1 \times \Delta^n$ the sequence

$$\Delta^1 \cong \Delta^{\{n, n+1\}} \rightarrow \Delta^{n+1} \rightarrow \Delta^1 \times \Delta^n$$

recovers the edge $\Delta^1 \times \{n\}$.

Now, since f is inner anodyne we can solve, in order, the lifting problems

$$\begin{array}{ccc} Y(i-1) & \longrightarrow & \mathcal{C} \\ \downarrow & \nearrow & \downarrow \\ Y(i) & \longrightarrow & S \end{array}$$

at all $0 < i \leq n$. For the final lifting problem, along the inclusion $Y(n) \rightarrow Y(n+1)$, we need to solve a lifting problem of the form

$$\begin{array}{ccc} \Lambda_{n+1}^{n+1} & \longrightarrow & \mathcal{C} \\ \downarrow & \nearrow & \downarrow \\ \Delta^{n+1} & \longrightarrow & S \end{array}$$

in which $\sigma|\Delta^{\{n,n+1\}}$ is of a constant value s in S . Since the fiber \mathcal{C}_s is a Kan complex, the morphism $\Delta^{\{n,n+1\}} \rightarrow \Lambda_{n+1}^{n+1} \rightarrow \mathcal{C}$ is an isomorphism in \mathcal{C} . We can therefore solve this final lifting problem, by Proposition I-4.33, and hence obtain the desired solution to the problem (19). \square

When applied to the case of a Kan fibration we have the following, which can also be deduced from Propositions I-3.30 and I-3.42.

Corollary 9.7. *A map between Kan complexes $f : \mathcal{X} \rightarrow \mathcal{Y}$ is a trivial Kan fibration if and only if it is a Kan fibration and, at each point $y : * \rightarrow \mathcal{Y}$, the fiber \mathcal{X}_y is contractible.*

9.3. Initial objects and undercategories.

prop:terminal_over

Proposition 9.8. *An object in an ∞ -category $x : * \rightarrow \mathcal{C}$ is initial if and only if the forgetful functor $\mathcal{C}_{x/} \rightarrow \mathcal{C}$ is a trivial Kan fibration. Dually, an object $y : * \rightarrow \mathcal{C}$ is terminal if and only if the functor $\mathcal{C}_{/y} \rightarrow \mathcal{C}$ is a trivial Kan fibration.*

Proof. If x is initial then all of the left pinched mapping spaces are contractible, so that all of the fibers of the left fibration $\mathcal{C}_{x/} \rightarrow \mathcal{C}$ are contractible. It follows that this map is a trivial Kan fibration. For the converse, we simply note that trivial Kan fibrations are stable under pullback. The arguments in the terminal case are similar. \square

Let us now give a technical lemma.

lem:1699

Lemma 9.9. *For each positive integer n , the map*

$$(\Delta^1 \star \partial \Delta^n) \coprod_{(\{0\} \star \partial \Delta^n)} \{0\} \star \Delta^n \rightarrow \Delta^1 \star \Delta^n \cong \Delta^{n+2}$$

induced by the respective inclusions is an isomorphism onto the horn Λ_0^{n+2} .

See [4] for the proof. We have the following characterization of isomorphisms via initial and terminal objects.

prop:isom_initial

Proposition 9.10. *For a map $\alpha : x \rightarrow y$ in an ∞ -category \mathcal{C} , the following are equivalent:*

- (a) α is an isomorphism.
- (b) α is initial when considered as an object in the undercategory $\mathcal{C}_{x/}$.
- (c) α is terminal when considered as an object in the overcategory.

Proof. We prove the equivalence between (a) and (b). The equivalence between (a) and (c) is obtained by taking opposites. The implication (b) \Rightarrow (a) just follows by considering maps between α and id_x in the undercategory. So suppose that α is an isomorphism. By Proposition 9.8, α is initial in $\mathcal{C}_{x/}$ if and only if the forgetful functor

$$\mathcal{C}_{\alpha/} \cong (\mathcal{C}_{x/})_{\alpha/} \rightarrow \mathcal{C}_{x/}$$

is a trivial Kan fibration. Now, via a consideration of the identification from Lemma 9.9, solving a lifting problem of the form

$$\begin{array}{ccc} \partial\Delta^n & \longrightarrow & \mathcal{C}_{\alpha/} \\ \downarrow & & \downarrow \\ \Delta^n & \longrightarrow & \mathcal{C}/_x \end{array}$$

is equivalent to solving the corresponding lifting problem

$$\begin{array}{ccc} \Lambda_0^{n+2} & \longrightarrow & \mathcal{C} \\ \downarrow & & \downarrow \\ \Delta^{n+2} & \longrightarrow & * \end{array}$$

in which the initial edge $\Delta^{\{0,1\}} \rightarrow \mathcal{C}$ is α . Such a problem admits a solution by Proposition I-4.33, so that the forgetful functor is seen to be a trivial Kan fibration. \square

We take a moment to discuss some examples before returning to the theoretical foundations of this topic.

9.4. Initial and terminal objects in simplicial nerves.

Definition 9.11. An object x in a simplicial category \underline{A} is called initial (resp. terminal) if, for each y in \underline{A} , the mapping complex $\underline{\mathrm{Hom}}_{\underline{A}}(x, y)$ (resp. $\underline{\mathrm{Hom}}_{\underline{A}}(y, x)$) is a contractible Kan complexes.

The easiest way for this to occur is if the relevant mapping complexes are just points. For example, one sees immediately that \emptyset and $*$ are initial and terminal in $\underline{\mathrm{Kan}}$, respectively.

For \underline{A} enriched in Kan complexes, and $\mathcal{A} = \mathrm{N}^{\mathrm{hc}}(\underline{A})$, the equivalence

$$\underline{\mathrm{Hom}}_{\underline{A}}(x, y) \xrightarrow{\sim} \mathrm{Hom}_{\mathcal{A}}^{\mathrm{L}}(x, y)$$

of Theorem 5.27 tells us that an object x is initial (resp. terminal) in \underline{A} if and only if x is initial (resp. terminal) when considered as an object in the ∞ -category \mathcal{A} . The analogous claim is seen to hold for terminal objects via a consideration of the opposite categories.

Lemma 9.12. *Let \underline{A} be a simplicial category which is enriched in Kan complexes. Then an object x is initial (resp. terminal) in \underline{A} if and only if the corresponding object x is initial (resp. terminal) in $\mathrm{N}^{\mathrm{hc}}(\underline{A})$.*

The following corollary is not an immediate consequence of triviality of the mapping categories $\mathrm{Fun}(\emptyset, \mathcal{C})$ and $\mathrm{Fun}(\mathcal{C}, *)$, when \mathcal{C} is an ∞ -category.

cor:1761

Corollary 9.13. *The empty set \emptyset is initial in both $\mathcal{K}\mathrm{an}$ and $\mathcal{C}\mathrm{at}_{\infty}$. The point $*$ is terminal in both $\mathcal{K}\mathrm{an}$ and $\mathcal{C}\mathrm{at}_{\infty}$.*

9.5. Zero objects in pointed spaces. Though we will not use the term explicitly, a zero object in an ∞ -category is an object which is simultaneously initial and terminal. Such objects are familiar from our studies of abelian categories. In the ∞ -setting, the theory of abelian categories is, to some extent and in an indirect manner, reflected in the theory of stable categories. In the stable setting one again demands the existence of a zero object.

Proposition 9.14. *If x is terminal in an ∞ -category \mathcal{C} , then x is both initial and terminal in the category $\mathcal{C}_{x/}$.*

By x in $\mathcal{C}_{x/}$ we mean any morphism $x \rightarrow x$. Since x is terminal, this lift of x to an object in $\mathcal{C}_{x/}$ is uniquely determined up to a contractible space. Practically speaking, we can just take this lift to be $id_x : x \rightarrow x$.

Proof. The fact that x is initial in $\mathcal{C}_{x/}$ follows by Proposition 9.10. For terminality, we consider the forgetful functor

$$\mathcal{C}_{x//x} \rightarrow \mathcal{C},$$

where $\mathcal{C}_{x//x} = (\mathcal{C}_{x/})_{/x} = (\mathcal{C}_{/x})_{x/}$. For any inclusion of simplicial sets $A \rightarrow B$, the existence of a solution to a lifting problem

$$\begin{array}{ccc} A & \longrightarrow & \mathcal{C}_{x//x} \\ \downarrow & & \downarrow \\ B & \longrightarrow & \mathcal{C}_{x/} \end{array}$$

is equivalent to the existence of a solution to the corresponding lifting problem

$$\begin{array}{ccc} \{x\} \star A & \longrightarrow & \mathcal{C}_{/x} \\ \downarrow & & \downarrow \\ \{x\} \star B & \longrightarrow & \mathcal{C}. \end{array}$$

By Proposition 9.8 a solution to the latter problem exists, since x is terminal in \mathcal{C} . It follows that the map $\mathcal{C}_{x//x} \rightarrow \mathcal{C}_{x/}$ is a trivial Kan fibration, and hence that x is terminal in $\mathcal{C}_{x/}$, by Proposition 9.8. \square

Recall from Corollary 9.13 that the 1-point space $*$ is terminal in \mathcal{Kan} .

Corollary 9.15. *The 1-point space $*$ is both initial and terminal in the ∞ -category $\mathcal{Kan}_{*/}$ of pointed Kan complexes.*

9.6. Zero objects in derived categories.

Definition 9.16. An object x in a dg category \mathbf{A} is said to be initial (resp. terminal) if, at each y in \mathbf{A} , the Hom complex $\mathrm{Hom}_{\mathbf{A}}^*(x, y)$ (resp. $\mathrm{Hom}_{\mathbf{A}}^*(y, x)$) is acyclic.

Recall our calculation of the mapping spaces in the dg nerve $\mathcal{A} = \mathrm{N}^{\mathrm{dg}}(\mathbf{A})$ via the Hom complexes in \mathbf{A} ,

$$\mathrm{Hom}_{\mathcal{A}}^L(x, y) \xrightarrow{\sim} K(\mathrm{Hom}_{\mathbf{A}}^*(x, y))$$

(Proposition I-11.7). By Theorem I-10.13, the above calculation tells us that the mapping $\mathrm{Hom}_{\mathcal{A}}^L(x, y)$ are contractible whenever the complex $\mathrm{Hom}_{\mathbf{A}}^*(x, y)$ is acyclic. So we observe the following.

lem:init_dg

Lemma 9.17. *Let \mathbf{A} be a dg category and take $\mathcal{A} = \mathbf{N}^{\mathrm{dg}}(\mathbf{A})$. If an object x is initial (resp. terminal) in \mathbf{A} , then the corresponding object x is initial (resp. terminal) in \mathcal{A} .*

Remark 9.18. The converse to Lemma 9.17 holds if we assume that our dg category \mathbf{A} has a good shift functor (see Section 11.1).

For any abelian category \mathbb{A} , the object 0 is both initial and terminal in the dg category $\mathrm{Ch}(\mathbb{A})$ of cochains over \mathbb{A} , and hence also in the subcategories of K -projective and K -injective complexes. We recall that the derived ∞ -category $\mathcal{D}(\mathbb{A})$ is defined by taking the dg nerve of the dg category of K -injective objects in $\mathrm{Ch}(\mathbb{A})$ when we have enough such objects, or K -projectives when we have enough such objects (see Section I-12).

Corollary 9.19. *For any Grothendieck abelian category \mathbb{A} , the zero complex 0 is both initial and terminal in the derived ∞ -category $\mathcal{D}(\mathbb{A})$.*

9.7. Initial objects and weak contractibility. We phrase all results below in terms of initial objects. The corresponding results hold for terminal objects via duality.

Lemma 9.20. *A Kan complex \mathcal{X} admits an initial object if and only if \mathcal{X} is contractible.*

Proof. If x is initial in \mathcal{X} , then every object in \mathcal{X} admits a morphism from x , and hence is isomorphic to x (since \mathcal{X} is a Kan complex). Since any object which is isomorphic to an initial object is initial, we conclude that \mathcal{X} consists entirely of initial objects. We conclude that \mathcal{X} is contractible by Lemma 9.2. \square

In the case of an ∞ -category \mathcal{C} we do not gain such a precise understanding of \mathcal{C} via the existence of an initial object. This is clear from the examples discussed above. We can, however, constrain certain relative phenomena between ∞ -categories via the preservation of initial objects. The remainder of this section is dedicated to an elaboration on this, somewhat cryptic, point.

lem:1936

Lemma 9.21. *An object x in \mathcal{C} is initial if and only if the forgetful functor $\mathcal{C}_{x/} \rightarrow \mathcal{C}$ admits a section $F : \mathcal{C} \rightarrow \mathcal{C}_{x/}$ with $F(x) = id_x$.*

Proof. If x is initial then the forgetful functor is a trivial Kan fibration, by Proposition 9.8. It follows that the lifting problem

$$\begin{array}{ccc} * & \xrightarrow{id_x} & \mathcal{C}_{x/} \\ \downarrow x & \nearrow & \downarrow \\ \mathcal{C} & \xrightarrow{id} & \mathcal{C} \end{array}$$

admits a solution $s : \mathcal{C} \rightarrow \mathcal{C}_{x/}$. This solution provides the desired section. Conversely, if we have such a section F then for each y in \mathcal{C} we can split the identity on the mapping space as

$$\mathrm{Hom}_{\mathcal{C}}(x, y) \xrightarrow{F} \mathrm{Hom}_{\mathcal{C}_{x/}}(id_x, F(y)) \xrightarrow{forget} \mathrm{Hom}_{\mathcal{C}}(x, y).$$

Since id_x is initial in $\mathcal{C}_{x/}$, by Proposition 9.10, each mapping space $\mathrm{Hom}_{\mathcal{C}_{x/}}(id_x, F(y))$ is contractible. Thus each mapping space $\mathrm{Hom}_{\mathcal{C}}(x, y)$ is a retract of a contractible space, and hence contractible itself. \square

We record a little lemma.

lem:1957

Lemma 9.22 ([4, 0196]). *If $i : A \rightarrow B$ is an inclusion of simplicial sets, then the induced map*

$$\{*\} \star i : \{*\} \star A \rightarrow \{*\} \star B$$

is left anodyne.

Proof. The class of i at which $\{*\} \star i$ is left anodyne is saturated. We need only show that it contains the inclusions $\partial\Delta^n \rightarrow \Delta^n$. But in this case the inclusion in question is identified with the left anodyne map $\Lambda_0^{n+1} \rightarrow \Delta^{n+1}$. \square

prop:init_lanodyne

Proposition 9.23. *An object x in \mathcal{C} is initial if and only if the inclusion $x : * \rightarrow \mathcal{C}$ is left anodyne. If y is terminal in \mathcal{C} , then the inclusion $y : * \rightarrow \mathcal{C}$ is right anodyne.*

Proof. We deal with the initial claim. If $x : * \rightarrow \mathcal{C}$ is left anodyne then we can solve the lifting problem

$$\begin{array}{ccc} * & \xrightarrow{id_x} & \mathcal{C}_{x/} \\ x \downarrow & \nearrow & \downarrow \\ \mathcal{C} & \xrightarrow{id} & \mathcal{C} \end{array}$$

and hence obtain a section $F : \mathcal{C} \rightarrow \mathcal{C}_{x/}$ as in Lemma 9.21. We conclude that x is initial in \mathcal{C} .

Conversely, if x is initial then the section $F : \mathcal{C} \rightarrow \mathcal{C}_{x/}$ of Lemma 9.21 provides a map $F' : \{*\} \star \mathcal{C} \rightarrow \mathcal{C}$ with $F'|_{\mathcal{C}} = id_{\mathcal{C}}$, $F'(*) = x$, and $F'(* \rightarrow x) = id_x$. In particular, F' is defined on each simplex outside of \mathcal{C} by taking

$$F'(\{*\} \star \Delta^m) = F(\Delta^m).$$

This map F' gives a diagram

$$\begin{array}{ccccc} * & \xrightarrow{x} & \{*\} \star \{x\} & \longrightarrow & * \\ x \downarrow & & \downarrow & & \downarrow x \\ \mathcal{C} & \longrightarrow & \{*\} \star \mathcal{C} & \xrightarrow{F} & \mathcal{C} \end{array}$$

so that the inclusion $\{x\} \rightarrow \mathcal{C}$ is a retract of the inclusion $\{*\} \star \{x\} \rightarrow \{*\} \star \mathcal{C}$. Since this latter inclusion is left anodyne, by Lemma 9.22, we conclude that the inclusion $\{x\} \rightarrow \mathcal{C}$ is left anodyne as well. \square

As a consequence of Proposition 9.23 we observe a kind of relative triviality for \mathcal{C} .

cor:initial_eval

Corollary 9.24. *Suppose $f : X \rightarrow S$ is a left fibration, and that \mathcal{C} admits an initial object $x : * \rightarrow \mathcal{C}$. Then the map*

$$\mathrm{Fun}(\mathcal{C}, X) \rightarrow X \times_S \mathrm{Fun}(\mathcal{C}, S), \quad F \mapsto (F|_x, fF). \quad (20)$$

eq:2002

is a trivial Kan fibration.

Proof. Immediate from Propositions 9.23 and 4.3. \square

Let's consider what Corollary 9.24 is telling, in semi-human terms. In the extreme case where $\mathcal{X} \rightarrow *$ is a Kan complex, the right hand side of (20) is just \mathcal{X} and we obtain a trivial Kan fibration

$$ev_x : \text{Fun}(\mathcal{C}, \mathcal{X}) \rightarrow \mathcal{X}$$

which just evaluates a functor F at $x : * \rightarrow \mathcal{C}$. This says that for any choice of a point $z : * \rightarrow \mathcal{X}$ there is a unique functor $F_z : \mathcal{C} \rightarrow \mathcal{X}$ which evaluates as $F_z(x) = z$. Indeed, we can just take the fiber of ev_x at z to obtain a space $\text{Fun}(\mathcal{C}, \mathcal{X})_z$ which parametrizes such functors, and observe that this space is contractible. In this way \mathcal{C} “looks like a point” relative to any Kan complex.

In the relative setting, we consider a left fibration $f : X \rightarrow S$ and see that for any choice of a functor $\bar{F} : \mathcal{C} \rightarrow S$, and a point z in X which lifts $F(x)$, there is a unique lift of \bar{F} to a functor $F : \mathcal{C} \rightarrow X$ with $F(x) = z$. Rather, we observe that any lifting problem of the form

$$\begin{array}{ccc} * & \longrightarrow & X \\ x \downarrow & \nearrow & \downarrow f \\ \mathcal{C} & \longrightarrow & S \end{array}$$

admits a unique solution.

9.8. Equivalences of fibrations via initial objects.

`prop:equiv_initial`

Proposition 9.25. *Let*

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{F} & \mathcal{D} \\ & \searrow & \swarrow \\ & \mathcal{T} & \end{array}$$

be a diagram of ∞ -categories in which both of the maps to \mathcal{T} are left fibrations. If \mathcal{C} admits an initial object x , then F is an equivalence if and only if $F(x)$ is initial in \mathcal{D} .

Proof. If F is an equivalence then it preserves initial objects, by Lemma 9.4. Suppose conversely that F is such a map, that x is initial in \mathcal{C} , and that the image $F(x)$ is initial in \mathcal{D} . Consider the lifting problem

$$\begin{array}{ccc} * & \xrightarrow{x} & \mathcal{C} \\ F(x) \downarrow & \nearrow & \downarrow \\ \mathcal{D} & \longrightarrow & \mathcal{T}. \end{array}$$

By Corollary 9.24 there exists a solution to this problem, and hence there exists a functor $G : \mathcal{D} \rightarrow \mathcal{C}$ over \mathcal{T} with $GF(x) = x$. By Corollary 9.24 again the composition GF is also seen to be isomorphic to the identity in $\text{Fun}_{\mathcal{T}}(\mathcal{C}, \mathcal{C})$. We also have $FG(Fx) = F(x)$ so that FG is isomorphic to the identity on \mathcal{D} . Thus F is an equivalence. \square

Again, one observes a corresponding statement for right fibrations and terminal objects, via the opposite duality.

At first glance this proposition seems ridiculous. Indeed, it suggests that if $f : \mathcal{E} \rightarrow \mathcal{C}$ is a left fibration of ∞ -categories, and e is an object in \mathcal{E} with image

x in \mathcal{C} , then the induced map on undercategories $F : \mathcal{C}_{e/} \rightarrow \mathcal{C}_{x/}$ is an equivalence. This is because F fits into a diagram

$$\begin{array}{ccc} \mathcal{C}_{e/} & \xrightarrow{\quad} & \mathcal{C}_{x/} \\ & \searrow \quad \swarrow & \\ & \mathcal{C} & \end{array}$$

and which both maps to \mathcal{C} are left fibrations, and F is seen to send the initial object id_e to the initial object id_x . However, one sees that this is as bad as it gets.

cor:2491

Corollary 9.26. *Let $\mathcal{C} \rightarrow \mathcal{T}$ be a left fibration, suppose that \mathcal{C} admits an initial object x , and let t denote the image of x in \mathcal{T} . Then there is an equivalence $F : \mathcal{C} \rightarrow \mathcal{T}_{t/}$ of left fibrations over \mathcal{T} which sends x to id_t .*

So Proposition 9.25, said another way, classifies left fibrations up to equivalence via isoclasses of objects in \mathcal{T} .

Proof. By Proposition 9.25 the map $\mathcal{C}_{x/} \rightarrow \mathcal{T}_{t/}$ is an equivalence of left fibrations which sends id_x to id_t . The proposed equivalence $\mathcal{C} \rightarrow \mathcal{T}_{t/}$ is obtained by composing the equivalence $\mathcal{C}_{x/} \rightarrow \mathcal{T}_{t/}$ with a section $F : \mathcal{C} \rightarrow \mathcal{C}_{x/}$ as in Lemma 9.21. \square

10. REPRESENTABLE AND COREPRESENTABLE FUNCTORS

Definition 10.1. Let \mathcal{C} be an ∞ -category. A functor $F : \mathcal{C} \rightarrow \mathcal{K}an$ is corepresented by an object x in \mathcal{C} if F is a transport functor for the left fibration $\mathcal{C}_{x/} \rightarrow \mathcal{C}$. We say F is corepresentable if it is corepresented by some object in \mathcal{C} .

We say a functor $G : \mathcal{C}^{\text{op}} \rightarrow \mathcal{K}an$ is represented by an object y in \mathcal{C} if it is corepresented by y when considered as an object in \mathcal{C}^{op} , i.e. if it is a contravariant transport functor for the right fibration $\mathcal{C}_{/y} \rightarrow \mathcal{C}$. A corepresentable functor is a functor which is corepresented by some object in \mathcal{C} .

We note that if F and F' are corepresented by an object x in \mathcal{C} , then F and F' are isomorphic, via the uniqueness of transport functors. We also see, by Corollary 9.26, that a functor $F : \mathcal{C} \rightarrow \mathcal{K}an$ is representable if the corresponding ∞ -category $\text{Un}(F) \cong \int_{\mathcal{C}} F$ has an initial object. It is clear from Corollary 9.26 that h is representable if, in some sense, it has an “initial object” in some fiber $F(x)$.

Definition 10.2. Given a functor $F : \mathcal{C} \rightarrow \mathcal{K}an$, we say an object $1_x : * \rightarrow F(x)$ is an initial object for F , over x , if at each y in \mathcal{C} the composite

$$\text{Hom}_{\mathcal{C}}(x, y) \xrightarrow{F} \text{Hom}_{\mathcal{K}an}(F(x), F(y)) \xrightarrow{\theta^{-1}} \text{Fun}_{\mathcal{K}an}(F(x), F(y)) \xrightarrow{1_x^*} F(y) \quad (21)$$

eq:initial_fun

is an isomorphism in $\mathbf{h}\mathcal{K}an$.

Note that this condition is really a restriction on the induced functor on enriched categories $\pi F : \pi\mathcal{C} \rightarrow \pi\mathcal{K}an$. Since the isomorphisms θ of Theorem 5.27 is seen to preserve identity morphisms, we see that the above composite at x ,

$$\text{Hom}_{\mathcal{C}}(x, x) \rightarrow F(x)$$

sends the identity $id_x : x \rightarrow x$ to 1_x . Since the θ^{-1} assemble into an equivalence of $\mathbf{H}\mathcal{K}an$ -enriched categories $\pi\mathcal{K}an \xrightarrow{\sim} \pi\mathbf{K}an$ (Proposition 7.6), we also observe at

any choice of $1_x : * \rightarrow F(x)$ a diagram

$$\begin{array}{ccc} F(x) & \xrightarrow{F(\alpha)} & F(y) \\ \uparrow & & \uparrow \\ \mathrm{Hom}_{\mathcal{C}}(x, x) & \xrightarrow{\alpha_*} & \mathrm{Hom}_{\mathcal{C}}(x, y). \end{array}$$

Our first aim is to prove the following.

thm:rep_funs

Theorem 10.3. *A functor $F : \mathcal{C} \rightarrow \mathcal{K}an$ is corepresented by an object x in \mathcal{C} if and only if F admits an initial object which lies over x , $1_x : * \rightarrow F(x)$.*

10.1. Left fibrations with initial transport. Consider a left fibration $q : \mathcal{E} \rightarrow \mathcal{C}$ with transport functor $F : \mathcal{C} \rightarrow \mathcal{K}an$. The induced functor $\pi F : \pi \mathcal{C} \rightarrow \pi \mathcal{K}an$ is determined by parametrized homotopy transport, according to Corollary 7.12. Hence the composite of (21) at any object $1_x : * \rightarrow F(x) \cong \mathcal{E}_x$ is identified with the composite

$$\mathrm{Hom}_{\mathcal{C}}(x, y) \times \{1_x\} \rightarrow \mathrm{Hom}_{\mathcal{C}}(x, y) \times \mathcal{E}_x \rightarrow \mathcal{E}_y, \quad (22)$$

eq:2540

where the final map is given by parametrized homotopy transport (Definition 7.7). More precisely, the adjunction

$$\mathrm{Adj} : \mathrm{Hom}_{\mathrm{Kan}}(\mathrm{Hom}(x, y), \mathrm{Fun}(*, \mathcal{E}_y)) \xrightarrow{\sim} \mathrm{Hom}_{\mathrm{Kan}}(\mathrm{Hom}(x, y) \times \{*\}, \mathcal{E}_y)$$

the map (21) is identified with the map (22). One sees directly that any map $\gamma : \mathrm{Hom}_{\mathcal{C}}(x, y) \rightarrow \mathrm{Fun}(*, \mathcal{E}_y) \cong \mathcal{E}_y$ fits into a diagram

$$\begin{array}{ccc} \mathrm{Hom}_{\mathcal{C}}(x, y) \times \{*\} & \xrightarrow{\mathrm{Adj}(\gamma)} & \mathcal{E}_y \\ \cong \uparrow & & \uparrow \cong \\ \mathrm{Hom}_{\mathcal{C}}(x, y) & \xrightarrow{\gamma} & \mathrm{Fun}(*, \mathcal{E}_y), \end{array}$$

so that γ is an equivalence if and only if $\mathrm{Adj}(\gamma)$ is an equivalence. So we see that an object $1_x : * \rightarrow F(x)$ is initial if and only if the maps (22) are all equivalences.

prop:2653

Proposition 10.4. *Let $F : \mathcal{C} \rightarrow \mathcal{K}an$ be any functor and take $q : \mathcal{E} = \int_{\mathcal{C}} F \rightarrow \mathcal{C}$ the corresponding left fibration. For any choice object x in \mathcal{C} , any object $1_x : * \rightarrow F(x)$, and let \tilde{x} be the image of 1_x in \mathcal{E}_x under the equivalence $\theta : F(x) \rightarrow \mathcal{E}_x$. Then \tilde{x} is initial in \mathcal{E} if and only if 1_x is an initial object for F .*

Proof. Via the equivalence $\mathcal{E}_x \xrightarrow{\sim} \{x\} \tilde{\times}_{\mathcal{C}} \mathcal{C}$ we see that the oriented fiber product admits an initial object. By the specific expression of this equivalence give in Section I-(9.5) one sees that this equivalence sends id_x to id_x , so that the identity specifically is seen to be initial in $\{x\} \tilde{\times}_{\mathcal{C}} \mathcal{C}$.

By Corollary 9.24 there is a unique map of left fibrations $t : \{x\} \tilde{\times}_{\mathcal{C}} \mathcal{C} \rightarrow \mathcal{E}$ which sends id_x to \tilde{x} . This map is an equivalence if and only if all of the induced maps on the fibers

$$t : \mathrm{Hom}_{\mathcal{C}}(x, y) \rightarrow \mathcal{E}_y$$

are isomorphisms.

Since enriched homotopy transport for the oriented fiber product is given by composition on $\text{Hom}_{\mathcal{C}}$ (Proposition ??), the diagrams

$$\begin{array}{ccc} \text{Hom}_{\mathcal{C}}(x, y) \times \mathcal{E}_x & \xrightarrow{\quad} & \mathcal{E}_y \\ \uparrow \text{id} \times t & & \uparrow t \\ \text{Hom}_{\mathcal{C}}(x, y) \times \text{Hom}_{\mathcal{C}}(x, x)^{\circ} & \xrightarrow{\quad} & \text{Hom}_{\mathcal{C}}(x, y) \end{array}$$

commute at all y in \mathcal{C} , where the top map is enriched homotopy transport for \mathcal{E} . Restricting to id_x in the second factor produces a diagram

$$\begin{array}{ccc} \text{Hom}_{\mathcal{C}}(x, y) \times \{\tilde{x}\} & \xrightarrow{\quad} & \mathcal{E}_y \\ \uparrow \text{id} & & \uparrow t_y \\ \text{Hom}_{\mathcal{C}}(x, y) \times \{\text{id}_x\} & \xrightarrow[\text{id}]{} & \text{Hom}_{\mathcal{C}}(x, y) \end{array}$$

from which we see that t_y is an equivalence at all y if and only if the maps (22) are all isomorphisms in $\mathcal{H}an$. Via the equivalence of enriched functors $\pi F \xrightarrow{\sim} q_!$ we observe that these maps are all equivalences if and only if 1_x is an initial object for F . \square

Now, for an arbitrary left fibration $q : \mathcal{E} \rightarrow \mathcal{C}$, with transport functor $F : \mathcal{C} \rightarrow \mathcal{H}an$, we have the equivalence of left fibrations $\mathcal{E} \xrightarrow{\sim} \int_{\mathcal{C}} F$ which is implicit in the assertion that F is a transport functor. We therefore see that \mathcal{E} admits an initial object if and only if $\int_{\mathcal{C}} F$ admits an initial object. So Proposition 10.4 implies the following.

Corollary 10.5. *Let $q : \mathcal{E} \rightarrow \mathcal{C}$ be a left fibration between ∞ -categories. Then \mathcal{E} admits an initial object \tilde{x} over a point x in \mathcal{C} if and only if the corresponding transport functor F admits an initial object $1_x : * \rightarrow F(x)$.*

We now observe the proof of Theorem 10.3.

Proof of Theorem 10.3. By Proposition 10.4 a functor F admits an initial object over x if and only if the corresponding left fibration $\int_{\mathcal{C}} F$ admits an initial object in the fiber over x . This occurs if and only if there is an equivalence of left fibrations

$$\mathcal{C}_{x/} \xrightarrow{\sim} \int_{\mathcal{C}} F$$

over \mathcal{C} . The existence of such an equivalence, by definition, characterizes F as a transport functor for the fibration $\mathcal{C}_{x/} \rightarrow \mathcal{C}$. \square

10.2. Corepresentable functors for simplicial categories. Let \underline{A} be a simplicial category which is enriched in Kan complexes. At any choice of an object x in \underline{A} we have the simplicial functor

$$\underline{\text{Hom}}_{\underline{A}}(x, -) : \underline{A} \rightarrow \underline{\text{Kan}}.$$

Taking homotopy coherent nerves then provides a functor

$$\underline{\text{Hom}}_{\text{N}^{\text{hc}}(\underline{A})}(x, -) : \text{N}^{\text{hc}}(\underline{A}) \rightarrow \mathcal{H}an.$$

prop:simplicial_corep

Proposition 10.6. *Let \underline{A} be a Kan-enriched simplicial category and take $\mathcal{A} = \text{N}^{\text{hc}}(\underline{A})$. At any object x in \mathcal{A} the functor $\underline{\text{Hom}}_{\mathcal{A}}(x, -) : \mathcal{A} \rightarrow \mathcal{H}an$ is corepresented by x .*

Proof. Since the map $\pi \underline{A} \rightarrow \pi \mathcal{A}$ induced by the equivalences of Theorem 5.27 is an equivalence of ∞ -categories, we have an identification of $\mathbf{h}\mathcal{K}an$ -enriched functors

$$\underline{\mathrm{Hom}}_{\mathcal{A}}(x, -) \cong \mathrm{Hom}_{\mathcal{A}}(x, -).$$

Since the functor $\mathrm{Hom}_{\mathcal{A}}(x, -)$ admits an initial object over x , so does $\underline{\mathrm{Hom}}_{\mathcal{A}}(x, -)$. It follows that $\underline{\mathrm{Hom}}_{\mathcal{A}}(x, -)$ is corepresented by x . \square

Corollary 10.7. *For \underline{A} and \mathcal{A} as in Proposition 10.6, a functor $F : \mathcal{A} \rightarrow \mathcal{K}an$ is corepresentable if and only if F admits an isomorphism*

$$\underline{\mathrm{Hom}}_{\mathcal{A}}(x, -) \xrightarrow{\sim} F$$

at some x in \underline{A} .

10.3. Aside: Dold-Kan at the ∞ -level. Let k be a commutative ring and consider the category \mathbf{sSet}_k of simplicial k -modules. This category is symmetric monoidal with the expected product $M \otimes_k N$, where

$$(M \otimes_k N)[r] := M[r] \otimes_k N[r].$$

This product admits inner-Homs, so that \mathbf{sSet}_k is naturally enriched in itself, and we obtain the corresponding simplicial category $\underline{\mathbf{sSet}}_k$. Explicitly the morphism complexes $\mathrm{Fun}_k(M, N)$ are the complexes with n -simplices

$$\mathrm{Fun}_k(M, N)[n] = \mathrm{Hom}_{\mathbf{sSet}_k}(k\Delta^n \otimes_k M, N).$$

Since all simplicial k -modules are Kan complexes, the simplicial category $\underline{\mathbf{sSet}}_k$ is Kan-enriched and we have the forgetful simplicial functor $\underline{\mathbf{sSet}}_k \rightarrow \underline{Kan}$.

Definition 10.8. We take $\mathcal{K}an_k := \mathbf{N}^{\mathrm{hc}}(\underline{\mathbf{sSet}}_k)$.

Via faithfulness of the functor $\underline{\mathbf{sSet}}_k \rightarrow \underline{Kan}$ we observe that the induced functor on ∞ -categories $\mathcal{K}an_k \rightarrow \mathcal{K}an$ is an inclusion of simplicial sets.

Recall that we have the Dold-Kan functor

$$K : \mathrm{Ch}(k) \rightarrow \mathbf{sSet}_k$$

which restricts to an equivalence $K^{\leq 0}$ from connective cochains. In particular, K factors through the truncation

$$\mathrm{Ch}(k) \rightarrow \mathrm{Ch}(k)^{\leq 0}, \quad X \mapsto (\cdots \rightarrow X^{-1} \rightarrow Z^0 X \rightarrow 0)$$

and is identified with the composite of this truncation and the equivalence $K^{\leq 0}$. We let $\mathbf{Ch}(k)$ denote the usual dg category of k -cochains.

Proposition 10.9 ([4, 00SD]). *The functor K admits a lax-monoidal structure $m_{V,W} : K(V) \otimes_k K(W) \rightarrow K(V \otimes_k W)$. Furthermore, at each pair of objects this morphism $m_{V,W}$ is a (non-linear) homotopy equivalence.*

Proof. This lax monoidal structure is adjoint to the monoidal structure on the normalized cochain functor provided by the Alexander-Whitney map [4, 00S6]. Since the Alexander-Whitney maps are quasi-isomorphism [4, 00SB], we conclude that each $m_{V,W}$ is a homotopy equivalence. In particular, $m_{V,W}$ is obtained as the composite

$$K(V) \otimes_k K(W) \xrightarrow{\sim} KN(K(V) \otimes_k K(W)) \xrightarrow{K(\mathrm{AW})} K(NK(V) \otimes_k NK(W)) \xrightarrow{\sim} K(V \otimes_k W).$$

\square

In the case that one of V or W is concentrated in degree 0 one observes natural identifications $K(V) \otimes_k K(W) \cong K(V \otimes_k W)$ and the aforementioned lax monoidal structure extends these identifications to arbitrary complexes. In the case where V is concentrated in degree 0, for example, we have $K(V)[n] = V$ at all n and the aforementioned identification is explicitly the map

$$V \otimes_k \operatorname{Hom}_k^*(N\Delta^n, W) \rightarrow \operatorname{Hom}_k^*(N\Delta^n, V \otimes_k W), \quad v \otimes f \mapsto (x \mapsto v \otimes f(x)).$$

Corollary 10.10. *For any dg category \mathbf{A} there is an associated simplicial category $K\mathbf{A}$ obtained by applying the lax monoidal functor K to the morphism complexes.*

The identifications $K(V)[0] = V^0$ induce an identification of homotopy categories

$$\mathbf{h}N^{\mathbf{hc}}(K\mathbf{A}) = \mathbf{h}K\mathbf{A} = H^0\mathbf{A} = \mathbf{h}N^{\mathbf{dg}}(\mathbf{A}).$$

It's shown in [3, 4] that this equivalence lifts to the ∞ -categorical level.

thm:dk_compare

Theorem 10.11 ([4, 00SV]). *For any dg category \mathbf{A} , there is a natural equivalence of ∞ -categories*

$$\mathfrak{Z}_{\mathbf{A}} : N^{\mathbf{hc}}(K\mathbf{A}) \xrightarrow{\sim} N^{\mathbf{dg}}(\mathbf{A})$$

which is furthermore a trivial Kan fibration and lifts the identification on homotopy categories $\mathbf{h}K\mathbf{A} = H^0\mathbf{A}$.

By naturality, we mean that any dg functor F fits into a diagram

$$\begin{array}{ccc} N^{\mathbf{dg}}(\mathbf{A}) & \xrightarrow{N^{\mathbf{dg}} F} & N^{\mathbf{dg}}(\mathbf{B}) \\ \mathfrak{Z}_{\mathbf{A}} \uparrow & & \uparrow \mathfrak{Z}_{\mathbf{B}} \\ N^{\mathbf{hc}}(K\mathbf{A}) & \xrightarrow{N^{\mathbf{hc}} KF} & N^{\mathbf{hc}}(K\mathbf{B}). \end{array}$$

Now, lax monoidality tells us that the Dold-Kan functor K enriches to a simplicial functor $K\operatorname{Ch}(k)^* \rightarrow \mathbf{sSet}_k$ which just sends a cochain V to the object KV and which is defined on morphisms via the unique map

$$K\operatorname{Hom}_k(V, W) \rightarrow \underline{\operatorname{Hom}}_{\mathbf{sSet}_k}(KV, KW)$$

which is compatible with evaluation.

thm:enriched_dk

Theorem 10.12. *Let $\mathbf{Ch}(k)$ denote the dg category of cochains. If k is a field, the simplicial functor*

$$K : K\mathbf{Ch}(k)^* \rightarrow \mathbf{sSet}_k$$

restricts to an equivalence on $K\mathbf{Ch}(k)^{\leq 0}$.

Proof. In this case a map in $\operatorname{Ch}(k)^{\leq 0}$ is a homotopy equivalence if and only if it is a quasi-isomorphism. It follows via the Dold-Kan equivalence, Theorems I-10.12 & I-10.13 and Proposition I-10.16, that a map in \mathbf{sSet}_k is linear homotopy equivalence if and only if it is a homotopy equivalence. In particular, the comparison map $m_{V,W} : K(V) \otimes_k K(W) \rightarrow K(V \otimes_k W)$ is a linear homotopy equivalence. It follows that K induces a monoidal euivalence on homotopy categories

$$\mathbf{h}K : D(k)^{\leq 0} \xrightarrow{\sim} \mathbf{h}\mathcal{K}an_k.$$

Now, both of the categories $D(k)$ and $\mathbf{h}\mathcal{K}an_k$ admit inner-Homs, which are just given by the inner-Homs at the pre-homotopical level Hom_k^* and $\underline{\operatorname{Hom}}_{\mathbf{sSet}_k}$. Since $\mathbf{h}K$ is an equivalence we now obtain a unique isomorphism of inner-Homs

$$K\operatorname{Hom}_k^*(V, W) \xrightarrow{\sim} \underline{\operatorname{Hom}}_{\mathbf{sSet}_k}(KV, KW)$$

in $h\mathcal{K}an_k$ which is compatible with evaluation. This unique isomorphism is the image of the corresponding map at the pre-homotopical level, from which we conclude that the original morphism is a homotopy equivalence. This implies that K is fully faithful, and essential surjectivity just follows from the fact that the non-enriched Dold-Kan functor is essentially surjective. \square

We refer to the enriched equivalence of Theorem 10.12 as the enriched Dold-Kan equivalence.

cor:infy_dk

Corollary 10.13. *Suppose k is a field. Then enriched Dold-Kan provides a functor between ∞ -categories $K : \mathcal{V}ect_k \rightarrow \mathcal{K}an_k$ which restricts to an equivalence of ∞ -categories*

$$K : \mathcal{V}ect_k^{\leq 0} \xrightarrow{\sim} \mathcal{K}an_k.$$

Here $\mathcal{V}ect_k := N^{\text{dg}}(\mathbf{Ch}(k))$ denotes the ∞ -category of connective cochains, and we've written simply K for the composite of equivalences

$$\mathcal{V}ect_k \xrightarrow{\sim} N^{\text{hc}}(K\mathbf{Ch}(k)) \xrightarrow{N^{\text{hc}}K} \mathcal{K}an_k$$

by an abuse of notation.

10.4. Corepresentable functors for dg categories. Let \mathbf{A} be a dg category, and $\mathcal{A} = N^{\text{dg}}(\mathbf{A})$. At any object x in \mathbf{A} we have the dg functor $\text{Hom}_{\mathbf{A}}^*(x, -) : \mathbf{A} \rightarrow \mathbf{Ch}(k)$.

lem:2851

Lemma 10.14. *Let \mathbf{A} be a dg category. At any object V in \mathbf{A} the simplicial functor*

$$K \circ \text{Hom}_{\mathbf{A}}^*(V, -) : K\mathbf{A} \rightarrow K\mathbf{Ch}(k) \xrightarrow{K} \underline{\text{sSet}}_k$$

is equal to the functor $\underline{\text{Hom}}_{K\mathbf{A}}(V, -) : K\mathbf{A} \rightarrow \underline{\text{sSet}}_k$.

Proof. Take $h_V = \text{Hom}_{\mathbf{A}}^*(V, -)$ and $\underline{h}_V = \underline{\text{Hom}}_{K\mathbf{A}}(V, -)$. On objects these functors are the same. For the composite, the original map

$$h_V : \text{Hom}_{\mathbf{A}}^*(W, W') \rightarrow \text{Hom}_k^*(h_V W, h_V W')$$

fits into, and is specified by, a diagram

$$\begin{array}{ccc} \text{Hom}_{\mathbf{A}}^*(W, W') \otimes_k h_V(W) & & \\ \downarrow h_V \otimes_k id & \searrow \circ & \\ \text{Hom}_k^*(h_V W, h_V W') \otimes_k h_V(W) & \xrightarrow{ev} & h_V(W'). \end{array}$$

Hence Kh_V fits into a diagram

$$\begin{array}{ccc} K \text{Hom}_{\mathbf{A}}^*(W, W') \otimes_k Kh_V(W) & & \\ \downarrow Kh_V \otimes_k id & \searrow K \circ & \\ K \text{Hom}_k^*(h_V W, h_V W') \otimes_k Kh_V(W) & \xrightarrow{ev} & Kh_V(W'). \end{array}$$

Taking inner-Homs for $\underline{\text{sSet}}_k$ now gives a diagram

$$\begin{array}{ccc} K \text{Hom}_{\mathbf{A}}^*(W, W') & & \\ \downarrow Kh_V & \searrow \underline{h}_V & \\ K \text{Hom}_k^*(h_V W, h_V W') & \xrightarrow{K} & \underline{\text{Hom}}_{\underline{\text{sSet}}_k}(Kh_V W, Kh_V W'). \end{array}$$

This implies an equality between these two functors on morphism complexes as well. \square

lem:2889

Lemma 10.15. *Given a diagram in \mathcal{Cat}_∞*

$$\begin{array}{ccc} \mathcal{C} & & \\ \xi \downarrow & \searrow F & \\ \mathcal{D} & \xrightarrow{G} & \mathcal{K}an \end{array}$$

in which ξ is an equivalence, then F is corepresentable by an object x if and only if G is corepresentable by $\xi(x)$.

Proof. We have an isomorphism of functors $F \cong G \circ \xi$. Since representability is stable under isomorphism, we may assume $F = G \circ \xi$. Fix arbitrary points $x, y : * \rightarrow \mathcal{C}$ and take $x' = \xi(x)$, $y' = \xi(y)$.

We consider the corresponding maps on π -enriched categories to observe a diagram

$$\begin{array}{ccccc} \mathrm{Hom}_{\mathcal{C}}(x, y) \times F(x) & \longrightarrow & \mathrm{Hom}_{\mathcal{K}an}(F(x), F(y)) \times F(x) & \xrightarrow{ev} & F(y) \\ \xi \times id \downarrow & & \downarrow = & & \downarrow = \\ \mathrm{Hom}_{\mathcal{D}}(x', y') \times G(\xi x) & \longrightarrow & \mathrm{Hom}_{\mathcal{K}an}(G(x'), G(y')) \times G(x') & \xrightarrow{ev} & G(y') \end{array}$$

in $\mathbf{h}\mathcal{K}an$ in which all vertical maps are isomorphisms. From this diagram we see that F admits an initial object if and only if G admits an initial object. Hence F is representable if and only if G is representable. \square

We understand that, at any dg category, the functor

$$\underline{\mathrm{Hom}}_{K\mathbf{A}}(V, -) : N^{\mathrm{hc}}(K\mathbf{A}) \rightarrow \mathcal{K}an_k \subseteq \mathcal{K}an$$

is corepresented by the given object V . This follows by Proposition 10.6. Naturality of the equivalence \mathfrak{J} from Theorem 10.11, in conjunction with Lemmas 10.14 and 10.15 above, tell us that the Hom-complexes for dg categories also provide representable functors, in the only way that makes sense.

prop:dg_corep

Proposition 10.16. *Let \mathbf{A} be any dg category with associated ∞ -category $\mathcal{A} = N^{\mathrm{dg}}(\mathbf{A})$. At any object V in \mathbf{A} the functor*

$$K \mathrm{Hom}_{\mathbf{A}}^*(V, -) : \mathcal{A} \rightarrow \mathcal{V}ect \rightarrow \mathcal{K}an$$

is corepresented by V .

Corollary 10.17. *Let \mathbf{A} be any dg category with associated ∞ -category $\mathcal{A} = N^{\mathrm{dg}}(\mathbf{A})$. A functor $F : \mathcal{A} \rightarrow \mathcal{K}an$ is corepresented by an object V in \mathcal{A} if and only if F is isomorphic to the functor $K \mathrm{Hom}_{\mathbf{A}}^*(V, -)$.*

10.5. Representable functors for simplicial and dg categories.

Lemma 10.18. *Let \underline{A} be a simplicial category which is enriched in Kan complexes. Then we have $N^{\mathrm{hc}}(\underline{A}^{\mathrm{op}}) = N^{\mathrm{hc}}(\underline{A})^{\mathrm{op}}$.*

Here the opposite category $\underline{A}^{\mathrm{op}}$ is simply the category obtained by applying the symmetry on \mathbf{sSet} to the morphisms. The identification of opposites follows from the identification

$$\mathrm{Fun}_{\mathbf{sCat}}(\mathrm{Path} \Delta^n, \underline{A}^{\mathrm{op}}) = \mathrm{Fun}((\mathrm{Path} \Delta^n)^{\mathrm{op}}, \underline{A}) = \mathrm{Fun}(\mathrm{Path}((\Delta^n)^{\mathrm{op}}), \underline{A})$$

We now consider the functor $\underline{\mathrm{Hom}}_{\underline{A}}(-, y)$, for y in \underline{A} as a functor from the opposite category

$$\underline{\mathrm{Hom}}_{\underline{A}}(-, y) : \underline{A}^{\mathrm{op}} \rightarrow \underline{\mathrm{sSet}}.$$

As a corollary to Proposition 10.6 we observe the following.

Corollary 10.19. *Let \underline{A} be a simplicial category which is enriched in Kan complexes. Then at any object y in \underline{A} , the functor*

$$\underline{\mathrm{Hom}}_{\underline{A}}(-, y) : \mathrm{N}^{\mathrm{hc}}(\underline{A}) \rightarrow \mathcal{K}an$$

is a representable functor which is represented by y .

In the dg setting we also have the opposite category \mathbf{A}^{op} . Here we employ the Koszul sign rule in the symmetric on $\mathrm{Ch}(k)$, so that composition for \mathbf{A}^{op} inherits a sign

$$f \cdot_{\mathrm{op}} g := (-1)^{\deg(f) \deg(g)} gf.$$

One can check the following.

lem:2965

Lemma 10.20. *For any dg category \mathbf{A} , the maps on n -simplices*

$$\mathrm{N}^{dg}(\mathbf{A})^{\mathrm{op}}[n] \rightarrow \mathrm{N}^{dg}(\mathbf{A}^{\mathrm{op}})[n], \quad \{f_I : I \subseteq [n]\} \mapsto \{(-1)^{|I|(|I|-1)/2} f_I : I \subseteq [n]\},$$

define an isomorphism of ∞ -categories $\mathrm{N}^{dg}(\mathbf{A})^{\mathrm{op}} \xrightarrow{\sim} \mathrm{N}^{dg}(\mathbf{A}^{\mathrm{op}})$.

From Proposition 10.16 and Lemma 10.15 we now observe the following.

Proposition 10.21. *For any dg category \mathbf{A} , and any object W in \mathbf{A} , the functor*

$$K \mathrm{Hom}_{\mathbf{A}}^*(-, W) : \mathrm{N}^{dg}(\mathbf{A})^{\mathrm{op}} \rightarrow \mathcal{K}an$$

is a representable functor which is represented by W .

Here of course we have abused notation to view the functor $\mathrm{Hom}_{\mathbf{A}}^*(-, W) : \mathrm{N}^{\mathrm{hc}}(\mathbf{A}^{\mathrm{op}}) \rightarrow \mathcal{V}ect$ as a functor from $\mathrm{N}^{dg}(\mathbf{A})^{\mathrm{op}}$, via the identification of Lemma 10.20.

11. TWISTED ARROWS AND BIFUNCTORIAL HOMS

11.1. The twisted arrows category.

Definition 11.1. Given a simplicial set \mathcal{C} , we define the twisted arrow category $\mathcal{T}w(\mathcal{C})$ as the simplicial set whose n -simplices are

$$\mathcal{T}w(\mathcal{C})[n] := \mathrm{Hom}_{\mathrm{sSet}}((\Delta^n)^{\mathrm{op}} \star \Delta^n, \mathcal{C}).$$

Restricting along the inclusions

$$(\Delta^n)^{\mathrm{op}} \rightarrow (\Delta^n)^{\mathrm{op}} \star \Delta^n \quad \text{and} \quad \Delta^n \rightarrow (\Delta^n)^{\mathrm{op}} \star \Delta^n$$

provides a natural map to the product

$$\lambda : \mathcal{T}w(\mathcal{C}) \rightarrow \mathcal{C}^{\mathrm{op}} \times \mathcal{C}. \tag{23}$$

eq:lambda

To get our heads on straight here, let's observe directly that an object in $\mathcal{T}w(\mathcal{C})$ is a choice of a morphism $\alpha : x \rightarrow y$ in \mathcal{C} . A morphism from an objects $\alpha : x \rightarrow y$ to some other $\alpha' : x' \rightarrow y'$ is a diagram of the form

$$\begin{array}{ccc} x & \xrightarrow{\alpha} & y \\ \uparrow & \searrow & \nearrow \\ & & \\ x' & \xrightarrow{\alpha'} & y' \end{array}$$

If we consider the fiber $\{(x, y)\} \times_{(\mathcal{C}^{\text{op}} \times \mathcal{C})} \mathcal{T}w(\mathcal{C})$, a simplex in this fiber can be visualized as some directed diagram from x to y which is “completely filled in”,

$$x \begin{array}{c} \xrightarrow{\cdots} \\ \xrightarrow{\cdots} \\ \xrightarrow{\cdots} \end{array} y.$$

We prove below that these fibers are a type of bifunctorial Hom space for \mathcal{C} , where bifunctoriality simply refers to the fact that one has two variables to tune in the base.

We note that the join $(\Delta^n)^{\text{op}} \star \Delta^n$ is identified with Δ^{2n+1} via the bijection

$$[2n+1] \rightarrow [n] \amalg [n], \quad i \mapsto \begin{cases} n-i & \text{in the first set if } i \leq n \\ i-n & \text{in the second set if } i \geq n. \end{cases}$$

prop:tw_inner

Proposition 11.2 ([4, 03JT]). *The restriction map $\lambda : \mathcal{T}w(\mathcal{C}) \rightarrow \mathcal{C}^{\text{op}} \times \mathcal{C}$ is a left fibration. More generally, if $\mathcal{C} \rightarrow S$ is an inner fibration of simplicial sets, then the restriction map $\mathcal{T}w(\mathcal{C}) \rightarrow (\mathcal{C}^{\text{op}} \times \mathcal{C}) \times_{(S^{\text{op}} \times S)} \mathcal{T}w(S)$ is a left fibration.*

We only outline the main points of the proof. The reader can find details in the cited text.

Proof outline. We wish to show that any lifting problem of the form

$$\begin{array}{ccc} \Lambda_{n-i}^n & \xrightarrow{\quad} & \mathcal{T}w(\mathcal{C}) \\ \downarrow & \nearrow & \downarrow \\ \Delta^n & \xrightarrow{\quad} & (\mathcal{C}^{\text{op}} \times \mathcal{C}) \times_{(S^{\text{op}} \times S)} \mathcal{T}w(S), \end{array}$$

with $i \leq n$, admits a solution. Such a lifting problem admits a solution if and only if the corresponding problem

$$\begin{array}{ccc} K_0 & \xrightarrow{\quad} & \mathcal{C} \\ \downarrow & \nearrow & \downarrow \\ \Delta^{2n+1} & \xrightarrow{\quad} & S \end{array} \tag{24} \quad \text{eq:1641}$$

admits a solution, where $K_0 \subseteq \Delta^{2n+1}$ is some subcomplex which we describe below.

Given a subset $J \subseteq [2n+1]$, the non-degenerate simplex $\Delta^J \subseteq \Delta^{2n+1}$ lies in K_0 if and only if J is contained in one of $[n]$ or $[2n+1] - [n]$, or J is contained in a subset $[2n+1] - \{j, 2n+1-j\}$ with $j \neq i$. It is argued in [4] that the inclusion $K_0 \rightarrow \Delta^{2n+1}$ is in fact anodyne, by factoring this map into a sequence of inclusions

$$K_0 \rightarrow K_1 \rightarrow \cdots \rightarrow K_m = \Delta^{2n+1}$$

in which each K_{i+1} is obtained from K_i by attaching a single non-degenerate simplex. Each such inclusion $K_i \rightarrow K_{i+1}$ is shown to be anodyne, so that the composition $K_0 \rightarrow \Delta^{2n+1}$ is in fact anodyne, and we find that the problem (24) admits a solution, as desired. \square

Since $\mathcal{C}^{\text{op}} \times \mathcal{C}$ is itself an ∞ -category whenever \mathcal{C} is an ∞ -category, we find that the twisted arrow category $\mathcal{T}w(\mathcal{C})$ is also an ∞ -category in this case.

Corollary 11.3. *If \mathcal{C} is an ∞ -category, the twisted arrow category $\mathcal{T}w(\mathcal{C})$ is also an ∞ -category.*

Via Proposition 11.2, and the general phenomena of transport for left fibrations (Proposition 6.16), we understand that the left fibration $\lambda : \mathcal{T}w(\mathcal{C}) \rightarrow \mathcal{C}^{\text{op}} \times \mathcal{C}$ identifies a associated transport functor

$$H : \mathcal{C}^{\text{op}} \times \mathcal{C} \rightarrow \mathcal{K}an.$$

This transport functor is uniquely determined, up to a contractible space of choices, by the assertion that H fits into a categorical pullback diagram

$$\begin{array}{ccc} \mathcal{T}w(\mathcal{C}) & \longrightarrow & \mathcal{K}an_{*/} \\ \downarrow & & \downarrow \\ \mathcal{C}^{\text{op}} \times \mathcal{C} & \xrightarrow{H} & \mathcal{K}an. \end{array}$$

Definition 11.4. A Hom-functor for an ∞ -category \mathcal{C} is a transport functor

$$H : \mathcal{C}^{\text{op}} \times \mathcal{C} \rightarrow \mathcal{K}an$$

for the left fibration $\lambda : \mathcal{T}w(\mathcal{C}) \rightarrow \mathcal{C}^{\text{op}} \times \mathcal{C}$.

The first aims of this section are to provide a calculation of the fibers of the twisted arrow fibration sufficient conditions which allow us to identify a Hom functor when we see one. Of interest are Hom functors for nerves of dg and simplicial categories (e.g. Hom functors for derived categories).

Let us note, as a bit of foreshadowing, that any Hom functor determines maps into the functor categories

$$H_* : \mathcal{C} \rightarrow \text{Fun}(\mathcal{C}^{\text{op}}, \mathcal{K}an) \quad \text{and} \quad H^* : \mathcal{C}^{\text{op}} \rightarrow \text{Fun}(\mathcal{C}, \mathcal{K}an).$$

We will eventually find that these functors are both fully faithful embeddings.

11.2. Fibers of the twisted arrows fibration. At any points in an ∞ -category $x : * \rightarrow \mathcal{C}$ we can restrict along the projection $(\Delta^n)^{\text{op}} \rightarrow *$ to obtain an inclusion into the twisted arrows category

$$\mathcal{C}_{x/} \rightarrow \{x\} \times_{\mathcal{C}^{\text{op}}} \mathcal{T}w(\mathcal{C}). \tag{25} \quad \text{eq:2199}$$

This map fits into a diagram over \mathcal{C} .

Lemma 11.5 ([4, 03JW]). *At any object $x : * \rightarrow \mathcal{C}^{\text{op}}$, and isomorphism $\alpha : x \rightarrow x'$, the map α is an initial object in the fiber $\{x\} \times_{\mathcal{C}^{\text{op}}} \mathcal{T}w(\mathcal{C})$.*

We note that we can take $x' = x$ and $\alpha = id_x$. In particular, we observe that the fiber of the twisted arrow category over any point in \mathcal{C}^{op} admits an initial object.

Outline of proof. We want to show that the forgetful map

$$(\{x\} \times_{\mathcal{C}^{\text{op}}} \mathcal{T}w(\mathcal{C}))_{\alpha/} \rightarrow \{x\} \times_{\mathcal{C}^{\text{op}}} \mathcal{T}w(\mathcal{C})$$

is a trivial Kan fibration. We note that solving the relevant lifting problem along an inclusion $\partial\Delta^{n-1} \rightarrow \Delta^{n-1}$ is equivalent to extending the boundary of an n -simplex $\bar{\sigma} : \partial\Delta^n \rightarrow \mathcal{T}w(\mathcal{C})$, with $n > 0$ and $\bar{\sigma}|_{\{0\}} = \alpha$, to an n -simplex $\sigma : \Delta^n \rightarrow \mathcal{T}w(\mathcal{C})$. This problem, in turn, is equivalent to solving a lifting problem of the form

$$\begin{array}{ccc} K & \xrightarrow{\tau_0} & \mathcal{C} \\ \downarrow & \nearrow & \downarrow \\ \Delta^{2n+1} & \longrightarrow & * \end{array} \tag{26} \quad \text{eq:2209}$$

with K the subcomplex in Δ^{2n+1} which is the union of the J -simplexes $\Delta^J \rightarrow \Delta^{2n+1}$, where $J \subseteq [2n+1]$ is any subset which is either contained in $[n]$ or $[2n+1] - [n]$ or $[2n+1] - \{i, 2n+1-i\}$ for some i . The assumption that $\bar{\sigma}$ lands in the fiber $\{x\} \times_{\mathcal{C}^{\text{op}}} \mathcal{T}w(\mathcal{C})$ forces $\tau_0|_{\Delta^n}$ to be of constant value x , and the assumption that $\bar{\sigma}|_{\{0\}} = \alpha$ forces $\tau_0|_{\Delta^{\{n,n+1\}}} = \alpha$

Now, one argues that there is a factoring of the inclusion $K \rightarrow \Delta^{2n+1}$ into a sequence of inclusions

$$K = K_0 \rightarrow K_1 \rightarrow \cdots \rightarrow K_m = \Delta^{2n+1}$$

with each K_{i+1} fitting into a pushout square

$$\begin{array}{ccc} \Lambda_{k_i}^{d_i} & \longrightarrow & K_i \\ \downarrow & & \downarrow \\ \Delta^{d_i} & \longrightarrow & K_{i+1} \end{array}$$

with each $k_i < d_i$, or $k_i = 0$, $d_i > 1$, and $\Delta^{\{0,1\}} \rightarrow \Lambda_0^{d_i} \rightarrow \Delta^{2n+1}$ landing in the 1-skeleton $\text{Sk}_1 \Delta^{n+1} \subseteq K$. This final condition implies that, in the case $k_i = 0$, and map $\tau_i : K_i \rightarrow \mathcal{C}$ extending $\tau_0 : K \rightarrow \mathcal{C}$ sends the initial vertex

$$\Delta^{\{0,1\}} \rightarrow \Lambda_0^{d_i} \rightarrow K_i \xrightarrow{\tau_i} \mathcal{C}$$

to an isomorphism in \mathcal{C} .

Using the above information, and Proposition I-4.33, we can produce sequential solutions to the lifting problems

$$\begin{array}{ccc} K_i & \xrightarrow{\tau_i} & \mathcal{C} \\ \downarrow & \nearrow \tau_{i+1} & \downarrow \\ K_{i+1} & \longrightarrow & * \end{array}$$

in order to produce the desired solution $\tau = \tau_m : \Delta^{2n+1} \rightarrow \mathcal{C}$ to the problem (26). \square

prop:2273

Proposition 11.6. *The maps (25) define equivalences of left and right fibrations*

$$\begin{array}{ccc} \mathcal{C}_x/ & \xrightarrow{\sim} & \{x\} \times_{\mathcal{C}^{\text{op}}} \mathcal{T}w(\mathcal{C}) \\ & \searrow & \swarrow \\ & \mathcal{C} & \end{array} .$$

Proof. The map $F : \mathcal{C}_x/ \rightarrow \{x\} \times_{\mathcal{C}^{\text{op}}} \mathcal{T}w(\mathcal{C})$ preserves the initial object id_x , and so is an equivalence of ∞ -categories by Proposition 9.25. It follows that F is an equivalence of cocartesian fibrations, by Proposition 3.1. \square

We recall that the fibers of any equivalence of isofibrations are again equivalences (Corollary I-5.24). So Proposition 11.6 tells us that the fibers of the twisted arrows category $\mathcal{T}w(\mathcal{C})$ are identified with the mapping spaces for \mathcal{C} .

cor:tw_fibers

Corollary 11.7. *Let \mathcal{C} be an ∞ -category. At any pair of points $x, y : * \rightarrow \mathcal{C}$ the natural map*

$$\text{Hom}_{\mathcal{C}}^L(x, y) = \mathcal{C}_x/ \times_{\mathcal{C}} \{y\} \rightarrow \{x\} \times_{\mathcal{C}^{\text{op}}} \mathcal{T}w(\mathcal{C}) \times_{\mathcal{C}} \{y\}$$

is an equivalence of Kan complexes.

Corollary 11.8. *For any Hom functor $H : \mathcal{C}^{\text{op}} \times \mathcal{C} \rightarrow \mathcal{K}an$, i.e. classifying functor for the twisted arrows fibration, the restriction*

$$\mathcal{C} \xrightarrow{x \times \text{id}} \mathcal{C} \times \mathcal{C} \xrightarrow{H} \mathcal{K}an$$

*at any point $x : * \rightarrow \mathcal{C}$ is a classifying functor for the forgetful functor $\mathcal{C}_{x/} \rightarrow \mathcal{C}$.*

11.3. Recognition for Hom functors.

12. HOM FUNCTORS FOR DG CATEGORIES

REFERENCES

- | | |
|--|--|
| <div style="border: 1px solid black; padding: 2px; display: inline-block;">joyal02</div>
<div style="border: 1px solid black; padding: 2px; display: inline-block;">htt</div>

<div style="border: 1px solid black; padding: 2px; display: inline-block;">ha</div>
<div style="border: 1px solid black; padding: 2px; display: inline-block;">kerodon</div>
<div style="border: 1px solid black; padding: 2px; display: inline-block;">stacks</div> | <ul style="list-style-type: none"> [1] A. Joyal. Quasi-categories and kan complexes. <i>J. Pure Appl. Algebra</i>, 175(1-3):207–222, 2002. [2] J. Lurie. <i>Higher topos theory</i>. Number 170. Princeton University Press, Princeton, New Jersey 08540, 2009. [3] J. Lurie. Higher algebra. www.math.ias.edu/~lurie/papers/HA.pdf, 2017. [4] J. Lurie. Kerodon. https://kerodon.net, 2024. [5] S. project authors. Stacks project. stacks.math.columbia.edu. |
|--|--|

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SOUTHERN CALIFORNIA, LOS ANGELES, CA 90007

Email address: `cnegron@usc.edu`