I ideals

~ I. Primes and maximal ideals

Throughout we let R denote a <u>commutative</u> ring.

Def$^{\underline{n}}$: An ideal $I \subseteq R$ is called prime if I is properly contained in R, and for each $a, b \in R$, $ab \in I \Leftrightarrow a \in I$ or $b \in I$.

We call I maximal if $I \subsetneq R$ and any intermediate ideal $I \subseteq I' \subseteq R$ we have either $I' = I$ or $I' = R$.

Lemma 1: i) An ideal $I \subseteq R$ is prime if and only if R/I is a (nonzero) domain.

ii) An ideal $I \subseteq R$ is maximal if and only if R/I is a (nonzero) field.

Proof: (i) Consider $\bar{a}, \bar{b} \in R/I$ with reps $a, b \in R$. Then $\bar{a} \cdot \bar{b} = 0$ if and only if $a \cdot b \in I$. Hence, if I is prime we have one of $\bar{a}$ or $\bar{b} = 0$. Conversely, if R/I is a domain then $\bar{a} \cdot \bar{b} = 0$ implies one of $\bar{a} = 0$ or $\bar{b} = 0$, so that for arbitrary elements $a \cdot b \in I$ implies $a \in I$

or $b \in I$. So I is prime.

(ii) If I is maximal R/I has no proper nonzero ideals [Lemma 11, Sample]. Hence for any nonzero $\bar{a} \in R/I$ $(\bar{a}) = R$ and hence there exists $\bar{b}$ such that $\bar{a} \cdot \bar{b} = 1$. So we see R/I is a field. Conversely, if R/I is a field then the quotient contains no proper nonzero ideals, and thus any intermediate ideal $I \subseteq I' \subseteq R$ has $I' = I$ or $I' = R$ [Lemma 11, Sample]. ∎

Corollary 2: Any maximal ideal $\mathfrak{m} \subseteq R$ is prime.

Proposition 3: Any proper ideal $I \subseteq R$ is contained in a maximal ideal $I \subseteq \mathfrak{m}$.

Proof: Zorn's Lemma. ∎

Example: For $(x) \subseteq k[x,y]$, we have the map $\phi: k[x,y] \longrightarrow k[y]$, $\phi(f(x,y)) = f(0,y)$, has kernel

$$\ker(\phi) = \{f(x,y) : x \text{ divides } f(x,y)\} = (x).$$

Hence, by the universal property of the quotient ring, we obtain an algebra isomorphism $k[x,y]/(x) \cong k[y]$.

Since $k[x]$ is a domain (x) is prime.

Example: Let $f(x_1, \dots, x_n) \in k[x_1, \dots, x_n]$ be irreducible. For $a, b \in k[x_1, \dots, x_n]$ we have

$$a \cdot b \in (f) \iff f \mid a \cdot b.$$

If one of a or b is a unit then either $f \mid a$ or $f \mid b$. If neither is a unit then we have unique decomps into products of irreducibles

$$a = p_1 \cdots p_s, \quad b = q_1 \cdots q_t, \quad a \cdot b = p_1 \cdots p_s q_1 \cdots q_t$$

and, via uniqueness, we have $u \cdot f = p_i$ or $u \cdot f = q_j$ for some unit u, giving either $f \mid a$ or $f \mid b$. Hence $a \in (f)$ or $b \in (f)$. We conclude that (f) is a prime ideal.

Proposition 4: For any UFD R, and nonzero $f \in R$, the ideal (f) is prime if and only if f is irreducible.

Proof: We've already agreed in the example above that (f) is prime when f is irreducible. Conversely, if (f) is prime then under any factoring $f = a_1 \cdot a_2$ we have $a_1 \cdot a_2 \in (f)$ and either $f \mid a_1$ or $f \mid a_2$. Supposing arbitrarily $f \mid a_1$, then $a_1 = f \cdot u$ giving $f = (a_2 \cdot u) \cdot f \implies$

$a_2 \cdot x = 1$, since R is a domain. So a_2 is a unit.
We conclude that f is irreducible. ∎

Example: Consider the poly $y^2 - x^3$ in $\mathbb{C}[x,y]$.

$y = \sqrt{x^3}$



Supposing $f_1(x,y) \cdot f_2(x,y) = y^2 - x^3$ with neither f_i a unit, we have

$$f_i(x,y) = c_i x^{a_i} + d_i y^{b_i}$$

due to the absence of mixed terms in $y^2 - x^3$, giving

$$f_1 \cdot f_2 = c_1 c_2 x^{a_1 + a_2} + d_1 d_2 y^{b_1 + b_2} + c_1 d_2 x^{a_1} y^{b_2} + c_2 d_1 x^{a_2} y^{b_1}.$$

$\Rightarrow$ $c_1 d_2 = c_2 d_1 = 0$ $\Rightarrow$ one of c_1 or $c_2 = 0$

$\Rightarrow$ $y^2 - x^3 = d_1 d_2 y^{b_1 + b_2}$, a contradiction.

So we see $y^2 - x^3$ is an irreducible polynomial in $\mathbb{C}[x,y]$, and $(y^2 - x^3)$ is prime in $\mathbb{C}[x,y]$, and

$$\mathbb{C}[x,y] / (y^2 - x^3) \text{ is a domain!}$$

Example: For $y = f(x)$



we can argue as in the previous example to find that $y - f(x)$ is irreducible. Hence $(y - f(x))$ is prime. Alternatively, we have the algebra map

$$\varphi : k[x,y] \to k[x], \quad \varphi(x) = x, \quad \varphi(y) = f(x)$$

with $\ker(\varphi) = (y - f(x))$. Let argue this point. We have the inclusion $k[x] \to k[x,y]$ for which composite

$$\begin{tikzcd} k[x] \arrow[r] \arrow[rr, bend left, "id"] & k[x,y] \arrow[r, "\varphi"'] & k[x] \end{tikzcd}$$

$$\Rightarrow \quad \begin{tikzcd} k[x] \arrow[r] \arrow[rr, bend left, "id"] & k[x,y]/(y - f(x)) \arrow[r, "\bar{\varphi}"'] & k[x] \end{tikzcd}$$

so that the induced map from the quotient $\bar{\varphi}$, which is obviously surjective, if and only if $\bar{\varphi}$ is injective. For this it suffices to show that all $\overline{g(x,y)} = \overline{h(x)}$ in the quotient, since in this case for some poly $h \in k[x]$

$$\bar{\varphi}(\bar{g}) = \bar{\varphi}(\overline{h(x)}) = h(x) \Rightarrow$$

$$\bar{\varphi}(\bar{g}) = 0 \text{ implies } \overline{h(x)} = 0 \text{ implies } \bar{g} = 0.$$

But now this is clear.

Indeed, in the quotient $\bar{y} = \overline{f(x)}$, giving $\bar{y}^m = \overline{f(x)}^m$ for all $m \geq 0$, giving finally

$$\overline{g(x,y)} = \sum_i \overline{g_i(y)}\,\bar{x}^i$$

$$= \sum_i \overline{g_i(f(x))}\,\bar{x}^i = h(x).$$

So we see

$\bar{\varphi}$ injective, and thus $\ker(\varphi) = (y - f(x))$.

Example: For any field k and $\alpha, \beta \in k$, $(x-\alpha, y-\beta)$ is a maximal ideal in $k[x,y]$.

Example: $(x^2 - y^2)$ is <u>not</u> prime, as $(x-y)\cdot(x+y) = x^2 - y^2$ while $x-y \notin (x^2-y^2)$ and $x+y \notin (x^2-y^2)$.



Example: (x^3, y) is <u>not</u> prime in $k[x,y]$.

Question: Is every non-maximal prime ideal $I \subseteq k[x,y]$ principle? I.e. of the form $I = (f)$ for some poly $f(x,y)$?

Question: Does there exist a principle ideal in $k[x,y]$ which is maximal?

Question: Are there maximal ideals in $k[x,y]$ other than those of the form $(x-\alpha, y-\beta)$?

II. Zariski's Lemma

Theorem (Zariski's Lemma): For any field k, and any maximal ideal $\mathfrak{m}$ in $k[x_1,\dots,x_n]$, the quotient $k[x_1,\dots,x_n]/\mathfrak{m}$ is a finite field extension of k.

II ½ Proving Zariski: pt 1: Rational functions

For any field k, we let $k(x_1,\dots,x_n)$ denote the field of "rational functions" over k. Elements in $k(x_1,\dots,x_n)$ are equiv classes of fractions f/g where $g \neq 0$ and $f_1/g_1 = f_2/g_2 \iff f_1 g_2 = f_2 g_1$; addition

$$f_1/g_1 + f_2/g_2 = (f_1 g_2 + f_2 g_1)/g_1 \cdot g_2$$

and mult.

$$(f_1/g_1)\cdot(f_2/g_2) = (f_1 \cdot f_2)/(g_1 \cdot g_2).$$

Lemma 5: For any field k, $k(x_1,\dots,x_n)$ is a well-defined ring. All nonzero elements in $k(x_1,\dots,x_n)$ are invertible, and the apparent map

$$k[x_1,\dots,x_n] \to k(x_1,\dots,x_n), \quad f \mapsto f/1,$$

is an injective ring homomorphism.

Proof: Exercise. ∎

Lemma 6: Take $R = k[x_1, \dots, x_n]$ and $K = k(x_1, \dots, x_n)$, and let $k \to F$ be any field extension. Restriction along the inclusion $R \to K$ induces a bijection

$$\mathrm{Hom}_{k\text{-alg}}(K, F) \to \{\text{Injective } k\text{-alg maps } R \to F\}$$

Proof: For $\varphi : K \to F$ and $\bar{\varphi} : R \to F$ given by restriction, we have at any nonzero poly f

$$1 = \varphi(1) = \varphi(f \cdot f^{-1}) = \bar{\varphi}(f) \cdot \varphi(f^{-1}) \Rightarrow \bar{\varphi}(f) \in F - \{0\}$$

so that $\bar{\varphi}$ has trivial kernel, and

$$\varphi(f/g) = \varphi(f \cdot g^{-1}) = \bar{\varphi}(f)\, \varphi(g^{-1}) = \bar{\varphi}(f)\, \bar{\varphi}(g)^{-1}$$

so that φ is determined uniquely by its restriction $\bar{\varphi}$. So we see that (*) is a well-defined injection.

For surjectivity, take any inj alg map $\psi : R \to F$. We propose a map

$$\tilde{\psi} : K \to F \text{ defined by}$$

$$\tilde{\psi}(f/g) := \psi(f) \cdot \psi(g)^{-1}.$$

When $f_1/g_1 = f_2/g_2$ we have $f_1 g_2 = f_2 g_1$ giving

$$\psi(f_1)\, \psi(g_2) = \psi(f_2)\, \psi(g_1)$$

$$\Rightarrow \psi(f_1) \cdot \psi(g_1)^{-1} = \psi(f_2) \cdot \psi(g_2)^{-1}.$$

So $\tilde{\varphi}$ is well-def as a map of sets. For addition and k-linearity

$$\tilde{\varphi}((f_1 g_2 + f_2 g_1)/g_1 g_2) = (\varphi(f_1 g_2) + \varphi(f_2 g_1)) \cdot \varphi(g_1 g_2)^{-1}$$

$$= (\varphi(f_1)\varphi(g_2) + \varphi(f_2)\varphi(g_1)) \cdot \varphi(g_1)^{-1} \cdot \varphi(g_2)^{-1}$$

$$= \varphi(f_1)\varphi(g_1)^{-1} + \varphi(f_2) \cdot \varphi(g_2)^{-1}$$

$$= \tilde{\varphi}(f_1/g_1) + \tilde{\varphi}(f_2/g_2),$$

and for any $c \in k$

$$\tilde{\varphi}(c \cdot f/g) = \varphi(c \cdot f) \cdot \varphi(g)^{-1}$$

$$= c \cdot \varphi(f)\varphi(g)^{-1} = c\,\tilde{\varphi}(f/g).$$

Considering the cases $c = \pm 1$ we see $\tilde{\varphi}$ is additive group map, and hence also a k-module map.

For the product, finally, we have

$$\tilde{\varphi}(f_1 f_2/g_1 g_2) = \varphi(f_1)\varphi(f_2)\varphi(g_1)^{-1}\varphi(g_2)^{-1}$$

$$= \varphi(f_1)\varphi(g_1)^{-1}\varphi(f_2)\varphi(g_2)^{-1}$$

$$= \tilde{\varphi}(f_1/g_1) \cdot \tilde{\varphi}(f_2/g_2). \qquad \blacksquare$$

Def$^{\underline{n}}$: For a field extension $k \to F$, we say elements $a_1, \dots, a_n$ are algebraically independent over k if the induced k-alg map

$$\phi_a : k[x_1, \dots, x_n] \longrightarrow F, \quad x_i \mapsto a_i,$$

is injective.

By Lemma 6 and freeness of the poly ring [Thm 6, Polyr] we see that there is a bijection:

$$\mathrm{Hom}_{k\text{-alg}}(k[x_1,\dots,x_n], F) \xrightarrow{\sim} \{n\text{-tuples of alg. indep. elem in } F\}$$

$$\phi \longmapsto (\phi(x_1),\dots,\phi(x_n)).$$

~ II. 2/2 Proving Zariski pt 2: Some lemmas

Lemma 7: Let k be a Noeth. comm. ring and $k \to R \xrightarrow{\phi} S$ be a sequence of comm ring maps for which ϕ is injective and the following hold:

(a) S is of finite type over k (fin. gen'd k-alg).

(b) S is finitely gen'd as an R-module.

Then R is of finite type over k.

Proof: Let $x_1,\dots,x_n$ be generators for S, as a k-algebra, and $y_1,\dots,y_m$ be gen for S as a R-module. Then we have

$$x_i = \sum_{j=1}^{m} a_{ij} y_j, \quad y_r y_s = \sum_t b_{rs}^t y_t \quad \text{for } a_{ij}, b_{rs}^t \in R.$$

Consider the subalgebra $R' = k[a_{ij}, b_{rs}^t : i,j,r,s,t] \subseteq R$.

Since k is Noetherian R' is a Noetherian ring, and we have that S is a finite module over R'.

It follows by Noetherianity that R is a finite module over R', and hence that R is a finite type alg over k. ∎

Lemma 8: Any PID R has infinitely many distinct irreducible elements.

To be more clear, for $\mathrm{Irred}(R)$ the collection of irreducibles, and $R^\times$ acting on $\mathrm{Irred}(R)$ by scaling, the set $\mathrm{Irred}(R)/R^\times$ is infinite.

Proof: For $p_1, \dots, p_n$ any finite collection of irreducibles and $p = u p_i$ for some $i \in \{1, \dots, n\}$ and unit u, if $p \mid (p_1 \cdots p_n - 1)$ then

$$p \cdot a = (ua) \cdot p_i = p_1 \cdots p_n - 1$$

$$\Rightarrow p_i (ua - p_1 \cdots p_{i-1} p_{i+1} \cdots p_n) = 1$$

$$\Rightarrow p_i \text{ is a unit, a contradiction.}$$

Hence there is an irreducible p which is not an associate to any of the p_i. ∎

~ II 3/2 Proof of Zariski

Recall the statement: For any maximal ideal $m \subseteq k[x_1, \dots, x_n]$, the quotient

$$k(m) = k[x_1, \dots, x_n]/m$$

is a finite field extension of k

$$\begin{tikzcd} & k[x_1,\dots,x_m] \arrow[dr] & \\ k \arrow[ur, dashed] \arrow[rr] & & k(m) \end{tikzcd}$$

Proof: This is equivalent to the claim that all elem in $k(m)$ are algebraic. Suppose this is not the case, and (after permutation) we assume that

$$k[x_1,\dots,x_m] \to k(m) =: K$$

is injective, giving a field injection $k(x_1,\dots,x_m) \to K$ by Lemma 6, and that all remaining

$$\alpha_i = \text{image } x_i \in K$$

are algebraic over the subfield $k(x_1,\dots,x_m) \subseteq K$. Then K is finite over $k(x_1,\dots,x_m)$, and since K is of finite type over k, $k(x_1,\dots,x_m)$ is of finite type over k by Lemma 7.

Take finite generators

$$\xi_1,\dots,\xi_t \in k(x_1,\dots,x_m), \quad \xi_i = f_i/g_i,$$

and let p be irreducible in $k[x_1,\dots,x_m]$ with $p \nmid g_i$ for all i. Such p exists by Lemma 8. Since $p^{-1} \in k(x_1,\dots,x_m)$ we have now

$$p^{-1} = \sum_{\vec{i}} c_{\vec{i}} \xi^{\vec{i}} = f(x_1,\dots,x_m) / g(x_1,\dots,x_m)$$

for $g(x_1,\dots,x_m)$ a product of the g_i, which gives

$$p \cdot f/g = 1 \Rightarrow p \cdot f = g \Rightarrow p \mid g \Rightarrow p \mid g_i,$$

a contradiction. Thus, K is algebraic and of fin.

type over k, giving K finite dim over k. ∎

As an immediate corollary we have the following.

Theorem (Alt. Zariski): For any field k, any finite type k-algebra R, and any maximal ideal $m \in R$, the quotient $k(m)$ is a finite field extension of k.

~ III. Zariski and algebraic closure

For any finite field extension K/k, we have finitely many $\alpha_1, \dots, \alpha_n$ in $\bar{k}$ for which

$$K \cong k(\alpha_1, \dots, \alpha_n) \subseteq \bar{k},$$

as a k-algebra.

Corollary 9: Every maximal ideal $m \subseteq k[x_1, \dots, x_n]$ appears as the kernel of an algebra map

$$\phi_\alpha : k[x_1, \dots, x_n] \to \bar{k}, \quad x_i \mapsto \alpha_i .$$

More generally for any max ideal $m \subseteq R$ in a finite type k-alg, m appears as the kernel of an algebra map

$$\phi : R \to \bar{k}.$$

Proof: The case of $R = k[x_1, \dots, x_n]$ is clear, by the above arguments. For general R we have a surj. alg map $\pi : k[x_1, \dots, x_n] \to R$ and for $\tilde{m} = \pi^{-1}(m)$ we have that $\tilde{m} \subseteq k[x_1, \dots, x_n]$ is maximal and we have an induced algebra isomorphism

$$k(\tilde{m}) \xrightarrow{\sim} R/m.$$

Hence from an algebra embedding $k(\tilde{m}) \to \bar{k}$ we obtain an algebra embedding $R/m \to \bar{k}$. ∎

Corollary 10: For any alg closed field $k = \bar{k}$, any maximal ideal $m \subseteq k[x_1, \dots, x_n]$ is of the form $m = (x_1 - a_1, \dots, x_n - a_n)$ for uniquely determined $a_i \in k$.

Proof: By Zariski / Cor 9 $m = \ker(\varphi)$ for an alg map $\varphi : k[x_1, \dots, x_n] \to k$, via alg closure. Hence for each x_i we have $x_i - a_i \in \ker \varphi = m$ for $a_i = \varphi(x_i)$. Thus

$$m_a = (x_i - a_i : 1 \le i \le n) \subseteq m.$$

Since m_a is already maximal this gives $m = m_a$. As for uniqueness, since the composite $k \to k[x_1, \dots, x_n] \to k$ is injective, if $x_i - a \in \ker(\pi) = m$ then

$$a = \pi(a) = \pi(x_i) = a_i.$$ ∎

Alt Proof: By Zariski, the induced map

$$k \to k[x_1, \dots, x_n]/m \qquad (*)$$

is an isomorphism, so that $\bar{x}_i = a_i$ for some $a_i \in k$, at each $i = 1, \dots, n$. Thus

$$\bar{x}_i - a_i = 0 \implies x_i - a_i \in m.$$

If $x_i - a \in m$ for some $a \in k$ then, for uniqueness, we get

image of $a - a_i$ in $k[x_1, \dots, x_n]/m$

$$= (\bar{x}_i - a_i) - (\bar{x}_i - a)$$

$$= 0.$$

By injectivity of $(*)$ we have $a = a_i$. ∎

Below we let $\mathbb{A}^n_k \overset{\text{a fancy not for}}{=} k^n$.

Corollary 11: For any alg closed field $k = \bar{k}$, like $\overline{\mathbb{F}_p}$, $\overline{\mathbb{Q}}$, $\mathbb{C}$, $\overline{\mathbb{C}(t)}$, etc., the set map

$$\mathbb{A}^n_k \longrightarrow \{\text{max ideals in } k[x_1, \dots, x_n]\}$$

$$\alpha = (a_1, \dots, a_n) \longmapsto m_\alpha = (x_i - a_i : 1 \le i \le n)$$

is a bijection!

~ IV The maximal spectrum

Def$^{\underline{n}}$: For any commutative ring R, define

$$\mathrm{mSpec}(R) = \left\{ \begin{array}{l} \text{The collection of all} \\ \text{max'l ideals } \mathfrak{m} \subseteq R \end{array} \right\}.$$

Take more generally,

$$\mathrm{Spec}(R) = \left\{ \begin{array}{l} \text{The collection of all} \\ \underline{\text{prime}} \text{ ideals } \mathfrak{p} \subseteq R \end{array} \right\}.$$

We just saw, for $k = \mathbb{C}$ for example,

$$\mathrm{mSpec}(\mathbb{C}[x_1, \ldots, x_n]) \cong \mathbb{C}^n.$$

Over $k = \mathbb{R}$, alternatively,

$$\mathrm{mSpec}(\mathbb{R}[x_1, \ldots, x_n]) = \mathbb{R}^n \sqcup \left\{ \begin{array}{l} \text{a bunch of} \\ \text{other stuff} \\ \text{having to do w/} \\ \mathbb{C} \end{array} \right\}.$$

For example

$$\mathrm{mSpec}(\mathbb{R}[x]) = \mathbb{R} \sqcup \left\{ \begin{array}{l} (x^2 - bx - a) \\ \text{such that} \\ \frac{1}{4} b^2 < a \end{array} \right\}$$

$=$



IV Further analysis of mSpec(R)

Let's look at some examples of quotients $k[x_1, \ldots, x_n]/I$.

Consider the quotient, for example,

$$R = \mathbb{C}[x,y] / (y^2 - x^3)$$

By Zariski each max ideal $m \subseteq R$ has $R/m = \mathbb{C}$ and restriction along the projection $\pi : \mathbb{C}[x,y] \to R$ gives an injection

$$\pi^* : \mathrm{mSpec}(R) \hookrightarrow \mathrm{mSpec}(\mathbb{C}[x,y]) = \mathbb{C}^2$$

$$m \longmapsto \pi^{-1}(m)$$

$$\mathrm{mSpec}(R) \subseteq \mathbb{C}^2.$$

Each max ideal $m_\alpha \subseteq \mathbb{C}[x,y]$ appears as the corresponding kernel of the evaluation map $ev_\alpha : \mathbb{C}[x,y] \to \mathbb{C}$.

Observe: A max ideal $m_\alpha \in \mathrm{mSpec}(\mathbb{C}[x,y])$ is in the image of the inclusion π^*

$\Leftrightarrow$ (a) $y^2 - x^3 \in \ker(ev_\alpha)$

$\Leftrightarrow$ (b) $\alpha_2^2 - \alpha_1^3 = 0$

$\Leftrightarrow$ (c) (α_1, α_2) is a point on the complex curve $y^2 - x^3 = 0$

$\subseteq \mathbb{C}^2$

Hence

$$\mathrm{mSpec}(\mathbb{C}[x,y] / (y^2 - x^3)) =$$



More generally, for any finite type k-algebra with $k = \bar{k}$ and a given presentation

$$\pi : k[x_1, \dots, x_n] \longrightarrow R, \quad R \cong k[x_1, \dots, x_n] / \ker \pi,$$

Noetherianity gives

$$\ker(\pi) = (f_1, \dots, f_r)$$

for some functions $f_i(x_1, \dots, x_n)$. We again have the inclusion

$$\pi^* : m\mathrm{Spec}(R) = m\mathrm{Spec}(k[x_1, \dots, x_n] / (f_1, \dots, f_r)) \hookrightarrow m\mathrm{Spec}(k[x_1, \dots, x_n]) = \mathbb{C}^n.$$

Theorem 12: Let R be a finite type k-alg over an algebraically closed field $k = \bar{k}$. For a chosen presentation $R \cong k[x_1, \dots, x_n] / (f_1, \dots, f_r)$ and corresp. surj. alg map $\pi : k[x_1, \dots, x_n] \longrightarrow R$, taking the preimage along π provides an inclusion

$$\pi^* : m\mathrm{Spec}(R) \hookrightarrow m\mathrm{Spec}(\mathbb{C}[x_1, \dots, x_n]) \underset{Zar}{=} k^n$$

$$m \longmapsto \pi^{-1}(m).$$

The image of this inclusion consists of all points $(\alpha_1, \dots, \alpha_n) \in k^n$ such that $f_i(\alpha_1, \dots, \alpha_n) = 0$ at all n,

$$m\mathrm{Spec}(R) \xrightarrow{\cong} \left\{ \begin{array}{l} \text{Simultaneous sols} \\ \text{to the equations} \\ f_i(x_1, \dots, x_n) = 0, \ i = 1, \dots, r \end{array} \right\} \subseteq k^n.$$

Warning! A general finite type k-alg does not come equipped with a specified presentation $R \cong$ quotient of a poly ring. So you can think of

$$\underbrace{\mathrm{mSpec}(R)}_{\text{A choice free, coord free object}} \xrightarrow{\cong} \underbrace{\left\{ \begin{array}{c} \text{Soln to eq's} \\ f_1(x_1, \dots, x_n) = 0 \\ \vdots \\ f_r(x_1, \dots, x_n) = 0 \end{array} \right\}}_{\text{Choosing coordinates}}$$

Warning! If $k \neq \bar{k}$, then $\mathrm{mSpec}(R)$ is still a thing, it's still perfectly fine to think about, but this geometric picture is no longer valid,

$$\mathrm{mSpec}(R) \neq \text{Vanishing loci of some poly's in } k^n \text{ (or just } \mathbb{C}^n).$$

This is due to the existence of field extensions nontrivial finite-fin type field extensions $k \subseteq K$.

Question: If I have some presentation

$$\pi : k[x_1, \dots, x_n] \to R,$$

how do I think about <u>non-maximal</u> primes in $\mathrm{Spec}(R)$, in geometric terms?

HW

1. Prove that an ideal $\mathfrak{p}$ in a commutative ring R is prime if and only if $\mathfrak{p}$ is the kernel of a ring map $\phi: R \to S$ to a domain S.

2. (a) Prove that the ideal (y^3-x^2, y^5-x^2) is not prime in $\mathbb{C}[x,y]$.

(b) Prove that the ideal $(z-x^2, z-y^2)$ is not prime in $\mathbb{Q}[x,y,z]$

(c) Prove that the ideal $(y-x^3-2x, x^4+x^3+x^2+x+1)$ is prime in $\mathbb{Q}[x,y]$.

(d) Prove that the ideal (z^2-xy) is prime in $\mathbb{F}_5[x,y,z]$. (Hint (z^2-xy) is the kernel of an algebra map $\mathbb{F}_5[x,y,z] \to \mathbb{F}_5[t,u]$.)

3. For an ideal I in a commutative ring R, define $\mathrm{Van}(I) = \{\text{all primes } \mathfrak{p} \subseteq R \text{ s.t. } I \subseteq \mathfrak{p}\} \subseteq \mathrm{Spec}(R)$ and $\mathrm{mVan}(I) = \{\text{all max ideals } \mathfrak{m} \subseteq R \text{ with } I \subseteq \mathfrak{m}\} \subseteq \mathrm{mSpec}(R)$. Prove that for any collection of ideals I_λ, $\lambda \in \Lambda$,

$$\bigcap_{\lambda \in \Lambda} \mathrm{Van}(I_\lambda) = \mathrm{Van}\left(\sum_\lambda I_\lambda\right)$$

and for any finite collection $I_1, I_2, \ldots, I_n$

$$\mathrm{Van}(I_1) \cup \cdots \cup \mathrm{Van}(I_n) = \mathrm{Van}\left(\prod_{j=1}^{n} I_j\right).$$

Convince yourself that the analogous equalities hold when we replace Van with mVan.

4. For $\mathcal{U}_I : \mathrm{Spec}(R) - \mathrm{Van}(I)$ and $m\mathcal{U}_I = m\mathrm{Spec}(R) - m\mathrm{Van}(I)$, prove that the collections $\{\mathcal{U}_I : I \text{ an ideal in } R\}$ and $\{m\mathcal{U}_I : I \text{ an ideal in } R\}$ are the open sets in topologies for $\mathrm{Spec}(R)$ and $m\mathrm{Spec}(R)$ respectively. Prove furthermore that the $\mathcal{U}_f := \mathcal{U}_{(f)}$ and $m\mathcal{U}_f = m\mathcal{U}_{(f)}$ provide bases for these topologies.

These topologies are called the Zariski topology on $\mathrm{Spec}(R)$ and $m\mathrm{Spec}(R)$

5. Describe the open sets in $m\mathrm{Spec}(\mathbb{C}[x])$. Prove that $m\mathrm{Spec}(\mathbb{C}[x])$ is a compact topological space. Prove also that $m\mathrm{Spec}(\mathbb{C}[x])$ is non-Hausdorff as a topological space.

6. Prove that $\mathrm{Spec}(R)$ is compact whenever R is Noetherian.