


Influence of simulated risk group turnover in STI epidemics with assortative mixing

Jesse Knight, Sharmistha Mishra

Institute of Medical Science
University of Toronto

Canadian Student Health Research Forum

2020 August 25



Disclosures

None

Acknowledgements



Background

¹Rowley et al. 2019.

²UNAIDS 2020.

³Mishra et al. 2014.

Background

STI — Sexually Transmitted Infections

- ▶ 1+ million new STI infections per day¹
- ▶ 1.7 million new HIV infections per year²

¹Rowley et al. 2019.

²UNAIDS 2020.

³Mishra et al. 2014.

Background

STI — Sexually Transmitted Infections

- ▶ 1+ million new STI infections per day¹
- ▶ 1.7 million new HIV infections per year²

tPAF — Transmission Population Attributable Fraction³

- ▶ based on epidemic simulation models
- ▶ prioritize risk groups with interventions

¹Rowley et al. 2019.

²UNAIDS 2020.

³Mishra et al. 2014.

Key Modelling Concepts

Key Modelling Concepts

Turnover:

movement between risk groups

Key Modelling Concepts

Turnover:

movement between risk groups

High


Med

Low

Key Modelling Concepts

Turnover:


movement between risk groups



Key Modelling Concepts

Turnover:


movement between risk groups



Key Modelling Concepts

Turnover:

movement between risk groups




Assortative Mixing:

like-with-like (vs random) partnerships

Key Modelling Concepts

Turnover:

movement between risk groups



Assortative Mixing:


like-with-like (vs random) partnerships



Key Modelling Concepts


Turnover:

movement between risk groups



Assortative Mixing:


like-with-like (vs random) partnerships



Key Modelling Concepts


Turnover:

movement between risk groups



Assortative Mixing:

like-with-like (vs random) partnerships



Research Questions

Influence of turnover on:

1. equilibrium **STI prevalence**
2. **tPAF** of High Risk group

...under **random** vs **assortative** mixing

Methods

Methods

- Susceptible, Infectious, Recovered (SIR)



Methods

- Susceptible, Infectious, Recovered (SIR)



- Stable turnover in 3 risk groups

Methods

- ▶ Susceptible, Infectious, Recovered (SIR)



- ▶ Stable turnover in 3 risk groups
- ▶ **STI prevalence** vs turnover

Methods

- ▶ Susceptible, Infectious, Recovered (SIR)



- ▶ **Calibrate** risk group partners per year to reproduce the **same STI prevalence**
- ▶ Stable turnover in 3 risk groups
- ▶ **STI prevalence** vs turnover

Methods

- Susceptible, Infectious, Recovered (SIR)



- ▶ Stable turnover in 3 risk groups

- STI prevalence vs turnover

- **Calibrate** risk group partners per year to reproduce the **same STI prevalence**

- ▶ 4 model variants:

Random
VS
Assortative

×

No Turnover
VS
Turnover

Methods

- Susceptible, Infectious, Recovered (SIR)



- ▶ Stable turnover in 3 risk groups
- ▶ **STI prevalence** vs turnover


- **Calibrate** risk group partners per year to reproduce the **same STI prevalence**

- ▶ 4 model variants:


Random No Turnover
VS × VS
Assortative Turnover

- ▶ **tPAF** for each variant

Random mix — turnover “homogenizes” STI prevalence




Random mix — turnover “homogenizes” STI prevalence



No Turnover (A)




Random mix — turnover “homogenizes” STI prevalence



No Turnover (A)



Turnover (B)



Random mix — infer larger risk ratio with turnover → higher tPAF+


	No Turnover	Turnover
STI prevalence *		
Partners per year *		
10-year tPAF (Cal)		

* Ratios = (High : Low) Risk; Pre → Post-Calibration

No Turnover (A)



Turnover (B)



Random mix — infer larger risk ratio with turnover → higher tPAF+


	No Turnover	Turnover
STI prevalence *	9.2 → 6.7	6.7 → 6.7
Partners per year *		
10-year tPAF (Cal)		

* Ratios = (High : Low) Risk; Pre → Post-Calibration

No Turnover (A)



Turnover (B)



Random mix — infer larger risk ratio with turnover → higher tPAF+


	No Turnover	Turnover
STI prevalence *	9.2 → 6.7	6.7 → 6.7
Partners per year *	25.0 → 15.2	25.0 → 23.9
10-year tPAF (Cal)		

* Ratios = (High : Low) Risk; Pre → Post-Calibration

No Turnover (A)



Turnover (B)



Random mix — infer larger risk ratio with turnover → higher tPAF+


	No Turnover	Turnover
STI prevalence *	9.2 → 6.7	6.7 → 6.7
Partners per year *	25.0 → 15.2	25.0 → 23.9
10-year tPAF (Cal)	0.759	0.804

* Ratios = (High : Low) Risk; Pre → Post-Calibration

No Turnover (A)



Turnover (B)



Random mix — infer larger risk ratio with turnover → higher tPAF+

	No Turnover	Turnover
STI prevalence *	9.2 → 6.7	6.7 → 6.7
Partners per year *	25.0 → 15.2	25.0 → 23.9
10-year tPAF (Cal)	0.759	0.804


* Ratios = (High : Low) Risk; Pre → Post-Calibration

► Ignore turnover → underestimate tPAF (5.6%)


No Turnover (A)




Turnover (B)



Assort mix — turnover allows infections to “escape” sexual networks




Assort mix — turnover allows infections to “escape” sexual networks



No Turnover (A)




Assort mix — turnover allows infections to “escape” sexual networks



No Turnover (A)



Turnover (B)



Assort mix — higher risk ratio & escaped infections → higher tPAF++


	No Turnover	Turnover
STI prevalence *		
Partners per year *		
10-year tPAF (Cal)		

* Ratios = (High : Low) Risk; Pre → Post-Calibration

No Turnover (A)



Turnover (B)



Assort mix — higher risk ratio & escaped infections → higher tPAF++


	No Turnover	Turnover
STI prevalence *	17.5 → 6.7	9.8 → 6.7
Partners per year *		
10-year tPAF (Cal)		

* Ratios = (High : Low) Risk; Pre → Post-Calibration

No Turnover (A)



Turnover (B)



Assort mix — higher risk ratio & escaped infections → higher tPAF++


	No Turnover	Turnover
STI prevalence *	17.5 → 6.7	9.8 → 6.7
Partners per year *	25.0 → 6.0	25.0 → 10.1
10-year tPAF (Cal)		

* Ratios = (High : Low) Risk; Pre → Post-Calibration

No Turnover (A)



Turnover (B)



Assort mix — higher risk ratio & escaped infections → higher tPAF++


	No Turnover	Turnover
STI prevalence *	17.5 → 6.7	9.8 → 6.7
Partners per year *	25.0 → 6.0	25.0 → 10.1
10-year tPAF (Cal)	0.505	0.643

* Ratios = (High : Low) Risk; Pre → Post-Calibration

No Turnover (A)



Turnover (B)





Assort mix — higher risk ratio & escaped infections → higher tPAF++

	No Turnover	Turnover
STI prevalence *	17.5 → 6.7	9.8 → 6.7
Partners per year *	25.0 → 6.0	25.0 → 10.1
10-year tPAF (Cal)	0.505	0.643

* Ratios = (High : Low) Risk; Pre → Post-Calibration

- ▶ Ignore turnover → underestimate tPAF (21.5%)



Implications

Implications

1. Influence of **turnover** on STI epidemics is **larger** under **assortative** mixing

Implications

1. Influence of **turnover** on STI epidemics is **larger** under **assortative** mixing
2. If turnover is **ignored**: we **underestimate** impact of reaching **high risk** groups

Implications

1. Influence of **turnover** on STI epidemics is **larger** under **assortative** mixing
2. If turnover is **ignored**: we **underestimate** impact of reaching **high risk** groups

Also relevant to non-STI epidemics ...

Implications

1. Influence of **turnover** on STI epidemics is **larger** under **assortative** mixing
2. If turnover is **ignored**: we **underestimate** impact of reaching **high risk** groups

Also relevant to non-STI epidemics ...like **COVID-19**

References

- Mishra, Sharmistha et al. (Feb. 2014). “Distinguishing sources of HIV transmission from the distribution of newly acquired HIV infections: Why is it important for HIV prevention planning?” In: *Sex Transm Infect* 90.1, pp. 19–25. URL: doi.org/10.1136/sextrans-2013-051250.
- Rowley, Jane et al. (2019). “Chlamydia, gonorrhoea, trichomoniasis and syphilis: global prevalence and incidence estimates, 2016”. In: *Bull World Health Organ* 97, pp. 548–562. URL: doi.org/10.2471/BLT.18.228486.
- UNAIDS (2020). *AIDSinfo*. URL: <http://aidsinfo.unaids.org/>.