

INFORMATION FOR THE ORGANIZERS
IT WILL NOT APPEAR IN THE BOOK OF ABSTRACT

Please delete the answers that do not correspond to your case.

Type of communication:

- oral presentation

A limited number of oral presentations are available and will be selected by the scientific committee

Preferred session(s):

- Chemical and Biophysical Tools for Investigation Across Scales

Participant:

- Other : large scale facility for research

Interested to contribute to the EPJE special issue:

- No

NEUTRONS FOR MEMBRANES: OPPORTUNITIES AND ACCESS AT THE ILL

Bruno DEMÉ

Institut Laue - Langevin, Large Scale Structures group, Grenoble, France

(corresponding author: deme@ill.fr)

Keywords: lamellar diffraction, skin lipids, plant lipids, myelin structure

ABSTRACT

In this short presentation I will introduce the range of neutron techniques available at the ILL for membrane science, including diffraction, small-angle neutron scattering (SANS), reflectometry, and spectroscopy. I will then illustrate the potential of membrane diffraction, a method dedicated to membrane structural studies, that has been pioneered at Brookhaven and at the ILL on the D16 instrument [1] in the 1970s, and has been continuously improved ever since. The method relies on the combination of neutron diffraction by solid supported, oriented lamellar stacks of membranes with precise control of temperature and humidity [2]. I will illustrate the results obtained with this technique with a selection of past studies from the ILL user program and from my own research. These span from simple synthetic models [3], skin lipids [4-7], plant lipids from thylakoids [9-11], to studies of the myelin structure [12-13]. Finally, I will outline the proposal process to apply for beamtime, the support available to first-time users, with the aim of lowering the barrier to entry for the membrane science community.

REFERENCES

- 1 J.-C. Buffet, et al., *Eur. Phys. J.*, **286**, 03010, (2023), DOI: /10.1051/epjconf/202328603010
- 2 J. Gonthier, et al., *J. Neut. Res.*, **21**, 65-76, (2019), DOI: /10.3233/JNR-190109
- 3 M. Kanduč, et al., *Nat. Comm.*, **8**, 1-9, (2017), DO : /10.1038/ncomms14899
- 4 T. Schmitt, et al., *Chem. Phys. Lip.*, **216**, 91-103, (2018), DOI: /10.1016/j.chemphyslip.2018.09.017
- 5 T. Schmitt, et al., *BBA Biomem.*, **1861**, 306-315, (2019), DOI : /10.1016/j.bbamem.2018.06.008
- 6 C. Beddoes, et al., *Langmuir*, **36**, 10270–10278, (2020), DOI: /10.1021/acs.langmuir.0c01992
- 7 C. Beddoes, et al., *BBA Biomem.*, **1864**, 183886, (2022), DOI: /10.1016/j.bbamem.2022.183886
- 8 F. Fandrei, et al., *J. Lip. Res.*, **64**, 1-17, (2023), DOI: /10.1016/j.jlir.2023.100356
- 9 B. Demé, et al., *FASEB J.*, **Volume**, 3373-3383, (2014), DOI: /10.1096/fj.13-247395
- 10 S. Bolik, et al., *BMC Biol.*, **21**, 1-16, (2023), DOI : /10.1186/s12915-023-01775-z
- 11 S. Bolik, et al., *BBA Biomem.*, **1864**, 184037, (2022), DOI: /10.1016/j.bbamem.2022.18403
- 12 A. R. Denninger, et al., *Acta Cryst. D*, **70**, 3198–3211 (2014), DOI:/10.1107/S1399004714023815
- 13 A. Baumann et al., *Curr. Res. in Struct. Biol.* **4** 231–245 (2022), DOI: /10.1016/j.crstbi.2022.06.003