
MOORE GRAPH WITH PARAMETERS (3250,57,0,1) DOES NOT EXIST *

A.A.Makhnev

Krasovskii Institute of Mathematics and Mechanics UB RAS,
Ural Federal University
makhnev@imm.uran.ru

ABSTRACT

If a regular graph of degree k and diameter d has v vertices then

$$v \leq 1 + k + k(k-1) + \dots + k(k-1)^{d-1}$$

Graphs with $v = 1 + k + k(k-1) + \dots + k(k-1)^{d-1}$ are called Moore graphs. Damerell proved that a Moore graph of degree $k \geq 3$ has diameter 2. If Γ is a Moore graph of diameter 2, then $v = k^2 + 1$, Γ is strongly regular with $\lambda = 0$ and $\mu = 1$, and one of the following statements holds: $k = 2$ and Γ is the pentagon, $k = 3$ and Γ is the Petersen graph, $k = 7$ and Γ is the Hoffman-Singleton graph, or $k = 57$. The existence of a Moore graph of degree 57 was unknown.

Jurishich and Vidali have proved that the existence of a Moore graph of degree $k > 3$ is equivalent to the existence of a distance-regular graph with intersection array $\{k-2, k-3, 2; 1, 1, k-3\}$ (in the case $k = 57$ we have a distance-regular graph with intersection array $\{55, 54, 2; 1, 1, 54\}$).

In this paper we prove that a distance-regular graph with intersection array $\{55, 54, 2; 1, 1, 54\}$ does not exist. As a corollary, we prove that a Moore graph of degree 57 does not exist. K

Keywords distance-regular graph · Moore graph

1 Intro

We consider undirected graphs without loops and multi-edges. If a, b are vertices of a graph Γ then we denote by $d(a, b)$ the distance between a and b . We denote by $\Gamma_i(a)$ a subgraph of the graph Γ induced by a set of vertices with the distance i from the vertex a . A subgraph $\Gamma_1(a)$ is called a *neighbourhood* of a vertex a and denoted by $[a]$.

The definition of the *distance-regular graph* with the intersection array $\{b_0, \dots, b_{d-1}; c_1, \dots, c_d\}$ can be found in [1].

For a graph Γ with the diameter d and $i \in \{2, \dots, d\}$ let's denote by Γ_i a graph defined on the same set of vertices where any two vertices u, w are adjacent if the distance between u and w in the graph Γ is i .

If a regular graph of degree k and diameter d has v vertices then the following inequality holds:

$$v \leq 1 + k + k(k-1) + \dots + k(k-1)^{d-1}$$

Graphs for which the non-strict inequality above turns into equality are called *Moore graphs* (1960). This definition belongs to Hoffman and Singleton who explored graphs with diameters of 2 and 3. The simplest example of Moore graph is a regular $(2d+1)$ -polygon.

Damerell [3] (see also [4]) has proven that a graph of degree $k \geq 3$ has diameter 2. In this case $v = k^2 + 1$ the graph is strongly regular with $\lambda = 0$ and $\mu = 1$ and one of the following statements holds: $k = 2$ and the graph is the

*Citation: Translation from Russian: **arXiv:2010.13443v2 [math.CO] 9 Nov 2020**

pentagon, $k = 3$ and the graph is the Petersen graph, $k = 7$ and the graph is the Hoffman-Singleton graph, or $k = 57$. The existence of a Moore graph of degree 57 was unknown.

Ashbacher [5] has proven that a Moore graph of degree 57 is not distance-transitive (the Moore graph of degree 57 sometimes called Ashbacher's graph). G. Higman ([6, Theorem 3.13]) has proven that a Moore graph of degree 57 is not a (vertex?)-symmetric graph. A.A. Makhnev and D.V. Paduchikh explored possible automorphisms of the Moore graph of degree 57.

Jurishich and Vidali [9] have noticed that the existence of a Moore graph of degree $k > 3$ is equivalent to the existence of a distance-regular graph with intersection array $\{k-2, k-3, 2; 1, 1, k-3\}$ (in the case $k = 57$ we have a distance-regular graph with intersection array $\{55, 54, 2; 1, 1, 54\}$).

In this paper we explore properties of the graph with intersection array $\{55, 54, 2; 1, 1, 54\}$ and claim:

Theorem 1. *A distance-regular graph with intersection array $\{55, 54, 2; 1, 1, 54\}$ does not exist.*

Corollary 1. *Strongly regular Moore graph with parameters $(3250, 57, 0, 1)$ does not exist.*

This result puts an end to 60 years of research on Moore graphs. The main methods used to prove the results are: the triple intersection number method applied to a graph without the Q-polynomial condition; method of symmetrization of an array of triple intersection numbers (proposed by A.A. Makhnev, the analogue of tensor symmetrization).

2 Auxiliary Results

Our notation is conventional and can be found in [1]. In the proof of the theorem we use *triple intersection numbers* [11].

Let Γ be a distance-regular graph with the diameter d . For vertices u_1, u_2, u_3 and non-negative integers Γ and r_1, r_2, r_3 are non-greater than d define as $\left\{ \begin{smallmatrix} u_1 u_2 u_3 \\ r_1 r_2 r_3 \end{smallmatrix} \right\}$ a set of vertices $w \in \Gamma$ such that $d(w, u_i) = r_i$.²

Let's denote as $\left[\begin{smallmatrix} u_1 u_2 u_3 \\ r_1 r_2 r_3 \end{smallmatrix} \right]$ the number of vertices in $\left\{ \begin{smallmatrix} u_1 u_2 u_3 \\ r_1 r_2 r_3 \end{smallmatrix} \right\}$. Numbers $\left[\begin{smallmatrix} u_1 u_2 u_3 \\ r_1 r_2 r_3 \end{smallmatrix} \right]$ we refer to as *triple intersection numbers*. For the fixed triplet of vertices u_1, u_2, u_3 instead of $\left[\begin{smallmatrix} u_1 u_2 u_3 \\ r_1 r_2 r_3 \end{smallmatrix} \right]$ we write $[r_1 r_2 r_3]$. Unfortunately there is no general way to compute $[r_1 r_2 r_3]$. But in [11] we propose a technique to compute some of $[r_1 r_2 r_3]$ numbers.

Let u, v, w be vertices of a graph Γ and $W = d(u, v)$, $U = d(v, w)$, $V = d(u, w)$. There is only one vertex $x = u$ such that $d(x, u) = 0$. Therefore $[0jh]$ is equal to 0 or 1. Thus, $[0jh] = \delta_{jW} \delta_{hW}$, $[i0h] = \delta_{iU} \delta_{hU}$, $[ij0] = \delta_{iV} \delta_{jV}$.

Another set of equations we can get by fixing the distance between any two vertices from $\{u, v, w\}$ and by counting the number of vertices with any distance to the third one:

$$\sum_{l=1}^d [ljh] = p_{jh}^U - [0jh], \sum_{l=1}^d [ilh] = p_{ih}^V - [i0h], \sum_{l=1}^d [ijl] = p_{ij}^W - [ij0] \quad (+)$$

Wherein some of the triplets disappear. When $|i - j| > W$ or $|i + j| < W$ we have $p_{ij}^W = 0$. Therefore $[ijh] = 0$ for all $h \in \{0, \dots, d\}$.

Let $S_{ijh}(u, v, w) = \sum_{r,s,t=0}^d Q_{ri} Q_{sj} Q_{th} \left[\begin{smallmatrix} uvw \\ rst \end{smallmatrix} \right]$. If Krein parameter $q_{ij}^h = 0$ then $S_{ijh}(u, v, w) = 0$.

Let's fix vertices u, v, w of a distance-regular graph Γ with the diameter 3 and define $\{ijh\} = \left\{ \begin{smallmatrix} uvw \\ ijh \end{smallmatrix} \right\}$, $[ijh] = \left[\begin{smallmatrix} uvw \\ ijh \end{smallmatrix} \right]$, $[ijh]' = \left[\begin{smallmatrix} uvw \\ ihj \end{smallmatrix} \right]$, $[ijh]^* = \left[\begin{smallmatrix} uvw \\ jih \end{smallmatrix} \right]$, $[ijh]^\sim = \left[\begin{smallmatrix} uvw \\ hji \end{smallmatrix} \right]$. In cases where $d(u, v) = d(u, w) = d(v, w) = 2$ or

²Translator's note: I found the definition from [11] to be more precise

$$\left\{ \begin{smallmatrix} u_1 u_2 u_3 \\ r_1 r_2 r_3 \end{smallmatrix} \right\} := \{v \in \text{vertices}(\Gamma) \mid d(v, u_1) = r_1, d(v, u_2) = r_2, d(v, u_3) = r_3\}$$

$d(u, v) = d(u, w) = d(v, w) = 3$ computation of the parameters $[ijh]' = \begin{bmatrix} uvw \\ ihj \end{bmatrix}$, $[ijh]^* = \begin{bmatrix} vuw \\ jih \end{bmatrix}$, $[ijh]^\sim = \begin{bmatrix} wvu \\ hji \end{bmatrix}$ (symmetrization of the array of triple intersection numbers) can give new ratios to prove the non-existence of the graph.

3 Properties of $\Gamma_3(u)$ graph

In this section the graph Γ is a distance-regular graph with intersection array $\{55, 54, 2; 1, 1, 54\}$. Then Γ has the spectrum $55^1, 7^{1617}, -1^{110}, -8^{1408}, 1 + 55 + 2970 + 110 = 3136$ vertices and dual eigenvalue matrix:

$$\begin{pmatrix} 1 & 1617 & 110 & 1408 \\ 1 & \frac{1029}{5} & -2 & -\frac{1024}{5} \\ 1 & -\frac{49}{15} & -2 & \frac{64}{15} \\ 1 & \frac{147}{5} & 54 & -\frac{128}{5} \end{pmatrix}$$

Then the graph Γ_3 is 56×56 lattice graph and the neighborhood of the vertex in Γ_3 is a union of two isolated 55-cliques.

Lemma 1. *The intersection numbers of the graph Γ are:*

1. $p_{11}^1 = 0, p_{12}^1 = 54, p_{22}^1 = 2808, p_{23}^1 = 108, p_{33}^1 = 2$
2. $p_{11}^2 = 1, p_{12}^2 = 52, p_{13}^2 = 2, p_{22}^2 = 2811, p_{23}^2 = 106, p_{33}^2 = 2$
3. $p_{12}^3 = 54, p_{13}^3 = 1, p_{22}^3 = 2862, p_{23}^3 = 54, p_{33}^3 = 54$

Proof. Direct computation according to [1, Lemma 4.1.7]. Let's fix the vertices u, v, w of the graph Γ and define $\{ijh\} = \begin{bmatrix} uvw \\ ijh \end{bmatrix}$.

Let $\Delta = \Gamma_2(u)$ and $\Lambda = \Delta_2$. Then Λ is a regular graph of degree $p_{22}^2 = 2811$ defined on $k_2 = 2970$ vertices. \square

Lemma 2. *Let $d(u, v) = 2, d(u, w) = d(v, w) = 1$. Then triple intersection numbers are:*

1. $[122] = 52, [132] = 2$
2. $[212] = 52, [221] = 53, [222] = r_1 + 2650, [223] = -r_1 + 108, [232] = -r_1 + 106, [233] = r_1$
3. $[312] = 2, [322] = -r_1 + 106, [323] = [332] = r_1, [333] = -r_1 + 2$

where $r_1 \in \{0, 1, 2\}$

Proof. Simplification of the set of equations +. \square

Lemma 3. *Let $d(u, v) = d(u, w) = 2, d(v, w) = 3$. Then triple intersection numbers are:*

1. $[112] = [121] = -r_6 + 1, [113] = [131] = r_6, [122] = r_7, [123] = [132] = r_6 - r_7 + 51, [133] = -2r_6 + r_7 - 49$
2. $[212] = [221] = 52, [213] = [231] = 0, [222] = r_5 + r_6 + 2756 - r_7 = r_9, [223] = [232] = -2r_6 + r_7 + 3, [233] = 2r_6 - r_7 + 102$
3. $[312] = [321] = r_6 + 1, [313] = [331] = 1 - r_6, [322] = 5 - 2r_6, [323] = [332] = r_6, [333] = 1$

where $r_6 \in \{0, 1\}, r_7 \in \{49, \dots, 52\}$. Then one of the followings is true:

$$[222] = 2706 \text{ and } [222] = 2r_6 + 50 = r_7$$

or

$$[222] = 2707 \text{ and } [222] = 2r_6 + 51 = r_7 \text{ and } r_6 = 0$$

Proof. Using computer aided simplification of the set of equations + we have

$$\begin{aligned} [112] &= -r_6 + 1, [113] = r_6, [121] = -r_4 - r_8 + 54, [122] = r_7, [123] = r_4 - r_7 + r_8 - 2, [131] = r_4 + r_8 - 53, [132] = \\ &r_6 - r_7 + 51, [133] = -r_4 - r_6 + r_7 - r_8 + 4; \\ [212] &= r_5 + r_6 - r_7 - r_9 + 2808, [213] = -r_5 - r_6 + r_7 + r_9 - 2756, [221] = r_4, [222] = r_9, [223] = \\ &-r_4 - r_9 + 2811, [231] = -r_4 + 52, [232] = -r_5 - r_6 + r_7 + 3, [233] = r_4 + r_5 + r_6 - r_7 + 50; \\ [312] &= -r_5 + r_7 + r_9 - 2755, [313] = r_5 - r_7 - r_9 + 2757, [321] = r_8, [322] = -r_7 - r_9 + 2861, [323] = \\ &r_7 - r_8 + r_9 - 2755, [331] = -r_8 + 2, [332] = r_5, [333] = -r_5 + r_8, \end{aligned}$$

where $r_4 \in \{51, 52\}$, $r_5 \in \{0, 1, 2\}$, $r_6 \in \{0, 1\}$, $r_7 \in \{49, \dots, 52\}$, $r_8 \in \{0, 1, 2\}$, $r_9 \in \{2705, \dots, 2710\}$. Thus $2705 \leq [222] = r_9 \leq 2710$.

In the 56×56 lattice Γ_3 a subgraph $\Gamma_3(u) \cap \Gamma_3(v)$ is 2-co-clique which has intersection with $\Gamma_3(w)$ exactly in one vertex. Therefore $[333] = -r_5 + r_8 = 1$ and $r_5 \in \{0, 1\}$.

Same, $\Gamma_3(v) \cap \Gamma_3(w)$ is 54-clique which has intersection just with $\Gamma_3(u)$ in no more then one vertex. Therefore $[133] = -r_4 - r_6 + r_7 - r_5 + 3 \leq 1$.

Symmetrization. For the vertex triplet (u, v, w) we have $[122] = r'_7 = r'_7, -r_6 + 1 = [112] = [121]' = -r'_4 - r'_5 + 53$ and $r_4 + r_5 = r'_6 + 52$. Then, $[222] = r_9 = r'_9$, $r_6 = [323] = [332]' = r'_5$ and $r_4 = 52$. Therefore $[133] = -49 - r_6 + r_7 - r_5 \leq 1$ and $r_7 \leq r_6 + r_5 + 50$.

Then we have $[313] = r_5 - r_7 - r_9 + 2757$, therefore $r_7 + r_9 \leq r_5 + 2757$ and $[213] = -r_5 - r_6 + r_7 + r_9 - 2756 \leq 1 - r_6$.

Assume $[213] = 1$. Then $r_6 = 0$, $[233] = r_5 - r_7 + 102$ and $r_5 = r'_5 = 0$, thus $r_7 + r_9 = 2757$.

Because $[133] = r_7 - r_8 - 48 = r'_7 - r'_8 - 48$ then $r_8 = r'_8$. Further, $1 = [112] = [121]' = r'_8$ and $r_8 = 1$. Same, $[233] = r_4 - r_7 + 50$, therefore, $r_4 = r'_4$. We have a contradiction because $51 = [212] = [221]' = r'_4 = 52$. Therefore $[213] = 0$ and $r_7 + r_9 = r_5 + r_6 + 2756$. If $r_6 = 0$, then $[233] = r_5 - r_7 + 102$ and $r_5 = r'_5 = 1$.

In case if $r_6 = 1$, then $[233] = r_5 - r_7 + 103$ and $r_5 = r'_5 = 1$. Thus in either cases we have $r_5 = r_6$ and we have symmetrized array:

$$\begin{aligned} [112] &= [121] = -r_6 + 1, [113] = [131] = r_6, [122] = r_7, [123] = [132] = r_6 - r_7 + 51, [133] = -2r_6 + r_7 - 49; \\ [212] &= [221] = 52, [213] = [231] = 0, [222] = r_5 + r_6 + 2756 - r_7 = r_9, [223] = [232] = -2r_6 + r_7 + 3, [233] = \\ &2r_6 - r_7 + 102; \\ [312] &= [321] = r_6 + 1, [313] = [331] = 1 - r_6, [322] = 5 - 2r_6, [323] = [332] = r_6, [333] = 1 \end{aligned}$$

where $r_6 \in \{0, 1\}$, $r_7 \in \{49, \dots, 52\}$.

If $[222] \geq 2708$ then $r_7 \leq 48 + 2r_6$ and the equation $[133] = -2r_6 + r_7 - 49$ gives a contradiction. If $[222] = 2705$, then $r_7 = 51 + 2r_6$, we have contradiction with the fact $[133] = -2r_6 + r_7 - 49 \leq 1$. So, either $[222] = 2706$ and $[222] = r_5 + r_6 + 50 = r_7$ or $[222] = 2707$, $r_5 + r_6 + 51 = r_7$ and $r_5 + r_6 = 0$.

□

Lemma 4. Let $d(u, v) = d(u, w) = 2$, $d(v, w) = 1$. Then triple intersection numbers are:

1. $[122] = 51, [123] = [132] = 0, [133] = 2$;
2. $[212] = [221] = 51, [222] = 2654, [223] = [232] = 106, [233] = 0$;
3. $[312] = [321] = 2, [322] = 102, [323] = [332] = 2, [333] = 0$.

Further, $[222] = 2654$.

Proof. Using simplification of the formula set + we get:

$$\begin{aligned} [122] &= r_3, [123] = [132] = -r_3 + 51, [133] = r_3 - 49; \\ [212] &= [221] = 51, [222] = -r_2 - r_3 + 2705, [223] = [232] = r_2 + r_3 + 55, [233] = -r_2 - r_3 + 51; \end{aligned}$$

$[312] = [321] = 2, [322] = r_2 + 102, [323] = [332] = -r_2 + 2, [333] = r_2$
where $r_2 \in \{0, 1, 2\}, r_3 \in \{49, 50, 51\}$.

In the 56×56 -lattice Γ_3 a subgraph $\Gamma_3(u) \cap \Gamma_3(w)$ is 2-co-clique. It has no intersection with $\Gamma_3(v)$. Therefore $[333] = r_2 = 0$.

Let's compute a number d of pairs of vertices (y, z) with the distance 3 in the graph Λ where $y \in \begin{Bmatrix} uv \\ 21 \end{Bmatrix}, z \in \begin{Bmatrix} uv \\ 23 \end{Bmatrix}$.

On the one hand we have $[213] = 0$ and $d = 0$ by the Lemma 2. On the other hand we have $d = 52[233] = 0$, therefore $[233] = 0$ and $r_3 = 51$. Thereby we have the equations from the lemma statement. \square

Lemma 5. *Let $d(u, v) = d(u, w) = (v, w) = 2$. Then triple intersection numbers are:*

$$[111] = -r_{16} - r_{11} + 1, [112] = r_{11}, [113] = r_{16}, [121] = r_{14}, [122] = -r_{12} - r_{14} + 52, [123] = r_{12}, [131] = -r_{14} + r_{16} + r_{11}, [132] = r_{12} + r_{14} - r_{11}, [133] = -r_{12} - r_{16} + 2;$$

$$[211] = r_{15} + r_{16} - r_{17} + 50, [212] = r_{17}, [213] = -r_{15} - r_{16} + 2, [221] = -r_{15} - r_{16} + r_{17} - r_{10} + 2, [222] = r_{12} + r_{16} - r_{17} + r_{10} + 2704, [223] = -r_{12} + r_{15} + 104, [231] = r_{10}, [232] = -r_{12} - r_{16} - r_{10} + 106, [233] = r_{12} + r_{16};$$

$$[311] = -r_{15} + r_{17} + r_{11} - 50, [312] = -r_{17} - r_{11} + 52, [313] = r_{15}, [321] = -r_{14} + r_{15} + r_{16} - r_{17} + r_{10} + 50, [322] = r_{14} - r_{16} + r_{17} - r_{10} + 54, [323] = -r_{15} + 2, [331] = r_{14} - r_{16} - r_{10} - r_{11} + 2, [332] = -r_{14} + r_{16} + r_{10} + r_{11}, [333] = 0$$

where $r_{10}, r_{12}, r_{15} \in \{0, 1, 2\}, r_{11}, r_{14}, r_{16} \in \{0, 1\}, r_{17} \in \{49, \dots, 52\}$.

Further, $2652 \leq [222] = r_{12} + r_{16} - r_{17} + r_{10} + 2704 \leq 2659$.

Proof. With computer-aided simplification of $+$ we have:

$$[111] = -r_{16} - r_{11} + 1, [112] = r_{11}, [113] = r_{16}, [121] = r_{14}, [122] = -r_{12} - r_{14} + 52, [123] = r_{12}, [131] = -r_{14} + r_{16} + r_{11}, [132] = r_{12} + r_{14} - r_{11}, [133] = -r_{12} - r_{16} + 2;$$

$$[211] = r_{15} + r_{16} - r_{17} + 50, [212] = r_{17}, [213] = -r_{15} - r_{16} + 2, [221] = -r_{15} - r_{16} + r_{17} - r_{10} + 2, [222] = r_{12} - r_{13} + r_{16} - r_{17} + r_{10} + 2704, [223] = -r_{12} + r_{13} + r_{15} + 104, [231] = r_{10}, [232] = -r_{12} + r_{13} - r_{16} - r_{10} + 106, [233] = r_{12} - r_{13} + r_{16};$$

$$[311] = -r_{15} + r_{17} + r_{11} - 50, [312] = -r_{17} - r_{11} + 52, [313] = r_{15}, [321] = -r_{14} + r_{15} + r_{16} - r_{17} + r_{10} + 50, [322] = r_{13} + r_{14} - r_{16} + r_{17} - r_{10} + 54, [323] = -r_{13} - r_{15} + 2, [331] = r_{14} - r_{16} - r_{10} - r_{11} + 2, [332] = -r_{13} - r_{14} + r_{16} + r_{10} + r_{11}, [333] = r_{13}$$

where $r_{10}, r_{12}, r_{13}, r_{15} \in \{0, 1, 2\}, r_{11}, r_{14}, r_{16} \in \{0, 1\}, r_{17} \in \{49, \dots, 52\}$.

In the 56×56 -lattice Γ_3 a subgraph $\Gamma_3(u) \cap \Gamma_3(w)$ is 2-co-clique. It has no intersection with $\Gamma_3(v)$. Therefore $[333] = r_{13} = 0$. Thereby we have the required equations.

Let's note that $r_{12} + r_{16} \leq 2$, therefore $2652 \leq [222] = r_{12} + r_{16} - r_{17} + r_{10} + 2704 \leq 2659$. Lemma is proved. \square

For number e of edges between $\Lambda(v)$ and $\Lambda_2(v)$ in a graph Λ the following inequalities hold: $424844 = 52 \cdot 2654 + 106 \cdot 2706 \leq e \leq 52 \cdot 2654 + 106 \cdot 2707 = 424950$. Therefore $151.136 \leq 2810 - \lambda \leq 151.174$ and $2658.826 \leq \lambda \leq 2658.864$, where λ is a mean value of the parameter $\lambda(\Lambda)$.

Let's note that the mean value $\lambda(\Lambda)$ is very close to the upper bound 2659.

4 Proof of the theorem

Let Γ be a distance-regular graph with an intersection array $\{55, 54, 2; 1, 1, 54\}$. Here we will prove the Theorem 1.

Let's fix (u, v, w) vertices of the graph Γ and define $\{ijh\} = \begin{Bmatrix} uvw \\ ijh \end{Bmatrix}, [ijh] = \begin{Bmatrix} uvw \\ ijh \end{Bmatrix}$. Let $\Delta = \Gamma_2(u)$ and $\Lambda = \Delta_2$.

Then Λ is a regular graph with the degree $p_{22}^2 = 2811$ defined on $k_2 = 2970$ vertices.

Lemma 6. *Let $d(u, v) = d(u, w) = (v, w) = 2$. Then the following statements hold:*

$$I. \ r_{11} = r'_{14}, r'_{10} = [213] = r_{12}^*, r_{15} + r_{16} + r'_{10} = 2 \text{ and } -r_{17} + 52 = r_{12}^* + r_{14}^*$$

2. if $r'_{10} = 2$ then $r_{10} = 2$ and one of the following cases is true:

either $r_{11} = 0$ and

$$\begin{aligned} [111] &= 1, [112] = [113] = [121] = [131] = 0, [122] = 50, [123] = [132] = 2, [133] = 0; \\ [211] &= 0, [212] = r_{17} = 50, [213] = [231] = 2, [221] = 50, [222] = 2658, [223] = [232] = 102, [233] = 2; \\ [311] &= 0, [312] = [321] = 2, [313] = 0, [322] = 102, [323] = [332] = 2, [331] = [333] = 0, \end{aligned}$$

or $r_{11} = 1$ and

$$\begin{aligned} [111] &= 0, [112] = [121] = 1, [113] = [131] = 0, [122] = 49, [123] = [132] = 2, [133] = 0; \\ [211] &= 1, [212] = [221] = r_{17} = 49, [213] = [231] = 2, [222] = 2659, [223] = [232] = 102, [233] = 2; \\ [311] &= 0, [312] = [321] = 2, [313] = [331] = 0, [322] = 102, [323] = [332] = 2, [333] = 0. \end{aligned}$$

Proof. Symmetrization. From $[111] = -r_{16} - r_{11} + 1$ we have $r_{16} + r_{11} = r'_{16} + r'_{11} = r^*_{16} + r^*_{11} = \tilde{r}_{16} + \tilde{r}_{11}$. Then $[112] = r_{11} = r^*_{11}$, $[113] = r_{16} = r^*_{16}$, $[121] = r_{14} = \tilde{r}_{14}$, $[212] = r_{17} = \tilde{r}_{17}$, $[313] = r_{15} = \tilde{r}_{15}$, $r_{11} = [112] = [121]' = r'_{14}$, $r'_{10} = [213] = r^*_{12}$.

From $[122] = -r_{12} - r_{14} + 52$ follows $r_{12} + r_{14} = r'_{12} + r'_{14}$, $[233] = r_{12} + r_{16} = r'_{12} + r'_{16}$, $r_{10} = [231] = [213]' = -r'_{15} - r'_{16} + 2$ and $r'_{10} + r_{15} + r_{16} = 2$.

Similarly, $-r_{17} - r_{11} + 52 = [312] = [132]^* = r^*_{12} + r^*_{14} - r^*_{11}$ and $-r_{17} + 52 = r^*_{12} + r^*_{14}$. The statement (1) is proved.

Let's assume $r'_{10} = 2$. Then $r^*_{12} = 2$, $r_{15} + r_{16} = 0$ and $r_{17} + r^*_{14} = 50$. Because $[131] = -r_{14} + r_{11}$ then in the case of $r_{11} = 0$ we have $r_{14} = 0$, $[132] = r_{12} + r_{14} - r_{11} = r_{12} = r'_{12}$, $[211] = -r_{17} + 50$, $[311] = r_{17} - 50$. Now $-r_{12} - r_{10} + 106 = [232] = [223]' = -r'_{12} + 104 = -r_{12} + 104$, therefore $r_{10} = 2$, $[212] = [221] = r_{17} = 50$, $r_{17} = [122] = -r_{12} + 52$, $r_{12} = 2$ and

$$\begin{aligned} [111] &= 1, [112] = [113] = [121] = [131] = 0, [122] = 50, [123] = [132] = 2, [133] = 0; \\ [211] &= 0, [212] = r_{17} = 50, [213] = [231] = 2, [221] = 50, [222] = 2658, [223] = [232] = 102, [233] = 2; \\ [311] &= 0, [312] = [321] = 2, [313] = 0, [322] = 102, [323] = [332] = 2, [331] = [333] = 0. \end{aligned}$$

In case where $r_{11} = 1$ we have $[233] = r_{12} = r'_{12}$, thereby $r_{12} = [123] = [132] = [132] = r_{12} + r_{14} - 1$, $r_{14} = 1$, $-r_{17} + 51 = [312] = [321]' = -r'_{17} + r'_{10} + 49$, $r_{17} = r'_{17}$, $r_{10} = 2$ and $-r_{17} + 51 = [312] = [132]^* = r^*_{12}$.

We have $[222] = r_{12} + 2657$, thereby $r_{12} = r^*_{12}$, $r_{12} + r_{17} = 51$ and

$$\begin{aligned} [111] &= 0, [112] = [121] = 1, [113] = [131] = 0, [122] = 49, [123] = [132] = 2, [133] = 0; \\ [211] &= 1, [212] = [221] = r_{17} = 49, [213] = [231] = 2, [222] = 2659, [223] = [232] = 102, [233] = 2; \\ [311] &= 0, [312] = [321] = 2, [313] = [331] = 0, [322] = 102, [323] = [332] = 2, [333] = 0. \end{aligned}$$

Now $[212] = [221] = r_{17} = r^*_{17}$, $r_{17} = [122] = -r_{12} + 51$ and $r_{12} = 2$. Lemma is proved. \square

The following idea from Vidali can be used to simplify the proof of the theorem. Let $d(u, v) = 2$. Then $[u]$ has one element in each row and each column of the 55×55 -lattice $\Gamma_3 - (\{u\} \cap \Gamma_3(u))$. Let $\{x, y\} = [u] \cap \Gamma_3(v)$. Then there is only one vertex in the subgraph $\Gamma_3(x) \cap \Gamma_3(y) - \{v\}$. We have a contradiction with the fact $[133] = 2$ (see Lemma 4) for any three vertices (u, v, w) with $d(u, v) = d(u, w) = 2$, $d(v, w) = 1$.

Theorem 1 is proved.

References

- [1] Brouwer A. E., Cohen A. M., Neumaier A. Distance-Regular Graphs. Berlin; Heidelberg; New York: Springer-Verlag, 1989.
- [2] Hoffman Alan J., Singleton Robert R. Moore graphs with diameter 2 and 3 // IBM Journal of Research and Development. 1960. V. 5, N 4. P. 497–504.
- [3] Damerell R.M. On Moore graphs // Math. Proc. Cambr. Phil. Soc., Vol. 74 (1973). P. 227–236.

- [4] Bannai E., Ito T. On finite Moore graphs // Journal of the Faculty of Science, the University of Tokyo. Sect. 1 A, Mathematics. 1973, V. 20. P. 191-208.
- [5] Ashbacher M. The nonexistence of rank 3 permutation group of degree 3250 and subdegree 57 // J.Algebra, Vol. 19, no. 3 (1971). P. 538-540.
- [6] Cameron P.J. Permutation Groups. London Math. Soc. Student Texts N 45: Cambridge Univ. Press, 1999
- [7] Makhnev A.A., Paduchikh D.V. On automorphisms of Aschbacher graph // Algebra i Logika, Vol. 40, no. 2 (2001). P. 125-134.
- [8] Makhnev A.A., Paduchikh D.V. On automorphisms group of Aschbacher graph // Doklady RAN, Vol. 426, no. 3 (2009). P. 310-313.
- [9] Jurishich A., Vidali J. The Sylvester graph and Moore graphs // Europ. J. Comb., Vol. 80 (2019). P. 184-193.
- [10] Makhnev A.A., Paduchikh D.V. The most Moore graph and distance-regular graph with intersection array $\{55, 54, 2; 1, 1, 54\}$ // Algebra i Logika, Vol. 59, no. 4 (2020).
- [11] Jurishich A., Vidali J. Extremal 1-codes in distance-regular graphs of diameter 3 // Des. Codes Cryptogr., Vol. 65 (2012). P. 29-47.

About Authors

MAKHNEV Alexander Alexeevich
 Russia, Ekaterinburg, S.Kovalevskoy st, 16, 620990,
 N.N. Krasovskii Institute of Mathematics and Mechanics
 Russia, Ekaterinburg, Mira st, 19, 620002,
 Ural Federal University
 makhnev@imm.uran.ru

About Translators

Selivanov Konstantin, konstantin.selivanov@gmail.com