

Special Focus Section: New Antifungal Compounds



Antifungal therapeutics for dimorphic fungal pathogens

Kristie D. Goughenour & Chad A. Rappleye

To cite this article: Kristie D. Goughenour & Chad A. Rappleye (2017) Antifungal therapeutics for dimorphic fungal pathogens, *Virulence*, 8:2, 211-221, DOI: [10.1080/21505594.2016.1235653](https://doi.org/10.1080/21505594.2016.1235653)

To link to this article: <https://doi.org/10.1080/21505594.2016.1235653>



© 2017 Taylor & Francis



Published online: 01 Nov 2016.



Submit your article to this journal [↗](#)



Article views: 3086



View related articles [↗](#)




View Crossmark data [↗](#)



Citing articles: 19 View citing articles [↗](#)

REVIEW

Antifungal therapeutics for dimorphic fungal pathogens

Kristie D. Goughenour and Chad A. Rappleye 

Department of Microbiology, Ohio State University, Columbus, OH, USA

ABSTRACT

Dimorphic fungi cause several endemic mycoses which range from subclinical respiratory infections to life-threatening systemic disease. Pathogenic-phase cells of *Histoplasma*, *Blastomyces*, *Paracoccidioides* and *Coccidioides* escape elimination by the innate immune response with control ultimately requiring activation of cell-mediated immunity. Clinical management of disease relies primarily on antifungal compounds; however, dimorphic fungal pathogens create a number of challenges for antifungal drug therapy. In addition to the drug toxicity issues known for current antifungals, barriers to efficient drug treatment of dimorphic fungal infections include natural resistance to the echinocandins, residence of fungal cells within immune cells, the requirement for systemic delivery of drugs, prolonged treatment times, potential for latent infections, and lack of optimized standardized methodology for in vitro testing of drug susceptibilities. This review will highlight recent advances, current therapeutic options, and new compounds on the horizon for treating infections by dimorphic fungal pathogens.

ARTICLE HISTORY

Received 28 July 2016
Revised 2 September 2016
Accepted 7 September 2016

KEYWORDS

antifungal; azoles; dimorphic fungi; echinocandins; nikkomycin; polyenes

Introduction

Within the Ascomycetes phylum, members of the genera *Histoplasma*, *Blastomyces*, *Coccidioides*, and *Paracoccidioides* are dimorphic fungi which are responsible for hundreds of thousands of infections, many thousands of which require clinical intervention. Given similarities in their pathogenesis, the disease manifestations they produce, their clinical management, and particularly the resemblance of their antifungal susceptibility profiles, these 4 genera will be the subject of this review. Two additional dimorphic fungi, *Talaromyces marneffe* and *Sporothrix schenckii*, will not be covered. Neither dimorphic fungi nor human pathogens are monophyletic groups within the Onygenales as closely related non-pathogenic and non-dimorphic species exist for several of these genera. While once designated as a single *Coccidioides* species, *Coccidioides immitis*, advances in sequence analyses and phylogenetics have separated it into 2 species: *Coccidioides immitis* and *Coccidioides posadasii*.¹ Other dimorphic fungi have similarly been separated into different species with *Paracoccidioides brasiliensis* now comprised of *Paracoccidioides brasiliensis* and *Paracoccidioides lutzi*² and *Blastomyces dermatitidis* as the 2 species *Blastomyces dermatitidis* and *Blastomyces gilchristii*.³ From a clinical standpoint, management of these diseases does not rely on knowing the precise species. Thus, throughout this review, the dimorphic fungal pathogens will be referred to

by their genus name without regard to the specific taxonomic species designation. Both current antifungals and potential new drugs for treating infections by dimorphic fungal pathogens will be discussed.

Dimorphism and endemicity

The hallmark of the dimorphic fungal pathogens is their dual, yet distinct lifestyles. These lifestyles are correlated with separate fungal morphologies with filamentous growth (i.e., hyphae) characterizing the saprobic phase in the environment and yeasts (or spherules in the case of *Coccidioides*) characterizing the human pathogenic phase. This contrasts with polymorphic fungi (e.g. *Candida albicans*) which exhibit both yeast and hyphal morphologies in human tissues. Temperature appears to be the central factor that determines the form/lifestyle of the dimorphic fungi with exposure to mammalian body temperature triggering adoption of the pathogenic mode.^{4–7} Yeasts of the dimorphic fungi (or *Coccidioides* spherules) are not efficiently eliminated by immune cells. Preventing the transition into yeasts through pharmacologic^{8,9} or genetic interventions^{10–13} renders dimorphic fungal pathogens avirulent indicating the necessity of the dimorphic transition for pathogenesis. Unlike many parasites which must cycle between human and non-human

environments to complete their lifecycle, the dimorphic fungi have no such requirement for infection of mammalian hosts. Rather, infection of mammals by the dimorphic fungi is accidental, yet when these fungi transition to the yeast/spherule phase, they express efficient mechanisms that enable their survival and proliferation within this secondary environment.

The dimorphic fungal pathogens are endemic to particular geographic regions rather than being found ubiquitously in the environment. These areas have been largely defined by clinical case prevalence,¹⁴ isolation from soils,¹⁵ or skin-reactivity tests to specific antigens of the dimorphic fungi.^{16–18} Ecological aspects of soils that favor growth of the hyphal forms and/or animal or bird patterns for dispersal are thought to underlie dimorphic fungal pathogen endemicity. *Histoplasma* is endemic to the Midwest and Eastern parts of the United States (chiefly around the Ohio and Mississippi River valleys),¹⁶ and is also found in Latin America (particularly Brazil, Venezuela, Colombia, and Argentina;¹⁹), parts of Africa,²⁰ and some cases have been reported in China^{21,22} and India.¹⁹ *Blastomyces* overlaps with many of the *Histoplasma*-endemic regions in the United States (Midwest, Southeast, and South-central) and also includes Canadian provinces around the Great Lakes^{23,24} and parts of Africa.²⁵ In contrast to the moist environments with decaying organic material supporting *Histoplasma* and *Blastomyces*, *Coccidioides* is found in the more arid environments of the Southwest of the United States (Southern California, Arizona, New Mexico, Texas;²⁶), Central America, and parts of South America (Northern Brazil, Venezuela, Argentina, and Paraguay²⁷). *Paracoccidioides* species are endemic to South America, particularly areas of Brazil, Colombia, Venezuela, Argentina, and Uruguay.²⁸

The annual incidence of infections by dimorphic fungi is likely inaccurate as under-diagnosis and under-reporting of infections is common. Furthermore, most infections are self limiting without requiring clinical intervention. Nevertheless, estimates of over a half million infections by *Histoplasma* and *Coccidioides* occur each year.^{29,30} *Paracoccidioides* and *Blastomyces* infection estimates are more difficult due to regional differences, however reports suggest the incidence ranges from 0.05 to 3 per 100,000 individuals in endemic regions.^{23,31–33} Together these 4 dimorphic fungal pathogens infect hundreds of thousands each year, a number which does not include veterinary infections. The number of clinical cases typically range from 5–30% of infections, although in some outbreaks over 50% of infections can result in clinical disease.^{34–36} One survey of hospital records in the United States tallied

over 6000 hospitalizations in one year: 3360 cases of histoplasmosis, 771 cases of blastomycosis, and 2194 cases of coccidioidomycosis.¹⁴ Less than half of these were in immunocompromised individuals. A case series analysis of Latin America countries showed over 750 annual reported cases of paracoccidioidomycosis.²⁸

Dimorphic fungal pathogen disease and treatment

Inhalation of conidia initiates infection by the dimorphic fungi^{28,37–39} The conidia (or arthroconidia in the case of *Coccidioides*) are produced solely by the environmental mycelia. The small size of these propagules facilitates aerosolization upon environmental disturbance and deposition into alveoli once inhaled. Thus, occupational and recreational activities that contribute to soil and environmental disruption are a primary risk factor for infection by dimorphic fungal pathogens.^{39–42} Most infection outbreaks can be traced back to specific events that led to conidia release from soils and their subsequent inhalation.^{34,35,43–47} Therefore, practices to avoid or reduce the potential for aerosolization and inhalation of conidia are prescribed for high risk areas^{34,41,48,49}

Within the lungs, conidia transition into pathogenic-phase cells which survive innate immune defenses. For *Histoplasma*, *Blastomyces*, and *Paracoccidioides*, the conidia germinate into yeasts. The yeasts can be internalized by host phagocytes (e.g., alveolar macrophages) to varying degrees with *Histoplasma* yeasts residing almost exclusively within these host cells. This intracellular residence creates an additional barrier to antifungal drug penetration which must be considered in antifungal development. For *Coccidioides*, the inhaled arthroconidia become spherules that are exclusively extracellular. The spherules enlarge into endosporeulating structures within the lung tissue and release endospores for propagation of the infection. Whether phagocytosed or not, the dimorphic fungal pathogens are not controlled by the innate axis of the immune system unlike the opportunistic fungal pathogens. Thus, *Histoplasma*, *Blastomyces*, *Coccidioides* species, and *Paracoccidioides* infections are not restricted to immunocompromised hosts, but also cause disease in immunocompetent individuals. Control of the infection requires activation of CD4⁺ cells and consequently individuals lacking aspects of cellular immunity (e.g., HIV, immunosuppression due to tissue or organ transplantation, TNF α blockade, etc.) typically progress to severe and disseminated disease. Elimination of symptoms has been assumed to indicate clearance of the infection, but evidence is now suggesting that at least in some individuals, the infection can enter a latent state, which can re-emerge later when the balance between pathogen and host immunity is altered (e.g., immunosuppression of the host).^{50–56}

Since inhalation is the route of exposure, mycoses caused by dimorphic fungal pathogens are initially pulmonary diseases. In immunocompetent individuals, mild disease is mostly subclinical, often going undiagnosed. Infection causes varying degrees of pneumonia and influenza-like symptoms. For the majority of individuals, symptoms typically resolve without requiring intervention. Roughly 5% of *Histoplasma* infections are estimated to require clinical management⁵⁷ and up to 30% for *Coccidioides* infections.⁵⁸ For individuals inhaling a larger inoculum, or those that have some deficiency in cellular immune response, disease is more severe and the infection typically disseminates to extrapulmonary sites via the hemolymphatic system. Extrapulmonary disease presentation varies, but can include oral and pharyngeal mucosa (*Paracoccidioides*), cutaneous lesions (*Histoplasma*, *Blastomyces*, *Paracoccidioides*, and *Coccidioides*), and bone (*Blastomyces*).^{31,38–40,59} The systemic nature of infections by dimorphic fungi thus precludes any topical antifungal options for management, and the required systemic drug administration significantly elevates the potential for problems with host toxicities.

Current antifungal options

In vitro activities of current antifungal drugs used clinically have been established for the dimorphic fungi. These include drugs of the polyene, azole, and echinocandin classes. In general, reference methods for antimicrobial susceptibility tests follow those established by the Clinical and Laboratory Standards Institute (CLSI). Notably missing from these reference methods are procedures specifically designated for dimorphic fungi leading to confusion as to whether macro- and microdilution test methods for yeasts

(M27-A3)⁶⁰ or for filamentous fungi (M38-A2)⁶¹ should be followed. This is not an insignificant question as the yeast and hyphal forms of the dimorphic fungi can have dramatically different susceptibilities.^{62–69} For example, early studies with echinocandins showed antifungal effects on the hyphal form of *Histoplasma*.⁶⁷ However, subsequent tests against the yeast form indicated that caspofungin was anywhere from 20- to 1000-fold less effective.^{64,68,69} It is now well recognized that the echinocandins have poor antifungal activity on pathogenic phase cells of the dimorphic fungi. This situation underscores the need to use the form most relevant to human infection when establishing the antifungal susceptibility profile of dimorphic fungal pathogens. Unfortunately, some antifungal studies continue to ignore the pathogenic forms and instead limit studies to mycelial-phase cells.^{70–72} Further complicating antifungal susceptibility profiling are the lack of standardized methods addressing complications for dimorphic fungi. While the CLSI methods for yeasts (M27-A3)⁶⁰ work well for *Candida* and *Cryptococcus* yeasts, they are inadequate for testing yeast-phase cells of the dimorphic fungal pathogens, which have longer generation times and require higher inocula for efficient and consistent growth in broth culture. A more appropriate microdilution method has recently been established for antifungal testing of *Histoplasma* yeasts,⁶⁹ and this methodology produces more accurate data for *Histoplasma* susceptibility.⁷³

Despite these caveats, a general profile of the antimicrobial susceptibilities of dimorphic fungi has been developed for amphotericin B and azole-class drugs. Each of the dimorphic fungal pathogens exhibit a similar susceptibility profile for current antifungals with both amphotericin B and azole drugs generally showing potent activity (low minimum inhibitory concentrations; MIC) with some variability in MICs reported for fluconazole (Table 1). Unfortunately,

Table 1. In vitro antifungal MICs for dimorphic fungal pathogens.

Drug class	Antifungal	MIC range ($\mu\text{g/mL}$)				References
		<i>Histoplasma</i>	<i>Blastomyces</i>	<i>Paracoccidioides</i>	<i>Coccidioides</i>	
Polyenes	Amphotericin B	Y: <0.03–2.0 M: 0.26–2.5	Y: <0.03–2.0 M:	Y: 0.06–2.0 M:	Y: 0.25–2.0 M: 0.03–0.50	68,69,109–119
Imidazoles	Ketoconazole	Y: M: 0.17	Y: <0.01–0.25 M: 0.1–0.4	Y: <0.01–0.03 M:	Y: M: 0.03–0.16	72,110,114–116,120
Triazoles	Fluconazole	Y: 0.25–8.0 M: 2.0–32	Y: 0.06–32 M: 0.06–32	Y: 0.13–0.50 M:	Y: M: 2.0–64	68,69,109,110,112–114,116,117,119–123
	Itraconazole	Y: <0.01–0.5 M: 0.03–1.0	Y: <0.01–0.13 M: 0.03–4	Y: <0.01–0.06 M:	Y: <0.03–0.50 M: 0.03–1.0	68,109–115,120–122
	Voriconazole	Y: 0.03–0.50 M: <0.01–2.0	Y: <0.03–0.25 M: 0.06–2.0	Y: M:	Y: <0.03–2.0 M: 0.03–1.0	109–111,113,121,123
	Posaconazole	Y: <0.01–0.50 M: 0.02–2.0	Y: <0.02–0.06 M: <0.02–2.0	Y: M:	Y: M: 0.06–1.0	67,109,112,121,123
Echinocandins	Micafungin	Y: >64 M: 0.03–0.06	Y: 32–64 M: <0.01–0.03	Y: >64 M: 4–16	Y: M: 0.02	68
	Caspofungin	Y: 8–32 ^a M: 0.02–4.0	Y: M: 0.5–8.0	Y: M:	Y: M: 8–64	67,69,113,117,118

Note. ^a2 studies reported low caspofungin MICs for yeast which disagree with the majority of studies based on clinical isolates in India (0.03–1.0 $\mu\text{g/mL}$)¹⁰⁹ and for a single laboratory *Histoplasma* strain (MIC <0.125 $\mu\text{g/mL}$).¹²²

the echinocandins, which have dramatically decreased host toxicity, have poor antifungal activity against the pathogenic-phase cells of dimorphic fungal pathogen yeast cells. The reasons underlying this natural resistance of yeasts to the β -glucan synthase inhibitors is currently unknown.

The in vitro antifungal susceptibilities for amphotericin B and azole-class drugs have been validated in murine models of dimorphic fungal disease and/or in clinical trials. As a consequence, the Infectious Disease Society of America has released treatment guidelines for infections with *Histoplasma*,⁷⁴ *Blastomyces*,⁷⁵ and *Coccidioides*.⁷⁶ Despite its potential for host toxicity, amphotericin B is recommended for severe disseminated disease, with one of the liposomal formulations preferred. For less severe situations, for follow up after amphotericin B, or for possible prophylaxis of highly susceptible individuals (e.g. AIDS), IDSA guidelines recommend treatment with an oral azole (i.e., itraconazole) with monitoring of serum concentrations to ensure sufficient absorption and bioavailability. To ensure sufficient clearance of the dimorphic fungal infection, treatments typically involve protracted regimens. For mild disease, treatment durations can range from several months to a year depending on the specific dimorphic fungus. Treatment of disseminated disease and disease in immunocompromised hosts can be a year or longer. Beyond the prolonged treatment times, the potential for latency, rather than clearance of the infection, further complicates antifungal management of dimorphic fungal infections.

New and alternative antifungal developments

With the known host toxicity risks of current antifungals,⁷⁷⁻⁷⁹ and the endogenous resistance of dimorphic fungi to the lower toxicity echinocandins, new and alternative antifungal drug options have been explored. The eukaryotic nature shared by both the dimorphic fungal pathogens and the host is recognized as a significant obstacle to antifungal drug development. Consequently, many molecules with promising antifungal activity fail due to lack of adequate selectivity.

One approach to improve the selectivity of existing antifungal molecules, is to test modifications to the structure that decrease drug affinity to host targets. The oral availability of azole-class antifungals is a great benefit to management of dimorphic fungal infections. However, the heme iron-binding action of the imidazoles and triazoles that inhibits fungal cytochrome CYP51 (Erg11) also targets host P450 cytochromes leading to hepatotoxicity and problematic drug-drug interactions. Exploration of other metal binding azole groups in combination with structure-guided improvement of affinity for fungal CYP51 outside of the active site led to the design of compound VT-1161.⁸⁰ As predicted from structure affinity calculations, the molecule has

low affinity for mammalian CYP3A4 but is highly potent against *C. albicans*. Subsequent testing against *Coccidioides* arthroconidia in vitro showed VT-1161 has an MIC around 1.5 $\mu\text{g/mL}$.⁸¹ Oral administration of VT-1161 in a murine model of coccidioidomycosis demonstrated VT-1161 reduced spherule burden in the lungs by 100- to 1000-fold, leading to improved mouse survival.⁸¹ While VT-1161 had similar effectiveness as fluconazole in vivo, it is hoped the better selectivity of the molecules will decrease potential host toxicity.

Following the goal of reduced host toxicity, some antibacterial compounds have been investigated due to their established safety profile to the host. Drugs targeting the folate pathway have been effective against microbes from bacteria to parasites,⁸² are orally available, and have a moderately safe host toxicity profile. These drugs have already been a means of prophylaxis against *Pneumocystis* fungal infections in HIV+ individuals. Sulfonamides in combination with dihydrofolate reductase inhibitors (e.g., cotrimoxazole) have activity against *Histoplasma* yeasts (8–14 $\mu\text{g/mL}$ and 2–3 $\mu\text{g/mL}$, respectively)⁶⁵ and *Coccidioides* (MICs of 1000 $\mu\text{g/mL}$ and 200 $\mu\text{g/mL}$, respectively),⁶⁶ although only the mycelial phase has been tested for *Coccidioides*. The sulfonamides/antifolate combination has been one of the traditional treatments for *Paracoccidioides*^{59,83-85} but they have not been advanced clinically for the other dimorphic fungal pathogens.

Using a similar rationale, ciprofloxacin, a fluoroquinolone antibacterial with very low host toxicity, has been tested for antifungal potential. The fluoroquinolones inhibit bacterial type II topoisomerases (e.g., DNA gyrase) and were suggested to potentially also inhibit fungal topoisomerase. While not effective against mycelial forms of *Histoplasma* and *Coccidioides*, tests against *Histoplasma* yeast showed ciprofloxacin has an in vitro MIC of 62.5–250 $\mu\text{g/mL}$.⁸⁶ While not considered a low MIC, the lack of significant host toxicity for ciprofloxacin creates a good selectivity index and raises its potential, at least as an adjunct therapy for fungal infections. Addition of ciprofloxacin to amphotericin B or itraconazole indicated some additive effects could be gained through drug combinations.

Anti-tuberculosis drugs show some, albeit low potency, antifungal activity for dimorphic fungal pathogens. When tested in vitro against *Coccidioides* mycelia, rifampicin, isoniazid, and ethambutol had very high MICs (8500 $\mu\text{g/mL}$, 500 $\mu\text{g/mL}$, and 2500 $\mu\text{g/mL}$, respectively⁷²). Effectiveness against spherules was not determined but tests against *Histoplasma* yeasts similarly had high MICs⁷¹ arguing these antibacterial compounds lack any significant antifungal utility. However, using isoniazid as the core, chemical structure modification improved its antifungal potential; 3 of 9 isoniazid derivatives had significantly lower MICs against

Histoplasma yeasts with one having inhibitory activity at concentrations as low as 7.8 $\mu\text{g/mL}$.⁶² Extending the in vitro tests of the isoniazid-hydrazone derivative to *Coccidioides* demonstrated it had an MIC of 50 $\mu\text{g/mL}$ against mycelia of this dimorphic fungal pathogen.⁸⁷ Although *Coccidioides* spherules were not tested, there is a reasonable possibility that the MIC will be lower since *Histoplasma* yeasts were 8–16X more sensitive to the isoniazid-hydrazone than were mycelia.⁶²

Antiretroviral protease inhibitors have inhibitory activity against *Histoplasma*. Saquinavir is active against both filamentous and yeast forms with an MIC around 0.4 $\mu\text{g/mL}$.⁶³ Another protease inhibitor, ritonavir, has an MIC of 1.0 $\mu\text{g/mL}$ against *Histoplasma* mycelia but is 7-fold more potent against yeast cells. As HIV+ individuals comprise a significant portion of clinical *Histoplasma* cases, this raises the possibility that management of HIV through protease inhibitor cocktails could simultaneously directly combat infection by dimorphic fungal pathogens.

Surprisingly, farnesol can inhibit growth and morphological transitions of *Paracoccidioides* and *Coccidioides*. Farnesol is an alcohol present in some essential oils but is also produced by *C. albicans* as a quorum sensing molecule. When tested on *Paracoccidioides* yeasts, farnesol had an MIC of 30 μM .⁸⁸ Farnesol also inhibited the transition from yeast cells to hyphae, although this morphological defect has questionable relevance to the situation in mammalian hosts. Against *Coccidioides* mycelia, farnesol was nearly 100-fold more potent (MIC approximately 0.3 μM ; ⁷⁰). Farnesol was also reported to inhibit *Histoplasma* yeast growth with an average MIC of 0.02 μM .⁸⁹ As a caution, tests with *Histoplasma* yeasts followed the CLSI methodology (M27-A3) and this has been shown to be inferior for reliable testing of *Histoplasma* yeast susceptibility.⁶⁹

Repurposing of anti-cancer drugs has also been fruitful in the search for new antifungal prospects. AR-12 is a celecoxib-derivative that has been shown to have antimicrobial properties against microbes ranging from bacteria to parasites to fungi and even viruses.^{90–94} Early mechanistic studies in cancer cells suggested AR-12 inhibited cellular phosphoinositide-dependent kinase-1 (Pdk1), however this does not appear to be the case in fungi. Recently, it was shown that AR-12 inhibits acetyl-CoA synthetase, an essential enzyme in fungi.⁹⁵ AR-12 has antifungal activity against a broad collection of fungi, including the dimorphic fungal pathogens at concentrations around 4–8 $\mu\text{g/mL}$.⁷³ In vitro, AR-12 is fungicidal against *Histoplasma* yeasts (Rapple CA, personal communication). These antifungal characteristics combined with AR-12s safety as established in Phase I trials make AR-12 an attractive antifungal candidate for further development.

Both phenotypic and target-based screening approaches have been used to identify novel compounds that inhibit dimorphic fungal pathogens. Genome analyses for potential antifungal targets focused on those genes that were (i) demonstrated to be essential for fungal growth and (ii) conserved among diverse fungi (Basidiomycetes to Ascomycetes;⁹⁶). From this thioredoxin reductase (Trr) was selected as a candidate target for computational screening of a library of 3000 compounds. Three compounds were identified and tests against purified *Paracoccidioides* Trr protein confirmed the compound inhibited Trr enzymatic activity.⁹⁷ Phenotype tests demonstrated that 2 compounds had significant biological activity against *Paracoccidioides* yeasts with MICs of 8–16 $\mu\text{g/mL}$ (compounds F1806-0122 and F3307-0100). This study validates target based screening as an approach to identification of potential antifungal compounds.

Phenotypic screening of small molecule libraries has identified a group of aminothiazoles with antifungal activity toward *Histoplasma*. To enable rapid quantitative screening of yeast growth, yeasts were engineered to express a red-fluorescent protein, which could be used as a surrogate indicator of the number of yeast cells.⁹⁸ Furthermore, fluorescence-based growth analysis facilitated efficient monitoring of *Histoplasma* within macrophages, and thus the determination of the effectiveness of compounds on inhibition of *Histoplasma* within its host cell environment. The top hit aminothiazole (41F5) was effective against yeasts in culture (MIC 2–4 μM) and effectively inhibited proliferation of yeasts resident within macrophages.⁹⁸ Importantly, the aminothiazole compound had low toxicity to cultured mammalian cells, including macrophages. Unfortunately, the aminothiazole was not active against *Blastomyces* yeasts despite the very close phylogenetic relationship between *Histoplasma* and *Blastomyces*. The molecular target of the aminothiazole remains to be determined, and this may facilitate structure optimization to broaden its activity for other dimorphic fungal pathogens.

The most advanced new antifungal option for clinical development to manage dimorphic fungal pathogens is nikkomycin Z. The polyoxins (from which the nikkomyces are derived) are peptide modified-nucleoside analogs which were originally identified by screening *Streptomyces* products for antifungal and insecticidal activities.⁹⁹ These compounds were later shown to be inhibitors of chitin synthesis.¹⁰⁰ Since chitin synthase is absent from mammalian cells, these compounds have very high selectivity for fungi. Following polyoxin studies on *C. albicans*, the nikkomyces were demonstrated to have good potency in vitro against dimorphic fungal pathogens,^{101–103} with the most potency against *Coccidioides* cells (MIC of 0.125 $\mu\text{g/mL}$ for spherules).¹⁰³ Examination of treated *Coccidioides* spherules showed lack of endosporulation and even lysis

of the spherules consistent with impairments in cell wall structure.¹⁰³ The nikkomycins are orally available, simplifying extension of studies to animal models of mycoses caused by dimorphic fungal pathogens. Administration of nikkomycin Z in murine models of lethal histoplasmosis resulted in decreased organ fungal burdens (3- to 28-fold) and a concomitant enhancement of mouse survival at doses from 20 to 100 mg/kg per day.¹⁰²⁻¹⁰⁴ Significant protection from lethal infection was realized, however, only at moderate inocula suggesting nikkomycin antifungal activity may not be sufficiently rapid to curb severe infections. Similar results were found for treatment of blastomycosis with reduction in fungal burdens ranging from 10-fold (at a dose of 50 mg/kg)¹⁰³ to apparent lung tissue sterilization (over 3500-fold reduction at higher doses of 400–1000 mg/kg).¹⁰⁵ The ability to safely administer such doses is a testament to the high selectivity gained by targeting the fungi-specific molecule chitin synthase. Nikkomycin Z is also effective in treating pulmonary coccidioidomycosis with over 10,000-fold reduction in lung fungal burdens and complete protection from lethal infection of mice at a dose of 20 mg/kg.¹⁰³ Recently, a small clinical trial in dogs with *Coccidioides* infection was completed. Dogs with mild to moderate disease had significant improvement but those with severe disease had little or poor responses to treatment, again suggesting nikkomycin may not act rapidly enough to quickly curtail disease progression in high fungal burden situations. Nonetheless, nikkomycin Z shows good promise as a highly selective (low host toxicity) antifungal for development against dimorphic fungal pathogen infections.

Conclusions

Current clinical management of infections by dimorphic fungal pathogens is limited to azole-class antifungal drugs and amphotericin B. While orally available, the azoles are not without host toxicity issues and the treatment course is lengthy for infections by dimorphic fungi. Development of resistance to azoles is not widespread, although treatment failures due to azole resistance have occurred.¹⁰⁶⁻¹⁰⁸ Unfortunately, the better tolerated echinocandin antifungals lack efficacy against the pathogenic-phase of the dimorphic fungal pathogens raising the need for alternative or second-line treatment options.

While a number of strides have been made in repurposing existing drugs and development of new inhibitors of fungal growth, careful attention must be paid to challenges posed by dimorphic fungi. As the yeasts/spherules of the dimorphic fungi are the state present within the mammalian host, antimicrobial susceptibilities need to be performed with these pathogenic-phase cells, not the mycelia which has led to erroneous conclusions. Testing

of the pathogenic-phase in vitro should follow recently optimized procedures as the CLSI methodology for yeasts is inadequate for the dimorphic fungi. Since yeast cells of the dimorphic fungi reside within host phagocytes, it is also advisable for in vitro tests to be followed with tests on drug effectiveness on intracellular yeasts, at least during initial drug development stages.

The overall selectivity of antifungal drug candidates is critical for progression of drugs through the development pipeline. Structure-guided rational design is one approach that has improved the selectivity of an azole structure (VT-1161). Many of the repurposed drugs have relatively high MICs (greater than 100 ug/mL) questioning their therapeutic utility, however if their selectivity is sufficiently high, formulations may be developed to facilitate sufficiently high serum and tissue levels. Lower MICs have been found for drugs targeting the folate pathway, an isoniazid-hydrazone derivative, antiretroviral protease inhibitors, and the anti-cancer drug AR-12, all of which are expected to be reasonably well-tolerated by the mammalian host. Novel drugs with good in vitro MICs and good selectivity include thioredoxin-reductase inhibitors, an aminothiazole compound, and nikkomycin Z. Since nikkomycin Z targets an enzyme absent from the host, nikkomycin has an excellent basis for high selectivity for fungi. In addition, nikkomycin Z has maintained antifungal effectiveness against multiple dimorphic fungal pathogens in animal models of disease. While the current antifungal armament is limited, there are exciting prospects on the horizon for treating dimorphic fungal infections.

Disclosure of potential conflicts of interest

No potential conflicts of interest were disclosed.

Funding

Studies on susceptibility to established drugs and development of new antifungals was supported, in part, by grant R21-AI109437 from the National Institute of Allergy and Infectious Diseases (NIAID).

ORCID

Chad A. Rappeye  <http://orcid.org/0000-0001-7880-5958>

References

- [1] Fisher MC, Koenig GL, White TJ, Taylor JW. Molecular and phenotypic description of *Coccidioides posadasii* sp. nov., previously recognized as the non-California population of *Coccidioides immitis*. *Mycologia* 2002; 94:73–84; PMID:21156479; <http://dx.doi.org/10.2307/3761847>

- [2] Teixeira MM, Theodoro RC, de Carvalho MJA, Fernandes L, Paes HC, Hahn RC, Mendoza L, Bagagli E, San-Blas G, Felipe MSS. Phylogenetic analysis reveals a high level of speciation in the *Paracoccidioides* genus. *Mol Phylogenet Evol* 2009; 52:273-83; PMID:19376249; <http://dx.doi.org/10.1016/j.ympev.2009.04.005>
- [3] Brown EM, McTaggart LR, Zhang SX, Low DE, Stevens DA, Richardson SE. Phylogenetic analysis reveals a cryptic species *Blastomyces gilchristii*, sp. nov. within the human pathogenic fungus *Blastomyces dermatitidis*. *PLoS One* 2013; 8:e59237; PMID:23533607; <http://dx.doi.org/10.1371/journal.pone.0059237>
- [4] Horwath MC, Fecher RA, Deepe GS. *Histoplasma capsulatum*, lung infection and immunity. *Future Microbiol* 2015; 10:967-75; PMID:26059620; <http://dx.doi.org/10.2217/fmb.15.25>
- [5] Klein BS, Newman SL. Role of cell-surface molecules of *Blastomyces dermatitidis* in host-pathogen interactions. *Trends Microbiol* 1996; 4:246-51; PMID:8795162; [http://dx.doi.org/10.1016/0966-842X\(96\)10028-7](http://dx.doi.org/10.1016/0966-842X(96)10028-7)
- [6] Kellner EM, Orsborn KI, Siegel EM, Mandel MA, Orbach MJ, Galgiani JN. *Coccidioides posadasii* contains a single 1,3- β -glucan synthase gene that appears to be essential for growth. *Eukaryotic Cell* 2005; 4:111-20; PMID:15643067; <http://dx.doi.org/10.1128/EC.4.1.111-120.2005>
- [7] Abreu e Silva MÀ de, Salum FG, Figueiredo MA, Cherubini K. Important aspects of oral paracoccidioidomycosis—a literature review. *Mycoses* 2013; 56:189-99; PMID:23088400; <http://dx.doi.org/10.1111/myc.12017>
- [8] Medoff G, Kobayashi GS, Painter A, Travis S. Morphogenesis and pathogenicity of *Histoplasma capsulatum*. *Infect Immun* 1987; 55:1355-8; PMID:3032799
- [9] Medoff G, Sacco M, Maresca B, Schlessinger D, Painter A, Kobayashi GS, Carratu L. Irreversible block of the mycelial-to-yeast phase transition of *Histoplasma capsulatum*. *Science* 1986; 231:476-9; PMID:3001938; <http://dx.doi.org/10.1126/science.3001938>
- [10] Nemecek JC, Wüthrich M, Klein BS. Global control of dimorphism and virulence in fungi. *Science* 2006; 312:583-8; PMID:16645097; <http://dx.doi.org/10.1126/science.1124105>
- [11] Nguyen VQ, Sil A. Temperature-induced switch to the pathogenic yeast form of *Histoplasma capsulatum* requires Ryp1, a conserved transcriptional regulator. *Proc Natl Acad Sci U S A* 2008; 105:4880-5; PMID:18339808; <http://dx.doi.org/10.1073/pnas.0710448105>
- [12] Beyhan S, Gutierrez M, Voorhies M, Sil A. A temperature-responsive network links cell shape and virulence traits in a primary fungal pathogen. *PLoS Biol* 2013; 11:e1001614; PMID:23935449; <http://dx.doi.org/10.1371/journal.pbio.1001614>
- [13] Boyce KJ, Schreider L, Kirszenblat L, Andrianopoulos A. The two-component histidine kinases DrkA and SlnA are required for in vivo growth in the human pathogen *Penicillium marneffei*. *Mol Microbiol* 2011; 82:1164-84; PMID:22059885; <http://dx.doi.org/10.1111/j.1365-2958.2011.07878.x>
- [14] Chu JH, Feudtner C, Heydon K, Walsh TJ, Zaoutis TE. Hospitalizations for endemic mycoses: a population-based national study. *Clin Infect Dis* 2006; 42:822-5; PMID:16477560; <http://dx.doi.org/10.1086/500405>
- [15] Ajello L. Coccidioidomycosis and histoplasmosis. A review of their epidemiology and geographical distribution. *Mycopathol Mycol Appl* 1971; 45:221-30; PMID:4950673; <http://dx.doi.org/10.1007/BF02051969>
- [16] Edwards LB, Acquaviva FA, Livesay VT, Cross FW, Palmer CE. An atlas of sensitivity to tuberculin, PPD-B, and histoplasmin in the United States. *Am Rev Respir Dis* 1969; 99(1-132); PMID:5767603
- [17] Mochi A, Edwards PQ. Geographical distribution of histoplasmosis and histoplasmin sensitivity. *Bull World Health Organ* 1952; 5:259-91; PMID:14935779
- [18] Kauffman CA. Histoplasmosis: a clinical and laboratory update. *Clin Microbiol Rev* 2007; 20:115-32; PMID:17223625; <http://dx.doi.org/10.1128/CMR.00027-06>
- [19] Bahr NC, Antinori S, Wheat LJ, Sarosi GA. Histoplasmosis infections worldwide: thinking outside of the Ohio River valley. *Curr Trop Med Rep* 2015; 2:70-80; PMID:26279969; <http://dx.doi.org/10.1007/s40475-015-0044-0>
- [20] Gugnani HC. Histoplasmosis in Africa: a review. *Indian J Chest Dis Allied Sci* 2000; 42:271-7; PMID:15597674
- [21] Zhao B, Xia X, Yin J, Zhang X, Wu E, Shi Y, Tong M, Sun S, Gui X, Zhao Z, et al. Epidemiological investigation of *Histoplasma capsulatum* infection in China. *Chin Med J* 2001; 114:743-6; PMID:11780341
- [22] Pan B, Chen M, Pan W, Liao W. Histoplasmosis: a new endemic fungal infection in China? Review and analysis of cases. *Mycoses* 2013; 56:212-21; PMID:23216676; <http://dx.doi.org/10.1111/myc.12029>
- [23] Smith JA, Gauthier G. New Developments in Blastomycosis. *Semin Respir Crit Care Med* 2015; 36:715-28; PMID:26398538; <http://dx.doi.org/10.1055/s-0035-1562898>
- [24] Klein BS, Vergeront JM, Davis JP. Epidemiologic aspects of blastomycosis, the enigmatic systemic mycosis. *Semin Respir Infect* 1986; 1:29-39; PMID:3317597
- [25] Carman WF, Frean JA, Crewe-Brown HH, Culligan GA, Young CN. Blastomycosis in Africa. *Mycopathologia* 1989; 107:25-32; PMID:2682252; <http://dx.doi.org/10.1007/BF00437587>
- [26] Edwards PQ, Palmer CE. Prevalence of sensitivity to coccidioidin, with special reference to specific and non-specific reactions to coccidioidin and to histoplasmin. *Dis Chest* 1957; 31:35-60; PMID:13384171; <http://dx.doi.org/10.1378/chest.31.1.35>
- [27] Laniado-Laborin R. Expanding understanding of epidemiology of coccidioidomycosis in the Western hemisphere. *Ann N Y Acad Sci* 2007; 1111:19-34; PMID:17395731; <http://dx.doi.org/10.1196/annals.1406.004>
- [28] Martinez R. Epidemiology of paracoccidioidomycosis. *Rev Inst Med Trop Sao Paulo* 2015; 57(19):11-20; PMID:26465364; <http://dx.doi.org/10.1590/S0036-46652015000700004>
- [29] Hammerman KJ, Powell KE, Tosh FE. The incidence of hospitalized cases of systemic mycotic infections. *Med Mycol* 1974; 12:33-45; <http://dx.doi.org/10.1080/00362177485380061>
- [30] Stevens DA. Coccidioidomycosis. *N Engl J Med* 1995; 332:1077-82; PMID:7898527; <http://dx.doi.org/10.1056/NEJM199504203321607>
- [31] Saccente M, Woods GL. Clinical and laboratory update on blastomycosis. *Clin Microbiol Rev* 2010; 23:367-81; PMID:20375357; <http://dx.doi.org/10.1128/CMR.00056-09>

- [32] Restrepo A, McEwen JG, Castañeda E. The habitat of *Paracoccidioides brasiliensis*: how far from solving the riddle? *Med Mycol* 2001; 39:233-41; PMID:11446526; <http://dx.doi.org/10.1080/714031028>
- [33] Colombo AL, Tobón A, Restrepo A, Queiroz-Telles F, Nucci M. Epidemiology of endemic systemic fungal infections in Latin America. *Med Mycol* 2011; 49:785-98; PMID:21539506
- [34] Cummings KC, McDowell A, Wheeler C, McNary J, Das R, Vugia DJ, Mohle-Boetani JC. Point-source outbreak of coccidioidomycosis in construction workers. *Epidemiol Infect* 2010; 138:507-11; PMID:19845993; <http://dx.doi.org/10.1017/S0950268809990999>
- [35] Chamany S, Mirza SA, Fleming JW, Howell JF, Lenhart SW, Mortimer VD, Phelan MA, Lindsley MD, Iqbal NJ, Wheat LJ, et al. A large histoplasmosis outbreak among high school students in Indiana, 2001. *Pediatr Infect Dis J* 2004; 23:909-14; PMID:15602189; <http://dx.doi.org/10.1097/01.inf.0000141738.60845.da>
- [36] Klein BS, Vergeront JM, Weeks RJ, Kumar UN, Mathai G, Varkey B, Kaufman L, Bradsher RW, Stoebig JF, Davis JP. Isolation of *Blastomyces dermatitidis* in soil associated with a large outbreak of blastomycosis in Wisconsin. *N Engl J Med* 1986; 314:529-34; PMID:3945290; <http://dx.doi.org/10.1056/NEJM198602273140901>
- [37] Kauffman CA. Histoplasmosis. *Clin Chest Med* 2009; 30:217-25; PMID:19375629; <http://dx.doi.org/10.1016/j.ccm.2009.02.002>
- [38] Stockamp NW, Thompson GR. Coccidioidomycosis. *Infect Dis Clin North Am* 2016; 30:229-46; PMID:26739609; <http://dx.doi.org/10.1016/j.idc.2015.10.008>
- [39] Castillo CG, Kauffman CA, Miceli MH. Blastomycosis. *Infect Dis Clin North Am* 2016; 30:247-64; PMID:26739607; <http://dx.doi.org/10.1016/j.idc.2015.10.002>
- [40] Wheat LJ, Azar MM, Bahr NC, Spec A, Relich RF, Hage C. Histoplasmosis. *Infect Dis Clin North Am* 2016; 30:207-27; PMID:26897068; <http://dx.doi.org/10.1016/j.idc.2015.10.009>
- [41] Johnson L, Gaab EM, Sanchez J, Bui PQ, Nobile CJ, Hoyer KK, Peterson MW, Ojcius DM. Valley fever: danger lurking in a dust cloud. *Microbes Infect* 2014; 16:591-600; PMID:25038397; <http://dx.doi.org/10.1016/j.micinf.2014.06.011>
- [42] Twarog M, Thompson GR. Coccidioidomycosis: Recent updates. *Semin Respir Crit Care Med* 2015; 36:746-55; PMID:26398540; <http://dx.doi.org/10.1055/s-0035-1562900>
- [43] Carlos WG, Rose AS, Wheat LJ, Norris S, Sarosi GA, Knox KS, Hage CA. Blastomycosis in Indiana: digging up more cases. *Chest* 2010; 138:1377-82; PMID:20558552; <http://dx.doi.org/10.1378/chest.10-0627>
- [44] Benedict K, Mody RK. Epidemiology of Histoplasmosis Outbreaks, United States, 1938-2013. *Emerging Infect Dis* 2016; 22:370-8; PMID:26890817; <http://dx.doi.org/10.3201/eid2203.151117>
- [45] Schneider E, Hajjeh RA, Spiegel RA, Jibson RW, Harp EL, Marshall GA, Gunn RA, McNeil MM, Pinner RW, Baron RC, et al. A coccidioidomycosis outbreak following the Northridge, Calif, earthquake. *JAMA* 1997; 277:904-8; PMID:9062329; <http://dx.doi.org/10.1001/jama.1997.03540350054033>
- [46] Crum N, Lamb C, Utz G, Amundson D, Wallace M. Coccidioidomycosis outbreak among United States Navy SEALs training in a *Coccidioides immitis*-endemic area—Coalinga, California. *J Infect Dis* 2002; 186:865-8; PMID:12198626; <http://dx.doi.org/10.1086/342409>
- [47] Wheat LJ, Slama TG, Eitzen HE, Kohler RB, French ML, Biesecker JL. A large urban outbreak of histoplasmosis: clinical features. *Ann Intern Med* 1981; 94:331-7; PMID:7224378; <http://dx.doi.org/10.7326/0003-4819-94-3-331>
- [48] Das R, McNary J, Fitzsimmons K, Dobraca D, Cummings K, Mohle-Boetani J, Wheeler C, McDowell A, Iossifova Y, Bailey R, et al. Occupational coccidioidomycosis in California: outbreak investigation, respirator recommendations, and surveillance findings. *J Occup Environ Med* 2012; 54:564-71; PMID:22504958; <http://dx.doi.org/10.1097/JOM.0b013e3182480556>
- [49] Histoplasmosis - Protecting Workers at Risk. DHHS (NIOSH) Publication 2004; No. 2005-109. Available from: <https://www.cdc.gov/niosh/docs/2005-109/>
- [50] Jain VV, Evans T, Peterson MW. Reactivation histoplasmosis after treatment with anti-tumor necrosis factor α in a patient from a nonendemic area. *Respir Med* 2006; 100:1291-3; PMID:16545951; <http://dx.doi.org/10.1016/j.rmed.2005.09.020>
- [51] Vergidis P, Avery RK, Wheat LJ, Dotson JL, Assi MA, Antoun SA, Hamoud KA, Burdette SD, Freifeld AG, McKinsey DS, et al. Histoplasmosis complicating tumor necrosis factor- α blocker therapy: a retrospective analysis of 98 cases. *Clin Infect Dis* 2015; 61:409-17; PMID:25870331; <http://dx.doi.org/10.1093/cid/civ299>
- [52] Arbeloa-Gutierrez L, Kuberski T, Johnson SM, Sagastibelza I, Alaez JI, Pappagianis D. Reactivation of coccidioidomycosis: a prosthetic joint infection in Spain. *Eur J Clin Microbiol Infect Dis* 2016; 35:183-6; PMID:26638217; <http://dx.doi.org/10.1007/s10096-015-2526-3>
- [53] Nakelchik M, Mangino JE. Reactivation of histoplasmosis after treatment with infliximab. *Am J Med* 2002; 112:78; PMID:11812415; [http://dx.doi.org/10.1016/S0002-9343\(01\)00945-7](http://dx.doi.org/10.1016/S0002-9343(01)00945-7)
- [54] Van Koeveeringe MP, Brouwer RE. Histoplasma capsulatum reactivation with haemophagocytic syndrome in a patient with chronic lymphocytic leukaemia. *Neth J Med* 2010; 68:418-21; PMID:21209468
- [55] Ankobiah WA, Vaidya K, Powell S, Carrasco M, Allam A, Chechani V, Kamholz SL. Disseminated histoplasmosis in AIDS. Clinicopathologic features in seven patients from a non-endemic area. *N Y State J Med* 1990; 90:234-8; PMID:2348942
- [56] Keckich DW, Blair JE, Vikram HR, Seville MT, Kusne S. Reactivation of coccidioidomycosis despite antifungal prophylaxis in solid organ transplant recipients. *Transplantation* 2011; 92:88-93; PMID:21516063; <http://dx.doi.org/10.1097/TP.0b013e31821c1df6>
- [57] Hage CA, Azar MM, Bahr N, Loyd J, Wheat LJ. Histoplasmosis: up-to-date evidence-based approach to diagnosis and management. *Semin Respir Crit Care Med* 2015; 36:729-45; PMID:26398539; <http://dx.doi.org/10.1055/s-0035-1562899>
- [58] Thompson GR. Pulmonary coccidioidomycosis. *Semin Respir Crit Care Med* 2011; 32:754-63; PMID:22167403; <http://dx.doi.org/10.1055/s-0031-1295723>

- [59] Marques SA. Paracoccidioidomycosis. *Clin Dermatol* 2012; 30:610-5; PMID:23068148; <http://dx.doi.org/10.1016/j.clindermatol.2012.01.006>
- [60] Clinical and Laboratory Standards Institute. Reference method for broth dilution antifungal susceptibility testing of yeasts; approved standard—3rd ed. CLSI document M27-A3. 3rd ed. Wayne PA: Clinical and Laboratory Standards Institute 2008.
- [61] Clinical and Laboratory Standards Institute. Reference method for broth dilution antifungal susceptibility testing of filamentous fungi; approved standard—2nd ed. CLSI document M38-A2. 2nd ed. Wayne, PA: Clinical and Laboratory Standards Institute 2008.
- [62] De Aguiar Cordeiro R, de Farias Marques FJ, de Aguiar Cordeiro R, da Silva MR, Donato Maia Malaquias A, Silva de Melo CV, Mafezoli J, Ferreira de Oliveira M da C, Nogueira Brilhante RS, Gadelha Rocha MF, et al. Synthesis and antifungal activity in vitro of isoniazid derivatives against *histoplasma capsulatum* var. *capsulatum*. *Antimicrob Agents Chemother* 2014; 58:2504-11; PMID:24514090; <http://dx.doi.org/10.1128/AAC.01654-13>
- [63] Brilhante RSN, Caetano ÉP, Riello GB, Guedes GM de M, Castelo-Branco D de SCM, Fechine MAB, Oliveira JS de, Camargo ZP de, Mesquita JRL de, Monteiro AJ, et al. Antiretroviral drugs saquinavir and ritonavir reduce inhibitory concentration values of itraconazole against *Histoplasma capsulatum* strains in vitro. *Braz J Infect Dis* 2016; 20:155-9; PMID:26748233; <http://dx.doi.org/10.1016/j.bjid.2015.11.003>
- [64] Hage CA, Connolly P, Horan D, Durkin M, Smedema M, Zarnowski R, Smith P, Wheat LJ. Investigation of the efficacy of micafungin in the treatment of histoplasmosis using two North American strains of *Histoplasma capsulatum*. *Antimicrob Agents Chemother* 2011; 55:4447-50; PMID:21670186; <http://dx.doi.org/10.1128/AAC.01681-10>
- [65] Brilhante RSN, Fechine MAB, Cordeiro R de A, Rocha MFG, Ribeiro JF, Monteiro AJ, de Lima RAC, Mesquita JRL, de Camargo ZP, Sidrim JJC. In vitro effect of sulfamethoxazole-trimethoprim against *Histoplasma capsulatum* var. *capsulatum*. *Antimicrob Agents Chemother* 2010; 54:3978-9; PMID:20566767; <http://dx.doi.org/10.1128/AAC.00793-10>
- [66] Cordeiro R de A, Astete-Medrano DJ, Marques FJ de F, Andrade HTL, Perdigão Neto LV, Tavares JL, de Lima RAC, Patoilo KKNR, Monteiro AJ, Brilhante RSN, et al. Cotrimoxazole enhances the in vitro susceptibility of *Coccidioides posadasii* to antifungals. *Mem Inst Oswaldo Cruz* 2011; 106:1045-8; PMID:22241131; <http://dx.doi.org/10.1590/S0074-02762011000800024>
- [67] Espinel-Ingroff A. Comparison of In vitro activities of the new triazole SCH56592 and the echinocandins MK-0991 (L-743,872) and LY303366 against opportunistic filamentous and dimorphic fungi and yeasts. *J Clin Microbiol* 1998; 36:2950-6; PMID:9738049
- [68] Nakai T, Uno J, Ikeda F, Tawara S, Nishimura K, Miyaji M. In vitro antifungal activity of Micafungin (FK463) against dimorphic fungi: comparison of yeast-like and mycelial forms. *Antimicrob Agents Chemother* 2003; 47:1376-81; PMID:12654673; <http://dx.doi.org/10.1128/AAC.47.4.1376-1381.2003>
- [69] Goughenour KD, Balada-Llasat JM, Rappleye CA. Quantitative microplate-based growth assay for determination of antifungal susceptibility of *histoplasma capsulatum* yeasts. *J Clin Microbiol* 2015; 53:3286-95; PMID:26246483; <http://dx.doi.org/10.1128/JCM.00795-15>
- [70] Brilhante RSN, de Lima RAC, Caetano EP, Leite JJC, Castelo-Branco D de SCM, Ribeiro JF, Bandeira T de JPG, Cordeiro R de A, Monteiro AJ, Sidrim JJC, et al. Effect of farnesol on growth, ergosterol biosynthesis, and cell permeability in *Coccidioides posadasii*. *Antimicrob Agents Chemother* 2013; 57:2167-70; PMID:23459491; <http://dx.doi.org/10.1128/AAC.02457-12>
- [71] Cordeiro R de A, Marques FJ de F, Brilhante RSN, Rocha de Castro E Silva K, Mourão CI, Caetano EP, Fechine MAB, Ribeiro JF, Castelo-Branco D de SCM, de Lima RAC, et al. Synergistic effect of antituberculosis drugs and azoles in vitro against *Histoplasma capsulatum* var. *capsulatum*. *Antimicrob Agents Chemother* 2011; 55:4482-4; PMID:21690286; <http://dx.doi.org/10.1128/AAC.01471-10>
- [72] De Aguiar Cordeiro R, Brilhante RSN, Rocha MFG, Fechine MAB, Camargo ZP de, Sidrim JJC. In vitro inhibitory effect of antituberculosis drugs on clinical and environmental strains of *Coccidioides posadasii*. *J Antimicrob Chemother* 2006; 58:575-9; PMID:16867997; <http://dx.doi.org/10.1093/jac/dkl290>
- [73] Koselny K, Green J, DiDone L, Halterman JP, Fothergill AW, Wiederhold NP, Patterson TF, Cushion MT, Rappleye CA, Wellington M, et al. The celecoxib derivative AR-12 has broad spectrum antifungal activity in vitro and improves the 1 activity of fluconazole in a murine model of cryptococcosis. *Antimicrob Agents Chemother* 2016; PMID:27645246; <http://dx.doi.org/10.1128/AAC.01061-16>
- [74] Wheat LJ, Freifeld AG, Kleiman MB, Baddley JW, McKinsey DS, Loyd JE, Kauffman CA. Clinical Practice Guidelines for the Management of Patients with Histoplasmosis: 2007 Update by the infectious diseases society of America. *Clin Infect Dis* 2007; 45:807-25; PMID:17806045; <http://dx.doi.org/10.1086/521259>
- [75] Chapman SW, Dismukes WE, Proia LA, Bradsher RW, Pappas PG, Threlkeld MG, Kauffman CA, Infectious Diseases Society of America. Clinical practice guidelines for the management of blastomycosis: 2008 update by the infectious diseases society of America. *Clin Infect Dis* 2008; 46:1801-12; PMID:18462107; <http://dx.doi.org/10.1086/588300>
- [76] Galgiani JN, Ampel NM, Blair JE, Catanzaro A, Johnson RH, Stevens DA, Williams PL, Infectious Diseases Society of America. Coccidioidomycosis. *Clin Infect Dis* 2005; 41:1217-23; PMID:16206093; <http://dx.doi.org/10.1086/496991>
- [77] Nett JE, Andes DR. Antifungal Agents: Spectrum of activity, pharmacology, and clinical indications. *Infect Dis Clin North Am* 2016; 30:51-83; PMID:26739608; <http://dx.doi.org/10.1016/j.idc.2015.10.012>
- [78] Cheng JT, Witty RT, Robinson RR, Yarger WE. Amphotericin B nephrotoxicity: increased renal resistance and tubule permeability. *Kidney Int* 1982; 22:626-33; PMID:7162036; <http://dx.doi.org/10.1038/ki.1982.221>
- [79] Bates DW, Su L, Yu DT, Chertow GM, Seger DL, Gomes DR, Dasbach EJ, Platt R. Mortality and costs of acute renal failure associated with amphotericin B therapy. *Clin Infect Dis* 2001; 32:686-93; PMID:11229835; <http://dx.doi.org/10.1086/319211>

- [80] Hoekstra WJ, Garvey EP, Moore WR, Rafferty SW, Yates CM, Schotzinger RJ. Design and optimization of highly-selective fungal CYP51 inhibitors. *Bioorg Med Chem Lett* 2014; 24:3455-8; PMID:24948565; <http://dx.doi.org/10.1016/j.bmcl.2014.05.068>
- [81] Shubitiz LF, Trinh HT, Galgiani JN, Lewis ML, Fothergill AW, Wiederhold NP, Barker BM, Lewis ERG, Doyle AL, Hoekstra WJ, et al. Evaluation of VT-1161 for Treatment of Coccidioidomycosis in Murine Infection Models. *Antimicrob Agents Chemother* 2015; 59:7249-54; PMID:26369964; <http://dx.doi.org/10.1128/AAC.00593-15>
- [82] Capasso C, Supuran CT. Sulfa and trimethoprim-like drugs - antimetabolites acting as carbonic anhydrase, dihydropteroate synthase and dihydrofolate reductase inhibitors. *J Enzyme Inhib Med Chem* 2014; 29:379-87; PMID:23627736; <http://dx.doi.org/10.3109/14756366.2013.787422>
- [83] Borges SRC, Silva GMS da, Chambela M da C, Oliveira R de VC de, Costa RLB, Wanke B, Valle ACF do. Itraconazole vs. trimethoprim-sulfamethoxazole: A comparative cohort study of 200 patients with paracoccidioidomycosis. *Med Mycol* 2014; 52:303-10; PMID:24577007; <http://dx.doi.org/10.1093/mmy/myt012>
- [84] Cavalcante R de S, Sylvestre TF, Levorato AD, de Carvalho LR, Mendes RP. Comparison between itraconazole and cotrimoxazole in the treatment of paracoccidioidomycosis. *PLoS Negl Trop Dis* 2014; 8:e2793; <http://dx.doi.org/10.1371/journal.pntd.0002793>
- [85] Restrepo A. Treatment of tropical mycoses. *J Am Acad Dermatol* 1994; 31:S91-102; PMID:8077517; [http://dx.doi.org/10.1016/S0190-9622\(08\)81277-7](http://dx.doi.org/10.1016/S0190-9622(08)81277-7)
- [86] Brilhante RSN, Caetano EP, Sidrim JJC, Cordeiro RA, Camargo ZP, Fachine MAB, Lima RAC, Castelo Branco DSCM, Marques FJF, Mesquita JRL, et al. Ciprofloxacin shows synergism with classical antifungals against *Histoplasma capsulatum* var. *capsulatum* and *Coccidioides posadasii*. *Mycoses* 2013; 56:397-401; PMID:23205615; <http://dx.doi.org/10.1111/myc.12025>
- [87] Cordeiro R de A, de Melo CVS, Marques FJ de F, Serpa R, Evangelista AJ de J, Caetano EP, Mafezoli J, de Oliveira M da CF, da Silva MR, Bandeira T de JPG, et al. Synthesis and in vitro antifungal activity of isoniazid-derived hydrazones against *Coccidioides posadasii*. *Microb Pathog* 2016; 98:1-5; PMID:27334293; <http://dx.doi.org/10.1016/j.micpath.2016.06.022>
- [88] Derengowski LS, De-Souza-Silva C, Braz SV, Mello-De-Sousa TM, Bão SN, Kyaw CM, Silva-Pereira I. Antimicrobial effect of farnesol, a *Candida albicans* quorum sensing molecule, on *Paracoccidioides brasiliensis* growth and morphogenesis. *Ann Clin Microbiol Antimicrob* 2009; 8:13; PMID:19402910; <http://dx.doi.org/10.1186/1476-0711-8-13>
- [89] Brilhante RSN, de Lima RAC, Marques FJ de F, Silva NF, Caetano EP, Castelo-Branco D de SCM, Bandeira T de JPG, Moreira JLB, Cordeiro R de A, Monteiro AJ, et al. *Histoplasma capsulatum* in planktonic and biofilm forms: in vitro susceptibility to amphotericin B, itraconazole and farnesol. *J Med Microbiol* 2015; 64:394-9; PMID:25657300; <http://dx.doi.org/10.1099/jmm.0.000030>
- [90] Baxter BK, DiDone L, Ogu D, Schor S, Krysan DJ. Identification, in vitro activity and mode of action of phosphoinositide-dependent-1 kinase inhibitors as antifungal molecules. *ACS Chem Biol* 2011; 6:502-10; PMID:21294551; <http://dx.doi.org/10.1021/cb100399x>
- [91] Hoang KV, Curry H, Collier MA, Borteh H, Bachelder EM, Schlesinger LS, Gunn JS, Ainslie KM. Needle-free delivery of acetalated Dextran-Encapsulated AR-12 protects mice from *Francisella tularensis* Lethal Challenge. *Antimicrob Agents Chemother* 2016; 60:2052-62; PMID:26787696; <http://dx.doi.org/10.1128/AAC.02228-15>
- [92] Collier MA, Peine KJ, Gautam S, Oghumu S, Varikuti S, Borteh H, Papenfuss TL, Sataoskar AR, Bachelder EM, Ainslie KM. Host-mediated *Leishmania donovani* treatment using AR-12 encapsulated in acetalated dextran microparticles. *Int J Pharm* 2016; 499:186-94; PMID:26768723; <http://dx.doi.org/10.1016/j.ijpharm.2016.01.004>
- [93] Chiu H-C, Kulp SK, Soni S, Wang D, Gunn JS, Schlesinger LS, Chen C-S. Eradication of intracellular *Salmonella enterica* serovar Typhimurium with a small-molecule, host cell-directed agent. *Antimicrob Agents Chemother* 2009; 53:5236-44; PMID:19805568; <http://dx.doi.org/10.1128/AAC.00555-09>
- [94] Mohr EL, McMullan LK, Lo MK, Spengler JR, Bergeron É, Albariño CG, Shrivastava-Ranjan P, Chiang CF, Nichol ST, Spiropoulou CF, et al. Inhibitors of cellular kinases with broad-spectrum antiviral activity for hemorrhagic fever viruses. *Antiviral Res* 2015; 120:40-7; PMID:25986249; <http://dx.doi.org/10.1016/j.antiviral.2015.05.003>
- [95] Koselny K, Green J, Favazzo L, Glazier VE, DiDone L, Ransford S, Krysan DJ. Antitumor/Antifungal Celecoxib Derivative AR-12 is a non-nucleoside inhibitor of the ANL-Family Adenylating Enzyme Acetyl CoA synthetase. *ACS Infect Dis* 2016; 2:268-80; PMID:27088128; <http://dx.doi.org/10.1021/acsinfecdis.5b00134>
- [96] Abadio AKR, Kioshima ES, Teixeira MM, Martins NF, Maigret B, Felipe MSS. Comparative genomics allowed the identification of drug targets against human fungal pathogens. *BMC Genomics* 2011; 12:75; PMID:21272313; <http://dx.doi.org/10.1186/1471-2164-12-75>
- [97] Abadio AKR, Kioshima ES, Leroux V, Martins NF, Maigret B, Felipe MSS. Identification of new antifungal compounds targeting thioredoxin reductase of *paracoccidioides* genus. *PLoS One* 2015; 10:e0142926; PMID:26569405; <http://dx.doi.org/10.1371/journal.pone.0142926>
- [98] Edwards JA, Kemski MM, Rappleye CA. Identification of an aminothiazole with antifungal activity against intracellular *Histoplasma capsulatum*. *Antimicrob Agents Chemother* 2013; 57:4349-59; PMID:23817367; <http://dx.doi.org/10.1128/AAC.00459-13>
- [99] Isono K, Asahi K, Suzuki S. Studies on polyoxins, antifungal antibiotics. 13. The structure of polyoxins. *J Am Chem Soc* 1969; 91:7490-505; PMID:5358618; <http://dx.doi.org/10.1021/ja01054a045>
- [100] Endo A, Misato T. Polyoxin D, a competitive inhibitor of UDP-N-acetylglucosamine: chitin N-acetylglucosaminyltransferase in *Neurospora crassa*. *Biochem Biophys Res Commun* 1969; 37:718-22; PMID:5353899; [http://dx.doi.org/10.1016/0006-291X\(69\)90870-5](http://dx.doi.org/10.1016/0006-291X(69)90870-5)
- [101] Garfoot AL, Shen Q, Wüthrich M, Klein BS, Rappleye CA. The Eng1 β -Glucanase Enhances *Histoplasma* Virulence by Reducing β -Glucan Exposure. *MBio* 2016; 7:e01388-15; PMID:27094334; <http://dx.doi.org/10.1128/mBio.01388-15>

- [102] Goldberg J, Connolly P, Schnitzlein-Bick C, Durkin M, Kohler S, Smedema M, Brizendine E, Hector R, Wheat J. Comparison of nikkomycin Z with amphotericin B and itraconazole for treatment of histoplasmosis in a murine model. *Antimicrob Agents Chemother* 2000; 44:1624-9; PMID:10817719; <http://dx.doi.org/10.1128/AAC.44.6.1624-1629.2000>
- [103] Hector RF, Zimmer BL, Pappagianis D. Evaluation of nikkomycins X and Z in murine models of coccidioidomycosis, histoplasmosis, and blastomycosis. *Antimicrob Agents Chemother* 1990; 34:587-93; PMID:2344165; <http://dx.doi.org/10.1128/AAC.34.4.587>
- [104] Graybill JR, Najvar LK, Bocanegra R, Hector RF, Luther MF. Efficacy of nikkomycin Z in the treatment of murine histoplasmosis. *Antimicrob Agents Chemother* 1998; 42:2371-4; PMID:9736565
- [105] Clemons KV, Stevens DA. Efficacy of nikkomycin Z against experimental pulmonary blastomycosis. *Antimicrob Agents Chemother* 1997; 41:2026-8; PMID:9303408
- [106] Kriesel JD, Sutton DA, Schulman S, Fothergill AW, Rinaldi MG. Persistent pulmonary infection with an azole-resistant *Coccidioides* species. *Med Mycol* 2008; 46:607-10; PMID:18608910; <http://dx.doi.org/10.1080/13693780802140923>
- [107] Wang CY, Jerng JS, Ko JC, Lin MF, Hsiao CH, Lee LN, Hsueh PR, Kuo SH. Disseminated coccidioidomycosis. *Emerging Infect Dis* 2005; 11:177-9; PMID:15714666; <http://dx.doi.org/10.3201/eid1101.040613>
- [108] Wheat LJ, Connolly P, Smedema M, Brizendine E, Hafner R, AIDS Clinical Trials Group and the Mycoses Study Group of the National Institute of Allergy and Infectious Diseases. Emergence of resistance to fluconazole as a cause of failure during treatment of histoplasmosis in patients with acquired immunodeficiency disease syndrome. *Clin Infect Dis* 2001; 33:1910-3; PMID:11692303; <http://dx.doi.org/10.1086/323781>
- [109] Kathuria S, Singh PK, Meis JF, Chowdhary A. In vitro antifungal susceptibility profile and correlation of mycelial and yeast forms of molecularly characterized *Histoplasma capsulatum* strains from India. *Antimicrob Agents Chemother* 2014; 58:5613-6; PMID:24982084; <http://dx.doi.org/10.1128/AAC.02973-14>
- [110] Ramani R, Chaturvedi V. Antifungal susceptibility profiles of *Coccidioides immitis* and *Coccidioides posadasii* from endemic and non-endemic areas. *Mycopathologia* 2007; 163:315-9; PMID:17484074; <http://dx.doi.org/10.1007/s11046-007-9018-7>
- [111] Li RK, Ciblak MA, Nordoff N, Pasarell L, Warnock DW, McGinnis MR. In vitro activities of voriconazole, itraconazole, and amphotericin B against *Blastomyces dermatitidis*, *Coccidioides immitis*, and *Histoplasma capsulatum*. *Antimicrob Agents Chemother* 2000; 44:1734-6; PMID:10817743; <http://dx.doi.org/10.1128/AAC.44.6.1734-1736.2000>
- [112] Sugar AM, Liu XP. In vitro and in vivo activities of SCH 56592 against *Blastomyces dermatitidis*. *Antimicrob Agents Chemother* 1996; 40:1314-6; PMID:8723494
- [113] Cordeiro RA, Brilhante RSN, Rocha MFG, Fechine MAB, Costa AKF, Camargo ZP, Sidrim JJC. In vitro activities of caspofungin, amphotericin B and azoles against *Coccidioides posadasii* strains from Northeast, Brazil. *Mycopathologia* 2006; 161:21-6; PMID:16389480; <http://dx.doi.org/10.1007/s11046-005-0177-0>
- [114] Cruz RC, Werneck SMC, Oliveira CS, Santos PC, Soares BM, Santos DA, Cisalpino PS. Influence of different media, incubation times, and temperatures for determining the MICs of seven antifungal agents against *Paracoccidioides brasiliensis* by microdilution. *J Clin Microbiol* 2013; 51:436-43; PMID:23175254; <http://dx.doi.org/10.1128/JCM.02231-12>
- [115] Andreu CMF, León AM, Medina YE, Machín GM, Lancha MRP, Zaragoza MTI. In vitro sensitivity of *Histoplasma capsulatum* var. *capsulatum* to amphotericin B, ketoconazole, itraconazole and fluconazole. *Rev Cubana Med Trop* 2003; 55:76-82; PMID:15849960
- [116] Zhanel GG, Karlowsky JA, Harding GA, Balko TV, Zelenitsky SA, Friesen M, Kabani A, Turik M, Hoban DJ. In vitro activity of a new semisynthetic echinocandin, LY-303366, against systemic isolates of *Candida* species, *Cryptococcus neoformans*, *Blastomyces dermatitidis*, and *Aspergillus* species. *Antimicrob Agents Chemother* 1997; 41:863-5; PMID:9087508
- [117] González GM, Tijerina R, Najvar LK, Bocanegra R, Luther M, Rinaldi MG, Graybill JR. Correlation between antifungal susceptibilities of *Coccidioides immitis* in vitro and antifungal treatment with caspofungin in a mouse model. *Antimicrob Agents Chemother* 2001; 45:1854-9; <http://dx.doi.org/10.1128/AAC.45.6.1854-1859.2001>
- [118] Kohler S, Wheat LJ, Connolly P, Schnitzlein-Bick C, Durkin M, Smedema M, Goldberg J, Brizendine E. Comparison of the echinocandin caspofungin with amphotericin B for treatment of histoplasmosis following pulmonary challenge in a murine model. *Antimicrob Agents Chemother* 2000; 44:1850-4; PMID:10858342; <http://dx.doi.org/10.1128/AAC.44.7.1850-1854.2000>
- [119] LeMonte AM, Washum KE, Smedema ML, Schnitzlein-Bick C, Kohler SM, Wheat LJ. Amphotericin B combined with itraconazole or fluconazole for treatment of histoplasmosis. *J Infect Dis* 2000; 182:545-50; PMID:10915087; <http://dx.doi.org/10.1086/315717>
- [120] Chapman SW, Rogers PD, Rinaldi MG, Sullivan DC. Susceptibilities of clinical and laboratory isolates of *Blastomyces dermatitidis* to ketoconazole, itraconazole, and fluconazole. *Antimicrob Agents Chemother* 1998; 42:978-80; PMID:9559827
- [121] González GM, Fothergill AW, Sutton DA, Rinaldi MG, Loebenberg D. In vitro activities of new and established triazoles against opportunistic filamentous and dimorphic fungi. *Med Mycol* 2005; 43:281-4; <http://dx.doi.org/10.1080/13693780500088416>
- [122] Van Duin D, Casadevall A, Nosanchuk JD. Melanization of *Cryptococcus neoformans* and *Histoplasma capsulatum* reduces their susceptibilities to amphotericin B and caspofungin. *Antimicrob Agents Chemother* 2002; 46:3394-400; PMID:12384341; <http://dx.doi.org/10.1128/AAC.46.11.3394-3400.2002>
- [123] Wheat LJ, Connolly P, Smedema M, Durkin M, Brizendine E, Mann P, Patel R, McNicholas PM, Goldman M. Activity of newer triazoles against *Histoplasma capsulatum* from patients with AIDS who failed fluconazole. *J Antimicrob Chemother* 2006; 57:1235-9; PMID:16627592; <http://dx.doi.org/10.1093/jac/dkl133>