

## SI PLANT BIOTIC INTERACTIONS

# Tools of the crook- infection strategies of fungal plant pathogens

Luis Rodriguez-Moreno<sup>1</sup>, Malaika K. Ebert<sup>1,2</sup>, Melvin D. Bolton<sup>2</sup> and Bart P.H.J. Thomma<sup>1,\*</sup> <sup>1</sup>Laboratory of Phytopathology, Wageningen University, Droevendaalsesteeg 1, 6708 PB Wageningen, The Netherlands, and<sup>2</sup>USDA – Agricultural Research Service, Red River Valley Agricultural Research Center, Fargo, ND, USA

Received 15 November 2017; revised 18 December 2017; accepted 18 December 2017; published online 26 December 2017.

\*For correspondence (e-mail bart.thomma@wur.nl).

L.R.-M., M.K.E., M.D.B. and B.P.H.J.T. contributed equally to this study.

## SUMMARY

Fungi represent an ecologically diverse group of microorganisms that includes plant pathogenic species able to cause considerable yield losses in crop production systems worldwide. In order to establish compatible interactions with their hosts, pathogenic fungi rely on the secretion of molecules of diverse nature during host colonization to modulate host physiology, manipulate other environmental factors or provide self-defence. These molecules, collectively known as effectors, are typically small secreted cysteine-rich proteins, but may also comprise secondary metabolites and sRNAs. Here, we discuss the most common strategies that fungal plant pathogens employ to subvert their host plants in order to successfully complete their life cycle and secure the release of abundant viable progeny.

**Keywords:** fungus, plant pathogen, effector, host immunity, virulence.


## INTRODUCTION

Fungi constitute an evolutionarily and ecologically diverse group of microorganisms that includes plant pathogenic species that cause considerable yield losses in agricultural production systems worldwide. Generally, the lifestyles of plant pathogenic fungi are differentiated depending on the strategies used to acquire nutrients from their hosts. As such, obligate biotrophic fungi comprise those species that can only feed on living host tissue to meet their nutritional requirements and complete their life cycle. At the complete opposite of the spectrum, necrotrophic fungi trigger cell death in the host to secure nutrient supply. In between these extremes is a wide array of hemibiotrophic fungi that start their compatible host interaction with an initial biotrophic phase that, at one point in time when the infection progressed sufficiently, is followed by a transition to a necrotrophic stage. A parasitic lifestyle that involves the extraction of sugars from other organisms is one of the ways in which non-heterotrophic organisms compensate for the inability to generate sugars through photosynthesis. Many biotrophic and hemibiotrophic fungi evolved haustoria, appendages of fungal hyphae that invaginate the host plasma membrane and grow inside host cells, to obtain these nutrients. Recently, it was demonstrated that

the obligate biotrophic powdery mildew fungus *Golovino-mycetes cichoracerum* requires lipids for colonization that it receives from the host plant (Jiang *et al.*, 2017).

Lifestyle differences largely determine the wide array of strategies that fungi use to evade, counteract or hijack plant defences in their effort to complete their life cycle and secure the production of viable progeny. Irrespective of their lifestyle, microbial pathogens are all believed to utilize so-called effectors, *in planta*-secreted molecules of various nature, to support host colonization, often, but not exclusively, through suppression of host immune responses (Rovenich *et al.*, 2014). Over the years it has become evident that haustoria are not only fungal feeding structures, but are also active sites for secretion and translocation of effectors into the host (Mendgen and Hahn, 2002; Whisson *et al.*, 2007; Rafiqi *et al.*, 2010, 2012).

Plants have developed an innate immune system to recognize and respond to microbes (Thomma *et al.*, 2001; Jones and Dangl, 2006; Cook *et al.*, 2015; Figure 1). This immune system relies on the presence of immune receptors that detect pathogen invasion through sensing of pathogen(-induced) ligands, collectively termed invasion patterns, to mount appropriate immune responses (Cook



**Figure 1.** Schematic representation of the 'Invasion Model' to describe the molecular basis of plant immunity against fungal pathogens. In this model, invasion pattern receptors, comprising any type of host receptor, detect invasion patterns, comprising externally encoded and modified-self ligands that announce invasion, to mount an effective immune response and halt the symbiosis. Fungal effectors may manipulate the induced response to tweak the symbiosis to their benefit.


*et al.*, 2015). Recognition of invasion patterns triggers both local and systemic reactions to respond in a quick and focussed manner to attempted microbial ingress (Thomma *et al.*, 2001; Jones and Dangl, 2006; Cook *et al.*, 2015). For example, the well-characterized invasion pattern chitin, an important constituent of fungal cell walls, is recognized by plants through plasma membrane-localized extracellular lysin motif (LysM)-containing receptor molecules (Felix *et al.*, 1993; Shibuya *et al.*, 1993). Pathogen recognition by plant immune receptors causes ion fluxes, the accumulation of reactive oxygen species (ROS), and a quick activation of defence-related mitogen-activated protein kinase (MAPKs) cascades that cause an extensive transcriptional reprogramming of the host (Altenbach and Robatzek, 2007; Boller and Felix, 2009; Bolton, 2009). Furthermore, pathogen perception leads to reinforcement of plant cell walls by callose deposition, changes in hormone biosynthesis and the production of antimicrobial compounds (Macho and Zipfel, 2014). In many cases, these defence responses collectively are sufficient to render the interaction between the plant and the invader incompatible, implying that pathogen ingress is halted or at least significantly slowed down. However, co-evolutionary processes have selected pathogens that employ a plethora of virulence strategies to overcome various mechanisms within plant immune

systems. In this review, we summarize the different virulence strategies that plant pathogenic fungi use to subvert their hosts. While there are excellent reviews that discuss individual strategies in detail, the aim of this review is to outline the broad diversity of known fungal virulence mechanisms (Figure 2).

### Fungal strategies for host penetration

One of the first barriers that fungal pathogens have to breach to gain entrance to their hosts is cell walls that are mainly composed of carbohydrates. Many plant pathogenic fungi utilize specialized infection structures, called appressoria, to generate focused turgor pressure to breach the cell wall by force (Ryder and Talbot, 2015). Depending on the fungal species, the turgor pressure is combined with the localized release of cell-wall-degrading enzymes (CWDEs; Ryder and Talbot, 2015). Furthermore, effectors are secreted from appressorial penetration pores prior to host invasion (Kleemann *et al.*, 2012).

Fungi typically produce an arsenal of so-called carbohydrate-active enzymes (CAZymes) that are grouped into five enzyme classes, namely glycoside hydrolases, glycosyltransferases, polysaccharide lyases, carbohydrate esterases and redox enzymes with auxiliary activities (Lombard *et al.*, 2014). Several of the polysaccharide lyases, glycoside hydrolases and carbohydrate esterases are known as CWDEs that are used to degrade host cell walls. Typically plant pathogenic species contain higher numbers of CAZyme genes than saprophytic and animal pathogenic strains (Zhao *et al.*, 2013). Whereas obligate biotrophs typically lack extensive catalogs of CWDE genes and likely only use such enzymes for subtle manipulations of host cell walls such as at the cellular entrance sites for haustoria, necrotrophic fungi were often thought of as 'brute-force' pathogens that rely on large CWDE catalogs to macerate host cell walls and initiate colonization (Bolton *et al.*, 2006). These enzymes occur in multiple isoforms that not only differ in isoelectric point and molecular weight, but also in timing of their production and processing, offering especially broad host-range necrotrophs particular flexibility to penetrate and colonize their hosts. Besides colonization, these enzymes also liberate nutrients for the pathogen. For example, hydrolysis of pectin by fungal pectinases weakens the cell wall to enable penetration while also providing the fungus with important carbon sources for growth (Alghisi and Favaron, 1995). Indeed, strategies to limit pectin degradation were explored by generating transgenic wheat lines expressing pectin methyl esterase inhibitors, which exhibited altered pectin methyl esterification that resulted in reduced activity of pathogen pectic enzymes and reduced disease from hemibiotrophic pathogens *Fusarium graminearum* and *Bipolaris sorokiniana* (Volpi *et al.*, 2011). Similarly, wheat lines expressing genes encoding a xylanase inhibitor and



**Figure 2.** Illustration of fungal pathogen strategies to surmount host plants.

(a) Secretion of effectors that perturb recognition by plant immune receptors.

(b) Secretion of effectors that subvert plant defence responses that are induced upon pathogen detection.

(c) Delivery of various types of molecules to hijack host metabolism.

polygalacturonase-inhibiting protein exhibited increased resistance to *Fusarium* head blight (Tundo *et al.*, 2016). However, *F. graminearum* single-gene deletion mutants for polygalacturonase or xylanase resulted in minor effects on virulence, while double-gene mutants were significantly reduced in virulence on soybean and wheat plants, highlighting the synergism between CWDEs (Paccanaro *et al.*, 2017).

Besides plant CWDEs, fungi secrete CWDEs to modulate their own cell walls and accommodate morphological changes. It was recently proposed that such activity facilitates pathogenesis of plants by enabling host colonization. A glycosyltransferase enzyme from the hemibiotrophic wheat pathogen *Zymoseptoria tritici* was reported to enable hyphal growth on solid surfaces that is essential for fungal disease of wheat plants (King *et al.*, 2017). Homologues of this particular enzyme are widespread in fungi, and mutants in the taxonomically unrelated *F. graminearum* were similarly impaired in hyphal growth and pathogenicity (King *et al.*, 2017).

### Fungal strategies preventing plant recognition

Plants evolved a plethora of plasma membrane-localized immune receptors for surveillance of the extracellular space for pathogen(-induced) ligands (Boller and Felix, 2009; Macho and Zipfel, 2014; Couto and Zipfel, 2016). The perception of these ligands is relayed into downstream signalling events that lead to the activation of plant defences (Bolton, 2009; Couto and Zipfel, 2016). Structural components of the fungal cell wall, such as glucans and chitin, are typically recognized as pathogen ligands by plant receptors (Felix *et al.*, 1993; Shibuya *et al.*, 1993; Cosio *et al.*, 1996; Cote *et al.*, 2000). As part of their defence system, plants secrete glucanases and chitinases to compromise the integrity of fungal cell walls and release oligomeric fragments that can act as ligands for extracellular

immune receptors (Sanchez-Vallet *et al.*, 2015). Fungi have evolved several strategies to overcome host immune responses that involve fungal cell walls, including alterations in cell wall compositions and the secretion of effectors to protect cell walls or perturb recognition of cell wall components (Rovenich *et al.*, 2014; Sanchez-Vallet *et al.*, 2015).

*Magnaporthe oryzae* is a hemibiotrophic fungal pathogen and causal agent of rice blast disease (Dean *et al.*, 2005). During infection, *M. oryzae* responds to the epidermal wax component 1,16-hexadecanediol by accumulating  $\alpha$ -1,3-glucans at the surface of the cell wall, resulting in inhibition of chitin degradation by plant chitinases (Fujikawa *et al.*, 2012). Accordingly, mutants that are unable to accumulate  $\alpha$ -1,3-glucans at the fungal cell surface trigger rapid activation of host defences (Fujikawa *et al.*, 2012). A similar strategy has been reported for the maize pathogen *Colletotrichum graminicola* that modifies the  $\beta$ -glucan composition of its biotrophic hyphae, as the content of  $\beta$ -1,3- and  $\beta$ -1,6-glucans is significantly reduced when compared with appressoria and necrotrophic hyphae (El Gueddari *et al.*, 2002; Oliveira-Garcia and Deising, 2013, 2016). Hence, *C. graminicola* strains that overexpress a  $\beta$ -1,3-glucan synthase in their biotrophic hyphae induce stronger host defence responses and display reduced virulence (Oliveira-Garcia and Deising, 2013). However, *C. graminicola* strains that are unable to produce  $\beta$ -1,6-glucans are defective in appressorium formation and thus non-pathogenic (Oliveira-Garcia and Deising, 2016).

Modification of cell walls is not the only strategy employed by fungal pathogens to prevent plant recognition. For instance, the tomato leaf mould fungus *Cladosporium fulvum* secretes the carbohydrate-binding effector protein Ecp6 that suppresses chitin-triggered host immunity. The chitin-binding capacity of Ecp6 is mediated by three LysMs (de Jonge *et al.*, 2010; Sanchez-Vallet *et al.*,

2013) that occur in proteins of a wide range of organisms to confer the ability to bind various types of polysaccharides, including peptidoglycan and chitin, through a conserved  $\beta\alpha\beta$ -fold (Buist *et al.*, 2008). Interestingly, two out of the three LysM domains of Ecp6 cooperate to form a groove that binds chitin fragments with ultra-high (pM) affinity that allows to outcompete host receptors for chitin binding (Sanchez-Vallet *et al.*, 2013). Besides Ecp6, *C. fulvum* also secretes the chitin-binding effector molecule Avr4 during host colonization. As opposed to LysMs, Avr4 binds chitin through an invertebrate chitin-binding module to protect the cell wall against hydrolysis by host enzymes (van den Burg *et al.*, 2006; van Esse *et al.*, 2007). In contrast to Avr4 homologues that only occur in a limited set of fungi that are closely related to *C. fulvum* (Stergiopoulos *et al.*, 2010), LysM effector proteins occur in a wide variety of fungi (de Jonge and Thomma, 2009), and have been shown to suppress chitin-triggered immunity on various plant hosts such as for *Z. tritici* on wheat (Marshall *et al.*, 2011), for *M. oryzae* on rice (Mentlak *et al.*, 2012), for *Colletotrichum higginsianum* on Arabidopsis (Takahara *et al.*, 2016), and for *Verticillium dahliae* on tomato (Kombrink *et al.*, 2017).

Whereas chitin perception in plants is relatively well understood (Rovenich *et al.*, 2016),  $\beta$ -glucan perception and signalling mechanisms remain poorly characterized (Fesel and Zuccaro, 2016). The root endophyte *Piriformospora indica* secretes the  $\beta$ -glucan-binding lectin effector FGB1 that suppresses  $\beta$ -glucan-triggered host immunity (Wawra *et al.*, 2016). Prevention of  $\beta$ -glucan detection by the plant seems important for successful fungal infection as overexpression of the *P. indica* FGB1 homologue in *Ustilago maydis* was shown to lead to an increase in virulence. Interestingly, FGB1 homologues are widespread in fungi (Wawra *et al.*, 2016).

A further strategy to protect fungal cell walls and prevent detection of cell wall components is through the secretion of proteases that affect hydrolytic host enzymes (Albersheim and Valent, 1974). Fungal chitinase-modifying proteins (CMPs) have been reported in several maize pathogens, including *Bipolaris zeicola* (Naumann *et al.*, 2009), *Stenocarpella maydis* (Naumann and Wicklow, 2010) and *Fusarium verticilloides* (Naumann *et al.*, 2011). Similarly, *Fusarium oxysporum* f. sp. *lycopersici*, *V. dahliae* and *Botrytis cinerea* were found to secrete CMPs that can degrade extracellular tomato chitinases (Jashni *et al.*, 2015a).

### Fungal strategies for inhibiting host defence responses

Pathogen recognition by plants results in a panoply of defence responses to hamper pathogen invasion. These responses comprise swift ion fluxes, pH changes, production of ROS, but also the production of local and systemic signalling molecules and antimicrobial compounds. Various

mechanisms are employed by fungal pathogens to subvert such responses.

**Subverting ROS damage.** Reactive oxygen species production is mostly due to the activity of membrane-bound NADPH-oxidases and cell-wall-associated peroxidases (POX; Bolwell *et al.*, 2002; Sasaki *et al.*, 2004; Bindschedler *et al.*, 2006). While relatively low concentrations of ROS have been reported to act as defence signalling molecules (Lamb and Dixon, 1997; Orozco-Cardenas *et al.*, 2001; Qi *et al.*, 2017), high concentrations of ROS are extremely harmful to cells as they have been shown to cause oxidative damage (Levine *et al.*, 1994; Wu *et al.*, 1997). The apoplastic effector Pep1 of the biotrophic maize pathogen *U. maydis* accumulates at sites where biotrophic hyphae move from cell to cell in maize tissue to inhibit the oxidative burst through inhibition of POX12, a type-III class heme-peroxidase that is highly induced after *U. maydis* penetration (Doehlemann *et al.*, 2009; Hemetsberger *et al.*, 2012). Pep1 only causes partial inhibition of the maize apoplastic peroxidase activity, suggesting that not all peroxidase-producing enzymes in the maize apoplast are targeted by Pep1 (Hemetsberger *et al.*, 2012).

**Manipulating tissue pH.** Many fungal pathogens induce a pH shift in the host tissue surrounding the infection site (Prusky and Yakoby, 2003). For instance, *Sclerotinia sclerotiorum* causes acidification of the infection area through the production of oxalic acid, leading to rapid death of host tissues (Bolton *et al.*, 2006). However, other pathogens induce alkalinization of host tissue (Prusky *et al.*, 2001; Masachis *et al.*, 2016). During host colonization, the vascular wilt pathogen *F. oxysporum* causes an increase of the extracellular pH from about 5 to 7 through the secretion of a peptide with homology to plant rapid alkalinizing factors (RALFs; Murphy and De Smet, 2014; Masachis *et al.*, 2016). Interestingly, *F. oxysporum* strains that are no longer able to produce this peptide trigger enhanced host defence, indicating a role in suppression of host immunity. Although this role in virulence has been challenged (Thynne *et al.*, 2017), RALF-encoding genes can be found in many fungal pathogens, suggesting a universal mechanism to alkalinize infection sites to suppress host immunity (Masachis *et al.*, 2016; Thynne *et al.*, 2017).

**Inhibition of host proteases.** Many of the molecules that fungal pathogens secrete in order to establish the parasitic interaction with their hosts are of a proteinaceous nature, and hence plants secrete proteases to undermine this pathogen strategy (van der Hoorn, 2008; Jashni *et al.*, 2015b). The apoplast of tomato and Arabidopsis contains various proteases that contribute to host defence (Shabab *et al.*, 2008; van Esse *et al.*, 2008). Among these, the tomato apoplast contains the extracellular cysteine



protease Rcr3 that plays a central role in resistance mediated by the Cf-2 immune receptor of tomato and that is activated by the *C. fulvum* effector Avr2 (Kruger *et al.*, 2002). The Avr2 effector inhibits the activity of Rcr3, likely causing a conformational change in the Rcr3 structure that is recognized by Cf-2 (Kruger *et al.*, 2002; Rooney *et al.*, 2005). Besides Rcr3, Avr2 inhibits various other host proteases that are required for pathogen defence (van Esse *et al.*, 2008). Other fungal pathogens also produce protease effectors to inhibit host proteases, such as the *U. maydis* Pit2 effector (Doehlemann *et al.*, 2011; Mueller *et al.*, 2013).

**Subverting hormone signalling.** Plant growth and their responses to environmental cues, including pathogens, are largely governed by phytohormones. Typically, salicylic acid (SA) signalling governs resistance against biotrophic pathogens, whereas a combination of jasmonic acid (JA) and ethylene (ET) signalling activates resistance against necrotrophic pathogens (Thomma *et al.*, 1998, 2001; Glazebrook, 2005). To a large extent, these signalling pathways act antagonistically and their balance needs to be governed carefully. Thus, it is not surprising that pathogens evolved various strategies to affect phytohormone signalling. For instance, *U. maydis* secretes the chorismate mutase Cmu1 into host cells to perturb SA production by affecting the production of its precursor (Djamei *et al.*, 2011). Likely, Cmu1 acts in combination with the maize chorismate mutase Cm1 to increase the flow of chorismate from the plastid to the cytosol to diminish the available substrate for SA biosynthesis in plastids in turn (Djamei *et al.*, 2011). Furthermore, *U. maydis* produces Shy1, a salicylate hydroxylase that degrades SA during host invasion (Rabe *et al.*, 2013). Together these results suggest that perturbation of SA-mediated immunity is crucial for *U. maydis* colonization. Chorismate mutases have been identified in many eukaryotic plant pathogens pointing towards a common strategy for host manipulation. Similar to *U. maydis*, also *V. dahliae* has been proposed to target SA biosynthesis by secreting effectors with isochorismatase activity to hydrolyse isochorismate (Liu *et al.*, 2014). Besides targeting SA signalling, fungal effectors that target JA signalling or ET signalling have been described as well (Kloppholz *et al.*, 2011; Plett *et al.*, 2014). For instance, the beneficial fungus *Laccaria bicolor* produces the Mycorrhiza-induced small secreted protein-7 (MiSSP7) during the interaction with its host *Populus trichocarpa* (Plett *et al.*, 2014). Intriguingly, MiSSP7 interacts with the plant JASMONATE ZIM-DOMAIN (JAZ)-6 protein to provoke blockage of the expression of JA-inducible genes in the host to promote fungal colonization (Plett *et al.*, 2014).

Besides the capacity to manipulate hormone balances in plant tissues, particular fungi appear to have the ability to produce hormone-mimicking compounds to promote host colonization (Reineke *et al.*, 2008; Yin *et al.*, 2014; Chanclud

and Morel, 2016; Chanclud *et al.*, 2016). For instance, *Fusarium pseudograminearum* produces cytokinin-like molecules that activate plant cytokinin signalling to reprogramme the host (Sorensen *et al.*, 2017).

### The molecules that do the job: fungal effectors

Typically, fungal effectors are described as small secreted, cysteine-rich proteins that are produced during host invasion (Stergiopoulos and de Wit, 2009). These fungal effectors can be divided into two types based on their extra- or intracellular localization in the host. Yet, how cytoplasmic effectors are translocated into host cells remains poorly understood (Kale and Tyler, 2011; Ribot *et al.*, 2013; Petre and Kamoun, 2014). Nevertheless, two distinct secretion systems to target effectors have been described for *M. oryzae*. Cytoplasmic effectors accumulate in a so-called biotrophic interfacial complex, a plant membrane-rich structure associated with invasive hyphae that involves exocyst and t-SNARE components (Khang *et al.*, 2010; Giraldo *et al.*, 2013). By contrast, apoplastic effectors are secreted from invasive hyphae via conventional secretion. In addition to proteinaceous effector molecules, other types of molecules are secreted by fungi with the aim to establish the parasitic relationship that therefore qualify to be labelled as effectors just as well.

**Secondary metabolites (SMs).** Secondary metabolites are small bioactive molecules that often play crucial roles in the establishment of specific ecological niches but, unlike primary metabolites, are not essential for fungal growth, development or reproduction. While fungal SMs are often known and valued for their anti-microbial activities, many fungi employ SMs to promote virulence. Traditionally SMs involved with virulence are classified as either host-specific toxins (HSTs; discussed below), because they have specific targets in the host, or non-HSTs that typically do not have a specific host target and are generally toxic to a wide range of organisms including the host instead (Wolpert *et al.*, 2002). Perylenequinones, for example, are a family of photosensitizing SMs for which the mode of action is well studied. The most prominent member of the family is cercosporin. This light-activated toxin is produced by most *Cercospora* spp. and has a very broad toxicity range to many organisms, including plants, animals, bacteria and most fungi. Due to its photosensitizing nature, cercosporin is able to absorb light energy and subsequently react with oxygen (Daub and Ehrenshaft, 2000). Products of this reaction are ROS that can cause protein and DNA damage and lipid peroxidation, and eventually lead to cell death of the host (Blokina *et al.*, 2003; Birben *et al.*, 2012). As necrosis development lays the ground for fungal spore formation, it is speculated that cercosporin secretion might facilitate cell wall breaching to enable conidiophore and conidia production (Weltmeier *et al.*, 2011). The cercosporin biosynthesis

gene cluster was recently shown to be found widespread in the *Colletotrichum* genus, implicating the role of cercosporin as a virulence factor in an important group of fungal plant pathogens (de Jonge *et al.*, 2017).

**HSTs.** Host-selective toxins are known to induce necrotic host tissue reactions to promote host susceptibility (Wolpert *et al.*, 2002). The effectiveness of HSTs depends on whether a plant possesses a corresponding toxin target, which may also define the host range of the producing pathogen. For example, maize lines harboring Texas cytoplasm for male sterility (Tcms) display extreme sensitivity to T-toxin and PM-toxin secreted by *Cochliobolus heterostrophus* race T and *Mycosphaerella zeae-maydis*, respectively (Levings III *et al.*, 1995; Wolpert *et al.*, 2002; Tsuge *et al.*, 2013). Here, host susceptibility is conferred by a single plant gene *T-urf13* that encodes URF13, a mitochondrial membrane protein to which either toxin can directly bind. Binding triggers URF13 to experience a conformational change that in turn results in the formation of a pore in the mitochondrial membrane. The ability to produce T-toxin is relevant for fungal virulence, as *C. heterostrophus* race O, a natural T-toxin lacking race and Tox1<sup>-</sup>-deficient mutants of race T show reduced virulence on Tcms-carrying maize (Yang *et al.*, 1994). Similarly, PM-toxin-deficient Tox<sup>-</sup> mutants of *M. zeae-maydis* lost the ability to infect Tcms maize (Yun *et al.*, 1998). Besides mitochondria, HSTs are also reported to target enzymes or other plant cell organelles like plasma membrane, chloroplast, endoplasmic reticulum, nucleus, vacuole and Golgi bodies, with the objective to suppress host defence responses and/or induce host cell death (Meena *et al.*, 2017).

Another HST toxin is victorin, a family of related, cyclized pentapeptides (Wolpert *et al.*, 1985, 1986) secreted by the necrotrophic fungus *Cochliobolus victoriae* that causes Victoria blight on susceptible oats (Meehan and Murphy, 1947). Fungal pathogenicity is solely attributed to the ability to produce victorin, as victorin-deficient mutants are entirely non-pathogenic (Scheffer *et al.*, 1967). While susceptibility of oats can be traced back to one dominant gene called *Vb* (Welsh *et al.*, 1954), it was later found that a single, dominant gene, called Locus Orchestrating Victorin Effects1 (LOV1), provides victorin susceptibility in Arabidopsis plants (Gilbert and Wolpert, 2013). Interestingly, only oat lines carrying *Pc2*, a resistance gene against crown rust, are susceptible to victorin-producing *C. victoriae* isolates (Litzenberger, 1948; Welsh *et al.*, 1954). As studies to create plants resistant to both Victoria blight and crown rust were unsuccessful, it was suggested that *Vb* and *Pc2* are the same gene conferring susceptibility and resistance, respectively (Litzenberger, 1948; Mayama *et al.*, 1995). Further evidence for this hypothesis was provided by the discovery that victorin perception triggers a defence

response in susceptible oats and Arabidopsis (Wolpert *et al.*, 2002; Gilbert and Wolpert, 2013), hinting that *C. victoriae* hijacks the classic gene-for-gene interaction needed to provide resistance against crown rust and utilizes victorin to elicit host cell death via the same defence mechanism to suit its necrotrophic lifestyle.

The necrotrophic fungus *Parastagonospora nodorum* (formerly *Stagonospora nodorum*) is the causal agent of the Septoria nodorum blotch (SNB) disease on wheat (King *et al.*, 1983; Oliver *et al.*, 2012). Besides the ability to produce CWDEs and non-specific toxins, *P. nodorum* has been characterized for its ability to produce a wide range of HSTs (also called necrotrophic effectors) that result in different levels of susceptibility depending on the wheat cultivar (Keller *et al.*, 1994; Wicki *et al.*, 1999a,b; Friesen and Faris, 2010). So far, a total of nine interactions between necrotrophic effectors of *P. nodorum* and corresponding wheat susceptibility genes have been found (Liu *et al.*, 2004a,b, 2006; Friesen *et al.*, 2006, 2007, 2008, 2009, 2012; Abeysekara *et al.*, 2009, 2012; Chu *et al.*, 2010; Zhang *et al.*, 2011b; Gao *et al.*, 2015; Shi *et al.*, 2015). Furthermore, it was reported that homologues of the necrotrophic *P. nodorum* effector gene *ToxA* have been acquired via horizontal gene transfer and interspecific hybridization by the wheat pathogens *Pyrenophora tritici-repentis*, *Phaeosphaeria avenaria triti* and *B. sorokiniana* (McDonald *et al.*, 2017). So far, two host targets *Snn1* and *Tsn1*, of *P. nodorum* necrotrophic effectors ToxA and Tox1, respectively, have been cloned (Faris *et al.*, 2010; Shi *et al.*, 2016). While *Tsn1* resembles a plant resistance gene structure as it harbors a serine/threonine protein kinase, a nucleotide binding and leucine-rich repeat domains (Faris *et al.*, 2010), *Snn1* is a wall-associated kinase with a predicted transmembrane domain (Shi *et al.*, 2016). However, in both cases interaction with a corresponding necrotrophic effector leads to a so-called necrotrophic effector-triggered susceptibility (Friesen *et al.*, 2007; Faris *et al.*, 2010; Liu *et al.*, 2012; Shi *et al.*, 2016) in opposition to the conventional effector-trigger immunity (ETI) observed in most of the biotrophic interactions. Taken together, these studies suggest that some necrotrophic fungal pathogens use effectors to subvert the host resistance mechanism for their own benefit (Wolpert *et al.*, 2002; Liu *et al.*, 2012, 2015).

**Non-typical effectors: sRNAs.** Small RNAs (sRNA) induce gene silencing by binding to Argonaute (AGO) proteins and directing the RNA-induced silencing complex to genes with complementary sequences (Castel and Martienssen, 2013). As regulatory molecules, sRNAs are involved in a wide range of biological processes, such as organ morphogenesis, genome modification, and adaptive responses to abiotic and biotic stresses (Carrington and Ambros, 2003; Ruiz-Ferrer and Voinnet, 2009; Katiyar-Agarwal and Jin, 2010). Both animals and plants have been reported to

exchange sRNAs with parasites, pathogens or symbiotic organisms in cross-kingdom sRNAs transfer (Wang *et al.*, 2016). It is generally assumed that sRNAs from plants are integral components of plant responses to adverse environmental conditions, including host-microbial interactions (Katiyar-Agarwal and Jin, 2010; Zhang *et al.*, 2011a). While host sRNAs play important roles in pathogen resistance, pathogens also encode sRNAs to manipulate host defence responses and mediate virulence (Weiberg *et al.*, 2013; Chaloner *et al.*, 2016; Wang *et al.*, 2016). The necrotrophic fungus *B. cinerea* infects almost all vegetable and fruit crops, causing major losses worldwide. Recently, it has been reported that some *B. cinerea* sRNAs (Bc-sRNAs) can silence Arabidopsis and tomato genes involved in immunity (Weiberg *et al.*, 2013). The produced Bc-sRNAs hijack the host RNA interference (RNAi) machinery by binding to Arabidopsis AGO1. Furthermore, Bc-sRNAs silence host target immunity genes in both Arabidopsis and tomato plants during fungal infection (Weiberg *et al.*, 2013). Cross-kingdom RNAi to suppress host immunity genes by hijacking host AGO1 has also been reported for *V. dahliae* (Wang *et al.*, 2016). Arabidopsis ago1-27 mutants were less susceptible to the infection with *V. dahliae* than wild-type plants in both soil and root culture conditions (Wang *et al.*, 2016). These results indicate that fungal pathogens and hosts utilize cross-kingdom RNAi to manipulate their interactions to their own benefit.

### Evolution of pathogen virulence

As effectors are pathogen molecules that are crucial for establishing the parasitic symbiosis, hosts continuously evolve to intercept pathogen effectors or their activities with their immune receptor repertoire to halt pathogen ingress (Cook *et al.*, 2015). To avoid or overcome such recognition, pathogens need to be able to swiftly purge or modify effectors that are intercepted by host immune systems, or evolve novel effectors to suppress the reinstated immune response, leading to an everlasting co-evolution between pathogen and host (Jones and Dangl, 2006; Cook *et al.*, 2015). Based on genomics of plant pathogenic species, it has been proposed that many pathogens possess a bipartite genome architecture where effector genes cluster in repeat-rich dynamic compartments, a phenomenon that has been coined a 'two-speed' genome (Croll and McDonald, 2012; Raffaele and Kamoun, 2012). These regions are typically repeat-rich, sometimes with active transposable elements (TEs), and often display increased structural polymorphism, increased point mutagenesis and positive selection (Raffaele *et al.*, 2010; Rouxel *et al.*, 2011; de Jonge *et al.*, 2013; Dong *et al.*, 2015; Faino *et al.*, 2016; Möller and Stukenbrock, 2017). TEs are likely to contribute to pathogen adaptation by facilitating the swift evolution of effector catalogues by establishing genetic variability (Faino *et al.*, 2016; Bao *et al.*, 2017), yet the underlying mechanisms

remain largely unknown (Seidl and Thomma, 2017). However, genomic analysis in *V. dahliae* revealed active and passive contributions of TEs, through TE activity, and through acting as substrate for homology-based double-strand repair pathways, respectively (Faino *et al.*, 2016).

To control the spread and activity of TEs, TE-rich genomic regions are often highly condensed in heterochromatin, which is directed by DNA methylation. As effector genes and other virulence-related genes, such as toxin biosynthesis genes, often reside in TE-rich regions, TEs can impact the expression of these genes (Connolly *et al.*, 2013; Chujo and Scott, 2014; Soyer *et al.*, 2014). Consequently, specific and differential methylation may be associated with adaptive evolution of two-speed pathogen genomes (Seidl *et al.*, 2016; Chen *et al.*, 2017; Seidl and Thomma, 2017). Thus, TEs drive genome and transcriptome variability that, in turn, impacts pathogen adaptation (Seidl and Thomma, 2017).

### CONCLUSION

While all plant pathogenic fungi come across common plant defence mechanisms during host colonization, they employ different strategies to bypass these. As the ongoing co-evolution with their hosts prompts pathogens to appropriately respond to modifications in host immunity in a timely manner, fungi need to continuously adapt their repertoire of virulence strategies to keep their parasitic relationships ongoing. A deep understanding of the molecular mechanisms underlying these virulence strategies and host-pathogen interactions will result in the identification of precise virulence targets in the host plant. Such knowledge is paramount to improve current crop protection strategies or to design novel measures for disease control.

### ACKNOWLEDGEMENTS

Work in the laboratory of B.P.H.J.T. is supported by the Research Council Earth and Life Sciences (ALW) of the Netherlands Organization of Scientific Research (NWO). The authors declare no conflict of interest.

### REFERENCES

- Abeysekara, N.S., Friesen, T.L., Keller, B. and Faris, J.D. (2009) Identification and characterization of a novel host-toxin interaction in the wheat-*Stagonospora nodorum* pathosystem. *Theor. Appl. Genet.* **120**, 117–126.
- Abeysekara, N.S., Faris, J.D., Chao, S., McClean, P.E. and Friesen, T.L. (2012) Whole-Genome QTL analysis of *Stagonospora nodorum* blotch resistance and validation of the SnTox4-Snn4 interaction in hexaploid wheat. *Phytopathology*, **102**, 94–104.
- Albersheim, P. and Valent, B.S. (1974) Host-pathogen interactions. VII. Plant pathogens secrete proteins which inhibit enzymes of the host capable of attacking the pathogen. *Plant Physiol.* **53**, 684–687.
- Alghisi, P. and Favaron, F. (1995) Pectin-degrading enzymes and plant-parasite interactions. *Eur. J. Plant Pathol.* **101**, 365–375.
- Altenbach, D. and Robatzek, S. (2007) Pattern recognition receptors: from the cell surface to intracellular dynamics. *Mol. Plant Microbe Interact.* **20**, 1031–1039.
- Bao, J., Chen, M., Zhong, Z. *et al.* (2017) PacBio sequencing reveals transposable elements as a key contributor to genomic plasticity and virulence variation in *Magnaporthe oryzae*. *Molec. Plant*, **10**, 1465–1468.



- Bindschedler, L.V., Dewdney, J., Blee, K.A. *et al.* (2006) Peroxidase-dependent apoplastic oxidative burst in *Arabidopsis* required for pathogen resistance. *Plant J.* **47**, 851–863.
- Birben, E., Sahiner, U.M., Sackesen, C., Erzurum, S. and Kalayci, O. (2012) Oxidative stress and antioxidant defense. *World Allergy Organ J.* **5**, 9–19.
- Blokhina, O., Virolainen, E. and Fagerstedt, K.V. (2003) Antioxidants, oxidative damage and oxygen deprivation stress: a review. *Ann. Bot.* **91** Spec No, 179–194.
- Boller, T. and Felix, G. (2009) A renaissance of elicitors: perception of microbe-associated molecular patterns and danger signals by pattern-recognition receptors. *Annu. Rev. Plant Biol.* **60**, 379–406.
- Bolton, M.D. (2009) Primary metabolism and plant defense-fuel for the fire. *Mol. Plant Microbe Interact.* **22**, 487–497.
- Bolton, M.D., Thomma, B.P. and Nelson, B.D. (2006) *Sclerotinia sclerotiorum* (Lib.) de Bary: biology and molecular traits of a cosmopolitan pathogen. *Mol. Plant Pathol.* **7**, 1–16.
- Bolwell, G.P., Bindschedler, L.V., Blee, K.A., Butt, V.S., Davies, D.R., Gardner, S.L., Gerrish, C. and Minibayeva, F. (2002) The apoplastic oxidative burst in response to biotic stress in plants: a three-component system. *J. Exp. Bot.* **53**, 1367–1376.
- Buist, G., Steen, A., Kok, J. and Kuipers, O.P. (2008) LysM, a widely distributed protein motif for binding to (peptido)glycans. *Mol. Microbiol.* **68**, 838–847.
- van den Burg, H.A., Harrison, S.J., Joosten, M.H., Vervoort, J. and de Wit, P.J. (2006) *Cladosporium fulvum* Avr4 protects fungal cell walls against hydrolysis by plant chitinases accumulating during infection. *Mol. Plant Microbe Interact.* **19**, 1420–1430.
- Carrington, J.C. and Ambros, V. (2003) Role of microRNAs in plant and animal development. *Science*, **301**, 336–338.
- Castel, S.E. and Martienssen, R.A. (2013) RNA interference in the nucleus: roles for small RNAs in transcription, epigenetics and beyond. *Nat. Rev. Genet.* **14**, 100–112.
- Chaloner, T., van Kan, J.A. and Grant-Downton, R.T. (2016) RNA 'information warfare' in pathogenic and mutualistic interactions. *Trends Plant Sci.* **21**, 738–748.
- Chanclud, E. and Morel, J.B. (2016) Plant hormones: a fungal point of view. *Mol. Plant Pathol.* **17**, 1289–1297.
- Chanclud, E., Kisiala, A., Emery, N.R., Chalvon, V., Ducasse, A., Romiti-Michel, C., Gravot, A., Kroj, T. and Morel, J.B. (2016) Cytokinin production by the rice blast fungus is a pivotal requirement for full virulence. *PLoS Pathog.* **12**, e1005457.
- Chen, K., Liu, J., Liu, S.X., Xia, M., Zhang, X.M., Han, D., Jiang, Y.M., Wang, C.M. and Cao, X.T. (2017) Methyltransferase SETD2-mediated methylation of STAT1 is critical for interferon antiviral activity. *Cell*, **170**, 492.
- Chu, C.-G., Faris, J.D., Xu, S.S. and Friesen, T.L. (2010) Genetic analysis of disease susceptibility contributed by the compatible Tsn1-SnToxA and Snn1-SnTox1 interactions in the wheat-*Stagonospora nodorum* pathosystem. *Theor. Appl. Genet.* **120**, 1451–1459.
- Chujo, T. and Scott, B. (2014) Histone H3K9 and H3K27 methylation regulates fungal alkaloid biosynthesis in a fungal endophyte-plant symbiosis. *Molec. Microbiol.* **92**, 413–434.
- Connolly, L.R., Smith, K.M. and Freitag, M. (2013) The *Fusarium graminearum* histone H3 K27 methyltransferase KMT6 regulates development and expression of secondary metabolite gene clusters. *PLoS Genet.* **9**, e1003916.
- Cook, D.E., Mesarich, C.H. and Thomma, B.P. (2015) Understanding plant immunity as a surveillance system to detect invasion. *Annu. Rev. Phytopathol.* **53**, 541–563.
- Cosio, E.G., Feger, M., Miller, C.J., Antelo, L. and Ebel, J. (1996) High-affinity binding of fungal  $\beta$ -glucan elicitors to cell membranes of species of the plant family *Fabaceae*. *Planta*, **200**, 92–99.
- Cote, F., Roberts, K.A. and Hahn, M.G. (2000) Identification of high-affinity binding sites for the hepta- $\beta$ -glucoside elicitor in membranes of the model legumes *Medicago truncatula* and *Lotus japonicus*. *Planta*, **211**, 596–605.
- Couto, D. and Zipfel, C. (2016) Regulation of pattern recognition receptor signalling in plants. *Nat. Rev. Immunol.* **16**, 537–552.
- Croll, D. and McDonald, B.A. (2012) The accessory genome as a cradle for adaptive evolution in pathogens. *PLoS Pathog.* **8**, e1002608.
- Daub, M.E. and Ehrenshaft, M. (2000) The photoactivated *Cercospora* toxin cercosporin: contributions to plant disease and fundamental biology. *Annu. Rev. Phytopathol.* **38**, 461–490.
- Dean, R.A., Talbot, N.J., Ebbole, D.J. *et al.* (2005) The genome sequence of the rice blast fungus *Magnaporthe grisea*. *Nature*, **434**, 980–986.
- Djamei, A., Schipper, K., Rabe, F. *et al.* (2011) Metabolic priming by a secreted fungal effector. *Nature*, **478**, 395–398.
- Doehlemann, G., van der Linde, K., Assmann, D., Schwammbach, D., Hof, A., Mohanty, A., Jackson, D. and Kahmann, R. (2009) Pep1, a secreted effector protein of *Ustilago maydis*, is required for successful invasion of plant cells. *PLoS Pathog.* **5**, e1000290.
- Doehlemann, G., Reissmann, S., Assmann, D., Fleckenstein, M. and Kahmann, R. (2011) Two linked genes encoding a secreted effector and a membrane protein are essential for *Ustilago maydis*-induced tumour formation. *Mol. Microbiol.* **81**, 751–766.
- Dong, Y.H., Li, Y., Zhao, M.M. *et al.* (2015) Global genome and transcriptome analyses of *Magnaporthe oryzae* epidemic isolate 98-06 uncover novel effectors and pathogenicity-related genes, revealing gene gain and loss dynamics in genome evolution. *PLoS Pathog.* **11**, e1004801.
- El Gueddari, N.E., Rauchhaus, U., Moerschbacher, B.M. and Deising, H.B. (2002) Developmentally regulated conversion of surface-exposed chitin to chitosan in cell walls of plant pathogenic fungi. *New Phytol.* **156**, 103–112.
- van Esse, H.P., Van't Klooster, J.W., Bolton, M.D., Yadeta, K.A., van Baarlen, P., Boeren, S., Vervoort, J., de Wit, P.J. and Thomma, B.P. (2008) The *Cladosporium fulvum* virulence protein Avr2 inhibits host proteases required for basal defense. *Plant Cell*, **20**, 1948–1963.
- van Esse, H.P., Bolton, M.D., Stergiopoulos, I., de Wit, P.J. and Thomma, B.P. (2007) The chitin-binding *Cladosporium fulvum* effector protein Avr4 is a virulence factor. *Mol. Plant Microbe Interact.* **20**, 1092–1101.
- Faino, L., Seidl, M.F., Shi-Kunne, X., Pauper, M., van den Berg, G.C.M., Wittenberg, A.H.J. and Thomma, B.P.H.J. (2016) Transposons passively and actively contribute to evolution of the two-speed genome of a fungal pathogen. *Genome Res.* **26**, 1091–1100.
- Faris, J.D., Zhang, Z., Lu, H. *et al.* (2010) A unique wheat disease resistance-like gene governs effector-triggered susceptibility to necrotrophic pathogens. *Proc. Natl Acad. Sci. USA*, **107**, 13544–13549.
- Felix, G., Regenass, M. and Boller, T. (1993) Specific perception of sub-nanomolar concentrations of chitin fragments by tomato cells: induction of extracellular alkalization, changes in protein phosphorylation, and establishment of a refractory state. *Plant J.* **4**, 307–316.
- Fessel, P.H. and Zuccaro, A. (2016)  $\beta$ -glucan: crucial component of the fungal cell wall and elusive MAMP in plants. *Fungal Genet. Biol.* **90**, 53–60.
- Friesen, T.L. and Faris, J.D. (2010) Characterization of the wheat-*Stagonospora nodorum* disease system: what is the molecular basis of this quantitative necrotrophic disease interaction? *Can. J. Plant Pathol.* **32**, 20–28.
- Friesen, T.L., Stukenbrock, E.H., Liu, Z., Meinhardt, S., Ling, H., Faris, J.D., Rasmussen, J.B., Solomon, P.S., McDonald, B.A. and Oliver, R.P. (2006) Emergence of a new disease as a result of interspecific virulence gene transfer. *Nat. Genet.* **38**, 953–956.
- Friesen, T.L., Meinhardt, S.W. and Faris, J.D. (2007) The *Stagonospora nodorum*-wheat pathosystem involves multiple proteinaceous host-selective toxins and corresponding host sensitivity genes that interact in an inverse gene-for-gene manner. *Plant J.* **51**, 681–692.
- Friesen, T.L., Zhang, Z., Solomon, P.S., Oliver, R.P. and Faris, J.D. (2008) Characterization of the interaction of a novel *Stagonospora nodorum* host-selective toxin with a wheat susceptibility gene. *Plant Physiol.* **146**, 682–693.
- Friesen, T.L., Chu, C.-G., Liu, Z.H., Xu, S.S., Halley, S. and Faris, J.D. (2009) Host-selective toxins produced by *Stagonospora nodorum* confer disease susceptibility in adult wheat plants under field conditions. *Theor. Appl. Genet.* **118**, 1489–1497.
- Friesen, T.L., Chu, C., Xu, S.S. and Faris, J.D. (2012) SnTox5-Snn5: a novel *Stagonospora nodorum* effector-wheat gene interaction and its relationship with the SnToxA-Tsn1 and SnTox3-Snn3-B1 interactions. *Molec. Plant Pathol.* **13**, 1101–1109.
- Fujikawa, T., Sakaguchi, A., Nishizawa, Y., Kouzai, Y., Minami, E., Yano, S., Koga, H., Meshi, T. and Nishimura, M. (2012) Surface  $\alpha$ -1,3-glucan facilitates fungal stealth infection by interfering with innate immunity in plants. *PLoS Pathog.* **8**, e1002882.



- Gao, Y., Faris, J.D., Liu, Z., Kim, Y.M., Syme, R.A., Oliver, R.P., Xu, S.S. and Friesen, T.L. (2015) Identification and characterization of the SnTox6-Snn6 interaction in the *Parastagonospora nodorum*-wheat pathosystem. *Molec. Plant-Microbe Interact.* **28**, 615–625.
- Gilbert, B.M. and Wolpert, T.J. (2013) Characterization of the LOV1-mediated, victorin-induced, cell-death response with virus-induced gene silencing. *Molec. Plant-Microbe Interact.* **26**, 903–917.
- Giraldo, M.C., Dagdas, Y.F., Gupta, Y.K., Mentlak, T.A., Yi, M., Martinez-Rocha, A.L., Saitoh, H., Terauchi, R., Talbot, N.J. and Valent, B. (2013) Two distinct secretion systems facilitate tissue invasion by the rice blast fungus *Magnaporthe oryzae*. *Nat. Commun.* **4**, 1996.
- Glazebrook, J. (2005) Contrasting mechanisms of defense against biotrophic and necrotrophic pathogens. *Annu. Rev. Phytopathol.* **43**, 205–227.
- Hemetsberger, C., Herrberger, C., Zechmann, B., Hillmer, M. and Doeblemann, G. (2012) The *Ustilago maydis* effector Pep1 suppresses plant immunity by inhibition of host peroxidase activity. *PLoS Pathog.* **8**, e1002684.
- van der Hoorn, R.A. (2008) Plant proteases: from phenotypes to molecular mechanisms. *Annu. Rev. Plant Biol.* **59**, 191–223.
- Jashni, M.K., Dols, I.H., Iida, Y., Boeren, S., Beenen, H.G., Mehrabi, R., Collemare, J. and de Wit, P.J. (2015a) Synergistic action of a metalloprotease and a serine protease from *Fusarium oxysporum* f. sp. *lycopersici* cleaves chitin-binding tomato chitinases, reduces their antifungal activity, and enhances fungal virulence. *Mol. Plant Microbe Interact.* **28**, 996–1008.
- Jashni, M.K., Mehrabi, R., Collemare, J., Mesarich, C.H. and de Wit, P.J. (2015b) The battle in the apoplast: further insights into the roles of proteases and their inhibitors in plant-pathogen interactions. *Front. Plant Sci.* **6**, 584.
- Jiang, Y., Wang, W., Xie, Q. et al. (2017) Plants transfer lipids to sustain colonization by mutualistic mycorrhizal and parasitic fungi. *Science*, **356**, 1172–1175.
- Jones, J.D. and Dangl, J.L. (2006) The plant immune system. *Nature*, **444**, 323–329.
- de Jonge, R. and Thomma, B.P. (2009) Fungal LysM effectors: extinguishers of host immunity? *Trends Microbiol.* **17**, 151–157.
- de Jonge, R., Ebert, M.K., Huit-Roehl, C.R. et al. (2017) Conservation of a gene cluster reveals novel cercosporin biosynthetic mechanisms and extends production to the genus *Colletotrichum*. *BioRxiv*. <https://doi.org/10.1101/100545>
- de Jonge, R., van Esse, H.P., Kombrink, A., Shinya, T., Desaki, Y., Bours, R., van der Krol, S., Shibuya, N., Joosten, M.H. and Thomma, B.P. (2010) Conserved fungal LysM effector Ecp6 prevents chitin-triggered immunity in plants. *Science*, **329**, 953–955.
- de Jonge, R., Bolton, M.D., Kombrink, A., van den Berg, G.C.M., Yadeta, K.A. and Thomma, B.P.H.J. (2013) Extensive chromosomal reshuffling drives evolution of virulence in an asexual pathogen. *Genome Res.* **23**, 1271–1282.
- Kale, S.D. and Tyler, B.M. (2011) Entry of oomycete and fungal effectors into plant and animal host cells. *Cell. Microbiol.* **13**, 1839–1848.
- Katiyar-Agarwal, S. and Jin, H. (2010) Role of small RNAs in host-microbe interactions. *Annu. Rev. Phytopathol.* **48**, 225–246.
- Keller, B., Winzeler, H., Winzeler, M. and Fried, P.M. (1994) Differential sensitivity of wheat embryos against extracts containing toxins of *Septoria nodorum*: first steps towards *in vitro* selection. *J. Phytopathol.* **141**, 233–240.
- Khang, C.H., Berruyer, R., Giraldo, M.C., Kankanala, P., Park, S.Y., Czymmek, K., Kang, S. and Valent, B. (2010) Translocation of *Magnaporthe oryzae* effectors into rice cells and their subsequent cell-to-cell movement. *Plant Cell*, **22**, 1388–1403.
- King, J.E., Cook, R.J. and Melville, S.C. (1983) A review of *Septoria* diseases of wheat and barley. *Ann. Appl. Biol.* **103**, 345–373.
- King, R., Urban, M., Lauder, R.P., Hawkins, N., Evans, M., Plummer, A., Halsey, K., Lovegrove, A., Hammond-Kosack, K. and Rudd, J.J. (2017) A conserved fungal glycosyltransferase facilitates pathogenesis of plants by enabling hyphal growth on solid surfaces. *PLoS Pathog.* **13**, e1006672.
- Kleemann, J., Rincon-Rivera, L.J., Takahara, H. et al. (2012) Sequential delivery of host-induced virulence effectors by appressoria and intracellular hyphae of the phytopathogen *Colletotrichum higginsianum*. *PLoS Pathog.* **8**, e1002643.
- Kloppholz, S., Kuhn, H. and Requena, N. (2011) A secreted fungal effector of *Glomus intraradices* promotes symbiotic biotrophy. *Curr. Biol.* **21**, 1204–1209.
- Kombrink, A., Rovenich, H., Shi-Kunne, X. et al. (2017) *Verticillium dahliae* LysM effectors differentially contribute to virulence on plant hosts. *Mol. Plant Pathol.* **18**, 596–608.
- Kruger, J., Thomas, C.M., Golstein, C., Dixon, M.S., Smoker, M., Tang, S., Mulder, L. and Jones, J.D. (2002) A tomato cysteine protease required for Cf-2-dependent disease resistance and suppression of autonecrosis. *Science*, **296**, 744–747.
- Lamb, C. and Dixon, R.A. (1997) The oxidative burst in plant disease resistance. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **48**, 251–275.
- Levine, A., Tenhaken, R., Dixon, R. and Lamb, C. (1994) H<sub>2</sub>O<sub>2</sub> from the oxidative burst orchestrates the plant hypersensitive disease resistance response. *Cell*, **79**, 583–593.
- Levings III, C.S., Rhoads, D.M. and Siedow, J.N. (1995) Molecular interactions of *Bipolaris maydis* T-toxin and maize. *Can. J. Botany*, **73**, 483–489.
- Litzenberger, S.C. (1948) Inheritance of resistance to specific races of crown rust, stem rust, and to Helminthosporium blight of oats. *Res. Bull.* **29**, Article 1.
- Liu, Z.H., Faris, J.D., Meinhardt, S.W., Ali, S., Rasmussen, J.B. and Friesen, T.L. (2004a) Genetic and physical mapping of a gene conditioning sensitivity in wheat to a partially purified host-selective toxin produced by *Stagonospora nodorum*. *Phytopathology*, **94**, 1056–1060.
- Liu, Z.H., Friesen, T.L., Rasmussen, J.B., Ali, S., Meinhardt, S.W. and Faris, J.D. (2004b) Quantitative trait loci analysis and mapping of seedling resistance to *Stagonospora nodorum* leaf blotch in wheat. *Phytopathology*, **94**, 1061–1067.
- Liu, Z., Friesen, T.L., Ling, H., Meinhardt, S.W., Oliver, R.P., Rasmussen, J.B. and Faris, J.D. (2006) The Tsn1-ToxA interaction in the wheat-*Stagonospora nodorum* pathosystem parallels that of the wheat-tan spot system. *Genome*, **49**, 1265–1273.
- Liu, Z., Zhang, Z., Faris, J.D. et al. (2012) The cysteine rich necrotrophic effector SnTox1 produced by *Stagonospora nodorum* triggers susceptibility of wheat lines harboring Snn1. *PLoS Pathog.* **8**, e1002467.
- Liu, T., Song, T., Zhang, X. et al. (2014) Unconventionally secreted effectors of two filamentous pathogens target plant salicylate biosynthesis. *Nat. Commun.* **5**, 4686.
- Liu, Z., Holmes, D.J., Faris, J.D., Chao, S., Brueggeman, R.S., Edwards, M.C. and Friesen, T.L. (2015) Necrotrophic effector-triggered susceptibility (NETS) underlies the barley-Pyrenophora teres f. teres interaction specific to chromosome 6H. *Mol. Plant Pathol.* **16**, 188–200.
- Lombard, V., Golaconda Ramulu, H., Drula, E., Coutinho, P.M. and Henrissat, B. (2014) The carbohydrate-active enzymes database (CAZy) in 2013. *Nucl. Acids Res.* **42**, 490–4955.
- Macho, A.P. and Zipfel, C. (2014) Plant PRRs and the activation of innate immune signaling. *Mol. Cell*, **54**, 263–272.
- Marshall, R., Kombrink, A., Motteram, J., Loza-Reyes, E., Lucas, J., Hammond-Kosack, K.E., Thomma, B.P. and Rudd, J.J. (2011) Analysis of two in planta expressed LysM effector homologs from the fungus *Mycosphaerella graminicola* reveals novel functional properties and varying contributions to virulence on wheat. *Plant Physiol.* **156**, 756–769.
- Masachis, S., Segorbe, D., Turra, D. et al. (2016) A fungal pathogen secretes plant alkalizing peptides to increase infection. *Nat. Microbiol.* **1**, 16043.
- Mayama, S., Bordin, A.P.A., Morikawa, T., Tanpo, H. and Kato, H. (1995) Association of avenalumin accumulation with co-segregation of victorin sensitivity and crown rust resistance in oat lines carrying the Pc-2 gene. *Physiol. Molec. Plant Pathol.* **46**, 263–274.
- McDonald, M.C., Ahren, D., Simpfendorfer, S., Milgate, A. and Solomon, P.S. (2017) The discovery of the virulence gene ToxA in the wheat and barley pathogen *Bipolaris sorokiniana*. *Mol. Plant Pathol.* <https://doi.org/10.1111/mpp.12535>
- Meehan, F. and Murphy, H. (1947) Differential phytotoxicity of metabolic by-products of *Helminthosporium victoriae*. *Science*, **106**, 270–271.
- Meena, M., Gupta, S.K., Swapnil, P., Zehra, A., Dubey, M.K. and Upadhyay, R.S. (2017) Alternaria toxins: potential virulence factors and genes related to pathogenesis. *Front. Microbiol.* **8**, 1451.
- Mendgen, K. and Hahn, M. (2002) Plant infection and the establishment of fungal biotrophy. *Trends Plant Sci.* **7**, 352–356.

- Mentlak, T.A., Kombrink, A., Shinya, T. *et al.* (2012) Effector-mediated suppression of chitin-triggered immunity by *Magnaporthe oryzae* is necessary for rice blast disease. *Plant Cell*, **24**, 322–335.
- Möller, M. and Stukenbrock, E.H. (2017) Evolution and genome architecture in fungal plant pathogens. *Nat. Rev. Microbiol.* **15**, 756–771.
- Mueller, A.N., Ziemann, S., Treitschke, S., Assmann, D. and Doehlemann, G. (2013) Compatibility in the *Ustilago maydis*-maize interaction requires inhibition of host cysteine proteases by the fungal effector Pit2. *PLoS Pathog.* **9**, e1003177.
- Murphy, E. and De Smet, I. (2014) Understanding the RALF family: a tale of many species. *Trends Plant Sci.* **19**, 664–671.
- Naumann, T.A. and Wicklow, D.T. (2010) Allozyme-specific modification of a maize seed chitinase by a protein secreted by the fungal pathogen *Stenocarpella maydis*. *Phytopathology*, **100**, 645–654.
- Naumann, T.A., Wicklow, D.T. and Kendra, D.F. (2009) Maize seed chitinase is modified by a protein secreted by *Bipolaris zeicola*. *Physiol. Mol. Plant Pathol.* **74**, 134–141.
- Naumann, T.A., Wicklow, D.T. and Price, N.P. (2011) Identification of a chitinase-modifying protein from *Fusarium verticillioides*: truncation of a host resistance protein by a fungicidal metalloprotease. *J. Biol. Chem.* **286**, 35358–35366.
- Oliveira-Garcia, E. and Deising, H.B. (2013) Infection structure-specific expression of  $\beta$ -1,3-glucan synthase is essential for pathogenicity of *Colletotrichum graminicola* and evasion of  $\beta$ -glucan-triggered immunity in maize. *Plant Cell*, **25**, 2356–2378.
- Oliveira-Garcia, E. and Deising, H.B. (2016) Attenuation of PAMP-triggered immunity in maize requires down-regulation of the key  $\beta$ -1,6-glucan synthesis genes *KRE5* and *KRE6* in biotrophic hyphae of *Colletotrichum graminicola*. *Plant J.* **87**, 355–375.
- Oliver, R.P., Friesen, T.L., Faris, J.D. and Solomon, P.S. (2012) *Stagonospora nodorum*: from pathology to genomics and host resistance. *Annu. Rev. Phytopathol.* **50**, 23–43.
- Orozco-Cardenas, M.L., Narvaez-Vasquez, J. and Ryan, C.A. (2001) Hydrogen peroxide acts as a second messenger for the induction of defense genes in tomato plants in response to wounding, systemin, and methyl jasmonate. *Plant Cell*, **13**, 179–191.
- Paccanaro, M.C., Sella, L., Castiglioni, C., Giacomello, F., Martinez-Rocha, A.L., D'Ovidio, R., Schafer, W. and Favaron, F. (2017) Synergistic effect of different plant cell wall-degrading enzymes is important for virulence of *Fusarium graminearum*. *Mol. Plant Microbe Interact.* **30**, 886–895.
- Petre, B. and Kamoun, S. (2014) How do filamentous pathogens deliver effector proteins into plant cells? *PLoS Biol.* **12**, e1001801.
- Plett, J.M., Daguerre, Y., Wittulsky, S. *et al.* (2014) Effector MiSSP7 of the mutualistic fungus *Laccaria bicolor* stabilizes the *Populus* JAZ6 protein and represses jasmonic acid (JA) responsive genes. *Proc. Natl Acad. Sci. USA*, **111**, 8299–8304.
- Prusky, D. and Yakoby, N. (2003) Pathogenic fungi: leading or led by ambient pH? *Mol. Plant Pathol.* **4**, 509–516.
- Prusky, D., McEvoy, J.L., Leverenz, B. and Conway, W.S. (2001) Local modulation of host pH by *Colletotrichum* species as a mechanism to increase virulence. *Mol. Plant Microbe Interact.* **14**, 1105–1113.
- Qi, J., Wang, J., Gong, Z. and Zhou, J.M. (2017) Apoplastic ROS signaling in plant immunity. *Curr. Opin. Plant Biol.* **38**, 92–100.
- Rabe, F., Ajami-Rashidi, Z., Doehlemann, G., Kahmann, R. and Djamei, A. (2013) Degradation of the plant defence hormone salicylic acid by the biotrophic fungus *Ustilago maydis*. *Mol. Microbiol.* **89**, 179–188.
- Raffaele, S. and Kamoun, S. (2012) Genome evolution in filamentous plant pathogens: why bigger can be better. *Nat. Rev. Microbiol.* **10**, 417–430.
- Raffaele, S., Farrer, R.A., Cano, L.M. *et al.* (2010) Genome evolution following host jumps in the Irish potato famine pathogen lineage. *Science*, **330**, 1540–1543.
- Rafiqi, M., Gan, P.H., Ravensdale, M., Lawrence, G.J., Ellis, J.G., Jones, D.A., Hardham, A.R. and Dodds, P.N. (2010) Internalization of flax rust avirulence proteins into flax and tobacco cells can occur in the absence of the pathogen. *Plant Cell*, **22**, 2017–2032.
- Rafiqi, M., Ellis, J.G., Ludowici, V.A., Hardham, A.R. and Dodds, P.N. (2012) Challenges and progress towards understanding the role of effectors in plant-fungal interactions. *Curr. Opin. Plant Biol.* **15**, 477–482.
- Reineke, G., Heinze, B., Schirawski, J., Buettner, H., Kahmann, R. and Basse, C.W. (2008) Indole-3-acetic acid (IAA) biosynthesis in the smut fungus *Ustilago maydis* and its relevance for increased IAA levels in infected tissue and host tumour formation. *Mol. Plant Pathol.* **9**, 339–355.
- Ribot, C., Cesari, S., Abidi, I., Chalvon, V., Bournaud, C., Vallet, J., Lebrun, M.H., Morel, J.B. and Kroj, T. (2013) The *Magnaporthe oryzae* effector AVR1-CO39 is translocated into rice cells independently of a fungal-derived machinery. *Plant J.* **74**, 1–12.
- Rooney, H.C., Van't Klooster, J.W., van der Hoorn, R.A., Joosten, M.H., Jones, J.D. and de Wit, P.J. (2005) Cladosporium Avr2 inhibits tomato Rcr3 protease required for Cf-2-dependent disease resistance. *Science*, **308**, 1783–1786.
- Rouxel, T., Grandaubert, J., Hane, J.K. *et al.* (2011) Effector diversification within compartments of the *Leptosphaeria maculans* genome affected by Repeat-Induced Point mutations. *Nat. Commun.* **2**, 202.
- Rovenich, H., Boshoven, J.C. and Thomma, B.P. (2014) Filamentous pathogen effector functions: of pathogens, hosts and microbiomes. *Curr. Opin. Plant Biol.* **20**, 96–103.
- Rovenich, H., Zuccaro, A. and Thomma, B.P.H.J. (2016) Convergent evolution of filamentous microbes towards evasion of glycan-triggered immunity. *New Phytol.* **212**, 896–901.
- Ruiz-Ferrer, V. and Voinnet, O. (2009) Roles of plant small RNAs in biotic stress responses. *Annu. Rev. Plant Biol.* **60**, 485–510.
- Ryder, L.S. and Talbot, N.J. (2015) Regulation of appressorium development in pathogenic fungi. *Curr. Opin. Plant Biol.* **26**, 8–13.
- Sanchez-Vallet, A., Saleem-Batcha, R., Kombrink, A., Hansen, G., Valkenburg, D.J., Thomma, B.P. and Mesters, J.R. (2013) Fungal effector Ecp6 outcompetes host immune receptor for chitin binding through intrachain LysM dimerization. *Elife*, **2**, e00790.
- Sanchez-Vallet, A., Mesters, J.R. and Thomma, B.P. (2015) The battle for chitin recognition in plant-microbe interactions. *FEMS Microbiol. Rev.* **39**, 171–183.
- Sasaki, K., Iwai, T., Hiraga, S. *et al.* (2004) Ten rice peroxidases redundantly respond to multiple stresses including infection with rice blast fungus. *Plant Cell Physiol.* **45**, 1442–1452.
- Scheffer, R.P., Nelson, R.R. and Ullstrup, A.J. (1967) Inheritance of toxin production and pathogenicity in *Cochliobolus carbonum* and *Cochliobolus victoriae*. *Phytopathology*, **57**, 1288.
- Seidl, M.F. and Thomma, B.P.H.J. (2017) Transposable elements direct the coevolution between plants and microbes. *Trends Genet.* **33**, 842–851.
- Seidl, M.F., Cook, D.E. and Thomma, B.P.H.J. (2016) Chromatin biology impacts adaptive evolution of filamentous plant pathogens. *PLoS Pathog.* **12**, e1005920.
- Shabab, M., Shindo, T., Gu, C., Kaschani, F., Pansuriya, T., Chintia, R., Harzen, A., Colby, T., Kamoun, S. and van der Hoorn, R.A.L. (2008) Fungal effector protein AVR2 targets diversifying defense-related Cys proteases of tomato. *Plant Cell*, **20**, 1169–1183.
- Shi, G., Friesen, T.L., Saini, J., Xu, S.S., Rasmussen, J.B. and Faris, J.D. (2015) The wheat Snn7 gene confers susceptibility on recognition of the *Parastagonospora nodorum* necrotrophic effector SnTox7. *Plant Genom.* **8**.
- Shi, G., Zhang, Z., Friesen, T.L. *et al.* (2016) The hijacking of a receptor kinase-driven pathway by a wheat fungal pathogen leads to disease. *Sci. Adv.* **2**, e1600822.
- Shibuya, N., Kaku, H., Kuchitsu, K. and Maliarik, M.J. (1993) Identification of a novel high-affinity binding site for N-acetylchitooligosaccharide elicitor in the membrane fraction from suspension-cultured rice cells. *FEBS Lett.* **329**, 75–78.
- Sorensen, J.L., Benfield, A.H., Wollenberg, R.D. *et al.* (2017) The cereal pathogen *Fusarium pseudograminearum* produces a new class of active cytokinins during infection. *Mol. Plant Pathol.*
- Soyer, J.L., El Ghalid, M., Glaser, N. *et al.* (2014) Epigenetic control of effector gene expression in the plant pathogenic fungus *Leptosphaeria maculans*. *PLoS Genet.* **10**, e1004227.
- Stergiopoulos, I. and de Wit, P.J. (2009) Fungal effector proteins. *Annu. Rev. Phytopathol.* **47**, 233–263.
- Stergiopoulos, I., van den Burg, H.A., Okmen, B., Beenen, H.G., van Liere, S., Kema, G.H. and de Wit, P.J. (2010) Tomato Cf resistance proteins mediate recognition of cognate homologous effectors from fungi pathogenic on dicots and monocots. *Proc. Natl Acad. Sci. USA*, **107**, 7610–7615.
- Takahara, H., Hacquard, S., Kombrink, A. *et al.* (2016) *Colletotrichum higginsianum* extracellular LysM proteins play dual roles in appressorial

- function and suppression of chitin-triggered plant immunity. *New Phytol.* **211**, 1323–1327.
- Thomma, B.P.H.J., Eggermont, K., Penninckx, I.A.M.A., Mauch-Mani, B., Vogelsang, R., Cammue, B.P.A. and Broekaert, W.F. (1998) Separate jasmonate-dependent and salicylate-dependent defense-response pathways in *Arabidopsis* are essential for resistance to distinct microbial pathogens. *Proc. Natl Acad. Sci. USA*, **95**, 15107–15111.
- Thomma, B.P.H.J., Tierens, K.F.M., Penninckx, I.A.M.A., Mauch-Mani, B., Broekaert, W.F. and Cammue, B.P.A. (2001) Different micro-organisms differentially induce *Arabidopsis* disease response pathways. *Plant Physiol. Bioch.* **39**, 673–680.
- Thynne, E., Saur, I.M.L., Simbaqueba, J. et al. (2017) Fungal phytopathogens encode functional homologues of plant rapid alkalization factor (RALF) peptides. *Mol. Plant Pathol.* **18**, 811–824.
- Tsuge, T., Harimoto, Y., Akimitsu, K., Ohtani, K., Kodama, M., Akagi, Y., Egusa, M., Yamamoto, M. and Otani, H. (2013) Host-selective toxins produced by the plant pathogenic fungus *Alternaria alternata*. *FEMS Microbiol. Rev.* **37**, 44–66.
- Tundo, S., Kalunke, R., Janni, M., Volpi, C., Lionetti, V., Bellincampi, D., Favaron, F. and D'Ovidio, R. (2016) Pyramiding PvPGIP2 and TAXI-III but not PvPGIP2 and PME1 enhances resistance against *Fusarium graminearum*. *Mol. Plant Microbe Interact.* **29**, 629–639.
- Volpi, C., Janni, M., Lionetti, V., Bellincampi, D., Favaron, F. and D'Ovidio, R. (2011) The ectopic expression of a pectin methyl esterase inhibitor increases pectin methyl esterification and limits fungal diseases in wheat. *Mol. Plant Microbe Interact.* **24**, 1012–1019.
- Wang, M., Weiberg, A., Lin, F.M., Thomma, B.P., Huang, H.D. and Jin, H. (2016) Bidirectional cross-kingdom RNAi and fungal uptake of external RNAs confer plant protection. *Nat. Plants*, **2**, 16151.
- Wawra, S., Fesel, P., Widmer, H. et al. (2016) The fungal-specific  $\beta$ -glucan-binding lectin FGB1 alters cell-wall composition and suppresses glucan-triggered immunity in plants. *Nat. Commun.* **7**, 13188.
- Weiberg, A., Wang, M., Lin, F.M., Zhao, H., Zhang, Z., Kaloshian, I., Huang, H.D. and Jin, H. (2013) Fungal small RNAs suppress plant immunity by hijacking host RNA interference pathways. *Science*, **342**, 118–123.
- Welsh, J.N., Peturson, B. and Machacek, J.E. (1954) Associated inheritance of reaction to races of crown rust, *Puccinia coronata avenae* Erikss, and to Victoria blight, *Helminthosporium victoriae* M. and M. in oats. *Can. J. Botany-Revue Canadienne De Botanique*, **32**, 55.
- Weltmeier, F., Maser, A., Menze, A., Hennig, S., Schad, M., Breuer, F., Schulz, B., Holtschulte, B., Nehls, R. and Stahl, D.J. (2011) Transcript profiles in sugar beet genotypes uncover timing and strength of defense reactions to *Cercospora beticola* infection. *Mol. Plant Microbe Interact.* **24**, 758–772.
- Whisson, S.C., Boevink, P.C., Moleleki, L. et al. (2007) A translocation signal for delivery of oomycete effector proteins into host plant cells. *Nature*, **450**, 115–118.
- Wicki, W., Messmer, M., Winzeler, M., Stamp, P. and Schmid, J.E. (1999a) *In vitro* screening for resistance against *Septoria nodorum* blotch in wheat. *Theor. Appl. Genet.* **99**, 1273–1280.
- Wicki, W., Winzeler, M., Schmid, J.E., Stamp, P. and Messmer, M. (1999b) Inheritance of resistance to leaf and glume blotch caused by *Septoria nodorum* Berk. in winter wheat. *Theor. Appl. Genet.* **99**, 1265–1272.
- Wolpert, T.J., Macko, V., Acklin, W., Jaun, B., Seibl, J., Meili, J. and Arigoni, D. (1985) Structure of Victorin-C, the major host-selective toxin from *Cochliobolus victoriae*. *Experientia*, **41**, 1524–1529.
- Wolpert, T.J., Macko, V., Acklin, W., Jaun, B. and Arigoni, D. (1986) Structure of minor host-selective toxins from *Cochliobolus victoriae*. *Experientia*, **42**, 1296–1299.
- Wolpert, T.J., Dunkle, L.D. and Ciuffetti, L.M. (2002) Host-selective toxins and avirulence determinants: what's in a name? *Annu. Rev. Phytopathol.* **40**, 251–285.
- Wu, G.S., Shortt, B.J., Lawrence, E.B., Leon, J., Fitzsimmons, K.C., Levine, E.B., Raskin, I. and Shah, D.M. (1997) Activation of host defense mechanisms by elevated production of  $H_2O_2$  in transgenic plants. *Plant Physiol.* **115**, 427–435.
- Yang, G., Turgeon, B.G. and Yoder, O.C. (1994) Toxin-deficient mutants from a toxin-sensitive transformant of *Cochliobolus heterostrophus*. *Genetics*, **137**, 751–757.
- Yin, C., Park, J.J., Gang, D.R. and Hulbert, S.H. (2014) Characterization of a tryptophan 2-monooxygenase gene from *Puccinia graminis* f. sp. *tritici* involved in auxin biosynthesis and rust pathogenicity. *Mol. Plant Microbe Interact.* **27**, 227–235.
- Yun, S.H., Turgeon, B.G. and Yoder, O.C. (1998) REMI-induced mutants of *Mycosphaerella zeae-maydis* lacking the polyketide PM-toxin are deficient in pathogenesis to corn. *Physiol. Mol. Plant Pathol.* **52**, 53–66.
- Zhang, X., Zhao, H., Gao, S., Wang, W.C., Katiyar-Agarwal, S., Huang, H.D., Raikhel, N. and Jin, H. (2011a) *Arabidopsis* Argonaute 2 regulates innate immunity via miRNA393(\*)-mediated silencing of a Golgi-localized SNARE gene, *MEMB12*. *Mol. Cell*, **42**, 356–366.
- Zhang, Z., Friesen, T.L., Xu, S.S., Shi, G., Liu, Z., Rasmussen, J.B. and Faris, J.D. (2011b) Two putatively homoeologous wheat genes mediate recognition of SnTox3 to confer effector-triggered susceptibility to *Stagonospora nodorum*. *Plant J.* **65**, 27–38.
- Zhao, Z., Liu, H., Wang, C. and Xu, J.R. (2013) Comparative analysis of fungal genomes reveals different plant cell wall degrading capacity in fungi. *BMC Genom.* **14**, 274.