


## ANNALS OF THE NEW YORK ACADEMY OF SCIENCES

Special Issue: *Antimicrobial Therapeutics Reviews*

REVIEW

# New pathogens, new tricks: emerging, drug-resistant fungal pathogens and future prospects for antifungal therapeutics

Jennifer Geddes-McAlister<sup>1,2</sup> and Rebecca S. Shapiro <sup>1</sup><sup>1</sup>Department of Molecular and Cellular Biology, University of Guelph, Guelph, Ontario, Canada. <sup>2</sup>Department of Proteomics and Signal Transduction, Max Planck Institute of Biochemistry, Munich, Germany

Address for correspondence: Rebecca S. Shapiro, Department of Molecular and Cellular Biology, University of Guelph, Guelph, ON N1G 2W1, Canada. shapiror@uoguelph.ca

Fungal pathogens are a growing threat to public health. As human immunodeficiency becomes increasingly common, fungal infections are becoming more prevalent. The use of antifungal agents for prophylaxis and treatment of fungal infections has favored the emergence of previously rare or unidentified species of drug-resistant fungal pathogens, including several *Candida* and *Cryptococcus* species, as well as mold pathogens. As these new and increasingly drug-resistant fungal pathogens continue to emerge, novel strategies for rapid identification and treatment are necessary to combat these life-threatening infections.

**Keywords:** fungal pathogens; emerging pathogens; antifungal drug resistance

## Introduction

Fungal pathogens are emerging as critically important threats to global health. Recent estimates indicate that more than 300 million people are affected by serious fungal diseases worldwide, resulting in 1.6 million deaths annually<sup>1,2</sup>—similar to the number of deaths caused by tuberculosis.<sup>2,3</sup> Fungal pathogens are a highly diverse group of infectious agents, including yeast, or yeast-like species such as *Candida* and *Cryptococcus* species, and molds, such as *Aspergillus* species. These pathogens are responsible for a range of diseases, with variable prevalence and clinical outcomes. Nearly, one quarter of the world's population (~1.7 billion people) is estimated to have superficial skin, hair, and nail fungal infections caused primarily by dermatophyte fungi.<sup>2–4</sup> Mucosal fungal infections of the oral and genital tracts are also exceedingly common, with ~75% of women experiencing at least one vulvovaginal fungal infection caused by *Candida* species during their lifetime.<sup>3,5</sup> *Aspergillus* species are responsible for ~3 million cases of

chronic lung disease, and are a substantial cause of fungal-associated asthma, affecting ~10 million people.<sup>2,3</sup> While significantly less common, invasive fungal diseases caused by *Cryptococcus*, *Candida*, *Aspergillus*, and *Pneumocystis* species are associated with exceedingly high mortality rates, ranging from 30% to 90%, depending on the fungal pathogen and patient group.<sup>6,7</sup> These species are currently the most common cause of fungal diseases; however, many new and emerging fungal pathogens are being identified, and are poised to significantly threaten human health.

Emerging pathogens may include newly identified species, as well as species that are dramatically increasing in prevalence (e.g., as a result of a growing population of vulnerable hosts) or spreading to new geographic niches.<sup>8,9</sup> Here, we define emerging fungal pathogens as recently identified species in clinical settings, and species undergoing a population expansion, change of ecological niche, or a geographic spread. However, for completeness and comprehensiveness, within our definition of “emerging,” we also consider pathogenic fungi with

a history of human and plant pathogenesis that presently demonstrate increased rates of occurrence attributed to a growing population of vulnerable hosts and to improved detection methods, awareness, and surveillance, including *Fusarium* spp., *Scedosporium* spp., and *Mucormycetes*. Overall, the focus of our review is to comprehensively address emerging fungal pathogens associated with high rates of antifungal drug resistance and to present novel strategies to combat drug resistance.

Several factors contribute to the escalating emergence of new human fungal pathogens in the modern era. Fungi are commonly opportunistic pathogens—exploiting susceptibilities presented by a host with a compromised immune system, an altered microbiome, or breached physical barriers, to cause infection. Specifically, the growing prevalence of fungal infections is directly associated with ever-increasing rates of medical vulnerabilities, including (1) immunosuppression, which is becoming more common as a result of HIV/AIDS,<sup>10,11</sup> hematologic cancers and cancer chemotherapeutics,<sup>12,13</sup> the use of immunosuppressive agents for organ and bone marrow transplantation,<sup>14,15</sup> and an aging population;<sup>16,17</sup> (2) pervasive dysbiosis of the human microbiome associated with exposure to broad-spectrum antimicrobials;<sup>18,19</sup> and (3) breached barriers linked to the frequent use of modern medical devices such as ventilators, stents, and catheters.<sup>20,21</sup> Aside from medical factors leading to expanding pervasiveness of fungal pathogens, environmental fluctuations such as global climate change are contributing to geographic spread of many fungal crop pathogens<sup>22</sup> and may be associated with the emergence of newly pathogenic fungal species.<sup>23</sup> Of particular concern is the emergence of previously unidentified or rare fungal pathogens with high rates of antifungal drug resistance, such as the recent discovery of *Candida auris*,<sup>24</sup> and the emergence of rare but highly drug-resistant molds. These emerging pathogens are likely selected due to the mounting clinical administration of antifungal drugs for both prophylaxis and disease treatment.<sup>25</sup>

In this review, we discuss emerging fungal pathogens that pose a growing threat to human health. In particular, we highlight the emergence of drug-resistant fungal pathogens—including less common *Candida* and *Cryptococcus* species and emerging filamentous fungal pathogens—and the

difficulties associated with treating their associated infections. Initially, we present an overview of existing antifungals and clinically relevant mechanisms of drug resistance, and then discuss specific emerging drug-resistant pathogens. Further, we discuss future prospects for the development of innovative antifungal drugs, new treatment regimens, and novel approaches to sensitize drug-resistant fungal infections.

## Antifungal agents and drug resistance

Given the growing importance of fungal diseases, novel treatment options are in high demand. As eukaryotic fungal cells share a close evolutionary relationship with their human hosts, antifungal drug development presents a unique challenge for the pharmaceutical industry. Specifically, antifungal agents must target the eukaryotic fungal cell while ensuring limited damage to human cellular function during treatment. In general, antifungal agents are classified as fungistatic drugs that inhibit fungal cell growth, or fungicidal drugs that lead to fungal cell death. These classifications are important for defining mechanisms of drug action and understanding treatment options and outcomes. Presently, four classes of antifungal therapies are routinely used in monotherapy or in combination: polyenes, azoles, pyrimidine analogs, and echinocandins.<sup>26</sup> Although these classes of antifungal agents are often active against many infections, limitations associated with off-target host toxicity, limited fungicidal activity, drug–drug interactions, prolonged treatment courses, and the emergence of drug resistance can significantly impede their applicability and efficacy.

Antimicrobial resistance is a growing threat for bacteria as well as fungal infections and is one of the most critical global health concerns faced today. For fungal infections, rates of resistance vary by geography, species, and available treatment options; however, the expansion of drug-resistant fungal species is universal. In general, intrinsic and acquired mechanisms of resistance of fungal pathogens are associated with decreasing the effective drug concentration, altering the drug target, or diverting antifungal toxicity by metabolic modification.<sup>27,28</sup> Here, we present each class of antifungal agents along with their mechanisms of action and occurrence of resistance, as well as an overview of laboratory methods to quantify antifungal drug resistance.

## Polyenes

Polyenes were the first antifungal drugs available for clinical use and are considered to be dose-dependent fungicidal agents with broad-spectrum activity.<sup>29,30</sup> They are macrocyclic organic molecules derived from natural products that interact and disrupt ergosterol-containing fungal cell membranes through pore formation. Specifically, exposure to polyenes results in the leakage of cytoplasmic contents and oxidative damage, leading to fungal cell death.<sup>31</sup> First-generation amphotericin B, which is active against most systemic infections caused by yeasts and filamentous fungi, binds ergosterol to form an extramembranous fungicidal sterol sponge for the destabilization of membrane function following intravenous administration.<sup>32</sup> Additionally, nystatin and natamycin (administered as topical agents due to low gut absorption and high toxicity) are used for the treatment of mucocutaneous candidiasis and fungal keratitis.<sup>33</sup> Polyenes also interact with cholesterol-containing membranes, albeit with a lower affinity than to ergosterols, resulting in high host toxicity.<sup>34</sup> For polyenes, resistance is often intrinsic and only observed in fungal species with nonsusceptible or mildly affected ergosterol membranes. However, prolonged exposure to polyenes can lead to the emergence of clinical resistance.<sup>35</sup> To combat challenges associated with drug efficacy, toxicity, and resistance, second- and third-generation polyenes focus on the applicability of lipid formulations to reduce toxicity. Specifically, for amphotericin B, structural alterations enabling the preferential formation of monomers<sup>36</sup> may improve penetrability to specific body compartments and reduce membrane toxicity.<sup>37</sup> Future generations may involve oral formulations,<sup>38</sup> nanoparticles,<sup>39</sup> and polysaccharide conjugation.<sup>40</sup>

## Azoles

Azoles are the most common antifungal drugs in clinical use today. They exert broad-spectrum activity through inhibition of lanosterol 14 $\alpha$ -demethylase (encoded by *ERG11/CYP51*), a cytochrome P450-dependent enzyme that converts lanosterol to ergosterol, a primary target in the fungal membrane that is absent in the host cell membrane.<sup>41</sup> First-generation azoles, including the imidazoles (clotrimazole, miconazole, and ketoconazole), displayed high host toxicity, severe side effects, and numerous off-target interactions.<sup>42</sup>

These were replaced with second-generation triazoles, such as fluconazole, which is active against several pathogenic yeast species, including *Candida* spp., *Cryptococcus neoformans*, and *Histoplasma*, but lacks activity against molds,<sup>43,44</sup> and itraconazole, which shows broader spectrum activity against yeasts as well as *Aspergillus* species.<sup>45</sup> Though widely employed, azoles show variable fungicidal activity against fungal pathogens. While several azole drugs display fungicidal activity against *Aspergillus* species,<sup>46</sup> they generally act in a concentration-independent fungistatic manner against yeast pathogens, including *Candida* and *Cryptococcus* species.<sup>29,30</sup> To overcome certain limitations of efficacy, third-generation azoles with extended-spectrum activity, including voriconazole, posaconazole, efinaconazole, and isavuconazole, have been developed through extensive structural modifications for improved activity, safety, formulations, and pharmacokinetic properties.<sup>47</sup>

The fungistatic nature of the azoles against common yeast pathogens can impose a strong selection pressure on surviving populations to evolve resistance to these antifungal agents.<sup>48,49</sup> Additionally, increased azole resistance associated with fungistatic versus fungicidal activity of the triazoles can limit their effectiveness against emerging fungal pathogens.<sup>50</sup> For azoles, intrinsic or acquired resistance is possible through genetic mutations to *ERG11/CYP51*, which prevents blockage of catalytic activity by the drug, as well as target gene amplification to overwhelm the inhibitory capacity of the drug.<sup>51</sup> Target mutations and amplifications are often associated with cross-resistance between certain azoles (i.e., fluconazole and voriconazole).<sup>52–55</sup> Another mechanism of azole resistance involves amplification or induction of efflux pumps for removal of azoles from the fungal cells, resulting in decreased drug efficacy.<sup>56</sup> Efflux-related resistance has also been associated with cross-resistance between azoles and amphotericin B.<sup>57</sup> Other resistance mutations, such as C-5,6-desaturase mutations that block the synthesis of ergosterol and lead to the accumulation of an alternate sterol in the fungal membrane, are associated with cross-resistance to amphotericin B, along with azole resistance.<sup>58,59</sup>

## Pyrimidine analogs

Pyrimidine analogs, including 5-flucytosine, exert their activity through conversion by cytosine

deaminase into the toxic compound 5-fluorouracil, which interferes with RNA and DNA biosynthesis in the fungal cell.<sup>60</sup> Flucytosine is primarily used to treat pathogenic yeasts, and has little activity against most filamentous fungi. Host toxicity is less severe than other antifungals due to little or no cytosine deaminase activity in mammalian cells; however, there are still side effects associated with flucytosine treatment. Development of antifungal drug resistance to flucytosine is rapid and common. Specifically, a point mutation in *FUR1* encoding uracil phosphoribosyltransferase leads to complete resistance to flucytosine and 5-fluorouracil in fungi.<sup>61</sup> Moreover, a second point mutation in the cytosine deaminase-encoding genes *FCY1* and *FCY2* results in flucytosine resistance in many *Candida* strains.<sup>62</sup> Therefore, flucytosine is rarely used in monotherapy, but rather in combinatory therapeutic approaches, such as in combination with amphotericin B for the treatment of cryptococcosis.<sup>63</sup> Indeed, amphotericin B and flucytosine have been shown to act synergistically; the combination of these drugs is significantly more effective than either amphotericin B or flucytosine monotherapy for reducing fungal burden in murine models of cryptococcosis.<sup>64,65</sup>

### Echinocandins

Echinocandins, a group of lipopeptides derived from natural products and the newest class of antifungal agents, target the fungal cell wall. This is achieved by noncompetitively inhibiting the plasma membrane-bound  $\beta$ -(1,3)-D-glucan synthase enzyme complex associated with synthesis of the structural polymer  $\beta$ -(1,3)-D-glucan, causing osmotic instability and fungal cell death.<sup>66</sup> Caspofungin, micafungin, and anidulafungin represent the three first-line echinocandins with broad-spectrum fungicidal activity against yeasts, specifically used to treat invasive systemic fungal infections including nosocomial candidemia and invasive candidiasis.<sup>43</sup> However, echinocandins are not active against many important fungal pathogens, including *Cryptococcus* and *Fusarium* species.<sup>67</sup> Interestingly, echinocandins display opposite fungicidal activity compared with azole drugs: they are generally fungicidal against pathogenic yeasts such as *Candida* species, but fungistatic against *Aspergillus* species.<sup>67</sup>

The echinocandins display good safety profiles, limited interactions with other drugs, and minimal host toxicity due to the absence of the enzyme in mammalian cells. Additionally, these drugs generally do not display cross-resistance to other existing antifungal agents, and are therefore broadly effective against azole-resistant pathogens.<sup>68</sup> For echinocandin resistance, mutations in the gene encoding the echinocandin target ( $\beta$ -(1,3)-D-glucan synthase (*FKS1*)) are by far the most common mechanisms of drug resistance.<sup>66</sup> These mutations, which typically occur in certain hot-spot regions of the gene *FKS1*, usually impart cross-resistance to all of the echinocandin drugs.<sup>68</sup>

### Measuring antifungal drug resistance

Antifungal drug resistance profiles for fungal pathogens are established using specific assays and protocols. Resistance is measured as the reduction of growth of fungal cells in the presence of the drug *in vitro*, compared with cells grown in the absence of drug. Such measurements can be established experimentally with antifungal drug gradients in liquid or on solid agar media.<sup>28,69,70</sup> In order to establish unified, reproducible methods of measuring antifungal susceptibility between laboratories, internationally recognized standard testing methods have been established from two agencies: the Clinical and Laboratory Standards Institute and the European Committee on Antimicrobial Susceptibility Testing.<sup>71</sup> In both cases, susceptibility measurements rely on broth microdilution testing along a drug concentration gradient in order to establish a minimum inhibitory concentration (MIC)—the lowest concentration of an antimicrobial that will inhibit fungal growth. A fungal strain is deemed drug resistant if its MIC value falls above clinical breakpoints that have been previously established for specific antifungals and specific fungal pathogens. In some cases, all strains of a given fungal species are not susceptible to a particular antifungal (without having been previously exposed to that drug) and these species are considered to be intrinsically resistant to that drug.<sup>28</sup>

There are certain issues that arise while determining antifungal susceptibility of fungal pathogens—and particularly, of emerging pathogens. Susceptibility testing relies on the use of established breakpoints, and if these have not been determined for a newly emerging pathogen, it can

be difficult to infer how *in vitro* susceptibility will reflect clinical *in vivo* drug resistance. Even for well-studied pathogens, drug resistance defined by *in vitro* analysis is often not sufficient to explain clinical outcomes.<sup>72</sup> Additionally, antifungal drug tolerance—defined as a pathogen's ability to withstand killing at drug concentrations above the MIC<sup>73</sup>—is a less well understood biological phenomenon, likely attributed to an epigenetic, rather than genetic, mechanism that facilitates fungal survival in the presence of drug.<sup>73</sup>

### Emerging threats of drug-resistant yeast fungal pathogens

Yeast and yeast-like species are a diverse group of fungal organisms that grow as single cells and divide through budding or binary fission. Some yeast species, such as *Candida albicans*, are considered dimorphic (or polymorphic) as they grow in both yeast morphology as well as elongated filamentous cells.<sup>74</sup> Yeast pathogens are among the most pervasive and deadly to humans, including *Candida* and *Cryptococcus* species. The emergence of rare or previously unidentified species of *Candida* and *Cryptococcus* is becoming more frequent and poses a challenge to human health.

#### Emerging drug-resistant *Candida* pathogens

*Candida* species can cause a range of human diseases, from superficial mucosal infection of the oral and urogenital tracts to life-threatening invasive infection in immunocompromised individuals.<sup>74,75</sup> *Candida* species are the fourth leading cause of hospital-acquired bloodstream infections and the most common fungal species isolated from medical device-associated biofilm infections.<sup>7,76</sup> The most commonly encountered of these yeast species is *C. albicans*, which accounts for more than half of all cases of invasive candidiasis.<sup>75</sup> However, despite the prevalence of *C. albicans*, newly emerging non-*albicans* *Candida* species are increasingly being documented in clinical settings.<sup>77</sup> Worryingly, many of these emerging *Candida* pathogens, including *Candida glabrata*, *Candida krusei*, *Candida lusitanae*, and the very newly emerging pathogen *C. auris*, are associated with greater resistance to antifungal therapeutics<sup>78,79</sup> and are a growing threat to public health.

Among non-*albicans* *Candida* pathogens, *C. glabrata* is the most common and of growing clin-

ical concern among elderly patients, HIV positive individuals, and those receiving solid organ transplants (SOTs) and bone marrow transplants.<sup>75,80</sup> *C. glabrata* is the second most common cause of invasive candidiasis, representing 15–30% of clinically isolated pathogens.<sup>75,80–82</sup> It is intrinsically less susceptible to fluconazole and amphotericin B;<sup>77,83,84</sup> ~10–15% of clinical isolates are resistant to fluconazole and voriconazole, and strains are likely to exhibit cross-resistance to multiple azole drugs.<sup>77</sup> Most clinical azole resistance in *C. glabrata* has been associated with upregulation of ATP-binding cassette (ABC) transporters, such as Cdr1 and Cdr2.<sup>85,86</sup> As a result of higher rates of innate and acquired azole resistance, *C. glabrata* is often found to cause infection in patients exposed to fluconazole as prophylaxis or for treatment of other fungal infections.<sup>77,87–89</sup> Notably, while most *Candida* species have relatively low frequency of resistance to echinocandins,<sup>90–92</sup> *C. glabrata* isolates have unusually high rates of resistance, ranging from less than 5% in 2001, to 12% in 2010,<sup>93</sup> often associated with mutations in the drug target *FKS* genes.<sup>92</sup> Of particular concern is the high frequency of multidrug-resistant *C. glabrata* isolates with reduced susceptibility to both azoles and echinocandins. One study indicates that 36% of echinocandin-resistant *C. glabrata* strains are also resistant to fluconazole,<sup>94</sup> limiting treatment options for this increasingly prevalent fungal pathogen. A recent study suggested that a prevalent “mutator” genotype—routinely identified among *C. glabrata* clinical isolates—may promote multidrug resistance.<sup>95</sup> Such mutator strains have mutations rendering the fungus defective in mismatch repair, leading to high rates of genetic mutations and a high propensity to develop multidrug resistance.<sup>95</sup>

Less frequently encountered emerging *Candida* pathogens, including *C. krusei*, *C. lusitanae*, and *C. haemulonii*, also have intrinsic antifungal drug resistance profiles that hinder successful antifungal treatment. Similar to *C. glabrata*, *C. krusei* is intrinsically resistant to fluconazole, mainly due to a diminished sensitivity of the target enzyme CYP51 to fluconazole inhibition.<sup>96</sup> As a result, *C. krusei* is frequently associated with infection in patients receiving prophylactic or therapeutic fluconazole treatment.<sup>97,98</sup> Additionally, *C. lusitanae* is generally susceptible to azoles, but is associated with intrinsic resistance to amphotericin B.<sup>99</sup> Moreover,



isolates of both *C. krusei* and *C. lusitaniae* have also been identified as multidrug resistant to azoles and echinocandins.<sup>100,101</sup> Finally, *C. haemulonii* isolates are frequently found to be resistant to both amphotericin B and azoles,<sup>102–104</sup> and azole-resistant *Candida parapsilosis* clinical isolates are increasingly being identified.<sup>105,106</sup>

### *Candida auris*

The appearance of newly emerging drug-resistant *Candida* species is ongoing. *C. auris* is the most recent and worrisome example of a previously unidentified pathogen that is highly refractory to existing antifungal treatment. Initially identified in Japan in 2009,<sup>107</sup> *C. auris* infections simultaneously and seemingly independently emerged in hospital settings in distinct geographical locations,<sup>24,108</sup> and have now been identified across five continents.<sup>24,109</sup> In some geographical regions, *C. auris* is becoming quite prevalent. For example, in India, a multicenter study of intensive care units found *C. auris* in 19 out of 27 units, accounting for ~5% of all candidemia cases.<sup>110</sup> Inaccuracies associated with the diagnosis of *C. auris* have resulted in misidentification of this pathogen and difficulties in quantifying the prevalence of these infections,<sup>111,112</sup> highlighting the need for accurate diagnostic tests. Although *C. auris* does not seem to form hyphae or pseudohyphae, some isolates display a cellular aggregation phenotype and are able to form biofilms.<sup>113</sup> *C. auris* also harbors virulence factors, including phospholipase activity and secreted proteinase activity.<sup>114</sup>

*C. auris* strains are frequently resistant to antifungal drugs, although currently there are no antifungal clinical breakpoints reported for this pathogen. A global study of *C. auris* isolates was conducted from 54 patients across three continents from 2012 to 2015.<sup>108</sup> This analysis found that an overwhelming majority of *C. auris* isolates are resistant to fluconazole (~93% of isolates<sup>108</sup>), and approximately half of the tested isolates are also resistant to voriconazole.<sup>108</sup> Additionally, 35% of *C. auris* isolates are resistant to amphotericin B, and 7% to echinocandins.<sup>108</sup> Multidrug resistance is common in *C. auris*, where 41% of strains are resistant to at least two classes of antifungals and 4% are resistant to all antifungal agents.<sup>108</sup> While *C. auris* research is still in its infancy, studies suggest that elevated drug resistance is due to a large number of genomically encoded efflux pumps,<sup>115,116</sup> enhanced efflux

pump activity,<sup>117</sup> and mutations in the ergosterol biosynthesis pathway.<sup>108</sup>

### Emerging drug-resistant *Cryptococcus* pathogens

*Cryptococcus* species are responsible for causing cryptococcosis, an important, primarily opportunistic fungal infection common among patients with HIV/AIDS. Infection begins with inhalation of desiccated yeast cells or spores from the environment, causing primary infection in the alveolar macrophages of the lungs, followed by dissemination to the central nervous system, leading to fungal meningitis, as well as bloodstream infections.<sup>118</sup> Cryptococcal meningitis is one of the most important HIV/AIDS-related infections, responsible for 15% of AIDS-related deaths.<sup>119</sup> *C. neoformans* and *Cryptococcus gattii* are responsible for an overwhelming majority of cryptococcal disease.<sup>118</sup> Specifically, the emergence of *C. gattii* since 1999 in the Pacific Northwest of the United States and on the west coast of Vancouver Island, Canada, has identified the species as a significant human pathogen causing infection in immunocompetent individuals.<sup>120</sup> While *C. gattii* is more susceptible to amphotericin B and flucytosine compared with *C. neoformans*, it is less susceptible to fluconazole and other azole drugs;<sup>121,122</sup> however, mechanisms of resistance have not been defined.<sup>123</sup> *Cryptococcus* species are not susceptible to the echinocandin class of antifungals;<sup>67,74</sup> therefore, additional antifungal drug resistance severely limits treatment options. Additionally, mismatch-repair defective mutator strains have been identified for *Cryptococcus* species and are associated with high rates of mutations and antifungal drug resistance.<sup>124–126</sup>

Aside from *C. neoformans*, the majority of *Cryptococcus* species were historically believed to be avirulent in humans. However, in recent years, less common non-*neoformans* *Cryptococcus* species have been increasingly identified as opportunistic human pathogens, causing infection primarily in patients with impaired cellular immunity due to hematological cancer, neutropenia, use of immunosuppressive drugs, or HIV/AIDS.<sup>127</sup> The rise in incidence of these pathogens may be due to improvements and availability of diagnostic tools. The most commonly identified emerging cryptococcal species associated with immunocompromised hosts are *Cryptococcus laurentii* and

*Cryptococcus albidus*, which account for 80% of non-*neoformans* and non-*gattii* *Cryptococcus* infections.<sup>127,128</sup> Susceptibility testing has revealed that these species are not susceptible to flucytosine, and 50% of *C. laurentii* and 80% of *C. albidus* have an elevated MIC to fluconazole ( $\geq 16$  mg/L),<sup>78</sup> although the mechanisms of resistance have not been conclusively identified. This mirrors the trend observed among emerging non-*albicans* *Candida* species with high rates of fluconazole resistance and suggests that widespread use of this antifungal may be responsible for the new emergence of diverse, fluconazole-resistant fungal pathogens. The majority of *Cryptococcus* isolates remain sensitive to amphotericin B, which is used to successfully treat these infections—but a trend toward emerging *Cryptococcus* species with expanded drug-resistance profiles results in a scarcity of therapeutic options.

### Emerging threats of drug-resistant filamentous fungal pathogens

Filamentous fungi or molds are ubiquitous within the environment and grow by apical extension of their filaments, or hyphae, with or without septation, leading to the formation of mycelia. The occurrence of invasive filamentous fungal infections has risen over the past few decades and these infections are often associated with high rates of morbidity and mortality. Moreover, in recent years, the increasing emergence of pathogenic fungi, including *Aspergillus* spp., Mucormycetes, *Fusarium* spp., and *Scedosporium* spp., that display intrinsic or acquired mechanisms of drug resistance demands our attention.

#### Emerging drug-resistant *Aspergillus* pathogens

*Aspergillus* species are saprotrophic fungi commonly found in soil and decaying organic material within the environment. *Aspergillus fumigatus* is the most common *Aspergillus* spp., causing invasive aspergillosis (IA) in immunocompromised individuals, particularly patients undergoing chemotherapy treatment, hematopoietic stem cell transplantations (HSCTs), SOTs, immunosuppressive drug treatment, or patients with HIV/AIDS.<sup>129</sup> Other species capable of causing disease include *Aspergillus flavus*, *Aspergillus niger*, *Aspergillus terreus*, *Aspergillus clavatus*, and *Aspergillus nidulans*.<sup>130</sup> Annually, *Aspergillus* causes

over 200,000 life-threatening infections, with mortality rates ranging from 30% to 95%.<sup>131</sup> Aside from IA infections, the occurrence of allergic bronchopulmonary aspergillosis affects 4.8 million people worldwide, with an estimated 400,000 with chronic pulmonary aspergillosis.<sup>132,133</sup> While *A. fumigatus* itself is not a new pathogen, the emergence of azole resistance in this species is relatively recent and concerning. Since azole-resistant strains were first identified in 1997, the rate of resistance has risen rapidly to account for ~5–6% of *A. fumigatus* isolates.<sup>134,135</sup> There is evidence to suggest that mounting rates of azole resistance are linked to the exposure of *A. fumigatus* to azole-based agricultural fungicides in the environment and the resultant selection for resistant strains.<sup>136–139</sup>

Given the clinical relevance of *Aspergillus* spp., the emergence of opportunistic *A. terreus*, the third leading cause of global IA, is of particular concern. For example, *A. terreus* presents a growing problem among patients with hematological malignancies (HMs),<sup>140,141</sup> as it causes severe infections and has a propensity to disseminate, resulting in systemic patient infections. Due to high rates of *in vitro* and *in vivo* resistance to antifungal agents,<sup>142,143</sup> including intrinsic resistance to amphotericin B,<sup>144</sup> *A. terreus* infections are associated with very poor patient outcomes.<sup>143,145</sup> Recently, evaluation of the mechanism of resistance to amphotericin B determined that ergosterol content plays a minor role in intrinsic resistance and that the occurrence of higher catalase production in resistant isolates may block oxidative damage caused during antifungal treatment.<sup>146</sup> In addition, reduced azole susceptibility, attributed to mutations in the target Cyp51 protein that are associated with acquired resistance,<sup>147</sup> has been reported in *A. terreus* infections in Spain, the United States, and Denmark.<sup>131,148,149</sup> Alternatively, posaconazole appears promising for *A. terreus* treatment based on its efficacy in animal models and clinical studies.<sup>150</sup>

#### Emerging drug-resistant Mucormycetes pathogens

Mucormycetes (formerly Zygomycetes) infections are fungal infections caused by pathogenic orders Mucorales and Entomophthorales. Here, we focus on Mucorales, the most common emerging cause of non-*Aspergillus* mold infections in humans, with reported increases in incidence across various

geographical regions.<sup>151–153</sup> Mucorales represent fast-growing, common saprobes that cause mucormycosis by invasion of blood vessels and tissue necrosis. Mucormycosis occurs primarily in immunocompromised patients and results in mortality rates of 40–70% despite antifungal treatments.<sup>154</sup> Among Mucorales, *Rhizopus* spp. are the most commonly identified in human infection, along with *Mucor*, *Rhizomucor*, *Lichtheimia*, and *Cunninghamella* spp., which are particularly aggressive in HSCT recipients, with a mortality rate of 80%.<sup>155</sup>

Treatment of infection is often challenging due to delayed diagnosis, acute disease progression, and intrinsic resistance to many currently available antifungal agents.<sup>156</sup> Treatment strategies often involve surgery at the pathogen-infected body sites, as well as antifungal therapies. Susceptibility to drug treatments varies significantly among Mucorales, and microenvironmental factors at the site of fungal growth, such as tissue hypoxia, may affect antifungal drug efficacy.<sup>157,158</sup> Mucorales are intrinsically resistant to fluconazole and highly resistant to voriconazole,<sup>159,160</sup> and, as a result, are commonly associated with infections of HSCT recipients following prophylactic voriconazole treatment.<sup>161</sup> Posaconazole and isavuconazole generally demonstrate significant anti-Mucorales activity and are routinely used for treatment of mucormycosis;<sup>162</sup> however, *Cunninghamella bertholletiae* is highly resistant to posaconazole treatment. Additionally, *in vitro* testing suggests that flucytosine and echinocandins have little activity against Mucorales.<sup>160,163,164</sup> Morbidity and mortality rates for mucormycosis remain high, most likely attributed to poor clinical drug efficacy of monotherapies, demonstrating the need for novel treatment strategies.

### Emerging drug-resistant *Fusarium* pathogens

*Fusarium* species are a large genus of environmentally ubiquitous hyphomycete fungi. *Fusarium* are well-known plant pathogens but are also classified as emerging, opportunistic human pathogens. Specifically, *Fusarium* spp. represent the second most common category of non-*Aspergillus* molds responsible for human infections.<sup>165</sup> Approximately 50% of reported infections are due to the most pathogenic species, *Fusarium solani*, followed by 20% of infections attributed to *Fusarium oxysporum*.<sup>166</sup> Primarily acquired from the environment, *Fusarium* spp.

cause keratitis and onychomycosis among immunocompetent hosts, and respiratory and disseminated infections in immunocompromised individuals due to the weakened defenses of pulmonary alveolar macrophages and neutrophils.<sup>167</sup> The occurrence of fusariosis, endemic in tropical and subtropical countries, is associated with high mortality and morbidity rates of >80% in patients with prolonged and severe neutropenia, HMs, and HSCT recipients.<sup>168,169</sup>

Intrinsic multidrug resistance to a broad range of antifungals, along with species- and isolate-dependent resistance, is characteristic of *Fusarium* species. For example, second-generation broad-spectrum triazoles (fluconazole, itraconazole, voriconazole, and posaconazole), flucytosine, and echinocandins (caspofungin, anidulafungin, and micafungin) have limited activity in almost all *Fusarium* spp., and cross-resistance has been observed among echinocandins.<sup>170,171</sup> Although most *Fusarium* species have reduced susceptibility to these diverse antifungals, *F. solani* is considered the most highly resistant to antifungal drugs.<sup>172,173</sup> While the basis for *Fusarium* species' near-universal intrinsic drug resistance profile is not fully understood, intrinsic echinocandin resistance may be attributed to mutations occurring in the gene for the Fks1 target protein.<sup>174</sup>

Beyond intrinsic resistance, secondary or acquired resistance to azoles develops among previously susceptible strains of *Fusarium* after exposure to an antifungal agent. The mechanism of resistance is usually dependent on altered expression of CYP51, consisting of three CYP51 paralogues (CYP51A, -B, and -C).<sup>175,176</sup> Owing to high levels of resistance, monotherapy for *Fusarium* infections is associated with poor patient prognosis. However, synergistic antifungal interactions between drug combinations, including natamycin and voriconazole, and amphotericin B with caspofungin, rifampin, 5-flucytosine, or voriconazole, hold promise for more effective treatment of these infections.<sup>177,178</sup> Taken together, emerging *Fusarium* species pose a major challenge for medicine and agriculture alike, with very limited treatment options available for patients and crops.

### Emerging drug-resistant *Scedosporium* pathogens

*Scedosporium*, including *Scedosporium apiospermum* and the related fungal species *Lomentospora*



*prolificans* (previously *Scedosporium prolificans*), are hyphomycete fungi of significant emerging medical importance. Although rare human pathogens, the incidence of *Scedosporium* infections has been rising significantly.<sup>179</sup> Globally, infection is acquired through inhalation of contaminated plant and soil residues and occurs in both immunocompetent and immunocompromised individuals.<sup>180</sup> Risk factors associated with infection include chronic obstructive or suppurative lung disease (including cystic fibrosis (CF)), HM, SOT or HSCTs, corticosteroid use, neutropenia, and diabetes mellitus.<sup>181,182</sup> *L. prolificans* is the second most abundant filamentous fungi colonizing the respiratory tract of CF patients after *Aspergillus* species.<sup>183,184</sup> Infection by *Scedosporium* species can cause a broad range of diseases, including localized skin and soft tissue infections, pneumonia, meningitis, and disseminated, multiorgan infection in immunocompromised hosts. For immunocompromised hosts, particularly those with neutropenia, the risk of *Scedosporium* infection dissemination is high, and associated mortality rates are >80%.<sup>185</sup>

*Scedosporium* species are among fungi with the highest degrees of resistance to antifungal compounds, and the majority of *Scedosporium* isolates have multidrug-resistance profiles.<sup>186,187</sup> *L. prolificans* in particular is one of the most highly drug-resistant fungal organisms identified, with high rates of resistance to polyenes,<sup>188</sup> azoles (including posaconazole and itraconazole),<sup>188,189</sup> and echinocandin antifungals.<sup>190</sup> *S. apiospermum* is also resistant to polyenes, echinocandins, and some azoles, but is susceptible to voriconazole and posaconazole.<sup>191</sup> While the mechanisms of *Scedosporium* resistance remain to be fully elucidated, echinocandin resistance has been largely attributed to hot spot mutations in the gene for the target protein Fks1, which render the fungi intrinsically resistant to echinocandin treatment.<sup>190</sup> Currently, the first-line treatment for *Scedosporium*-associated infection is voriconazole, as resistance against this compound is rare.

### Future prospects for emerging fungal disease: diagnosis and treatment

Given the prevalence of intrinsic drug resistance among emerging fungal pathogens, it is critical that these less common species are correctly identified,

as misidentification can lead to inappropriate therapeutic treatment and poor patient prognosis. This is of particular concern in resource-limited hospital environments, where species-level pathogen diagnostics may not be readily available. Furthermore, since fluconazole is by far the most readily available and inexpensive antifungal, the ability to acquire more expensive and more effective alternatives is limited in developing countries—presenting an additional hurdle to treating these emerging pathogens, many of which are intrinsically resistant to this ubiquitous antifungal. Difficulties associated with diagnosis and treatment of emerging pathogens may explain why these infections are associated with significantly longer hospital stays and higher inpatient costs compared with other infections.<sup>28,192–194</sup>

### Future outlook for diagnosis of emerging fungal disease

Accurate identification of newly emerging fungal species is critical. In some cases, fungal pathogens classified as emerging are not in fact new pathogens but have been historically misclassified or mistaken for a more common species. For instance, a retrospective review of *Candida* isolates found a misidentified strain of *C. auris* from 1996<sup>195</sup>—13 years before *C. auris* was first identified as an emerging fungal pathogen. Additionally, intrinsic drug-resistance associated with many emerging fungi means that accurate, species-level diagnosis is crucial to ensure that proper treatment is administered. Most current fungal diagnostic techniques, such as culturing strains, microscopy-based staining, or antigen-based tests are time-consuming, may not provide species-level detection specificity, and/or require specialized and costly equipment and reagents. Therefore, novel tools for the rapid and accurate detection of fungal pathogens are imperative, and, given the prevalence of fungal infections in resource-limited environments, low-cost and low-tech diagnostic tools would significantly improve detection of these organisms in the developing world.<sup>196</sup>

Identification of fungal species based on genetic signatures provides accurate detection of pathogens and distinguishes between closely related species based on unique genomic identifiers. PCR-based diagnostics are often used for fungal

identification,<sup>197</sup> but require the use of specialized instrumentation. However, new, rapid, low-cost PCR-based technologies<sup>198–200</sup> may improve global access to these diagnostic techniques. New technologies are focusing on whole-genome sequencing for pathogen detection.<sup>201</sup> For example, portable genomic sequencers, such as the minION from Oxford Nanopore Technologies,<sup>202</sup> have been successfully used for rapid and relatively low-cost next-generation sequencing for the identification of viral<sup>203–205</sup> and bacterial<sup>206</sup> diseases in resource-limited settings, and could similarly be employed for the diagnosis of fungal disease. Whole-genome sequencing has the additional benefit of detecting not only a particular species, but also any potential drug-resistance mutations. Additionally, cutting-edge technologies for nucleic acid detection, including paper-based synthetic gene networks<sup>207,208</sup> and CRISPR-based diagnostic platforms,<sup>209</sup> are rapid, inexpensive, and sensitive methods for infectious disease detection that could be applied to fungal pathogens.

In addition to novel genomic technologies, other newer diagnostics focus on biomarkers of fungal infection. Matrix-assisted laser desorption/ionization time-of-flight mass spectrometry has been successfully exploited for species-level fungal pathogen identification based on unique proteomic signatures.<sup>210,211</sup> This approach has been particularly useful in the identification of rare and commonly misidentified fungal species, including unusual *Aspergillus* pathogens,<sup>212,213</sup> and the identification of antifungal drug-resistant *Candida* strains.<sup>214,215</sup> Moreover, quantitative bottom-up proteomic analysis combined with selected reaction monitoring has discovered secreted cryptococcal proteins in infected tissue samples, suggesting the utility of such techniques for detection of diagnostic biomarkers of fungal infection.<sup>216</sup> Finally, compelling recent work has exploited engineered *Saccharomyces cerevisiae* as a biosensor capable of detecting fungal pathogen-derived peptides in order to develop a low-cost and nontechnical fungal diagnostic.<sup>217</sup>

### **New antifungal drugs for emerging infections**

New antifungal drugs and treatment strategies are urgently needed to combat ever-emerging and increasingly drug-resistant fungal

infections,<sup>26,218,219</sup> and several novel promising treatment strategies are currently being researched and developed (Table 1). For example, improved versions of existing antifungals, such as potent new azoles for candidiasis (VT-1161),<sup>220</sup> cryptococcosis (VT-1129),<sup>221</sup> and coccidioidomycosis (VT-1598, whose development has recently been fast-tracked by the Food and Drug Administration),<sup>222</sup> and CD101, a novel echinocandin with an improved safety profile,<sup>223–225</sup> are under investigation or development. Another important new antifungal, ASP2397, displaying potent fungicidal activity against several mold species, disrupts intracellular fungal biochemical machinery through uptake by a siderophore iron transporter.<sup>226</sup> In addition, several existing pharmaceutical agents are being repurposed as antifungals, such as the use of the selective serotonin reuptake inhibitor sertraline as an adjuvant to potentiate azole activity in cryptococcal infections,<sup>227</sup> which is currently in phase III clinical trials,<sup>26</sup> and the anticancer drug bortezomib, which interferes with virulence factor production and disease caused by *C. neoformans*.<sup>228</sup> Finally, new antifungal drugs with novel targets are also under development, such as APX001 (currently completed phase II clinical trials), a prodrug targeting glycosylphosphatidylinositol synthesis that has broad-spectrum activity against diverse yeast and mold pathogens, including highly drug-resistant emerging pathogens such as *Scedosporium* species.<sup>229–231</sup>

In addition to these antifungals currently in development, ongoing research is continually identifying new compounds with antifungal activity or new fungal factors that may serve as targets for antifungal therapy. Recent screens of bio-active molecules identified drugs with antifungal activity<sup>232–235</sup> that may serve as promising leads for antifungal drug development. Other work has focused on identifying and characterizing prospective antifungal drug targets, including key fungal stress response factors, such as the molecular chaperone Hsp90 (see Refs. 236–238), and the protein phosphatase calcineurin.<sup>236,239–241</sup> Antifungal drug target identification has also benefited from computational methods used to compare human and fungal proteomes, in order to identify fungal-specific domains with potential to bind small molecules.<sup>242</sup> These avenues lay the groundwork for future antifungal drug development pipelines.

**Table 1.** Emerging fungal pathogens: mechanisms of antifungal drug resistance profiles and future therapeutic prospects

	Pathogen	Resistance profile	Future therapeutic prospects	Reference				
Candida species	Candida glabrata	Decreased intrinsic susceptibility to fluconazole, amphotericin B	New azoles (i.e., VT-1161); new echinocandins (i.e., CD101)	Garvey <i>et al.</i> , <sup>220</sup> Ong <i>et al.</i> , <sup>223</sup> Pfaller <i>et al.</i> <sup>225</sup>				
		High rates of acquired resistance to fluconazole, voriconazole, and echinocandins	New antifungal APX001 targeting GPI synthesis	Watanabe <i>et al.</i> , <sup>229</sup> Hata <i>et al.</i> <sup>230</sup>				
	Candida krusei	Intrinsic resistance to fluconazole	Targeting fungal stress response	Cowen <i>et al.</i> , <sup>236</sup> Juvvadi <i>et al.</i> <sup>239</sup>				
	Candida lusitaniae	Intrinsic resistance to amphotericin B	Computational methods for antifungal drug target discovery	Barrera <i>et al.</i> <sup>242</sup>				
	Candida haemulonii	High rates of resistances to azoles and amphotericin B	New antifungal adjuvants	Pfaller <i>et al.</i> , <sup>254</sup> Fiori <i>et al.</i> , <sup>255</sup> Liu <i>et al.</i> , <sup>257</sup> Shekhar-Guturja <i>et al.</i> , <sup>258</sup> Spitzer <i>et al.</i> , <sup>259</sup> Robbins <i>et al.</i> <sup>260</sup>				
			Cytokine therapy as antifungal adjuvants	Delsing <i>et al.</i> <sup>270</sup>				
	Candida auris	High rates of resistance to fluconazole, voriconazole, and amphotericin B  Very high prevalence of multidrug-resistant strains (including strains resistant to all existing antifungals)	Monoclonal antibody therapy	Pachl <i>et al.</i> , <sup>238</sup> Bugli <i>et al.</i> <sup>277</sup>				
			Novel techniques in immuno-modulation	Zhao <i>et al.</i> <sup>281</sup>				
			Fungal vaccine	Xin <i>et al.</i> , <sup>290</sup> Xin <i>et al.</i> <sup>291</sup>				
			Targeting fungal virulence	Zhang <i>et al.</i> , <sup>294</sup> Romo <i>et al.</i> , <sup>295</sup> Vila <i>et al.</i> , <sup>296</sup> Fazly <i>et al.</i> , <sup>297</sup> Shareck and Belhumeur, <sup>298</sup> Murzyn <i>et al.</i> , <sup>300</sup> Mayer and Kronstad <sup>301</sup>				
Cryptococcus species	Cryptococcus gattii	Resistance to azoles	New azoles (i.e., VT-1129)	Lockhart <i>et al.</i> <sup>221</sup>				
					Cryptococcus laurentii	Intrinsic resistance to flucytosine High rates of resistance to fluconazole	SSRI sertraline as azole adjuvant New antifungal APX001 targeting GPI synthesis	Rhein <i>et al.</i> <sup>227</sup> Schell <i>et al.</i> <sup>231</sup>
					Cytokine therapy as antifungal adjuvants Monoclonal antibody therapy Fungal vaccine	Pappas <i>et al.</i> , <sup>268</sup> Jarvis <i>et al.</i> <sup>272</sup> Larsen <i>et al.</i> , <sup>274</sup> Rachini <i>et al.</i> <sup>276</sup> Chow and Casadevall, <sup>286</sup> Upadhyia <i>et al.</i> , <sup>287</sup> Mor <i>et al.</i> , <sup>288</sup> Wormley <i>et al.</i> <sup>289</sup>		
	Aspergillus terreus	Intrinsic resistance to amphotericin B Acquired azole resistance	New echinocandins (i.e., CD101) New antifungal APX001 targeting GPI synthesis	Pfaller <i>et al.</i> <sup>225</sup> Hata <i>et al.</i> <sup>230</sup>				
					Mucorales species	Intrinsic resistance to fluconazole, highly resistant to voriconazole Resistant to flucytosine and echinocandins	New antifungal ASP2397 Targeting fungal stress response	Nakamura <i>et al.</i> <sup>226</sup> Cöwen <i>et al.</i> , <sup>236</sup> Juvvadi <i>et al.</i> , <sup>239</sup> Nambu <i>et al.</i> <sup>241</sup>
	Fusarium species	High rates of resistance to azoles, echinocandins, and flucytosine	Computational methods for antifungal drug target discovery	Barrera <i>et al.</i> <sup>242</sup>				
	Scedosporium species	High rates of resistance to azoles, echinocandins, and polyenes	New antifungal adjuvants	Pfaller <i>et al.</i> , <sup>225</sup> Shekhar-Guturja <i>et al.</i> , <sup>258</sup> Robbins <i>et al.</i> <sup>260</sup>				
			Cytokine therapy as antifungal adjuvants	Shao <i>et al.</i> , <sup>269</sup> Delsing <i>et al.</i> , <sup>270</sup> Armstrong-James <i>et al.</i> , <sup>271</sup> Goldman <i>et al.</i> <sup>273</sup>				
			Monoclonal antibody therapy	Rollin-Pinheiro <i>et al.</i> <sup>278</sup>				
Novel techniques in immuno-modulation			Kumaresan <i>et al.</i> <sup>280</sup>					

*Prospects for combination antifungal drug therapy*

Combination drug therapy is another promising therapeutic strategy for the treatment of fungal infections, as well as many other infectious and noncommunicable diseases.<sup>243–245</sup> Combination

therapies can improve the effectiveness of drug treatment and may limit or prevent the emergence of drug resistance,<sup>246–250</sup> making this strategy of particular interest for emerging and highly drug-resistant fungal pathogens. Combination therapy has been proposed as a treatment option

for highly drug-resistant fungal pathogens, including *Fusarium* and *Scedosporium* species, where treatment with combinations of existing antifungals has shown promising results as effective therapeutics,<sup>177,178,251</sup> as well as combination therapy between antifungals and antiparasitics.<sup>252</sup>

In addition to combinations of existing antifungals, many new promising antifungal adjuvants have been described and are in various stages of clinical development, including: histone deacetylase inhibitors, which synergize with azoles and echinocandins;<sup>253,254</sup> iron and calcium homeostasis inhibitors,<sup>255–257</sup> which enhance azole activity; and beauvericin, which enhances the efficacy of azoles and blocks the emergence of antifungal resistance.<sup>258</sup> Additionally, several compelling novel screening platforms have identified new molecules that act to enhance the activity of existing antifungals<sup>259–261</sup> and may serve as a template for future screening efforts to identify antifungal adjuvants.

In addition to drug screening, an alternative strategy for target identification in combination therapy is to use chemical-genomic approaches or genetic interaction analysis. Chemical-genomic screening has been successfully used to identify fungal genetic mutants with altered susceptibility to chemical perturbations, and further exploitation of this chemogenomic information represents a powerful tool to predict antifungal drug synergies in *Cryptococcus*.<sup>262</sup> Genetic interaction analysis can also be utilized to identify drug target synergies by identifying synthetic lethal interactions,<sup>263</sup> where simultaneous deletion of perturbation of two genes results in lethality to the cell.<sup>264</sup> Innovative tools for genetic interaction and synthetic lethal screening in fungal pathogens<sup>265</sup> may facilitate such identification of fungal targets for combination antifungal therapy.

### *New alternatives for antifungal drug therapy*

Given the prevalence of antifungal drug resistance among emerging fungal pathogens, unconventional therapeutic strategies may be essential for treating these infections. One alternative treatment strategy for fungal diseases aims to manipulate the host immune system to better combat infection. This includes cell-based therapies focused on replenishing immune cells<sup>266,267</sup> and the use of cytokines as adjunctive antifungal therapy,<sup>268–272</sup> which

has shown promising results for the treatment of invasive fungal infections in organ transplant patients,<sup>271</sup> for HIV/AIDS-associated *Cryptococcus* infections,<sup>268,272</sup> and for highly drug-resistant *Scedosporium* infection.<sup>273</sup> Additionally, there is growing interest in the use of monoclonal antibodies to target fungal pathogens<sup>238,274–278</sup> or for use in radioimmunotherapy.<sup>279</sup> Other innovative approaches to immune-based antifungal therapy include a bioengineering strategy to genetically modify T cells to redirect their specificity toward *Aspergillus*,<sup>280</sup> and a targeted kinase approach in mammalian cells that negatively regulates the host antifungal immune response in order to boost innate immunity against fungal infection.<sup>281</sup> Finally, there has been substantial interest in the development of antifungal vaccines.<sup>282–285</sup> There is mounting evidence for the efficacy of antifungal vaccines in preventing infection in animal models,<sup>286–291</sup> and a vaccine for recurrent vulvovaginal candidiasis has completed phase II clinical trials.

Another new paradigm for antimicrobial therapy is targeting virulence factors.<sup>292,293</sup> This strategy may be of particular interest for treating emerging drug-resistant fungi, as antivirulence therapeutics are predicted to impose less selection pressure toward drug resistance. Several recent studies have identified molecules that reduce virulence of *Candida* pathogens by preventing fungal adhesion, cellular morphogenesis, and/or biofilm formation.<sup>294–298</sup> Other factors targeting fungal virulence have also been identified through interactions with competing microbial organisms. The probiotic yeast *Saccharomyces boulardii* inhibits *C. albicans* filamentation, adhesion, and biofilm formation,<sup>299,300</sup> and a *Bacillus* bacterial species is able to inhibit virulence factor production and biofilm formation in both *C. neoformans* and *C. albicans*,<sup>301</sup> suggesting that investigating natural products from other microbial organisms may be a promising strategy to identify novel ways to target fungal virulence.

Finally, targeting antifungal resistance mechanisms holds potential as a powerful strategy to eliminate drug-resistant emerging fungal pathogens. Targeting drug efflux pumps has been the primary tool employed to overcome antifungal drug resistance. Inhibitors of ABC<sup>258,302–304</sup> and major facilitator superfamily<sup>305</sup> transporters, as well as broad-spectrum efflux pump inhibitors,<sup>306</sup>

have been identified, with the ability to reverse antifungal drug resistance in fungal pathogens, including drug-resistant *C. glabrata* species. Recent work exploited a high-throughput screening platform to identify a small-molecule inhibitor of a critical protein–protein interaction between the Pdr1 efflux pump in *C. glabrata* and the Mediator complex.<sup>307</sup> This new inhibitor blocks efflux pump activity and successfully reduced *C. glabrata* resistance to azole antifungals in *in vitro* and *in vivo* models of fungal infection,<sup>307</sup> demonstrating a compelling and innovative strategy to desensitize drug-resistant fungal pathogens.

## Conclusion

Pathogenic fungi are a leading cause of human mortality, particularly among an ever-increasing population of immunocompromised individuals. As newly identified fungal pathogens continue to emerge and previously rare pathogens increase in both prevalence and resistance to antifungal agents, combating these infections is becoming a pressing challenge for global health. Despite the significant burden of fungal infections, fungal diseases have been generally neglected by many public health initiatives<sup>308,309</sup> and require our renewed attention. The development of innovative diagnostics and therapeutics for emerging drug-resistant fungal pathogens holds promise for the future of treating fungal disease, and must continue to be emphasized.

## Acknowledgment

We would like to thank L. Certain for helpful discussions.

## Competing interests

The authors declare no competing interests.

## References

1. Denning, D.W. 2017. Calling upon all public health mycologists. *Eur. J. Clin. Microbiol. Infect. Dis.* **36**: 923–924.
2. Bongomin, F., S. Gago, R. Oladele, *et al.* 2017. Global and multi-national prevalence of fungal diseases—estimate precision. *J. Fungi* **3**: 57.
3. Brown, G.D., D.W. Denning, N.A.R. Gow, *et al.* 2012. Hidden killers: human fungal infections. *Sci. Transl. Med.* **4**: 165rv13.
4. Havlickova, B., V.A. Czaika & M. Friedrich. 2008. Epidemiological trends in skin mycoses worldwide. *Mycoses* **51**(Suppl. 4): 2–15.
5. Sobel, J.D. 2007. Vulvovaginal candidosis. *Lancet* **369**: 1961–1971.

6. Pfaller, M.A., S.A. Messer, L. Boyken, *et al.* 2005. Global trends in the antifungal susceptibility of *Cryptococcus neoformans* (1990 to 2004). *J. Clin. Microbiol.* **43**: 2163–2167.
7. Pfaller, M.A. & D.J. Diekema. 2007. Epidemiology of invasive candidiasis: a persistent public health problem. *Clin. Microbiol. Rev.* **20**: 133–163.
8. Morse, S.S. 1995. Factors in the emergence of infectious diseases. *Emerg. Infect. Dis.* **1**: 7–15.
9. Nucci, M. & K.A. Marr. 2005. Emerging fungal diseases. *Clin. Infect. Dis.* **41**: 521–526.
10. Denning, D.W. 2016. Minimizing fungal disease deaths will allow the UNAIDS target of reducing annual AIDS deaths below 500 000 by 2020 to be realized. *Philos. Trans. R. Soc. B.* **371**: 20150468.
11. Armstrong-James, D., G. Meintjes & G.D. Brown. 2014. A neglected epidemic: fungal infections in HIV/AIDS. *Trends Microbiol.* **22**: 120–127.
12. Bodey, G., B. Bueltmann, W. Duguid, *et al.* 1992. Fungal infections in cancer patients: an international autopsy survey. *Eur. J. Clin. Microbiol. Infect. Dis.* **11**: 99–109.
13. Pagano, L., M. Akova, G. Dimopoulos, *et al.* 2011. Risk assessment and prognostic factors for mould-related diseases in immunocompromised patients. *J. Antimicrob. Chemother.* **66**(Suppl. 1): i5–i14.
14. Pappas, P.G., B.D. Alexander, D.R. Andes, *et al.* 2010. Invasive fungal infections among organ transplant recipients: results of the Transplant-Associated Infection Surveillance Network (TRANSNET). *Clin. Infect. Dis.* **50**: 1101–1111.
15. Shoham, S. & K.A. Marr. 2012. Invasive fungal infections in solid organ transplant recipients. *Future Microbiol.* **7**: 639–655.
16. Kauffman, C.A. 2001. Fungal infections in older adults. *Clin. Infect. Dis.* **33**: 550–555.
17. Powell, K. 2012. Candidiasis: fungal infections in older adults. *Nurs. Resident. Care* **14**: 637–641.
18. Krčmery, V., F. Matejička, E. Pichňová, *et al.* 1999. Documented fungal infections after prophylaxis or therapy with wide spectrum antibiotics: relationship between certain fungal pathogens and particular antimicrobials? *J. Chemother.* **11**: 385–390.
19. Van Boeckel, T.P., S. Gandra, A. Ashok, *et al.* 2014. Global antibiotic consumption 2000 to 2010: an analysis of national pharmaceutical sales data. *Lancet Infect. Dis.* **14**: 742–750.
20. Kojic, E.M. & R.O. Darouiche. 2004. Candida infections of medical devices. *Clin. Microbiol. Rev.* **17**: 255–267.
21. Percival, S.L., L. Suleman, C. Vuotto, *et al.* 2015. Healthcare-associated infections, medical devices and biofilms: risk, tolerance and control. *J. Med. Microbiol.* **64**: 323–334.
22. Bebbler, D.P., M.A.T. Ramotowski & S.J. Gurr. 2013. Crop pests and pathogens move polewards in a warming world. *Nat. Clim. Change* **3**: 985.
23. Garcia-Solache, M.A. & A. Casadevall. 2010. Global warming will bring new fungal diseases for mammals. *mBio* **1**: e00061-10.
24. Spivak, E.S. & K.E. Hanson. 2018. *Candida auris*: an emerging fungal pathogen. *J. Clin. Microbiol.* **56**: pii: e01588-17.



25. Vazquez, L. 2016. Antifungal prophylaxis in immunocompromised patients. *Mediterr. J. Hematol. Infect. Dis.* **8**: e2016040.
26. Perfect, J.R. 2017. The antifungal pipeline: a reality check. *Nat. Rev. Drug Discov.* **16**: 603–616.
27. Kanafani, Z.A. & J.R. Perfect. 2008. Antimicrobial resistance: resistance to antifungal agents: mechanisms and clinical impact. *Clin. Infect. Dis.* **46**: 120–128.
28. Sanglard, D. 2016. Emerging threats in antifungal-resistant fungal pathogens. *Front. Med.* **3**: 11.
29. Klepser, M.E., E.J. Wolfe, R.N. Jones, *et al.* 1997. Antifungal pharmacodynamic characteristics of fluconazole and amphotericin B tested against *Candida albicans*. *Antimicrob. Agents Chemother.* **41**: 1392–1395.
30. Ernst, E.J., M.E. Klepser & M.A. Pfaller. 2000. Postantifungal effects of echinocandin, azole, and polyene antifungal agents against *Candida albicans* and *Cryptococcus neoformans*. *Antimicrob. Agents Chemother.* **44**: 1108–1111.
31. Gray, K.C., D.S. Palacios, I. Dailey, *et al.* 2012. Amphotericin primarily kills yeast by simply binding ergosterol. *Proc. Natl. Acad. Sci. USA* **109**: 2234–2239.
32. Anderson, T.M., M.C. Clay, A.G. Cioffi, *et al.* 2014. Amphotericin forms an extramembranous and fungicidal sterol sponge. *Nat. Chem. Biol.* **10**: 400–406.
33. Zotchev, S.B. 2003. Polyene macrolide antibiotics and their applications in human therapy. *Curr. Med. Chem.* **10**: 211–223.
34. Lemke, A., A.F. Kiderlen & O. Kayser. 2005. Amphotericin B. *Appl. Microbiol. Biotechnol.* **68**: 151–162.
35. Rodero, L., S. Córdoba, P. Cahn, *et al.* 2000. *In vitro* susceptibility studies of *Cryptococcus neoformans* isolated from patients with no clinical response to amphotericin B therapy. *J. Antimicrob. Chemother.* **45**: 239–242.
36. Janout, V., W.A. Schell, D. Thévenin, *et al.* 2015. Taming amphotericin B. *Bioconjug. Chem.* **26**: 2021–2024.
37. Ickowicz, D.E., S. Farber, E. Sionov, *et al.* 2014. Activity, reduced toxicity, and scale-up synthesis of amphotericin B-conjugated polysaccharide. *Biomacromolecules* **15**: 2079–2089.
38. Delmas, G., S. Park, Z.W. Chen, *et al.* 2002. Efficacy of orally delivered cochleates containing amphotericin B in a murine model of aspergillosis. *Antimicrob. Agents Chemother.* **46**: 2704–2707.
39. Yang, Z., M. Liu, J. Chen, *et al.* 2014. Development and characterization of amphotericin B nanosuspensions for oral administration through a simple top-down method. *Curr. Pharm. Biotechnol.* **15**: 569–576.
40. Halperin, A., Y. Shadkhan, E. Pisarevsky, *et al.* 2016. Novel water-soluble amphotericin B-PEG conjugates with low toxicity and potent *in vivo* efficacy. *J. Med. Chem.* **59**: 1197–1206.
41. Sheehan, D.J., C.A. Hitchcock & C.M. Sibley. 1999. Current and emerging azole antifungal agents. *Clin. Microbiol. Rev.* **12**: 40–79.
42. Morita, T. & Y. Nozawa. 1985. Effects of antifungal agents on ergosterol biosynthesis in *Candida albicans* and *Trichophyton mentagrophytes*: differential inhibitory sites of naphthiomate and miconazole. *J. Invest. Dermatol.* **85**: 434–437.
43. Pappas, P.G., C.A. Kauffman, D.R. Andes, *et al.* 2016. Clinical practice guideline for the management of candidiasis: 2016 update by the Infectious Diseases Society of America. *Clin. Infect. Dis.* **62**: e1–e50.
44. Lass-Flörl, C. 2011. Triazole antifungal agents in invasive fungal infections: a comparative review. *Drugs* **71**: 2405–2419.
45. Andriole, V.T. 2000. Current and future antifungal therapy: new targets for antifungal therapy. *Int. J. Antimicrob. Agents* **16**: 317–321.
46. Manavathu, E.K., J.L. Cutright & P.H. Chandrasekar. 1998. Organism-dependent fungicidal activities of azoles. *Antimicrob. Agents Chemother.* **42**: 3018–3021.
47. Allen, D., D. Wilson, R. Drew, *et al.* 2015. Azole antifungals: 35 years of invasive fungal infection management. *Expert Rev. Anti-Infect. Ther.* **13**: 787–798.
48. Cowen, L.E. & W.J. Steinbach. 2008. Stress, drugs, and evolution: the role of cellular signaling in fungal drug resistance. *Eukaryot. Cell* **7**: 747–764.
49. Anderson, J.B. 2005. Evolution of antifungal-drug resistance: mechanisms and pathogen fitness. *Nat. Rev. Microbiol.* **3**: 547–556.
50. Peyton, L.R., S. Gallagher & M. Hashemzadeh. 2015. Triazole antifungals: a review. *Drugs Today* **51**: 705–718.
51. Sanglard, D. & F.C. Odds. 2002. Resistance of *Candida* species to antifungal agents: molecular mechanisms and clinical consequences. *Lancet Infect. Dis.* **2**: 73–85.
52. Xiang, M.-J., J.-Y. Liu, P.-H. Ni, *et al.* 2013. Erg11 mutations associated with azole resistance in clinical isolates of *Candida albicans*. *FEMS Yeast Res.* **13**: 386–393.
53. Wang, B., L.-H. Huang, J.-X. Zhao, *et al.* 2015. *ERG11* mutations associated with azole resistance in *Candida albicans* isolates from vulvovaginal candidosis patients. *Asian Pac. J. Trop. Biomed.* **5**: 909–914.
54. Forastiero, A., A.C. Mesa-Arango, A. Alastruey-Izquierdo, *et al.* 2013. *Candida tropicalis* antifungal cross-resistance is related to different azole target (Erg11p) modifications. *Antimicrob. Agents Chemother.* **57**: 4769–4781.
55. White, T.C., S. Holleman, F. Dy, *et al.* 2002. Resistance mechanisms in clinical isolates of *Candida albicans*. *Antimicrob. Agents Chemother.* **46**: 1704–1713.
56. Cannon, R.D., E. Lamping, A.R. Holmes, *et al.* 2009. Efflux-mediated antifungal drug resistance. *Clin. Microbiol. Rev.* **22**: 291–321.
57. Joseph-Horne, T., D. Hollomon, R.S. Loeffler, *et al.* 1995. Cross-resistance to polyene and azole drugs in *Cryptococcus neoformans*. *Antimicrob. Agents Chemother.* **39**: 1526–1529.
58. Kelly, S.L., D.C. Lamb, D.E. Kelly, *et al.* 1997. Resistance to fluconazole and cross-resistance to amphotericin B in *Candida albicans* from AIDS patients caused by defective sterol  $\Delta 5,6$ -desaturation. *FEBS Lett.* **400**: 80–82.
59. Vandeputte, P., G. Tronchin, T. Bergès, *et al.* 2007. Reduced susceptibility to polyenes associated with a missense mutation in the *ERG6* gene in a clinical isolate of *Candida glabrata* with pseudohyphal growth. *Antimicrob. Agents Chemother.* **51**: 982–990.
60. Waldorf, A.R. & A. Polak. 1983. Mechanisms of action of 5-fluorocytosine. *Antimicrob. Agents Chemother.* **23**: 79–85.

61. Edlind, T.D. & S.K. Katiyar. 2010. Mutational analysis of flucytosine resistance in *Candida glabrata*. *Antimicrob. Agents Chemother.* **54**: 4733–4738.
62. Vandeputte, P., L. Pineau, G. Larcher, *et al.* 2011. Molecular mechanisms of resistance to 5-fluorocytosine in laboratory mutants of *Candida glabrata*. *Mycopathologia* **171**: 11–21.
63. Day, J.N., T.T.H. Chau, M. Wolbers, *et al.* 2013. Combination antifungal therapy for cryptococcal meningitis. *N. Engl. J. Med.* **368**: 1291–1302.
64. Schwarz, P., F. Dromer, O. Lortholary, *et al.* 2006. Efficacy of amphotericin B in combination with flucytosine against flucytosine-susceptible or flucytosine-resistant isolates of *Cryptococcus neoformans* during disseminated murine cryptococcosis. *Antimicrob. Agents Chemother.* **50**: 113–120.
65. Loyse, A., F. Dromer, J. Day, *et al.* 2013. Flucytosine and cryptococcosis: time to urgently address the worldwide accessibility of a 50-year-old antifungal. *J. Antimicrob. Chemother.* **68**: 2435–2444.
66. Denning, D.W. 2002. Echinocandins: a new class of antifungal. *J. Antimicrob. Chemother.* **49**: 889–891.
67. Denning, D.W. 2003. Echinocandin antifungal drugs. *Lancet* **362**: 1142–1151.
68. Perlin, D.S. 2007. Resistance to echinocandin-class antifungal drugs. *Drug Resist. Updat.* **10**: 121–130.
69. Pfaller, M.A. & D.J. Diekema. 2012. Progress in antifungal susceptibility testing of *Candida* spp. by use of Clinical and Laboratory Standards Institute broth microdilution methods, 2010 to 2012. *J. Clin. Microbiol.* **50**: 2846–2856.
70. Arendrup, M.C., G. Garcia-Effron, C. Lass-Flörl, *et al.* 2010. Echinocandin susceptibility testing of *Candida* species: comparison of EUCAST EDef 7.1, CLSI M27-A3, Etest, disk diffusion, and agar dilution methods with RPMI and isosensitest media. *Antimicrob. Agents Chemother.* **54**: 426–439.
71. Cuenca-Estrella, M. & J.L. Rodriguez-Tudela. 2010. The current role of the reference procedures by CLSI and EUCAST in the detection of resistance to antifungal agents *in vitro*. *Expert Rev. Anti-Infect. Ther.* **8**: 267–276.
72. Doern, G.V. & S.M. Brecher. 2011. The clinical predictive value (or lack thereof) of the results of *in vitro* antimicrobial susceptibility tests. *J. Clin. Microbiol.* **49**: S11–S14.
73. Delarze, E. & D. Sanglard. 2015. Defining the frontiers between antifungal resistance, tolerance and the concept of persistence. *Drug Resist. Updat.* **23**: 12–19.
74. Shapiro, R.S., N. Robbins & L.E. Cowen. 2011. Regulatory circuitry governing fungal development, drug resistance, and disease. *Microbiol. Mol. Biol. Rev.* **75**: 213–267.
75. McCarty, T.P. & P.G. Pappas. 2016. Invasive candidiasis. *Infect. Dis. Clin. North Am.* **30**: 103–124.
76. Nobile, C.J. & A.D. Johnson. 2015. *Candida albicans* biofilms and human disease. *Annu. Rev. Microbiol.* **69**: 71–92.
77. Pfaller, M.A. & D.J. Diekema. 2004. Rare and emerging opportunistic fungal pathogens: concern for resistance beyond *Candida albicans* and *Aspergillus fumigatus*. *J. Clin. Microbiol.* **42**: 4419–4431.
78. Bernal-Martinez, L., A. Gomez-Lopez, M.V. Castelli, *et al.* 2010. Susceptibility profile of clinical isolates of non-*Cryptococcus neoformans*/non-*Cryptococcus gattii* *Cryptococcus* species and literature review. *Med. Mycol.* **48**: 90–96.
79. Kołaczowska, A. & M. Kołaczowski. 2016. Drug resistance mechanisms and their regulation in non-*albicans* *Candida* species. *J. Antimicrob. Chemother.* **71**: 1438–1450.
80. Rodrigues, C.F., S. Silva & M. Henriques. 2014. *Candida glabrata*: a review of its features and resistance. *Eur. J. Clin. Microbiol. Infect. Dis.* **33**: 673–688.
81. Diekema, D.J., S.A. Messer, A.B. Brueggemann, *et al.* 2002. Epidemiology of candidemia: 3-year results from the emerging infections and the epidemiology of Iowa organisms study. *J. Clin. Microbiol.* **40**: 1298–1302.
82. Papon, N., V. Courdavault, M. Clastre, *et al.* 2013. Emerging and emerged pathogenic *Candida* species: beyond the *Candida albicans* paradigm. *PLoS Pathog.* **9**: e1003550.
83. Pfaller, M.A., S.A. Messer, L. Boyken, *et al.* 2004. Geographic variation in the susceptibilities of invasive isolates of *Candida glabrata* to seven systemically active antifungal agents: a global assessment from the ARTEMIS Antifungal Surveillance Program conducted in 2001 and 2002. *J. Clin. Microbiol.* **42**: 3142–3146.
84. Pfaller, M.A., D.J. Diekema, D.L. Gibbs, *et al.* 2010. Results from the ARTEMIS DISK Global Antifungal Surveillance Study, 1997 to 2007: a 10.5-year analysis of susceptibilities of *Candida* species to fluconazole and voriconazole as determined by CLSI standardized disk diffusion. *J. Clin. Microbiol.* **48**: 1366–1377.
85. Bennett, J.E., K. Izumikawa & K.A. Marr. 2004. Mechanism of increased fluconazole resistance in *Candida glabrata* during prophylaxis. *Antimicrob. Agents Chemother.* **48**: 1773–1777.
86. Sanglard, D., F. Ischer, D. Calabrese, *et al.* 1999. The ATP binding cassette transporter gene *CgCDR1* from *Candida glabrata* is involved in the resistance of clinical isolates to azole antifungal agents. *Antimicrob. Agents Chemother.* **43**: 2753–2765.
87. Marr, K.A., K. Seidel, T.C. White, *et al.* 2000. Candidemia in allogeneic blood and marrow transplant recipients: evolution of risk factors after the adoption of prophylactic fluconazole. *J. Infect. Dis.* **181**: 309–316.
88. Vazquez, J.A., J.D. Sobel, G. Peng, *et al.* 1999. Evolution of vaginal *Candida* species recovered from human immunodeficiency virus-infected women receiving fluconazole prophylaxis: the emergence of *Candida glabrata*? Terry Bein Community Programs for Clinical Research in AIDS (CPCRA). *Clin. Infect. Dis.* **28**: 1025–1031.
89. Pfaller, M.A., D.J. Diekema, R.N. Jones, *et al.* 2002. Trends in antifungal susceptibility of *Candida* spp. isolated from pediatric and adult patients with bloodstream infections: SENTRY Antimicrobial Surveillance Program, 1997 to 2000. *J. Clin. Microbiol.* **40**: 852–856.
90. Kuse, E.-R., P. Chetchotisakd, C.A. da Cunha, *et al.* 2007. Micafungin versus liposomal amphotericin B for candidaemia and invasive candidosis: a phase III randomised double-blind trial. *Lancet* **369**: 1519–1527.
91. Pappas, P.G., C.M.F. Rotstein, R.F. Betts, *et al.* 2007. Micafungin versus caspofungin for treatment of candidemia

- and other forms of invasive candidiasis. *Clin. Infect. Dis.* **45**: 883–893.
92. Perlin, D.S. 2015. Echinocandin resistance in *Candida*. *Clin. Infect. Dis.* **61**: S612–S617.
  93. Alexander, B.D., M.D. Johnson, C.D. Pfeiffer, *et al.* 2013. Increasing echinocandin resistance in *Candida glabrata*: clinical failure correlates with presence of *FKS* mutations and elevated minimum inhibitory concentrations. *Clin. Infect. Dis.* **56**: 1724–1732.
  94. Pham, C.D., N. Iqbal, C.B. Bolden, *et al.* 2014. Role of *FKS* mutations in *Candida glabrata*: MIC values, echinocandin resistance, and multidrug resistance. *Antimicrob. Agents Chemother.* **58**: 4690–4696.
  95. Healey, K.R., Y. Zhao, W.B. Perez, *et al.* 2016. Prevalent mutator genotype identified in fungal pathogen *Candida glabrata* promotes multi-drug resistance. *Nat. Commun.* **7**: 11128.
  96. Orozco, A.S., L.M. Higginbotham, C.A. Hitchcock, *et al.* 1998. Mechanism of fluconazole resistance in *Candida krusei*. *Antimicrob. Agents Chemother.* **42**: 2645.
  97. Abi-Said, D., E. Anaissie, O. Uzun, *et al.* 1997. The epidemiology of hematogenous candidiasis caused by different *Candida* species. *Clin. Infect. Dis.* **24**: 1122–1128.
  98. Singh, N. 2001. Trends in the epidemiology of opportunistic fungal infections: predisposing factors and the impact of antimicrobial use practices. *Clin. Infect. Dis.* **33**: 1692–1696.
  99. Atkinson, B.J., R.E. Lewis & D.P. Kontoyiannis. 2008. *Candida lusitanae* fungemia in cancer patients: risk factors for amphotericin B failure and outcome. *Med. Mycol.* **46**: 541–546.
  100. Hakkı, M., J.F. Staab & K.A. Marr. 2006. Emergence of a *Candida krusei* isolate with reduced susceptibility to caspofungin during therapy. *Antimicrob. Agents Chemother.* **50**: 2522–2524.
  101. Asner, S.A., S. Giulieri, M. Diezi, *et al.* 2015. Acquired multidrug antifungal resistance in *Candida lusitanae* during therapy. *Antimicrob. Agents Chemother.* **59**: 7715–7722.
  102. Rodero, L., M. Cuenca-Estrella, S. Córdoba, *et al.* 2002. Transient fungemia caused by an amphotericin B-resistant isolate of *Candida haemulonii*. *J. Clin. Microbiol.* **40**: 2266–2269.
  103. Kim, M., J.H. Shin, H. Sung, *et al.* 2009. *Candida haemulonii* and closely related species at 5 university hospitals in Korea: identification, antifungal susceptibility, and clinical features. *Clin. Infect. Dis.* **48**: e57–e61.
  104. Ruan, S.-Y., Y.-W. Kuo, C.-T. Huang, *et al.* 2010. Infections due to *Candida haemulonii*: species identification, antifungal susceptibility and outcomes. *Int. J. Antimicrob. Agents* **35**: 85–88.
  105. Sarvikivi, E., O. Lyytikäinen, D.R. Soll, *et al.* 2005. Emergence of fluconazole resistance in a *Candida parapsilosis* strain that caused infections in a neonatal intensive care unit. *J. Clin. Microbiol.* **43**: 2729–2735.
  106. Cleveland, A.A., M.M. Farley, L.H. Harrison, *et al.* 2012. Changes in incidence and antifungal drug resistance in candidemia: results from population-based laboratory surveillance in Atlanta and Baltimore, 2008–2011. *Clin. Infect. Dis.* **55**: 1352–1361.
  107. Satoh, K., K. Makimura, Y. Hasumi, *et al.* 2009. *Candida auris* sp. nov., a novel ascomycetous yeast isolated from the external ear canal of an inpatient in a Japanese hospital. *Microbiol. Immunol.* **53**: 41–44.
  108. Lockhart, S.R., K.A. Etienne, S. Vallabhaneni, *et al.* 2017. Simultaneous emergence of multidrug-resistant *Candida auris* on 3 continents confirmed by whole-genome sequencing and epidemiological analyses. *Clin. Infect. Dis.* **64**: 134–140.
  109. Chowdhary, A., C. Sharma & J.F. Meis. 2017. *Candida auris*: a rapidly emerging cause of hospital-acquired multidrug-resistant fungal infections globally. *PLoS Pathog.* **13**: e1006290.
  110. Chakrabarti, A., P. Sood, S.M. Rudramurthy, *et al.* 2015. Incidence, characteristics and outcome of ICU-acquired candidemia in India. *Intensive Care Med.* **41**: 285–295.
  111. Sarma, S. & S. Upadhyay. 2017. Current perspective on emergence, diagnosis and drug resistance in *Candida auris*. *Infect. Drug Resist.* **10**: 155–165.
  112. Kathuria, S., P.K. Singh, C. Sharma, *et al.* 2015. Multidrug-resistant *Candida auris* misidentified as *Candida haemulonii*: characterization by matrix-assisted laser desorption/ionization-time of flight mass spectrometry and DNA sequencing and its antifungal susceptibility profile variability by vitek 2, CLSI broth microdilution, and Etest method. *J. Clin. Microbiol.* **53**: 1823–1830.
  113. Borman, A.M., A. Szekeley & E.M. Johnson. 2016. Comparative pathogenicity of United Kingdom isolates of the emerging pathogen *Candida auris* and other key pathogenic *Candida* species. *mSphere* **1**: pii: e00189-16.
  114. Larkin, E., C. Hager, J. Chandra, *et al.* 2017. The emerging pathogen *Candida auris*: growth phenotype, virulence factors, activity of antifungals, and effect of SCY-078, a novel glucan synthesis inhibitor, on growth morphology and biofilm formation. *Antimicrob. Agents Chemother.* **61**: e02396-16.
  115. Sharma, C., N. Kumar, J.F. Meis, *et al.* 2015. Draft genome sequence of a fluconazole-resistant *Candida auris* strain from a candidemia patient in India. *Genome Announc.* **3**: pii: e00722-15.
  116. Chatterjee, S., S.V. Alampalli, R.K. Nageshan, *et al.* 2015. Draft genome of a commonly misdiagnosed multidrug resistant pathogen *Candida auris*. *BMC Genomics* **16**: 686.
  117. Ben-Ami, R., J. Berman, A. Novikov, *et al.* 2017. Multidrug-resistant *Candida haemulonii* and *C. auris*, Tel Aviv, Israel. *Emerg. Infect. Dis.* **23**: 195–203.
  118. Derek, J. & V.P. Sloan. 2014. Cryptococcal meningitis: epidemiology and therapeutic options. *Clin. Epidemiol.* **6**: 169.
  119. Rajasingham, R., R.M. Smith, B.J. Park, *et al.* 2017. Global burden of disease of HIV-associated cryptococcal meningitis: an updated analysis. *Lancet Infect. Dis.* **17**: 873–881.
  120. Bartlett, K.H., S.E. Kidd & J.W. Kronstad. 2008. The emergence of *Cryptococcus gattii* in British Columbia and the Pacific Northwest. *Curr. Infect. Dis. Rep.* **10**: 58–65.
  121. Harris, J.R., S.R. Lockhart, E. Debess, *et al.* 2011. *Cryptococcus gattii* in the United States: clinical aspects of infection

- with an emerging pathogen. *Clin. Infect. Dis.* **53**: 1188–1195.
122. Gomez-Lopez, A., O. Zaragoza, M. Dos Anjos Martins, *et al.* 2008. *In vitro* susceptibility of *Cryptococcus gattii* clinical isolates. *Clin. Microbiol. Infect.* **14**: 727–730.
  123. Gast, C.E., L.R. Basso, Jr., I. Bruzual, *et al.* 2013. Azole resistance in *Cryptococcus gattii* from the Pacific Northwest: investigation of the role of ERG11. *Antimicrob. Agents Chemother.* **57**: 5478–5485.
  124. Billmyre, R.B., S.A. Clancey & J. Heitman. 2017. Natural mismatch repair mutations mediate phenotypic diversity and drug resistance in *Cryptococcus deuterogattii*. *Elife* **6**: pii: e28802.
  125. Rhodes, J., M.A. Beale, M. Vanhove, *et al.* 2017. A population genomics approach to assessing the genetic basis of within-host microevolution underlying recurrent cryptococcal meningitis infection. *G3* **7**: 1165–1176.
  126. Boyce, K.J., Y. Wang, S. Verma, *et al.* 2017. Mismatch repair of DNA replication errors contributes to microevolution in the pathogenic fungus *Cryptococcus neoformans*. *mBio* **8**: e00595–17.
  127. Khawcharoenporn, T., A. Apisarnthanarak & L.M. Mundy. 2007. Non-*neoformans* cryptococcal infections: a systematic review. *Infection* **35**: 51–58.
  128. Smith, N., M. Sehring, J. Chambers, *et al.* 2017. Perspectives on non-*neoformans* cryptococcal opportunistic infections. *J. Community Hosp. Intern. Med. Perspect.* **7**: 214–217.
  129. van de Veerdonk, F.L., M.S. Gresnigt, L. Romani, *et al.* 2017. *Aspergillus fumigatus* morphology and dynamic host interactions. *Nat. Rev. Microbiol.* **15**: 661–674.
  130. Walsh, T.J. & A.H. Groll. 2001. Overview: non-*fumigatus* species of *Aspergillus*: perspectives on emerging pathogens in immunocompromised hosts. *Curr. Opin. Investig. Drugs* **2**: 1366–1367.
  131. Mortensen, K.L., H.K. Johansen, K. Fuursted, *et al.* 2011. A prospective survey of *Aspergillus* spp. in respiratory tract samples: prevalence, clinical impact and antifungal susceptibility. *Eur. J. Clin. Microbiol. Infect. Dis.* **30**: 1355–1363.
  132. Denning, D.W., A. Pleuvry & D.C. Cole. 2013. Global burden of allergic bronchopulmonary aspergillosis with asthma and its complication chronic pulmonary aspergillosis in adults. *Med. Mycol.* **51**: 361–370.
  133. Stevens, D.A., R.B. Moss, V.P. Kurup, *et al.* 2003. Allergic bronchopulmonary aspergillosis in cystic fibrosis—state of the art: Cystic Fibrosis Foundation Consensus Conference. *Clin. Infect. Dis.* **37**(Suppl. 3): S225–S264.
  134. Snelders, E., H.A.L. van der Lee, J. Kuijpers, *et al.* 2008. Emergence of azole resistance in *Aspergillus fumigatus* and spread of a single resistance mechanism. *PLoS Med.* **5**: e219.
  135. Howard, S.J., D. Cerar, M.J. Anderson, *et al.* 2009. Frequency and evolution of azole resistance in *Aspergillus fumigatus* associated with treatment failure. *Emerg. Infect. Dis.* **15**: 1068–1076.
  136. Snelders, E., R.A.G. Huis In 't Veld, A.J.M.M. Rijs, *et al.* 2009. Possible environmental origin of resistance of *Aspergillus fumigatus* to medical triazoles. *Appl. Environ. Microbiol.* **75**: 4053–4057.
  137. Berger, S., Y. El Chazli, A.F. Babu, *et al.* 2017. Azole resistance in *Aspergillus fumigatus*: a consequence of antifungal use in agriculture? *Front. Microbiol.* **8**: 1024.
  138. Snelders, E., S.M.T. Camps, A. Karawajczyk, *et al.* 2012. Triazole fungicides can induce cross-resistance to medical triazoles in *Aspergillus fumigatus*. *PLoS One* **7**: e31801.
  139. Chowdhary, A., S. Kathuria, J. Xu, *et al.* 2013. Emergence of azole-resistant *Aspergillus fumigatus* strains due to agricultural azole use creates an increasing threat to human health. *PLoS Pathog.* **9**: e1003633.
  140. Steinbach, W.J., K.A. Marr, E.J. Anaissie, *et al.* 2012. Clinical epidemiology of 960 patients with invasive aspergillosis from the PATH Alliance registry. *J. Infect.* **65**: 453–464.
  141. Hachem, R.Y., D.P. Kontoyiannis, M.R. Boktour, *et al.* 2004. *Aspergillus terreus*: an emerging amphotericin B-resistant opportunistic mold in patients with hematologic malignancies. *Cancer* **101**: 1594–1600.
  142. Pfaller, M.A., S.A. Messer, R.J. Hollis, *et al.* 2002. Antifungal activities of posaconazole, ravuconazole, and voriconazole compared to those of itraconazole and amphotericin B against 239 clinical isolates of *Aspergillus* spp. and other filamentous fungi: report from SENTRY Antimicrobial Surveillance Program, 2000. *Antimicrob. Agents Chemother.* **46**: 1032–1037.
  143. Hachem, R., M.Z.R. Gomes, G. El Helou, *et al.* 2014. Invasive aspergillosis caused by *Aspergillus terreus*: an emerging opportunistic infection with poor outcome independent of azole therapy. *J. Antimicrob. Chemother.* **69**: 3148–3155.
  144. Van Der Linden, J.W.M., A. Warris & P.E. Verweij. 2011. *Aspergillus* species intrinsically resistant to antifungal agents. *Med. Mycol.* **49**: S82–S89.
  145. Ramos, E.R., Y. Jiang, R. Hachem, *et al.* 2011. Outcome analysis of invasive aspergillosis in hematologic malignancy and hematopoietic stem cell transplant patients: the role of novel antimold azoles. *Oncologist* **16**: 1049–1060.
  146. Blum, G., C. Hortnagl, E. Jukic, *et al.* 2013. New insight into amphotericin B resistance in *Aspergillus terreus*. *Antimicrob. Agents Chemother.* **57**: 1583–1588.
  147. Arendrup, M.C., R.H. Jensen, K. Grif, *et al.* 2012. *In vivo* emergence of *Aspergillus terreus* with reduced azole susceptibility and a Cyp51a M217I alteration. *J. Infect. Dis.* **206**: 981–985.
  148. Gomez-Lopez, A., G. Garcia-Effron, E. Mellado, *et al.* 2003. *In vitro* activities of three licensed antifungal agents against Spanish clinical isolates of *Aspergillus* spp. *Antimicrob. Agents Chemother.* **47**: 3085–3088.
  149. Pfaller, M.A., F. Duncanson, S.A. Messer, *et al.* 2011. *In vitro* activity of a novel broad-spectrum antifungal, E1210, tested against *Aspergillus* spp. determined by CLSI and EUCAST broth microdilution methods. *Antimicrob. Agents Chemother.* **55**: 5155–5158.
  150. Salas, V., F.J. Pastor, M.M. Rodríguez, *et al.* 2011. *In vitro* activity and *in vivo* efficacy of posaconazole in treatment of murine infections by different isolates of the *Aspergillus terreus* complex. *Antimicrob. Agents Chemother.* **55**: 676–679.
  151. Slavin, M., S. van Hal, T.C. Sorrell, *et al.* 2015. Invasive infections due to filamentous fungi other than *Aspergillus*:



- epidemiology and determinants of mortality. *Clin. Microbiol. Infect.* **21**: 490.e1-10.
152. Pfaller, M.A. & D.J. Diekema. 2010. Epidemiology of invasive mycoses in North America. *Crit. Rev. Microbiol.* **36**: 1–53.
  153. Bitar, D., O. Lortholary, Y. Le Strat, *et al.* 2014. Population-based analysis of invasive fungal infections, France, 2001–2010. *Emerg. Infect. Dis.* **20**: 1163–1169.
  154. Roden, M.M., T.E. Zaoutis, W.L. Buchanan, *et al.* 2005. Epidemiology and outcome of zygomycosis: a review of 929 reported cases. *Clin. Infect. Dis.* **41**: 634–653.
  155. Strasfeld, L., L. Espinosa-Aguilar, J.L. Gajewski, *et al.* 2013. Emergence of *Cunninghamella* as a pathogenic invasive mold infection in allogeneic transplant recipients. *Clin. Lymphoma Myeloma Leuk.* **13**: 622–628.
  156. Katragkou, A., T.J. Walsh & E. Roilides. 2014. Why is mucormycosis more difficult to cure than more common mycoses? *Clin. Microbiol. Infect.* **20**(Suppl. 6): 74–81.
  157. Grahl, N., S. Puttikamonkul, J.M. Macdonald, *et al.* 2011. *In vivo* hypoxia and a fungal alcohol dehydrogenase influence the pathogenesis of invasive pulmonary aspergillosis. *PLoS Pathog.* **7**: e1002145.
  158. Maurer, E., U. Binder, M. Sparber, *et al.* 2015. Susceptibility profiles of amphotericin B and posaconazole against clinically relevant mucorales species under hypoxic conditions. *Antimicrob. Agents Chemother.* **59**: 1344–1346.
  159. Sun, Q.-N., A.W. Fothergill, D.I. McCarthy, *et al.* 2002. *In vitro* activities of posaconazole, itraconazole, voriconazole, amphotericin B, and fluconazole against 37 clinical isolates of zygomycetes. *Antimicrob. Agents Chemother.* **46**: 1581–1582.
  160. Vitale, R.G., G.S. de Hoog, P. Schwarz, *et al.* 2012. Antifungal susceptibility and phylogeny of opportunistic members of the order mucorales. *J. Clin. Microbiol.* **50**: 66–75.
  161. Imhof, A., S.A. Balajee, D.N. Fredricks, *et al.* 2004. Break-through fungal infections in stem cell transplant recipients receiving voriconazole. *Clin. Infect. Dis.* **39**: 743–746.
  162. Marty, F.M., L. Ostrosky-Zeichner, O.A. Cornely, *et al.* 2016. Isavuconazole treatment for mucormycosis: a single-arm open-label trial and case-control analysis. *Lancet Infect. Dis.* **16**: 828–837.
  163. Pfaller, M.A., F. Marco, S.A. Messer, *et al.* 1998. *In vitro* activity of two echinocandin derivatives, LY303366 and MK-0991 (L-743,792), against clinical isolates of *Aspergillus*, *Fusarium*, *Rhizopus*, and other filamentous fungi. *Diagn. Microbiol. Infect. Dis.* **30**: 251–255.
  164. Ibrahim, A.S., J.C. Bowman, V. Avanesian, *et al.* 2005. Caspofungin inhibits *Rhizopus oryzae* 1,3-beta-D-glucan synthase, lowers burden in brain measured by quantitative PCR, and improves survival at a low but not a high dose during murine disseminated zygomycosis. *Antimicrob. Agents Chemother.* **49**: 721–727.
  165. Park, B.J., P.G. Pappas, K.A. Wannemuehler, *et al.* 2011. Invasive non-*Aspergillus* mold infections in transplant recipients, United States, 2001–2006. *Emerg. Infect. Dis.* **17**: 1855–1864.
  166. Nucci, M. & E. Anaissie. 2007. *Fusarium* infections in immunocompromised patients. *Clin. Microbiol. Rev.* **20**: 695–704.
  167. Dignani, M.C. & E. Anaissie. 2004. Human fusariosis. *Clin. Microbiol. Infect.* **10**(Suppl. 1): 67–75.
  168. Nucci, M., K.A. Marr, F. Queiroz-Telles, *et al.* 2004. *Fusarium* infection in hematopoietic stem cell transplant recipients. *Clin. Infect. Dis.* **38**: 1237–1242.
  169. Nucci, M., E.J. Anaissie, F. Queiroz-Telles, *et al.* 2003. Outcome predictors of 84 patients with hematologic malignancies and *Fusarium* infection. *Cancer* **98**: 315–319.
  170. Al-Hatmi, A.M., F. Hagen, S.B. Menken, *et al.* 2016. Global molecular epidemiology and genetic diversity of *Fusarium*, a significant emerging group of human opportunists from 1958 to 2015. *Emerg. Microbes Infect.* **5**: e124.
  171. Alastruay-Izquierdo, A., M. Cuenca-Estrella, A. Monzón, *et al.* 2008. Antifungal susceptibility profile of clinical *Fusarium* spp. isolates identified by molecular methods. *J. Antimicrob. Chemother.* **61**: 805–809.
  172. Pujol, I., J. Guarro, J. Gené, *et al.* 1997. *In-vitro* antifungal susceptibility of clinical and environmental *Fusarium* spp. strains. *J. Antimicrob. Chemother.* **39**: 163–167.
  173. Azor, M., J. Gené, J. Cano, *et al.* 2007. Universal *in vitro* antifungal resistance of genetic clades of the *Fusarium solani* species complex. *Antimicrob. Agents Chemother.* **51**: 1500–1503.
  174. Katiyar, S.K. & T.D. Edlind. 2009. Role for Fks1 in the intrinsic echinocandin resistance of *Fusarium solani* as evidenced by hybrid expression in *Saccharomyces cerevisiae*. *Antimicrob. Agents Chemother.* **53**: 1772–1778.
  175. Heyn, K., A. Tredup, S. Salvenmoser, *et al.* 2005. Effect of voriconazole combined with micafungin against *Candida*, *Aspergillus*, and *Scedosporium* spp. and *Fusarium solani*. *Antimicrob. Agents Chemother.* **49**: 5157–5159.
  176. Fan, J., M. Urban, J.E. Parker, *et al.* 2013. Characterization of the sterol 14 $\alpha$ -demethylases of *Fusarium graminearum* identifies a novel genus-specific CYP51 function. *New Phytol.* **198**: 821–835.
  177. Spader, T.B., T.P. Venturini, L. Rossato, *et al.* 2013. Synergism of voriconazole or itraconazole with other antifungal agents against species of *Fusarium*. *Rev. Iberoam. Micol.* **30**: 200–204.
  178. Al-Hatmi, A.M.S., J. Meletiadis, I. Curfs-Breuker, *et al.* 2016. *In vitro* combinations of natamycin with voriconazole, itraconazole and micafungin against clinical *Fusarium* strains causing keratitis. *J. Antimicrob. Chemother.* **71**: 953–955.
  179. Lamaris, G.A., G. Chamilos, R.E. Lewis, *et al.* 2006. *Scedosporium* infection in a tertiary care cancer center: a review of 25 cases from 1989–2006. *Clin. Infect. Dis.* **43**: 1580–1584.
  180. Cortez, K.J., E. Roilides, F. Quiroz-Telles, *et al.* 2008. Infections caused by *Scedosporium* spp. *Clin. Microbiol. Rev.* **21**: 157–197.
  181. Heath, C.H., M.A. Slavin, T.C. Sorrell, *et al.* 2009. Population-based surveillance for scedosporiosis in Australia: epidemiology, disease manifestations and emergence of *Scedosporium aurantiacum* infection. *Clin. Microbiol. Infect.* **15**: 689–693.
  182. Gilgado, F., J. Cano, J. Gené, *et al.* 2008. Molecular and phenotypic data supporting distinct species statuses for *Scedosporium apiospermum* and *Pseudallescheria boydii* and



- the proposed new species *Scedosporium dehoogii*. *J. Clin. Microbiol.* **46**: 766–771.
183. Cimon, B., J. Carrère, J.F. Vinatier, *et al.* 2000. Clinical significance of *Scedosporium apiospermum* in patients with cystic fibrosis. *Eur. J. Clin. Microbiol. Infect. Dis.* **19**: 53–56.
  184. Blyth, C.C., P.G. Middleton, A. Harun, *et al.* 2010. Clinical associations and prevalence of *Scedosporium* spp. in Australian cystic fibrosis patients: identification of novel risk factors? *Med. Mycol.* **48**(Suppl. 1): S37–S44.
  185. Rodriguez-Tudela, J.L., J. Berenguer, J. Guarro, *et al.* 2009. Epidemiology and outcome of *Scedosporium prolificans* infection, a review of 162 cases. *Med. Mycol.* **47**: 359–370.
  186. Gilgado, F., C. Serena, J. Cano, *et al.* 2006. Antifungal susceptibilities of the species of the *Pseudallescheria boydii* complex. *Antimicrob. Agents Chemother.* **50**: 4211–4213.
  187. Lackner, M., G.S. de Hoog, P.E. Verweij, *et al.* 2012. Species-specific antifungal susceptibility patterns of *Scedosporium* and *Pseudallescheria* species. *Antimicrob. Agents Chemother.* **56**: 2635–2642.
  188. Sabatelli, F., R. Patel, P.A. Mann, *et al.* 2006. *In vitro* activities of posaconazole, fluconazole, itraconazole, voriconazole, and amphotericin B against a large collection of clinically important molds and yeasts. *Antimicrob. Agents Chemother.* **50**: 2009–2015.
  189. Heng, S.-C., M.A. Slavin, S.C.-A. Chen, *et al.* 2012. Hospital costs, length of stay and mortality attributable to invasive scedosporiosis in haematology patients. *J. Antimicrob. Chemother.* **67**: 2274–2282.
  190. Johnson, M.E., S.K. Katiyar & T.D. Edlind. 2011. New Fks hot spot for acquired echinocandin resistance in *Saccharomyces cerevisiae* and its contribution to intrinsic resistance of *Scedosporium* species. *Antimicrob. Agents Chemother.* **55**: 3774–3781.
  191. Cuenca-Estrella, M., A. Alastruey-Izquierdo, L. Alcazar-Fuoli, *et al.* 2008. *In vitro* activities of 35 double combinations of antifungal agents against *Scedosporium apiospermum* and *Scedosporium prolificans*. *Antimicrob. Agents Chemother.* **52**: 1136–1139.
  192. Moran, C., C.A. Grussemeyer, J.R. Spalding, *et al.* 2009. *Candida albicans* and non-*albicans* bloodstream infections in adult and pediatric patients: comparison of mortality and costs. *Pediatr. Infect. Dis. J.* **28**: 433.
  193. Moran, D.C., M.C.A. Grussemeyer, D.J.R. Spalding, *et al.* 2010. Comparison of costs, length of stay, and mortality associated with *Candida glabrata* and *Candida albicans* bloodstream infections. *Am. J. Infect. Control* **38**: 78–80.
  194. Pound, M.W., M.L. Townsend, V. Dimondi, *et al.* 2011. Overview of treatment options for invasive fungal infections. *Med. Mycol.* **49**: 561–580.
  195. Lee, W.G., J.H. Shin, Y. Uh, *et al.* 2011. First three reported cases of nosocomial fungemia caused by *Candida auris*. *J. Clin. Microbiol.* **49**: 3139–3142.
  196. Urdea, M., L.A. Penny, S.S. Olmsted, *et al.* 2006. Requirements for high impact diagnostics in the developing world. *Nature* **444**(Suppl. 1): 73–79.
  197. Arvanitis, M., T. Anagnostou, B.B. Fuchs, *et al.* 2014. Molecular and nonmolecular diagnostic methods for invasive fungal infections. *Clin. Microbiol. Rev.* **27**: 490.
  198. Chan, K., P.-Y. Wong, P. Yu, *et al.* 2016. A rapid and low-cost PCR thermal cycler for infectious disease diagnostics. *PLoS One* **11**: e0149150.
  199. Xiang, Q., B. Xu, R. Fu, *et al.* 2005. Real time PCR on disposable PDMS chip with a miniaturized thermal cycler. *Biomed. Microdevices* **7**: 273–279.
  200. Wong, G., I. Wong, K. Chan, *et al.* 2015. A rapid and low-cost PCR thermal cycler for low resource settings. *PLoS One* **10**: e0131701.
  201. Gardy, J.L. & N.J. Loman. 2018. Towards a genomics-informed, real-time, global pathogen surveillance system. *Nat. Rev. Genet.* **19**: 9–20.
  202. Jain, M., H.E. Olsen, B. Paten, *et al.* 2016. The Oxford Nanopore MinION: delivery of nanopore sequencing to the genomics community. *Genome Biol.* **17**: 239.
  203. Quick, J., N.J. Loman, S. Duraffour, *et al.* 2016. Real-time, portable genome sequencing for Ebola surveillance. *Nature* **530**: 228–232.
  204. Hoenen, T., A. Groseth, K. Rosenke, *et al.* 2016. Nanopore sequencing as a rapidly deployable Ebola outbreak tool. *Emerg. Infect. Dis.* **22**: 331–334.
  205. Faria, N.R., E.C. Sabino, M.R.T. Nunes, *et al.* 2016. Mobile real-time surveillance of Zika virus in Brazil. *Genome Med.* **8**: 97.
  206. Votintseva, A.A., P. Bradley, L. Pankhurst, *et al.* 2017. Same-day diagnostic and surveillance data for tuberculosis via whole-genome sequencing of direct respiratory samples. *J. Clin. Microbiol.* **55**: 1285–1298.
  207. Pardee, K., A.A. Green, T. Ferrante, *et al.* 2014. Paper-based synthetic gene networks. *Cell* **159**: 940–954.
  208. Pardee, K., A.A. Green, M.K. Takahashi, *et al.* 2016. Rapid, low-cost detection of Zika virus using programmable biomolecular components. *Cell* **165**: 1255–1266.
  209. Gootenberg, J.S., O.O. Abudayyeh, J.W. Lee, *et al.* 2017. Nucleic acid detection with CRISPR-Cas13a/C2c2. *Science* **356**: 438–442.
  210. Bader, O. 2017. Fungal species identification by MALDI-ToF mass spectrometry. In *Human Fungal Pathogen Identification*. T. Lion, Ed.: Vol. 1508. Methods in Molecular Biology. New York: Humana Press.
  211. Bader, O. 2013. MALDI-TOF-MS-based species identification and typing approaches in medical mycology. *Proteomics* **13**: 788–799.
  212. Alanio, A., J.L. Beretti, B. Dauphin, *et al.* 2011. Matrix-assisted laser desorption ionization time-of-flight mass spectrometry for fast and accurate identification of clinically relevant *Aspergillus* species. *Clin. Microbiol. Infect.* **17**: 750–755.
  213. Gautier, M., A.-C. Normand & S. Ranque. 2016. Previously unknown species of *Aspergillus*. *Clin. Microbiol. Infect.* **22**: 662–669.
  214. Vella, A., E. De Carolis, E. Mello, *et al.* 2017. Potential use of MALDI-ToF mass spectrometry for rapid detection of antifungal resistance in the human pathogen *Candida glabrata*. *Sci. Rep.* **7**: 9099.
  215. Vella, A., E. De Carolis, L. Vaccaro, *et al.* 2013. Rapid antifungal susceptibility testing by matrix-assisted laser desorption ionization-time of flight mass spectrometry analysis. *J. Clin. Microbiol.* **51**: 2964–2969.

216. Geddes, J.M.H., D. Croll, M. Caza, *et al.* 2015. Secretome profiling of *Cryptococcus neoformans* reveals regulation of a subset of virulence-associated proteins and potential biomarkers by protein kinase A. *BMC Microbiol.* **15**: 206.
217. Ostrov, N., M. Jimenez, S. Billerbeck, *et al.* 2017. A modular yeast biosensor for low-cost point-of-care pathogen detection. *Sci. Adv.* **3**: e1603221.
218. Perfect, J.R. 2016. "Is there an emerging need for new antifungals?" *Expert Opin. Emerg. Drugs* **21**: 129–131.
219. Krysan, D.J. 2017. The unmet clinical need of novel antifungal drugs. *Virulence* **8**: 135–137.
220. Garvey, E.P., W.J. Hoekstra, R.J. Schotzinger, *et al.* 2015. Efficacy of the clinical agent VT-1161 against fluconazole-sensitive and -resistant *Candida albicans* in a murine model of vaginal candidiasis. *Antimicrob. Agents Chemother.* **59**: 5567–5573.
221. Lockhart, S.R., A.W. Fothergill, N. Iqbal, *et al.* 2016. The investigational fungal Cyp51 inhibitor VT-1129 demonstrates potent *in vitro* activity against *Cryptococcus neoformans* and *Cryptococcus gattii*. *Antimicrob. Agents Chemother.* **60**: 2528–2531.
222. Wiederhold, N.P., H.P. Patterson, B.H. Tran, *et al.* 2018. Fungal-specific Cyp51 inhibitor VT-1598 demonstrates *in vitro* activity against *Candida* and *Cryptococcus* species, endemic fungi, including *Coccidioides* species, *Aspergillus* species and *Rhizopus arrhizus*. *J. Antimicrob. Chemother.* **73**: 404–408.
223. Ong, V., G. Hough, M. Schlosser, *et al.* 2016. Preclinical evaluation of the stability, safety, and efficacy of CD101, a novel echinocandin. *Antimicrob. Agents Chemother.* **60**: 6872–6879.
224. James, K.D., C.P. Laudeman, N.B. Malkar, *et al.* 2017. Structure–activity relationships of a series of echinocandins and the discovery of CD101, a highly stable and soluble echinocandin with distinctive pharmacokinetic properties. *Antimicrob. Agents Chemother.* **61**: e01541-16.
225. Pfaller, M.A., S.A. Messer, P.R. Rhomberg, *et al.* 2017. CD101, a long-acting echinocandin, and comparator antifungal agents tested against a global collection of invasive fungal isolates in the SENTRY 2015 Antifungal Surveillance Program. *Int. J. Antimicrob. Agents* **50**: 352–358.
226. Nakamura, I., S. Yoshimura, T. Masaki, *et al.* 2017. ASP2397: a novel antifungal agent produced by *Acremonium persicinum* MF-347833. *J. Antibiot.* **70**: 45–51.
227. Rhein, J., B.M. Morawski, K.H. Hullsiek, *et al.* 2016. Efficacy of adjunctive sertraline for the treatment of HIV-associated cryptococcal meningitis: an open-label dose-ranging study. *Lancet Infect. Dis.* **16**: 809–818.
228. Geddes, J.M.H., M. Caza, D. Croll, *et al.* 2016. Analysis of the protein kinase A-regulated proteome of *Cryptococcus neoformans* identifies a role for the ubiquitin-proteasome pathway in capsule formation. *mBio* **7**: e01862-15.
229. Watanabe, N.-A., M. Miyazaki, T. Horii, *et al.* 2012. E1210, a new broad-spectrum antifungal, suppresses *Candida albicans* hyphal growth through inhibition of glycosylphosphatidylinositol biosynthesis. *Antimicrob. Agents Chemother.* **56**: 960–971.
230. Hata, K., T. Horii, M. Miyazaki, *et al.* 2011. Efficacy of oral E1210, a new broad-spectrum antifungal with a novel mechanism of action, in murine models of candidiasis, aspergillosis, and fusariosis. *Antimicrob. Agents Chemother.* **55**: 4543–4551.
231. Schell, W.A., C. Giamberardino, K.J. Shaw, *et al.* 2017. Efficacy of oral APX001 in a murine model of cryptococcal meningitis. *Open Forum Infect. Dis.* **4**: S478.
232. Robbins, N., M. Spitzer, W. Wang, *et al.* 2016. Discovery of ibomycin, a complex macrolactone that exerts antifungal activity by impeding endocytic trafficking and membran function. *Cell Chem. Biol.* **23**: 1383–1394.
233. Tebbets, B., D. Stewart, S. Lawry, *et al.* 2012. Identification and characterization of antifungal compounds using a *Saccharomyces cerevisiae* reporter bioassay. *PLoS One* **7**: e36021.
234. Roemer, T., D. Xu, S.B. Singh, *et al.* 2011. Confronting the challenges of natural product-based antifungal discovery. *Chem. Biol.* **18**: 148–164.
235. Breger, J., B.B. Fuchs, G. Aperis, *et al.* 2007. Antifungal chemical compounds identified using a *C. elegans* pathogenicity assay. *PLoS Pathog.* **3**: e18.
236. Cowen, L.E., S.D. Singh, J.R. Köhler, *et al.* 2009. Harnessing Hsp90 function as a powerful, broadly effective therapeutic strategy for fungal infectious disease. *Proc. Natl. Acad. Sci. USA* **106**: 2818–2823.
237. Cowen, L.E. 2013. The fungal Achilles' heel: targeting Hsp90 to cripple fungal pathogens. *Curr. Opin. Microbiol.* **16**: 377–384.
238. Pacht, J., P. Svoboda, F. Jacobs, *et al.* 2006. A randomized, blinded, multicenter trial of lipid-associated amphotericin B alone versus in combination with an antibody-based inhibitor of heat shock protein 90 in patients with invasive candidiasis. *Clin. Infect. Dis.* **42**: 1404–1413.
239. Juvvadi, P.R., S.C. Lee, J. Heitman, *et al.* 2017. Calcineurin in fungal virulence and drug resistance: prospects for harnessing targeted inhibition of calcineurin for an antifungal therapeutic approach. *Virulence* **8**: 186–197.
240. Blankenship, J.R., W.J. Steinbach, J.R. Perfect, *et al.* 2003. Teaching old drugs new tricks: reincarnating immunosuppressants as antifungal drugs. *Curr. Opin. Investig. Drugs* **4**: 192–199.
241. Nambu, M., J.A. Covel, M. Kapoor, *et al.* 2017. A calcineurin antifungal strategy with analogs of FK506. *Bioorg. Med. Chem. Lett.* **27**: 2465–2471.
242. Barrera, A., A. Alastruey-Izquierdo, M.J. Martín, *et al.* 2014. Analysis of the protein domain and domain architecture content in fungi and its application in the search of new antifungal targets. *PLoS Comput. Biol.* **10**: e1003733.
243. Lehár, J., A.S. Krueger, W. Avery, *et al.* 2009. Synergistic drug combinations tend to improve therapeutically relevant selectivity. *Nat. Biotechnol.* **27**: 659–666.
244. Spitzer, M., N. Robbins & G.D. Wright. 2017. Combinatorial strategies for combating invasive fungal infections. *Virulence* **8**: 169–185.
245. Butts, A., G.E. Palmer & P.D. Rogers. 2017. Antifungal adjuvants: preserving and extending the antifungal arsenal. *Virulence* **8**: 198–210.

246. Lazar, V., G. Pal Singh, R. Spohn, *et al.* 2014. Bacterial evolution of antibiotic hypersensitivity. *Mol. Syst. Biol.* **9**: 700.
247. Imamovic, L. & M.O.A. Sommer. 2013. Use of collateral sensitivity networks to design drug cycling protocols that avoid resistance development. *Sci. Transl. Med.* **5**: 204ra132.
248. Dhawan, A., D. Nichol, F. Kinose, *et al.* 2017. Collateral sensitivity networks reveal evolutionary instability and novel treatment strategies in ALK mutated non-small cell lung cancer. *Sci. Rep.* **7**: 1232.
249. Pluchino, K.M., M.D. Hall, A.S. Goldsborough, *et al.* 2012. Collateral sensitivity as a strategy against cancer multidrug resistance. *Drug Resist. Updat.* **15**: 98–105.
250. Hill, J.A. & L.E. Cowen. 2015. Using combination therapy to thwart drug resistance. *Future Microbiol.* **10**: 1719–1726.
251. Rodriguez, M.M., E. Calvo, C. Serena, *et al.* 2009. Effects of double and triple combinations of antifungal drugs in a murine model of disseminated infection by *Scedosporium prolificans*. *Antimicrob. Agents Chemother.* **53**: 2153–2155.
252. Biswas, C., T.C. Sorrell, J.T. Djordjevic, *et al.* 2013. *In vitro* activity of miltefosine as a single agent and in combination with voriconazole or posaconazole against uncommon filamentous fungal pathogens. *J. Antimicrob. Chemother.* **68**: 2842–2846.
253. Pfaller, M.A., P.R. Rhomberg, S.A. Messer, *et al.* 2015. *In vitro* activity of a Hos2 deacetylase inhibitor, MGCD290, in combination with echinocandins against echinocandin-resistant *Candida* species. *Diagn. Microbiol. Infect. Dis.* **81**: 259–263.
254. Pfaller, M.A., S.A. Messer, N. Georgopapadakou, *et al.* 2009. Activity of MGCD290, a Hos2 histone deacetylase inhibitor, in combination with azole antifungals against opportunistic fungal pathogens. *J. Clin. Microbiol.* **47**: 3797–3804.
255. Fiori, A. & P. Van Dijck. 2012. Potent synergistic effect of doxycycline with fluconazole against *Candida albicans* is mediated by interference with iron homeostasis. *Antimicrob. Agents Chemother.* **56**: 3785–3796.
256. Shi, W., Z. Chen, X. Chen, *et al.* 2010. The combination of minocycline and fluconazole causes synergistic growth inhibition against *Candida albicans*: an *in vitro* interaction of antifungal and antibacterial agents. *FEMS Yeast Res.* **10**: 885–893.
257. Liu, S., L. Yue, W. Gu, *et al.* 2016. Synergistic effect of fluconazole and calcium channel blockers against resistant *Candida albicans*. *PLoS One* **11**: e0150859.
258. Shekhar-Guturja, T., G.M.K.B. Gunaherath, E.M.K. Wijeratne, *et al.* 2016. Dual action antifungal small molecule modulates multidrug efflux and TOR signaling. *Nat. Chem. Biol.* **12**: 867–875.
259. Spitzer, M., E. Griffiths, K.M. Blakely, *et al.* 2011. Cross-species discovery of syncretic drug combinations that potentiate the antifungal fluconazole. *Mol. Syst. Biol.* **7**: 499.
260. Robbins, N., M. Spitzer, T. Yu, *et al.* 2015. An antifungal combination matrix identifies a rich pool of adjuvant molecules that enhance drug activity against diverse fungal pathogens. *Cell Rep.* **13**: 1481–1492.
261. Zhang, L., K. Yan, Y. Zhang, *et al.* 2007. High-throughput synergy screening identifies microbial metabolites as combination agents for the treatment of fungal infections. *Proc. Natl. Acad. Sci. USA* **104**: 4606–4611.
262. Brown, J.C.S., J. Nelson, B. VanderSluis, *et al.* 2014. Unraveling the biology of a fungal meningitis pathogen using chemical genetics. *Cell* **159**: 1168–1187.
263. Han, K., E.E. Jeng, G.T. Hess, *et al.* 2017. Synergistic drug combinations for cancer identified in a CRISPR screen for pairwise genetic interactions. *Nat. Biotechnol.* **35**: 463–474.
264. O'Neil, N.J., M.L. Bailey & P. Hieter. 2017. Synthetic lethality and cancer. *Nat. Rev. Genet.* **18**: 613–623.
265. Shapiro, R.S., A. Chavez, C.B.M. Porter, *et al.* 2018. A CRISPR-Cas9-based gene drive platform for genetic interaction analysis in *Candida albicans*. *Nat. Microbiol.* **3**: 73–82.
266. Deo, S.S. & D.J. Gottlieb. 2015. Adoptive T-cell therapy for fungal infections in haematology patients. *Clin. Transl. Immunology* **4**: e40.
267. Loreto, E.S., J.S.M. Tondolo, S.H. Alves & J.M. Santurio. 2017. Immunotherapy for fungal infections. In *Immunotherapy—Myths, Reality, Ideas, Future*. K. Metodiev, Ed.: 291–322. Rijeka, Croatia: InTech.
268. Pappas, P.G., B. Bustamante, E. Ticona, *et al.* 2004. Recombinant interferon-gamma 1b as adjunctive therapy for AIDS-related acute cryptococcal meningitis. *J. Infect. Dis.* **189**: 2185–2191.
269. Shao, C., J. Qu, L. He, *et al.* 2005. Transient overexpression of gamma interferon promotes *Aspergillus* clearance in invasive pulmonary aspergillosis. *Clin. Exp. Immunol.* **142**: 233–241.
270. Delsing, C.E., M.S. Gresnigt, J. Leentjens, *et al.* 2014. Interferon-gamma as adjunctive immunotherapy for invasive fungal infections: a case series. *BMC Infect. Dis.* **14**: 166.
271. Armstrong-James, D., I.A. Teo, S. Shrivastava, *et al.* 2010. Exogenous interferon- $\gamma$  immunotherapy for invasive fungal infections in kidney transplant patients. *Am. J. Transplant.* **10**: 1796–1803.
272. Jarvis, J.N., G. Meintjes, K. Rebe, *et al.* 2012. Adjunctive interferon- $\gamma$  immunotherapy for the treatment of HIV-associated cryptococcal meningitis: a randomized controlled trial. *AIDS* **26**: 1105–1113.
273. Goldman, C., M.J. Akiyama, J. Torres, *et al.* 2016. *Scedosporium apiospermum* infections and the role of combination antifungal therapy and GM-CSF: a case report and review of the literature. *Med. Mycol. Case Rep.* **11**: 40–43.
274. Larsen, R.A., P.G. Pappas, J. Perfect, *et al.* 2005. Phase I evaluation of the safety and pharmacokinetics of murine-derived anticryptococcal antibody 18B7 in subjects with treated cryptococcal meningitis. *Antimicrob. Agents Chemother.* **49**: 952–958.
275. Karwa, R. & K.A. Wargo. 2009. Efungumab: a novel agent in the treatment of invasive candidiasis. *Ann. Pharmacother.* **43**: 1818–1823.
276. Rachini, A., D. Pietrella, P. Lupo, *et al.* 2007. An anti- $\beta$ -glucan monoclonal antibody inhibits growth and capsule formation of *Cryptococcus neoformans* *in vitro* and exerts therapeutic, anticryptococcal activity *in vivo*. *Infect. Immun.* **75**: 5085–5094.
277. Bugli, F., M. Cacaci, C. Martini, *et al.* 2013. Human monoclonal antibody-based therapy in the treatment of invasive candidiasis. *Clin. Dev. Immunol.* **2013**: 403121.

278. Rollin-Pinheiro, R., L.C. Liporagi-Lopes, J.V. de Meirelles, *et al.* 2014. Characterization of *Scedosporium apiospermum* glucosylceramides and their involvement in fungal development and macrophage functions. *PLoS One* **9**: e98149.
279. Bryan, R.A., A.J. Guimaraes, S. Hopcraft, *et al.* 2011. Toward developing a universal treatment for fungal disease using radioimmunotherapy targeting common fungal antigens. *Mycopathologia* **173**: 463–471.
280. Kumaresan, P.R., P.R. Manuri, N.D. Albert, *et al.* 2014. Bio-engineering T cells to target carbohydrate to treat opportunistic fungal infection. *Proc. Natl. Acad. Sci. USA* **111**: 10660–10665.
281. Zhao, X., Y. Guo, C. Jiang, *et al.* 2017. JNK1 negatively controls antifungal innate immunity by suppressing CD23 expression. *Nat. Med.* **23**: 337–346.
282. Casadevall, A. & L.-A. Pirofski. 2005. Feasibility and prospects for a vaccine to prevent cryptococcosis. *Med. Mycol.* **43**: 667–680.
283. Cutler, J.E., G.S. Deepe, Jr. & B.S. Klein. 2007. Advances in combating fungal diseases: vaccines on the threshold. *Nat. Rev. Microbiol.* **5**: 13–28.
284. Leopold Wager, C.M. & F.L. Wormley, Jr. 2015. Is development of a vaccine against *Cryptococcus neoformans* feasible? *PLoS Pathog.* **11**: e1004843.
285. Caballero Van Dyke, M.C. & F.L. Wormley. 2017. A call to arms: quest for a cryptococcal vaccine. *Trends Microbiol.* **26**: 436–446.
286. Chow, S.-K. & A. Casadevall. 2011. Evaluation of *Cryptococcus neoformans* galactoxylomannan–protein conjugate as vaccine candidate against murine cryptococcosis. *Vaccine* **29**: 1891–1898.
287. Upadhyay, R., W.C. Lam, B. Maybruck, *et al.* 2016. Induction of protective immunity to cryptococcal infection in mice by a heat-killed, chitosan-deficient strain of *Cryptococcus neoformans*. *mBio* **7**: pii: e00547–16.
288. Mor, V., A.M. Farnoud, A. Singh, *et al.* 2016. Glucosylceramide administration as a vaccination strategy in mouse models of cryptococcosis. *PLoS One* **11**: e0153853.
289. Wormley, F.L., Jr., J.R. Perfect, C. Steele, *et al.* 2007. Protection against cryptococcosis by using a murine gamma interferon-producing *Cryptococcus neoformans* strain. *Infect. Immun.* **75**: 1453–1462.
290. Xin, H., S. Dziadek, D.R. Bundle, *et al.* 2008. Synthetic glycopeptide vaccines combining beta-mannan and peptide epitopes induce protection against candidiasis. *Proc. Natl. Acad. Sci. USA* **105**: 13526–13531.
291. Xin, H., J. Cartmell, J.J. Bailey, *et al.* 2012. Self-adjuvanting glycopeptide conjugate vaccine against disseminated candidiasis. *PLoS One* **7**: e35106.
292. Clatworthy, A.E., E. Pierson & D.T. Hung. 2007. Targeting virulence: a new paradigm for antimicrobial therapy. *Nat. Chem. Biol.* **3**: 541–548.
293. Gauwery, K., C. Borelli & H.C. Kortling. 2009. Targeting virulence: a new paradigm for antifungals. *Drug Discov. Today* **14**: 214–222.
294. Zhang, M., W. Chang, H. Shi, *et al.* 2017. Biatrisporin D displays anti-virulence activity through decreasing the intracellular cAMP levels. *Toxicol. Appl. Pharmacol.* **322**: 104–112.
295. Romo, J.A., C.G. Pierce, A.K. Chaturvedi, *et al.* 2017. Development of anti-virulence approaches for candidiasis via a novel series of small-molecule inhibitors of *Candida albicans* filamentation. *mBio* **8**: e01991–17.
296. Vila, T., J.A. Romo, C.G. Pierce, *et al.* 2017. Targeting *Candida albicans* filamentation for antifungal drug development. *Virulence* **8**: 150–158.
297. Fazly, A., C. Jain, A.C. Dehner, *et al.* 2013. Chemical screening identifies filastatin, a small molecule inhibitor of *Candida albicans* adhesion, morphogenesis, and pathogenesis. *Proc. Natl. Acad. Sci. USA* **110**: 13594–13599.
298. Shareck, J. & P. Belhumeur. 2011. Modulation of morphogenesis in *Candida albicans* by various small molecules. *Eukaryot. Cell* **10**: 1004–1012.
299. Krasowska, A., A. Murzyn, A. Dyjankiewicz, *et al.* 2009. The antagonistic effect of *Saccharomyces boulardii* on *Candida albicans* filamentation, adhesion and biofilm formation. *FEMS Yeast Res.* **9**: 1312–1321.
300. Murzyn, A., A. Krasowska, P. Stefanowicz, *et al.* 2010. Capric acid secreted by *S. boulardii* inhibits *C. albicans* filamentous growth, adhesion and biofilm formation. *PLoS One* **5**: e12050.
301. Mayer, F.L. & J.W. Kronstad. 2017. Disarming fungal pathogens: *Bacillus safensis* inhibits virulence factor production and biofilm formation by *Cryptococcus neoformans* and *Candida albicans*. *mBio* **8**: e01537–17.
302. Niimi, K., D.R.K. Harding, A.R. Holmes, *et al.* 2012. Specific interactions between the *Candida albicans* ABC transporter Cdr1p ectodomain and a D-octapeptide derivative inhibitor. *Mol. Microbiol.* **85**: 747–767.
303. Niimi, K., D.R.K. Harding, R. Parshot, *et al.* 2004. Chemosensitization of fluconazole resistance in *Saccharomyces cerevisiae* and pathogenic fungi by a D-octapeptide derivative. *Antimicrob. Agents Chemother.* **48**: 1256–1271.
304. Lee, M.D., J.L. Galazzo, A.L. Staley, *et al.* 2001. Microbial fermentation-derived inhibitors of efflux-pump-mediated drug resistance. *Farmaco* **56**: 81–85.
305. Keniya, M.V., E. Fleischer, A. Klinger, *et al.* 2015. Inhibitors of the *Candida albicans* major facilitator superfamily transporter Mdr1p responsible for fluconazole resistance. *PLoS One* **10**: e0126350.
306. Holmes, A.R., M.V. Keniya, I. Ivnitski-Steele, *et al.* 2012. The monoamine oxidase A inhibitor clorgyline is a broad-spectrum inhibitor of fungal ABC and MFS transporter efflux pump activities which reverses the azole resistance of *Candida albicans* and *Candida glabrata* clinical isolates. *Antimicrob. Agents Chemother.* **56**: 1508–1515.
307. Nishikawa, J.L., A. Boeszoermyenyi, L.A. Vale-Silva, *et al.* 2016. Inhibiting fungal multidrug resistance by disrupting an activator–mediator interaction. *Nature* **530**: 485–489.
308. Editorial. 2017. Stop neglecting fungi. *Nat. Microbiol.* **2**: 17120.
309. Denning, D.W. 2017. Calling upon all public health mycologists: to accompany the country burden papers from 14 countries. *Eur. J. Clin. Microbiol. Infect. Dis.* **36**: 923–924.