

STA 522, Spring 2021
Introduction to Theoretical Statistics II

Lecture 13

Department of Biostatistics
University at Buffalo

26 April, 2021

AGENDA

- ▶ Asymptotic Evaluations

Consistent Estimators

Definition: A sequence of estimators $W_n = W_n(\underline{X})$ is a (weakly) **consistent sequence of estimators** of the parameter θ if and only if for every $\theta \in \Theta$, $W_n \xrightarrow{P} \theta$ i.e., for every $\varepsilon > 0$,

$$\lim_{n \rightarrow \infty} P_{\theta} (|W_n - \theta| \geq \varepsilon) = 0,$$

or, equivalently,

$$\lim_{n \rightarrow \infty} P_{\theta} (|W_n - \theta| < \varepsilon) = 1.$$

Definition: A sequence of estimators $W_n = W_n(\underline{X})$ is a **strongly consistent sequence of estimators** of the parameter θ if and only if $W_n \xrightarrow{a.s.} \theta$, for every $\theta \in \Theta$, i.e.,

$$P \left(\lim_{n \rightarrow \infty} W_n = \theta \right) = 1.$$

Example: Let $X_1, X_2, \dots, X_n \sim \text{iid } N(\theta, 1)$. Then \bar{X}_n is a consistent sequence of estimators of θ .

Recall that $\bar{X}_n \sim N(\theta, 1/n)$. So,

$$\begin{aligned} P_\theta(|\bar{X}_n - \theta| < \varepsilon) &= \int_{\theta-\varepsilon}^{\theta+\varepsilon} \left(\frac{n}{2\pi}\right)^{1/2} e^{-(n/2)(\bar{x}_n - \theta)^2} d\bar{x}_n \\ &= \int_{-\varepsilon}^{\varepsilon} \left(\frac{n}{2\pi}\right)^{1/2} e^{-(n/2)y^2} dy && (y = \bar{x}_n - \theta) \\ &= \int_{-\varepsilon\sqrt{n}}^{\varepsilon\sqrt{n}} \left(\frac{1}{2\pi}\right)^{1/2} e^{-(1/2)t^2} dt && (t = y\sqrt{n}) \\ &= P(-\varepsilon\sqrt{n} < Z < \varepsilon\sqrt{n}) && (Z \sim N(0, 1)) \\ &\rightarrow 1 \quad \text{as } n \rightarrow \infty \end{aligned}$$

This shows $\bar{X}_n \xrightarrow{P} \theta$.

How to Verify Consistency for a Sequence of Estimators

Theorem 10.1.3

Let W_n be a sequence of estimators of a parameter θ satisfying

(a) $\lim_{n \rightarrow \infty} \text{Var}_{\theta}(W_n) = 0$ and

(b) $\lim_{n \rightarrow \infty} \text{Bias}_{\theta}(W_n) = 0$

for every $\theta \in \Theta$.

Then W_n is a consistent sequence of estimators of θ .

Example (contd.): $X_1, X_2, \dots, X_n \sim \text{iid } N(\theta, 1)$, consider the estimator \bar{X}_n of θ .

We have $E_{\theta}(\bar{X}_n) = \theta$ for all θ , i.e., $\text{Bias}_{\theta}(\bar{X}_n) = 0$, and $\text{Var}_{\theta}(\bar{X}_n) = \frac{1}{n} \rightarrow 0$. Hence, from the above theorem, it follows that \bar{X}_n is consistent for θ .

Theorem 10.1.6 (Consistency of MLEs)

Let $X_1, X_2, \dots, X_n \sim \text{iid } f(x | \theta)$.

Let $L(\theta | \underline{x}) = \prod_{i=1}^n f(x_i | \theta)$ be the likelihood function.

Let $\hat{\theta}$ denote the MLE of θ .

Let $\tau(\theta)$ be a continuous function of θ .

Under certain regularity conditions on $f(x | \theta)$ (see Miscellanea 10.6.2; these hold, e.g., for the regular exponential family) and, hence, $L(\theta | \underline{x})$, for every $\varepsilon > 0$ and every $\theta \in \Theta$,

$$\lim_{n \rightarrow \infty} P_{\theta} \left(|\tau(\hat{\theta}) - \tau(\theta)| \geq \varepsilon \right) = 0.$$

In other words, $\tau(\hat{\theta})$ is a consistent estimator of $\tau(\theta)$, i.e., MLEs are weakly consistent (converge in probability).

Asymptotic Variance & Efficiency

Definition: For an estimator T_n , suppose that $k_n(T_n - \tau(\theta)) \xrightarrow{d} N(0, \sigma^2)$, where $\{k_n\}$ is a sequence of constants. The parameter σ^2 is called the **asymptotic variance** or **variance of the limit distribution** of T_n .

Definition: A sequence of estimators W_n is **asymptotically efficient** for a parameter $\tau(\theta)$ if and only if

$$\sqrt{n}(W_n - \tau(\theta)) \xrightarrow{d} W \sim N(0, v(\theta))$$

and

$$v(\theta) = \frac{[\tau'(\theta)]^2}{I_1(\theta)} = \frac{[\tau'(\theta)]^2}{E_\theta \left[\left(\frac{\partial}{\partial \theta} \log f(X | \theta) \right)^2 \right]};$$

that is, the asymptotic variance of W_n (almost) achieves the Cramér–Rao Lower Bound. $I_1(\theta)$ is the *unit* Fisher information (i.e., Fisher information computed for *one* random observation).

Notes on Asymptotic Efficiency

- (a) There is no n in the denominator of $v(\theta)$, as it got moved to the left-hand side.
- (b) This definition does not mean

$$W_n \xrightarrow{d} N(\tau(\theta), \text{CRLB}),$$

since

$$\text{CRLB} = \frac{[\tau'(\theta)]^2}{n E_{\theta} \left[\left(\frac{\partial}{\partial \theta} \log f(X | \theta) \right)^2 \right]}.$$

$v(\theta)$ is written without the n because we are letting $n \rightarrow \infty$

- (c) In practical terms, what this does mean is that W_n is approximately normally distributed with mean $\tau(\theta)$ and variance equal to the CRLB, evaluated at $\hat{\theta}$. This is often denoted as $W_n \stackrel{a}{\sim} N(\tau(\theta), \text{CRLB}(\hat{\theta}))$.

Asymptotic distribution of a function of a sequence of random variables

Theorem 5.5.24 (Delta Method)

Suppose Y_n is a sequence of random variables that satisfies

$\sqrt{n}(Y_n - \theta) \xrightarrow{d} N(0, \sigma^2)$. For a given function g and a specific value of θ , suppose that $g'(\theta)$ exists and is not 0. Then

$$\sqrt{n}[g(Y_n) - g(\theta)] \xrightarrow{d} N(0, \sigma^2[g'(\theta)]^2)$$

Heuristic Proof: The Taylor expansion of $g(Y_n)$ around $Y_n = \theta$ is

$$g(Y_n) = g(\theta) + g'(\theta)(Y_n - \theta) + R_n$$

where R_n is such that $\sqrt{n}R_n \xrightarrow{P} 0$ as $n \rightarrow \infty$. Therefore,

$$\sqrt{n}[g(Y_n) - g(\theta)] = g'(\theta)\sqrt{n}(Y_n - \theta) + \sqrt{n}R_n \xrightarrow{d} N(0, \sigma^2[g'(\theta)]^2)$$

by Slutsky's theorem.

Theorem 10.1.12 (Asymptotic Efficiency of MLEs)

Let $X_1, X_2, \dots, X_n \sim \text{iid } f(x | \theta)$. Let $\hat{\theta}$ denote the MLE of θ . Under certain regularity conditions (see Miscellanea 10.6.2) on $f(x | \theta)$ and, hence, on $L(\theta | \underline{x})$,

$$\sqrt{n}(\hat{\theta} - \theta) \xrightarrow{d} N\left(0, \frac{1}{I_1(\theta)}\right),$$

where $I_1(\theta)$ is the unit Fisher information.

Corollary

Suppose $\tau(\theta)$ is a differentiable function of θ . If $\hat{\theta}$ denotes the MLE of θ , then $\tau(\hat{\theta})$ is the MLE of $\tau(\theta)$. The asymptotic distribution of $\tau(\hat{\theta})$ is obtained using the delta method:

$$\sqrt{n}(\tau(\hat{\theta}) - \tau(\theta)) \xrightarrow{d} N(0, v(\theta))$$

In other words, $\tau(\hat{\theta})$ is a consistent and asymptotically efficient estimator of $\tau(\theta)$.

Example: Suppose $X_1, X_2, \dots, X_n \sim \text{iid Bernoulli}(p)$. The MLE of p is $\hat{p} = \frac{1}{n} \sum_{i=1}^n X_i$. We have seen an asymptotic distribution of \bar{X}_n via de Moivre-Laplace CLT.

To apply the asymptotic normality of the MLE, note that here $I_1(\theta) = \frac{1}{p(1-p)}$.

Hence

$$\sqrt{n} (\hat{p} - p) \xrightarrow{d} N(0, p(1-p)) \implies \frac{\sqrt{n} (\hat{p} - p)}{\sqrt{p(1-p)}} \xrightarrow{d} N(0, 1)$$

The standard deviation $\sqrt{p(1-p)}$ has MLE $\sqrt{\hat{p}(1-\hat{p})}$, and due to consistency of \hat{p}

$$\sqrt{\hat{p}(1-\hat{p})} \xrightarrow{P} \sqrt{p(1-p)}$$

Therefore using Slutsky's theorem

$$\frac{\sqrt{n} (\hat{p} - p)}{\sqrt{\hat{p}(1-\hat{p})}} \xrightarrow{d} N(0, 1)$$

In practical terms this means $\hat{p} \overset{a}{\sim} N\left(p, \frac{p(1-p)}{n}\right)$.

Asymptotic Relative Efficiency (ARE)

If two estimators W_n and V_n satisfy

$$\sqrt{n} [W_n - \tau(\theta)] \xrightarrow{d} N(0, \sigma_W^2)$$

$$\sqrt{n} [V_n - \tau(\theta)] \xrightarrow{d} N(0, \sigma_V^2)$$

then the asymptotic relative efficiency (ARE) of V_n with respect to W_n is

$$\text{ARE}(V_n, W_n) = \frac{\sigma_W^2}{\sigma_V^2}$$

Example (AREs of Poisson Estimators): Suppose that X_1, X_2, \dots are iid $\text{Poisson}(\lambda)$, and we are interested in estimating $\tau(\lambda) = P_\lambda(X = 0) = e^{-\lambda}$. Consider two estimators of λ , viz.,

(a) $V_n = \frac{1}{n} \sum_{i=1}^n I(X_i = 0) = \frac{1}{n} Y_i =: \bar{Y}_n$ and

(b) $W_n = \tau(\bar{X}) = e^{-\bar{X}_n}$. Note that \bar{X}_n is the MLE of λ .

and find the ARE of V_n relative to W_n .

For (a), note that $Y_i = I(X_i = 0) \sim \text{iid Binomial}(1, \tau(\lambda))$. Therefore,

$$E_\lambda(V_n) = e^{-\lambda} \quad \text{and} \quad \text{Var}_\lambda(V_n) = \frac{e^{-\lambda}(1 - e^{-\lambda})}{n}$$

For (b), using delta method approximation we have (HW)

$$E_\lambda(W_n) \approx e^{-\lambda} \quad \text{and} \quad \text{Var}_\lambda(W_n) = \frac{\lambda e^{-2\lambda}}{n}$$

Since

$$\begin{aligned}\sqrt{n} [V_n - e^{-\lambda}] &\xrightarrow{d} N(0, e^{-\lambda}(1 - e^{-\lambda})) \\ \sqrt{n} [W_n - e^{-\lambda}] &\xrightarrow{d} N(0, \lambda e^{-2\lambda})\end{aligned}$$

Therefore

$$\text{ARE}(V_n, W_n) = \frac{\lambda e^{-2\lambda}}{e^{-\lambda}(1 - e^{-\lambda})} = \frac{\lambda}{1 - e^{-\lambda}}$$

The ARE is strictly decreasing with a maximum of 1 attained at $\lambda = 0$.

Note that the UMVUE for $e^{-\lambda}$ is

$$U_n = \left(\frac{n-1}{n}\right)^{\sum_{i=1}^n X_i} = \left[\left(1 - \frac{1}{n}\right)^n\right]^{\bar{X}_n} \approx e^{-\bar{X}_n} = V_n \text{ for large } n.$$

Asymptotic Distribution of LRT

Suppose $X_1, X_2, \dots, X_n \sim \text{iid poisson}(\lambda)$, and we want to construct a level α test of

$$H_0 : \lambda = \lambda_0$$

$$H_1 : \lambda \neq \lambda_0.$$

A level α test is obtained using rejection region

$$R = \{\underline{x} : -2 \log \lambda(\underline{x}) > \chi_{1,\alpha}^2\},$$

where $\chi_{1,\alpha}^2$ is the χ_1^2 value with area α to its right.

Example (Poisson Testing): Suppose that X_1, X_2, \dots are iid $\text{Poisson}(\lambda)$, and we are interested in testing $H_0 : \lambda = \lambda_0$ vs. $H_1 : \lambda \neq \lambda_0$.

We have

$$-2 \log \lambda(\underline{x}) = -2 \log \left(\frac{e^{-n\lambda_0} \lambda_0^{\sum_{i=1}^n x_i}}{e^{-n\hat{\lambda}} \hat{\lambda}^{\sum_{i=1}^n x_i}} \right) = 2n \left[(\lambda_0 - \hat{\lambda}) - \hat{\lambda} \log(\lambda_0/\hat{\lambda}) \right]$$

where $\hat{\lambda} = \bar{x}$ is the MLE of λ .

The asymptotic theory based test would be to reject H_0 at level α if $-2 \log \lambda(\underline{x}) > \lambda_{1,\alpha}$

Asymptotic normality based tests

Suppose X_1, X_2, \dots, X_n is a random sample from some population $f_\theta(x)$.

Wald test

Let $\hat{\theta}_n = \hat{\theta}_n(X_1, X_2, \dots, X_n)$ be the MLE of θ . Then using the asymptotic normality of $\hat{\theta}_n$ (holds under certain regularity conditions):

$$\sqrt{n}(\hat{\theta}_n - \theta) \xrightarrow{d} N\left(0, \frac{1}{I_1(\theta)}\right)$$

one can perform tests of hypotheses about the real valued parameter θ .

Score test

The *score statistic* is defined as

$$S(\theta) = \frac{\partial}{\partial \theta} \log f(\underline{X} \mid \theta) = \frac{\partial}{\partial \theta} \log L(\theta \mid \underline{X})$$

We know that $E_{\theta}(S(\theta)) = 0$ for all θ . Furthermore

$$\text{Var}_{\theta}(S(\theta)) = E_{\theta} \left[\left(\frac{\partial}{\partial \theta} \log L(\theta \mid \underline{X}) \right)^2 \right] = -E_{\theta} \left(\frac{\partial^2}{\partial \theta^2} \log L(\theta \mid \underline{X}) \right) = I_n(\theta)$$

where $I_1(\theta)$ is the Fisher information obtained from one random observations. Tests of hypothesis such as $H_0 : \theta = \theta_0$ vs. $H_1 : \theta \neq \theta_0$ can be performed using the asymptotic normality of the score statistic:

$$\frac{S(\theta)}{\sqrt{I_n(\theta)}} \xrightarrow{d} N(0, 1)$$

Homework

- ▶ Read p. 467 – 481, 488 – 495.
- ▶ Exercises: TBA.