## **Operon Promoter Landscape**

Operon	Strand	Operon start	Operon end
ssrS-fau	+	3054005	3054811






Figure 1: **Promoter activity in rich media (LB) surrounding query operon**. 17,767 previously reported TSSs were evaluated by measuring the promoter activity (right Y-axis) of the 150 bp surrounding the TSS (-120 to +30) to determine which were active or inactive. The genome-wide promoter activity (left Y-axis) was determined by measuring expression of over 300,000 genomic fragments spanning the *E. coli* genome and averaging promoter activity at all nucleotide positions in a strand-specific fashion. Genome coordinates corresponds to *E. coli* genome version U00096.2.

## TSS Summary

TSS name	TSS position	Strand	TSS activity	Category
TSS_11503_regulondb	3053509	+	1.5261833	active
TSS_11504_regulondb	3053573	+	4.2830854	active
TSS_11506_regulondb	3053779	+	2.1626631	active
TSS_11508_storz_regulondb	3053996	+	13.5773967	active
TSS_11507_regulondb	3053921	+	0.7810445	inactive

## TSS Scanning Mutagenesis




Figure 2: **(Left)** Mutagenesis profile(s) of active TSS(s). Sequences surrounding active TSSs were systematically mutated to identify regions controlling expression. Bar height indicates the relative change in promoter activity as a result of scrambling nucleotides within 10 bp regions at 5 bp intervals spanning the promoter. Bar color identifies the region as a putative activator (yellow) or repressor (purple). **(Right)** Dashed line indicates the expression of the indicated TSS relative to all tested TSS sequences. The distributions of expression is shown for all tested TSSs (black), 500 negative controls (red), and a set of constitutive promoters from the BioBrick registry (blue).