

Operon Promoter Landscape

Operon	Strand	Operon start	Operon end
gnd	-	2099292	2097886




Figure 1: **Promoter activity in rich media (LB) surrounding query operon.** 17,767 previously reported TSSs were evaluated by measuring the promoter activity (right Y-axis) of the 150 bp surrounding the TSS (-120 to +30) to determine which were active or inactive. The genome-wide promoter activity (left Y-axis) was determined by measuring expression of over 300,000 genomic fragments spanning the *E. coli* genome and averaging promoter activity at all nucleotide positions in a strand-specific fashion. Genome coordinates corresponds to *E. coli* genome version U00096.2.

TSS Summary

TSS name	TSS position	Strand	TSS activity	Category
TSS_8086_storz	2099691	-	0.7632222	inactive
TSS_8083_regulondb	2099460	-	0.8701478	inactive
TSS_8081_regulondb	2099344	-	2.0154637	active
TSS_8085_storz	2099637	-	0.9932795	active
TSS_8084_storz	2099496	+	0.6442522	inactive

TSS Scanning Mutagenesis




Figure 2: **(Left)** Mutagenesis profile(s) of active TSS(s). Sequences surrounding active TSSs were systematically mutated to identify regions controlling expression. Bar height indicates the relative change in promoter activity as a result of scrambling nucleotides within 10 bp regions at 5 bp intervals spanning the promoter. Bar color identifies the region as a putative activator (yellow) or repressor (purple). **(Right)** Dashed line indicates the expression of the indicated TSS relative to all tested TSS sequences. The distributions of expression is shown for all tested TSSs (black), 500 negative controls (red), and a set of constitutive promoters from the BioBrick registry (blue).