Operon Promoter Landscape

Operon	Strand	Operon start	Operon end
secE-nusG	+	4175381	4176311






Figure 1: **Promoter activity in rich media (LB) surrounding query operon**. 17,767 previously reported TSSs were evaluated by measuring the promoter activity (right Y-axis) of the 150 bp surrounding the TSS (-120 to +30) to determine which were active or inactive. The genome-wide promoter activity (left Y-axis) was determined by measuring expression of over 300,000 genomic fragments spanning the *E. coli* genome and averaging promoter activity at all nucleotide positions in a strand-specific fashion. Genome coordinates corresponds to *E. coli* genome version U00096.2.

TSS Summary

TSS name	TSS position	Strand	TSS activity	Category
TSS_16332_storz_regulondb	4174998	+	0.9514336	active
TSS_16334_regulondb	4175236	+	2.6428466	active
TSS_16333_storz_regulondb	4175200	+	2.5933348	active
TSS_16336_regulondb	4175301	+	0.7410552	inactive
TSS_16335_regulondb	4175266	+	2.3765478	active
TSS_16337_regulondb	4175352	+	1.2759458	active

TSS Scanning Mutagenesis




Figure 2: (Left) Mutagenesis profile(s) of active TSS(s). Sequences surrounding active TSSs were systematically mutated to identify regions controlling expression. Bar height indicates the relative change in promoter activity as a result of scrambling nucleotides within 10 bp regions at 5 bp intervals spanning the promoter. Bar color identifies the region as

a putative activator (yellow) or repressor (purple). (**Right)** Dashed line indicates the expression of the indicated TSS relative to all tested TSS sequences. The distributions of expression is shown for all tested TSSs (black), 500 negative controls (red), and a set of

1

constitutive promoters from the BioBrick registry (blue).