Operon Promoter Landscape

Operon	Strand	Operon start	Operon end
htrL	-	3791706	3790849







Figure 1: **Promoter activity in rich media (LB) surrounding query operon**. 17,767 previously reported TSSs were evaluated by measuring the promoter activity (right Y-axis) of the 150 bp surrounding the TSS (-120 to +30) to determine which were active or inactive. The genome-wide promoter activity (left Y-axis) was determined by measuring expression of over 300,000 genomic fragments spanning the *E. coli* genome and averaging promoter activity at all nucleotide positions in a strand-specific fashion. Genome coordinates corresponds to *E. coli* genome version U00096.2.

TSS Summary

TSS name	TSS position	Strand	TSS activity	Category
TSS_14696_regulondb	3791958	+	1.4271504	active
TSS_14694_storz	3791887	-	0.6909279	inactive
TSS_14693_regulondb	3791857	-	0.6652528	inactive
TSS_14692_storz	3791815	-	0.8764632	inactive
TSS_14697_regulondb	3791993	+	1.4588814	active
TSS_14695_regulondb	3791888	+	0.4140789	inactive

TSS Scanning Mutagenesis



TSSs were systematically mutated to identify regions controlling expression. Bar height indicates the relative change in promoter activity as a result of scrambling nucleotides within 10 bp regions at 5 bp intervals spanning the promoter. Bar color identifies the region as a putative activator (yellow) or repressor (purple). **(Right)** Dashed line indicates the expression of the indicated TSS relative to all tested TSS sequences. The distributions of expression is shown for all tested TSSs (black), 500 negative controls (red), and a set of constitutive promoters from the BioBrick registry (blue).