

# 18-660: Numerical Methods for Engineering Design and Optimization

Xin Li  
Department of ECE  
Carnegie Mellon University  
Pittsburgh, PA 15213



Slide 1

## Overview


### ■ Monte Carlo Analysis

- ▼ Random variable
- ▼ Probability distribution
- ▼ Random sampling

Slide 2

## Random Variables

- A **random variable** is a real-valued function of the outcome of the experiment



We get different results from different experiments (i.e., the output is random)

Slide 3


## Probability Distribution

- A continuous random variable  $x$  is defined by its probability distribution function

- Probability density function (PDF)

▼  $\text{pdf}_x(t_x)$  denotes the probability per unit length near  $x = t_x$


$$\int_a^b \text{pdf}_x(y) dy = P[a \leq x \leq b]$$



- Cumulative distribution function (CDF)

▼  $\text{cdf}_x(t_x)$  equals the probability of  $x \leq t_x$

$$\begin{aligned} \text{cdf}(t_x) &= P(x \leq t_x) \\ &= \int_{-\infty}^{t_x} \text{pdf}_x(y) dy \end{aligned}$$



Slide 4

## Expectation

- Given a random variable  $x$  and a function  $f(x)$ , the expectation of  $f(x)$  is the weighted average of the possible values of  $f(x)$

$$E[f(x)] = \int_{-\infty}^{+\infty} f(y) \cdot p_{df_x}(y) dy$$

- A useful equation for expected value calculation

$$E[f(x) + g(x)] = E[f(x)] + E[g(x)]$$

Slide 5

## Mean, Variance and Standard Deviation

- Mean

$$E[x]$$

- Variance

$$\text{VAR}[x] = E[(x - E[x])^2]$$



- Standard deviation

$$\text{STD}[x] = \sqrt{\text{VAR}(x)}$$



Slide 6

## Mean, Variance and Standard Deviation

- Mean measures the “average position” of  $x$



- Variance measures the “spread” of the distribution



$$\int_{-\infty}^{+\infty} p d x (y) d y = 1$$

Slide 7

## Moments and Central Moments

- k-th order moment

$$E[x^k]$$

- ▼ Mean is the first order moment ( $k=1$ )

- k-th order central moments

$$E[(x - E[x])^k]$$

- ▼ Variance is the second order central moment ( $k=2$ )

Slide 8




## Normal Distribution

- A random variable  $x$  is Normal if

$$pdf_x(t) = \frac{1}{\sigma\sqrt{2\pi}} \cdot e^{-\frac{(t-\mu)^2}{2\sigma^2}}$$

- ▼  $\mu$ : mean
- ▼  $\sigma$ : standard deviation
- ▼ Denoted as  $N(\mu, \sigma^2)$



Standard Normal distribution


- If  $\mu = 0$  and  $\sigma = 1$ , it is called standard Normal distribution

Why is Normal distribution important to us?

Slide 9

## Normal Distribution

- Many physical variations are Normal
- **Central limit theorem**: the variation caused by a large number of independent random factors is "almost" Normal



Assume that all  $x_i$ 's are independent and have the same uniform distribution

Slide 10

## Multiple Random Variables

- Two continuous random variables  $x$  and  $y$  are defined by their joint probability distribution
- Joint probability density function

$$\int_c^d \int_a^b pdf_{x,y}(\tau_x, \tau_y) \cdot d\tau_x \cdot d\tau_y = P(a \leq x \leq b, c \leq y \leq d)$$

$$x \in [a, b]$$

$$y \in [c, d]$$

- Joint cumulative distribution function


$$cdf_{x,y}(t_x, t_y) = P(x \leq t_x, y \leq t_y) = \int_{-\infty}^{t_x} \int_{-\infty}^{t_y} pdf_{x,y}(\tau_x, \tau_y) \cdot d\tau_x \cdot d\tau_y$$

Applicable to more than two random variables


Slide 11

## Joint Probability Distribution

- Example: bivariate Normal distribution



Joint probability density function



Joint cumulative distribution function

Slide 12

## Marginal Distribution Function

### ■ Marginal probability density function

$$pdf_x(t_x) = \int_{-\infty}^{+\infty} pdf_{x,y}(t_x, \tau_y) \cdot d\tau_y$$
$$pdf_y(t_y) = \int_{-\infty}^{+\infty} pdf_{x,y}(\tau_x, t_y) \cdot d\tau_x$$


### ■ Marginal cumulative distribution function

$$cdf_x(t_x) = P(x \leq t_x, y \leq +\infty) = \lim_{t_y \rightarrow +\infty} cdf_{x,y}(t_x, t_y)$$
$$cdf_y(t_y) = P(x \leq +\infty, y \leq t_y) = \lim_{t_x \rightarrow +\infty} cdf_{x,y}(t_x, t_y)$$

Slide 13

## Marginal Distribution Function

### ■ Example: bivariate Normal distribution



Slide 14



## Covariance and Correlation

### ■ Covariance

$$\text{Cov}[x, y] = E[(x - E[x]) \cdot (y - E[y])]$$

- ▼ If  $\text{COV}[x, y] = 0$ , then  $x$  and  $y$  are **uncorrelated**

### ■ Covariance matrix

$$\Sigma = \begin{bmatrix} \text{Cov}[x, x] & \text{Cov}[x, y] \\ \text{Cov}[y, x] & \text{Cov}[y, y] \end{bmatrix}$$

↓  
 $\text{Var}[y]$

- ▼  $\Sigma$  is always symmetric
- ▼ Diagonal components are corresponding to variance values
- ▼  $\Sigma$  is diagonal if  $x$  and  $y$  are uncorrelated


Slide 15

## Covariance and Correlation

### ■ Correlation (normalized covariance)

$$\text{COR}[x, y] = \frac{\text{COV}[x, y]}{\text{STD}[x] \cdot \text{STD}[y]}$$

- ▼ Correlation between two random variables can be visualized by **scatter plot**




Slide 16



## Covariance and Correlation

### ■ Example: correlated random variables




Slide 17

## Monte Carlo Analysis

### ■ Problem definition

- ▼ Find probability distribution and/or moments of



### ■ In general, the distribution and/or moments of $f$ cannot be calculated analytically, because

- ▼  $f(X)$  is nonlinear
- ▼  $f(X)$  may not have closed-form expression (we can only numerically calculate  $f$  for a given  $X$  value)

Slide 18

## Monte Carlo Analysis

### ■ Monte Carlo analysis for $f(X)$

- ▼ Randomly select  $M$  samples for  $X$
- ▼ Evaluate function  $f(X)$  at each sampling point
- ▼ Estimate distribution of  $f$  using these  $M$  samples



Slide 19

## Monte Carlo Analysis Example

### ■ Example: estimate the probability distribution of

$$y = \exp(x)$$

- ▼  $x \sim N(0,1)$  (standard Normal distribution)

Slide 20

## Monte Carlo Analysis Example

- Step 1: draw random samples for x

Samples	1	2	3	4	5	6	...
x	-0.4326	-1.6656	0.1253	0.2877	-1.1465	1.1909	...

M random samples for x

- Step 2: calculate y at each sampling point


Samples	1	2	3	4	5	6	...
y	0.6488	0.1891	1.1335	1.3333	0.3178	3.2901	...

M random samples for y

Slide 21

## Monte Carlo Analysis Result

- Monte Carlo result is typically represented by a histogram
  - ▼ A big table of data is not intuitive



Histogram of y based on 1000 random samples


**QUESTION:** how accurate is Monte Carlo analysis?

Slide 22



## Monte Carlo Analysis Accuracy


- Monte Carlo analysis is not deterministic
  - ▼ We cannot get identical results when running MC twice
  - ▼ The analysis error is not deterministic
- Monte Carlo accuracy depends on the number of samples
  - ▼ Examples: histogram of  $y$



Slide 23

## Monte Carlo Analysis Accuracy

- Example: bivariate Normal distribution
  - ▼  $x$  and  $y$  are independent and jointly standard Normal



Slide 24

## Monte Carlo Analysis Accuracy

- Statistical methods exist to analyze Monte Carlo accuracy
- Example: Monte Carlo accuracy analysis

$x \sim N(0,1)$       Standard Normal distribution

- ▼ Estimate the mean value  $\mu_x$  by Monte Carlo analysis
- ▼ Our question: how accurate is the estimated  $\mu_x$  (dependent on the number of Monte Carlo samples)?

Slide 25

## Monte Carlo Analysis Accuracy

- Monte Carlo analysis for the mean value  $\mu_x$ 
  - ▼ Randomly draw M sampling points  $\{x^{(1)}, x^{(2)}, \dots, x^{(M)}\}$
  - ▼ Estimate  $\mu_x$  by the following equation

$$\mu_x = \frac{x^{(1)} + x^{(2)} + \dots + x^{(M)}}{M}$$

↓  
Estimator

- Assumptions in our accuracy analysis
  - ▼ Each  $x^{(i)}$  is random and satisfies standard Normal distribution – it is randomly created for  $x \sim N(0,1)$
  - ▼ All  $x^{(i)}$ 's are mutually independent – samples from a good random number generator should be independent
- $\mu_x$  is a function of  $\{x^{(1)}, x^{(2)}, \dots, x^{(M)}\}$ , which is a random variable

Slide 26

## Monte Carlo Analysis Accuracy

### ■ Mean of $\mu_x$

$$\begin{aligned} E[\mu_x] &= E\left[\frac{x^{(1)} + x^{(2)} + \dots + x^{(M)}}{M}\right] \\ &= \frac{1}{M} [E[x^{(1)}] + E[x^{(2)}] + \dots + E[x^{(M)}]] \\ &= \frac{1}{M} [0 + 0 + 0 + \dots + 0] = 0 \end{aligned}$$

### ■ Variance of $\mu_x$

$$\begin{aligned} E[\mu_x^2] &= E\left[\left(\frac{x^{(1)} + x^{(2)} + \dots + x^{(M)}}{M}\right)^2\right] \\ &= E\left\{\frac{x^{(1)2} + x^{(2)2} + \dots + x^{(M)2}}{M^2}\right\} \\ &= \frac{1 + 1 + \dots + 1}{M^2} = \frac{M}{M^2} = \frac{1}{M} \end{aligned}$$

$E[x^{(i)} x^{(j)}] = E[x^{(i)}] E[x^{(j)}] = 0 \text{ (i} \neq \text{j)}$

Slide 27


## Monte Carlo Analysis Accuracy

### ■ $E\{\mu_x\} = 0$

- ▼  $\mu_x$  is an **unbiased estimator**
- ▼ Otherwise, if the estimator mean is not equal to the actual mean, it is called a **biased estimator**

### ■ $E\{\mu_x^2\} = 1/M$

- ▼ Variance decreases as M increases
- ▼ Distributions of  $\mu_x$  for different M values



$\mu_x$  is a Normal distribution  $N(0, 1/M)$

Slide 28



## Monte Carlo Analysis Accuracy

- “Average” estimation accuracy is better when using larger M

- In this  $\mu_x$  example

- ▼ If we require that  $\pm 3$  sigma of  $\mu_x$  is within  $[-0.1, 0.1]$

$$\sqrt{\frac{1}{M}} \quad \frac{3}{\sqrt{M}} \leq 0.1 \quad \Rightarrow \quad M \geq 900$$

- ▼ If we require that  $\pm 3$  sigma of  $\mu_x$  is within  $[-0.01, 0.01]$

$$\frac{3}{\sqrt{M}} \leq 0.01 \quad \Rightarrow \quad M \geq 90000$$

Slide 29

## Monte Carlo Analysis Accuracy

- Accuracy is improved by **10x** if the number of samples is increased by **100x**
- 1K ~ 10K sampling points are typically required to achieve reasonable accuracy
- However, even if you use 10K sampling points, an accurate result is not guaranteed!
  - ▼ Monte Carlo analysis is random, and you can be unlucky (e.g., going beyond  $\pm 3$  sigma range)

Slide 30

## Summary

### ■ Monte Carlo analysis

- ▼ Random variable
- ▼ Probability distribution
- ▼ Random sampling

Slide 31

# 18-660: Numerical Methods for Engineering Design and Optimization

Xin Li

Department of ECE  
Carnegie Mellon University  
Pittsburgh, PA 15213



Slide 1

## Overview

### ■ Monte Carlo Analysis

- ▼ Latin hypercube sampling
- ▼ Importance sampling



Slide 2



## Latin Hypercube Sampling (LHS)


- A great number of samples are typically required in traditional Monte Carlo to achieve good accuracy
- Various techniques exist to improve Monte Carlo accuracy
- Controlling sampling points is the key
  - ▼ Latin hypercube sampling is a widely-used method to generate controlled random samples
  - ▼ The basic idea is to make sampling point distribution close to probability density function (PDF)

M. McKay, R. Beckman and W. Conover, "A comparison of three methods for selecting values of input variables in the analysis of output from a computer code," *Technometrics*, vol. 21, no. 2, pp. 239-245, May. 1979

Slide 3

## Latin Hypercube Sampling (LHS)

- One dimensional Latin hypercube sampling
  - ▼ Evenly partition CDF into N regions
  - ▼ Randomly pick up one sampling point in each region




Slide 4

## Latin Hypercube Sampling (LHS)

### ■ Two dimensional Latin hypercube sampling

- ▼  $x_1$  and  $x_2$  must be independent
- ▼ Generate one-dimensional LHS samples for  $x_1$
- ▼ Generate one-dimensional LHS samples for  $x_2$
- ▼ Randomly combine the LHS samples to two-dimensional pairs



- One sample in each row and each column
- Sampling is random in each grid
- Higher-dimensional LHS samples can be similarly generated

Slide 5

## Latin Hypercube Sampling (LHS)

### ■ Matlab code for LHS sampling of independent standard Normal distributions

```
data = rand(NSample,NVar);
for i = 1:NVar
    index = randperm(NSample);
    prob = (index'-data(:,i))/NSample;
    data(:,i) = sqrt(2)*erfinv(2*prob-1);
end;
```

- ▼ NVar: # of random variables
- ▼ NSample: # of samples
- ▼ data: LHS sampling points


Slide 6

## Latin Hypercube Sampling (LHS)

### ■ Compare Monte Carlo accuracy for a simple example


- ▾  $x \sim N(0,1)$  (standard Normal distribution)
- ▾ Repeatedly estimate the mean value by Monte Carlo analysis

Accuracy is improved by increasing sampling #



Random sampling

Monte Carlo is not deterministic



Latin hypercube sampling

Slide 7

## Importance Sampling

### ■ Even with Latin hypercube sampling, Monte Carlo analysis requires a **HUGE** number of sampling points

### ■ Example: rare event estimation

$x \sim N(0,1)$  Standard Normal distribution

Estimate  $P(x \leq -5) = ???$

- ▾ The theoretical answer for  $P(x \leq -5)$  is equal to  $2.87 \times 10^{-7}$
- ▾ ~100M sampling points are required if we attempt to estimate this probability by random sampling or LHS

Slide 8




## Importance Sampling

### ■ Key idea:

- ▼ Do not generate random samples from  $\text{pdf}_x(t)$
- ▼ Instead, find a good distorted  $\text{pdf}_y(t)$  to improve Monte Carlo sampling accuracy

### ■ Example: if $x \sim N(0,1)$ , what is $P(x \leq -5)$ ?

- ▼ Intuitively, if we draw sampling points based on  $\text{pdf}_y(t)$ , more samples will fall into the grey area



Slide 9

## Importance Sampling

### ■ Assume that we want to estimate the following expected value

$$E[f(x)] = \int_{-\infty}^{+\infty} f(t) \cdot \text{pdf}_x(t) \cdot dt$$

### ■ Example: if we want to estimate $P(x \leq -5)$ , then

$$\begin{aligned}
 f(x) &= \begin{cases} 1 & (x \leq -5) \\ 0 & (x > -5) \end{cases} \\
 E[f(x)] &= \int_{-\infty}^{+\infty} f(t) \cdot \text{pdf}_x(t) dt \\
 &= \int_{-\infty}^{-5} \underbrace{f(t)}_{=1} \cdot \text{pdf}_x(t) dt + \int_{-5}^{+\infty} \underbrace{f(t)}_{=0} \cdot \text{pdf}_x(t) dt \\
 &= P(x \leq -5)
 \end{aligned}$$

Slide 10

## Importance Sampling

- Estimate  $E[f(x)]$  where  $x \sim \text{pdf}_x(t)$  by importance sampling

$$\begin{aligned}
 E[f(x)] &= \int_{-\infty}^{+\infty} f(t) \cdot \text{pdf}_x(t) dt \\
 &= \int_{-\infty}^{+\infty} f(t) \cdot \text{pdf}_x(t) \cdot \frac{\text{pdf}_y(t)}{\text{pdf}_y(t)} dt \\
 &= \int_{-\infty}^{+\infty} f(t) \cdot \underbrace{\frac{\text{pdf}_x(t)}{\text{pdf}_y(t)}}_{g(t)} \cdot \text{pdf}_y(t) dt \\
 &= E[g(x)] \quad x \sim \text{pdf}_x(x)
 \end{aligned}$$

Slide 11

## Importance Sampling

- Estimate  $E[f(x)]$  where  $x \sim \text{pdf}_x(t)$  by traditional sampling
  - ▼ Step 1: draw  $M$  random samples  $\{t_1, t_2, \dots, t_M\}$  based on  $\text{pdf}_x(t)$
  - ▼ Step 2: calculate  $f_m = f(t_m)$  at each sampling point  $m = 1, 2, \dots, M$
  - ▼ Step 3: calculate  $E[f] \approx (f_1 + f_2 + \dots + f_M)/M$
- Estimate  $E[f(x)]$  where  $x \sim \text{pdf}_x(t)$  by importance sampling
  - ▼ Step 1: draw  $M$  random samples  $\{t_1, t_2, \dots, t_M\}$  based on  $\text{pdf}_y(t)$
  - ▼ Step 2: calculate  $g_m = f(t_m) \cdot \text{pdf}_x(t_m) / \text{pdf}_y(t_m)$  at each sampling point  $m = 1, 2, \dots, M$
  - ▼ Step 3: calculate  $E[f] \approx (g_1 + g_2 + \dots + g_M)/M$

How do we decide the optimal  $\text{pdf}_y(t)$  to achieve minimal Monte Carlo analysis error?

Slide 12

## Importance Sampling

- Determine optimal pdf<sub>y</sub>(t) for importance sampling

$$E[f] \approx \mu_f = \frac{1}{M} \cdot \sum_{m=1}^M f(t_m) \cdot \underbrace{\frac{pdf_x(t_m)}{pdf_y(t_m)}}_{g(t)}$$

↓  
Estimator

- The accuracy of an estimator can be quantitatively measured by its variance

$$Error \sim VAR[\mu_f]$$

- ▼ To improve Monte Carlo analysis accuracy, we should minimize VAR[μ<sub>f</sub>]

Slide 13

## Importance Sampling

- Determine optimal pdf<sub>y</sub>(t) for importance sampling

$$E[f] \approx \mu_f = \frac{1}{M} \cdot \sum_{m=1}^M f(t_m) \cdot \frac{pdf_x(t_m)}{pdf_y(t_m)}$$

- We achieve the minimal VAR[μ<sub>f</sub>] = 0 if

$$f(t) = \frac{pdf_x(t)}{pdf_y(t)} = y(t) = k \text{ constant}$$

$$pdf_y(t) = \frac{f(t) \cdot pdf_x(t)}{k}$$

Slide 14



## Importance Sampling

$$pdf_y(t) = \frac{f(t) \cdot pdf_x(t)}{k}$$

- How do we decide the value  $k$ ?
- $k$  cannot be arbitrarily selected
  - ▼  $pdf_y(t)$  must be a valid PDF that satisfies the following condition

$$\int_{-\infty}^{+\infty} pdf_y(t) dt = \int_{-\infty}^{+\infty} \frac{f(t) \cdot pdf_x(t)}{k} dt = 1$$

$$k = \int_{-\infty}^{+\infty} f(t) \cdot pdf_x(t) dt = E[f]$$

must know  $E[f]$ !!!

Slide 15

## Importance Sampling

- In practice, such an optimal  $pdf_y(t)$  cannot be easily applied
- Instead, we typically look for a sub-optimal solution that satisfies the following constraints
  - ▼ Easy to construct – we do not have to know  $k = E[f]$
  - ▼ Easy to sample – not all random distributions can be easily sampled by a random number generator
  - ▼ Minimal estimator variance – the sub-optimal  $pdf_y(t)$  is close to the optimal case as much as possible

Slide 16


## Importance Sampling

- Finding the right  $\text{pdf}_y(t)$  is nontrivial for practical problems
  - ▼ No magic equation exists in general
  - ▼ Engineering approach is based on heuristics
  - ▼ Sometimes require a lot of human experience and numerical optimization
- The criterion to choose  $\text{pdf}_y(t)$  is also application-dependent

Slide 17

## Importance Sampling

- Example: if  $x \sim N(0,1)$ , what is  $P(x \leq -5)$ ?



Several possible choices for  $\text{pdf}_y(t)$

Slide 18

## Summary

### ■ Monte Carlo analysis

- ▼ Latin hypercube sampling
- ▼ Importance sampling

## Importance Sampling

Example: If  $x \sim N(0,1)$ , what is  $P(x \leq -2)$ ?

