

REFERENCE LIST – PROJECT 1 – BIOMARKERS OF EPILEPTOGENESIS AFTER EXPERIMENTAL TRAUMATIC BRAIN INJURY

1. Ahmed F, Plantman S, Cernak I, Agoston D V. The Temporal Pattern of Changes in Serum Biomarker Levels Reveals Complex and Dynamically Changing Pathologies after Exposure to a Single Low-Intensity Blast in Mice. *Front Neurol*. 2015;6:114. doi:10.3389/fneur.2015.00114.
2. Annegers JF, Hauser WA, Coan SP, Rocca WA. A population-based study of seizures after traumatic brain injuries. *N Engl J Med*. 1998;338(1):20-24. doi:10.1056/NEJM199801013380104.
3. Bartel DP. MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell*. 2004;116(2):281-297. <http://www.ncbi.nlm.nih.gov/pubmed/14744438>. Accessed July 17, 2014.
4. Di Battista AP, Buonora JE, Rhind SG, et al. Blood Biomarkers in Moderate-To-Severe Traumatic Brain Injury: Potential Utility of a Multi-Marker Approach in Characterizing Outcome. *Front Neurol*. 2015;6:110. doi:10.3389/fneur.2015.00110.
5. Bragin A, Azizyan A, Almajano J, Engel J. The cause of the imbalance in the neuronal network leading to seizure activity can be predicted by the electrographic pattern of the seizure onset. *J Neurosci*. 2009;29(11):3660-3671. doi:10.1523/JNEUROSCI.5309-08.2009.
6. Bragin A, Wilson CL, Almajano J, Mody I, Engel J. High-frequency oscillations after status epilepticus: epileptogenesis and seizure genesis. *Epilepsia*. 2004;45(9):1017-1023. doi:10.1111/j.0013-9580.2004.17004.x.
7. Bragin A, Wilson CL, Staba RJ, Reddick M, Fried I, Engel J. Interictal high-frequency oscillations (80-500 Hz) in the human epileptic brain: entorhinal cortex. *Ann Neurol*. 2002;52(4):407-415. doi:10.1002/ana.10291.
8. Bragin A. Pathological electrographic changes after experimental traumatic brain injury. *Epilepsia*. 2016:in press.
9. Burwell RD, Witter MP, Amaral DG. Perirhinal and postrhinal cortices of the rat: a review of the neuroanatomical literature and comparison with findings from the monkey brain. *Hippocampus*. 1995;5(5):390-408. doi:10.1002/hipo.450050503.
10. CDC. No Title. <http://www.cdc.gov/traumaticbraininjury/>.
11. CenterTBI. No Title. <https://www.center-tbi.eu/>.
12. Csicsvari J, Hirase H, Czurkó A, Mamiya A, Buzsáki G. Oscillatory coupling of hippocampal pyramidal cells and interneurons in the behaving Rat. *J Neurosci*. 1999;19(1):274-287. <http://www.ncbi.nlm.nih.gov/pubmed/9870957>. Accessed February 22, 2016.

13. D'Ambrosio R, Eastman CL, Darvas F, et al. Mild passive focal cooling prevents epileptic seizures after head injury in rats. *Ann Neurol*. 2013;73(2):199-209. doi:10.1002/ana.23764.
14. Engel JJ, Pitkanen A, Loeb JA, et al. Epilepsy biomarkers. *Epilepsia*. 2013;54 Suppl 4:61-69. doi:10.1111/epi.12299.
15. Falcone T, Janigro D, Lovell R, et al. S100B blood levels and childhood trauma in adolescent inpatients. *J Psychiatr Res*. 2015;62:14-22. doi:10.1016/j.jpsychires.2014.12.002.
16. Farquharson S, Tournier J-D, Calamante F, et al. White matter fiber tractography: why we need to move beyond DTI. *J Neurosurg*. 2013;118(6):1367-1377. doi:10.3171/2013.2.JNS121294.
17. Fisher RS, van Emde Boas W, Blume W, et al. Epileptic seizures and epilepsy: definitions proposed by the International League Against Epilepsy (ILAE) and the International Bureau for Epilepsy (IBE). *Epilepsia*. 2005;46(4):470-472. doi:10.1111/j.0013-9580.2005.66104.x.
18. Gardner AB, Worrell GA, Marsh E, Dlugos D, Litt B. Human and automated detection of high-frequency oscillations in clinical intracranial EEG recordings. *Clin Neurophysiol*. 2007;118(5):1134-1143. doi:10.1016/j.clinph.2006.12.019.
19. Giacobbo S, Bramanti P, Barresi M, et al. Predictive biomarkers of recovery in traumatic brain injury. *Neurocrit Care*. 2012;16(3):470-477. doi:10.1007/s12028-012-9707-z.
20. Gupta PK, Sayed N, Ding K, et al. Subtypes of post-traumatic epilepsy: clinical, electrophysiological, and imaging features. *J Neurotrauma*. 2014;31(16):1439-1443. doi:10.1089/neu.2013.3221.
21. Hudak AM, Trivedi K, Harper CR, et al. Evaluation of seizure-like episodes in survivors of moderate and severe traumatic brain injury. *J Head Trauma Rehabil*. 19(4):290-295.
<http://www.ncbi.nlm.nih.gov/pubmed/15263856>. Accessed February 22, 2016.
22. Immonen R, Kharatishvili I, Gröhn O, Pitkänen A. MRI biomarkers for post-traumatic epileptogenesis. *J Neurotrauma*. 2013;30(14):1305-1309. doi:10.1089/neu.2012.2815.
23. Immonen RJ, Kharatishvili I, Niskanen J-P, Gröhn H, Pitkänen A, Gröhn OHJ. Distinct MRI pattern in lesional and perilesional area after traumatic brain injury in rat--11 months follow-up. *Exp Neurol*. 2009;215(1):29-40. doi:10.1016/j.expneurol.2008.09.009.
24. Jack CR, Holtzman DM. Biomarker modeling of Alzheimer's disease. *Neuron*. 2013;80(6):1347-1358. doi:10.1016/j.neuron.2013.12.003.
25. Jandó G, Carpi D, Kandel A, et al. Spike-and-wave epilepsy in rats: sex differences and inheritance of physiological traits. *Neuroscience*. 1995;64(2):301-317.
<http://www.ncbi.nlm.nih.gov/pubmed/7700522>. Accessed February 22, 2016.

26. Jones NC, Cardamone L, Williams JP, Salzberg MR, Myers D, O'Brien TJ. Experimental traumatic brain injury induces a pervasive hyperanxious phenotype in rats. *J Neurotrauma*. 2008;25(11):1367-1374. doi:10.1089/neu.2008.0641.
27. Kharatishvili I, Nissinen JP, McIntosh TK, Pitkänen A. A model of posttraumatic epilepsy induced by lateral fluid-percussion brain injury in rats. *Neuroscience*. 2006;140(2):685-697. doi:10.1016/j.neuroscience.2006.03.012.
28. Kharatishvili I, Pitkänen A. Association of the severity of cortical damage with the occurrence of spontaneous seizures and hyperexcitability in an animal model of posttraumatic epilepsy. *Epilepsy Res*. 2010;90(1-2):47-59. doi:10.1016/j.eplepsyres.2010.03.007.
29. Kheiri F, Bragin A, Engel J, Almajano J, Winden E. Non-linear classification of heart rate parameters as a biomarker for epileptogenesis. *Epilepsy Res*. 2012;100(1-2):59-66. doi:10.1016/j.eplepsyres.2012.01.008.
30. Kochanek PM, Bramlett H, Dietrich WD, et al. A novel multicenter preclinical drug screening and biomarker consortium for experimental traumatic brain injury: operation brain trauma therapy. *J Trauma*. 2011;71(1 Suppl):S15-S24. doi:10.1097/TA.0b013e31822117fe.
31. Liu YR, Cardamone L, Hogan RE, et al. Progressive metabolic and structural cerebral perturbations after traumatic brain injury: an in vivo imaging study in the rat. *J Nucl Med*. 2010;51(11):1788-1795. doi:10.2967/jnumed.110.078626.
32. Marchi N, Granata T, Janigro D. Inflammatory pathways of seizure disorders. *Trends Neurosci*. 2014;37(2):55-65. doi:10.1016/j.tins.2013.11.002.
33. McIntosh TK, Vink R, Noble L, et al. Traumatic brain injury in the rat: characterization of a lateral fluid-percussion model. *Neuroscience*. 1989;28(1):233-244. <http://www.ncbi.nlm.nih.gov/pubmed/2761692>. Accessed February 22, 2016.
34. Messori A, Polonara G, Carle F, Gesuita R, Salvolini U. Predicting posttraumatic epilepsy with MRI: prospective longitudinal morphologic study in adults. *Epilepsia*. 2005;46(9):1472-1481. doi:10.1111/j.1528-1167.2005.34004.x.
35. Nissinen J, Halonen T, Koivisto E, Pitkänen A. A new model of chronic temporal lobe epilepsy induced by electrical stimulation of the amygdala in rat. *Epilepsy Res*. 2000;38(2-3):177-205. <http://www.ncbi.nlm.nih.gov/pubmed/10642046>. Accessed February 22, 2016.
36. Nowak JS, Michlewski G. miRNAs in development and pathogenesis of the nervous system. *Biochem Soc Trans*. 2013;41(4):815-820. doi:10.1042/BST20130044.
37. Pearce PS, Friedman D, Lafrancois JJ, et al. Spike-wave discharges in adult Sprague-Dawley rats and their implications for animal models of temporal lobe epilepsy. *Epilepsy Behav*. 2014;32:121-131. doi:10.1016/j.yebeh.2014.01.004.

38. Pitkanen A, Lukasiuk K. Mechanisms of epileptogenesis and potential treatment targets. *Lancet Neurol.* 2011;10(2):173-186. doi:10.1016/S1474-4422(10)70310-0.
39. Pitkanen A, Nehlig A, Brooks-Kayal AR, et al. Issues related to development of antiepileptogenic therapies. *Epilepsia.* 2013;54 Suppl 4:35-43. doi:10.1111/epi.12297.
40. Pitkänen A, Immonen RJ, Gröhn OHJ, Kharatishvili I. From traumatic brain injury to posttraumatic epilepsy: what animal models tell us about the process and treatment options. *Epilepsia.* 2009;50 Suppl 2:21-29. doi:10.1111/j.1528-1167.2008.02007.x.
41. Pitkänen A, Lukasiuk K, Dudek FE, Staley KJ. Epileptogenesis. *Cold Spring Harb Perspect Med.* 2015;5(10). doi:10.1101/cshperspect.a022822.
42. Redell JB, Moore AN, Ward NH, Hergenroeder GW, Dash PK. Human traumatic brain injury alters plasma microRNA levels. *J Neurotrauma.* 2010;27(12):2147-2156. doi:10.1089/neu.2010.1481.
43. Saatman KE, Duhaime A-C, Bullock R, Maas AIR, Valadka A, Manley GT. Classification of traumatic brain injury for targeted therapies. *J Neurotrauma.* 2008;25(7):719-738. doi:10.1089/neu.2008.0586.
44. Shultz SR, Cardamone L, Liu YR, et al. Can structural or functional changes following traumatic brain injury in the rat predict epileptic outcome? *Epilepsia.* 2013;54(7):1240-1250. doi:10.1111/epi.12223.
45. Sierra A, Laitinen T, Gröhn O, Pitkänen A. Diffusion tensor imaging of hippocampal network plasticity. *Brain Struct Funct.* 2015;220(2):781-801. doi:10.1007/s00429-013-0683-7.
46. Staba RJ, Stead M, Worrell GA. Electrophysiological biomarkers of epilepsy. *Neurotherapeutics.* 2014;11(2):334-346. doi:10.1007/s13311-014-0259-0.
47. Staba RJ, Wilson CL, Bragin A, Fried I, Engel J. Quantitative analysis of high-frequency oscillations (80-500 Hz) recorded in human epileptic hippocampus and entorhinal cortex. *J Neurophysiol.* 2002;88(4):1743-1752. <http://www.ncbi.nlm.nih.gov/pubmed/12364503>. Accessed February 22, 2016.
48. Staba RJ, Wilson CL, Bragin A, Jhung D, Fried I, Engel J. High-frequency oscillations recorded in human medial temporal lobe during sleep. *Ann Neurol.* 2004;56(1):108-115. doi:10.1002/ana.20164.
49. Tournier J-D, Calamante F, Connelly A. Determination of the appropriate b value and number of gradient directions for high-angular-resolution diffusion-weighted imaging. *NMR Biomed.* 2013;26(12):1775-1786. doi:10.1002/nbm.3017.
50. Truettner JS, Alonso OF, Bramlett HM, Dietrich WD. Therapeutic hypothermia alters microRNA responses to traumatic brain injury in rats. *J Cereb Blood Flow Metab.* 2011;31(9):1897-1907. doi:10.1038/jcbfm.2011.33.

51. Le Van Quyen M, Bragin A. Analysis of dynamic brain oscillations: methodological advances. *Trends Neurosci.* 2007;30(7):365-373. doi:10.1016/j.tins.2007.05.006.
52. Wang J, Tan L, Tan L, et al. Circulating microRNAs are promising novel biomarkers for drug-resistant epilepsy. *Sci Rep.* 2015;5:10201. doi:10.1038/srep10201.
53. Wang J, Yu J-T, Tan L, et al. Genome-wide circulating microRNA expression profiling indicates biomarkers for epilepsy. *Sci Rep.* 2015;5:9522. doi:10.1038/srep09522.
54. Wang X, Sun Y, Tan Z, et al. Serum MicroRNA-4521 is a Potential Biomarker for Focal Cortical Dysplasia with Refractory Epilepsy. *Neurochem Res.* December 2015. doi:10.1007/s11064-015-1773-0.
55. Wheeler BM, Heimberg AM, Moy VN, et al. The deep evolution of metazoan microRNAs. *Evol Dev.* 11(1):50-68. doi:10.1111/j.1525-142X.2008.00302.x.
56. Yokobori S, Hosein K, Burks S, Sharma I, Gajavelli S, Bullock R. Biomarkers for the clinical differential diagnosis in traumatic brain injury--a systematic review. *CNS Neurosci Ther.* 2013;19(8):556-565. doi:10.1111/cns.12127.