

REFERENCES – PROJECT 3 – BIOMARKERS OF HUMAN EPILEPTOGENESIS AFTER TRAUMATIC BRAIN INJURY

1. Alkawadri R, Gaspard N, Goncharova I et al. The spatial and signal characteristics of physiologic high frequency oscillations. *Epilepsia*. 2014 Dec;55(12):1986-95.
2. Annegers JF, Hauser WA, Coan SP et al. A population-based study of seizures after traumatic brain injuries. *N Engl J Med*. 1998;338(1):20-4.
3. Arango JI, Deibert CP, Brown D et al. Posttraumatic seizures in children with severe traumatic brain injury. *Childs Nerv Syst*. 2012;28(11):1925-9.
4. Arndt DH, Lerner JT, Matsumoto JH et al. Subclinical early posttraumatic seizures detected by continuous EEG monitoring in a consecutive pediatric cohort. *Epilepsia*. 2013;54:1780-8.
5. Bonilha L, Edwards JC, Kinsman SL et al. Extrahippocampal gray matter loss and hippocampal deafferentation in patients with temporal lobe epilepsy. *Epilepsia*. 2010 Apr;51(4):519-28.
6. Bolkvadze T and Pitkänen A. Development of post-traumatic epilepsy after controlled cortical impact and lateral fluid-percussion-induced brain injury in the mouse. *J Neurotrauma*. 2012;29:789-812.
7. Bragin A, Lin L, Almajano J et al. Pathological electrographic changes after experimental traumatic brain injury. *Epilepsia*. 2016, (in press).
8. Bragin A, Engel J Jr., Wilson CL et al. High-frequency oscillations in human brain. *Hippocampus* 1999;9:137-42.
9. Bragin A, Wilson CL, Almajano J et al. High-frequency oscillations after status epilepticus: epileptogenesis and seizure genesis. *Epilepsia*. 2004;45:1017-1023.
10. Burnham KP and Anderson DR. Model selection and multimodel inference: a practical information-theoretic approach. Springer. (2002; revised 2013); Chapters 1-7.
11. Chamberlain AC, Viventi J, Blanco JA et al. Millimeter-scale epileptiform spike patterns and their relationship to seizures. *Conf Proc IEEE Eng Med Biol Soc*. 2011;761-4.
12. Chauvière L, Doublet T, Ghestem A et al. Changes in interictal spike features precede the onset of temporal lobe epilepsy. *Ann Neurol*. 2012;71:805-14.
13. Christensen J, Pedersen MG, Pedersen CB et al. Long-term risk of epilepsy after traumatic brain injury in children and young adults: a population-based cohort study. *Lancet*. 2009;373:1105-10.
14. Chu A1, Cui J, Dinov ID. SOCR Analyses - an Instructional Java Web-based Statistical Analysis Toolkit. *J Online Learn Teach*. 2009;5:1-18.
15. Claassen J, Mayer SA, Kowalski RG et al. Detection of electrographic seizures with continuous EEG monitoring in critically ill patients. *Neurology*. 2004;62:1743-8
16. Coulter DA, Rafiq A, Shumate M et al. Brain injury-induced enhanced limbic epileptogenesis: anatomical and physiological parallels to an animal model of temporal lobe epilepsy. *Epilepsy Res*. 1996;26(1):81-91.
17. Dinkelacker V, Valabregue R, Thivard L et al. Hippocampal-thalamic wiring in medial temporal lobe epilepsy: Enhanced connectivity per hippocampal voxel. *Epilepsia*. 2015 Aug;56(8):1217-26.
18. Dinov ID, Petrosyan P, Liu Z et al. High-throughput neuroimaging-genetics computational infrastructure. *Front Neuroinform*. 2014;23;8:41.
19. Dinov ID, Petrosyan P, Liu Z et al. Alzheimer's Disease Neuroimaging Initiative. The perfect neuroimaging-genetics-computation storm: collision of petabytes of data, millions of hardware devices and thousands of software tools. *Brain Imaging Behav*. 2014;8:311-22.
20. Greicius MD, Krasnow B, Reiss AL et al. Functional connectivity in the resting brain: a network analysis of the default mode hypothesis. *Proceedings of the National Academy of Sciences*. 2003;100(1):253-8.

21. Gupta RK, Saksena S, Agarwal A et al. Diffusion tensor imaging in late posttraumatic epilepsy. *Epilepsia*. 2005;46:1465-71.
22. Haneef Z, Lenartowicz A, Yeh HJ et al. Functional connectivity of hippocampal networks in temporal lobe epilepsy. *Epilepsia*. 2014 Jan;55(1):137-45.
23. Haneef Z, Chiang S, Yeh HJ, Engel J Jr, Stern JM. (2015) Functional connectivity homogeneity correlates with duration of temporal lobe epilepsy. *Epilepsy Behav* 46:227-33.9.
24. Chiang S, Cassese A, Guindani M et al. Time-dependence of graph theory metrics in functional connectivity analysis. *Neuroimage*. 2016 Jan 15;125:601-15.
25. Hirsch LJ, LaRoche SM, Gaspard N et al. American Clinical Neurophysiology Society's Standardized Critical Care EEG Terminology: 2012 version. *J Clin Neurophysiol*. 2013 Feb;30(1):1-27.
26. Huneau C, Benquet P, Dieuset G et al. Shape features of epileptic spikes are a marker of epileptogenesis in mice. *Epilepsia*. 2013;54(12):2219-27
27. Hunt RF, Boychuk JA, Smith BN. Neural circuit mechanisms of post-traumatic epilepsy. *Front Cell Neurosci*. 2013;18;7:89.
28. Immonen R, Kharatishvili I, Gröhn O et al. MRI biomarkers for post-traumatic epileptogenesis. *J Neurotrauma*. 2013;30:1305-9.
29. Jacobs J, Vogt C, LeVan P et al. The identification of distinct high-frequency oscillations during spikes delineates the seizure onset zone better than high-frequency spectral power changes. *Clin Neurophysiol*. 2016 Jan;127(1):129-42.
30. Irimia A, Goh SY, Torgerson CM et al. Structural and connectomic neuroimaging for the personalized study of longitudinal alterations in cortical shape, thickness and connectivity after traumatic brain injury. *J Neurosurg Sci*. 2014;58(3):129.
31. Jennett B. Early traumatic epilepsy. Incidence and significance after nonmissile injuries. *Arch Neurol*. 1974;30:394-8.
32. Lamar CD, Hurley RA, Rowland JA et al. Post-traumatic epilepsy: review of risks, pathophysiology, and potential biomarkers. *J Neuropsychiatry Clin Neurosci*. 2014;26:113.
33. Liliang PC, Liang CL, Weng HC et al. Tau proteins in serum predict outcome after severe traumatic brain injury. *J Surg Res*. 2010 May 15;160(2):302-7
34. Lin JJ, Mula M, Hermann BP. Uncovering the neurobehavioural comorbidities of epilepsy over the lifespan. *Lancet*. 2012 380(9848):1180-92.
35. Shultz SR, Wright DK, Zheng P et al. Sodium selenate reduces hyperphosphorylated tau and improves outcomes after traumatic brain injury. *Brain*. 2015 May;138(5):1297-313
36. Löscher W and Brandt C. Prevention or modification of epileptogenesis after brain insults: experimental approaches and translational research. *Pharmacol Rev*. 2010 62(4):668-700.
37. Lutkenhoff ES, McArthur DL, Hua X et al. Thalamic atrophy in antero-medial and dorsal nuclei correlates with six-month outcome after severe brain injury. *Neuroimage Clin*. 2013;3:396-404.
38. Lutkenhoff E.S., Rosenberg M., Chiang J et al. (Optimized brain extraction for pathological brains (optiBET)). *PloS one*. 2014;9(12):e115551.
39. Lutkenhoff ES, Chiang J, Tshibanda L et al. Thalamic and extrathalamic mechanisms of consciousness after severe brain injury. *Annals of neurology*, 2014;78(1):68-76.
40. Marquez de la Plata CD, Garces J, Kojori ES et al. Deficits in functional connectivity of hippocampal and frontal lobe circuits after traumatic axonal injury. *Arch Neurol*. 2011 Jan;68(1):74-84.

41. Monti MM, Lutkenhoff ES, Rubinov M et al. Dynamic change of global and local information processing in propofol-induced loss and recovery of consciousness. *PLoS computational biology*. 2013;9(10):e1003271.
42. Mishra AM, Bai X, Sanganahalli BG et al. Decreased resting functional connectivity after traumatic brain injury in the rat. *PLoS One*. 2014 Apr 18;9(4):e95280.
43. Monti MM, Rosenberg M, Finoia P et al. Thalamo-frontal connectivity mediates top-down cognitive functions in disorders of consciousness. *Neurology*. 2015;84(2):167-73.
44. Ottman R, Barker-Cummings C, Leibson CL et al. Validation of a brief screening instrument for the ascertainment of epilepsy. *Epilepsia*. 2010;51:191-7.
45. Pitkänen A and Immonen R. Epilepsy related to traumatic brain injury. *Neurotherapeutics*. 2014 Apr;11(2):286-96.
46. Power JD, Mitra A, Laumann TO et al. Methods to detect, characterize, and remove motion artifact in resting state fMRI. *Neuroimage*. 2014;84:320-41.
47. Salazar AM, Jabbari B, Vance SC, Grafman J, Amin D, Dillon JD. Epilepsy after penetrating head injury. I. Clinical correlates: a report of the Vietnam Head Injury Study. *Neurology*. 1985;35(10):1406.
48. Shahim P, Tegner Y, Wilson DH et al. Blood biomarkers for brain injury in concussed professional ice hockey players. *JAMA Neurol*. 2014 Jun;71(6):684-92.
49. Staba RJ, Wilson CL, Bragin A et al. High-frequency oscillations recorded in human medial temporal lobe during sleep. *Ann Neurol* 204;56:108-115.
50. Temkin NR, Dikmen SS, Wilensky AJ et al. A randomized, double-blind study of phenytoin for the prevention of post-traumatic seizures. *N Engl J Med*. 1990;323(8):497-502.
51. Trinka E and Brigo F. Antiepileptogenesis in humans: disappointing clinical evidence and ways to move forward. *Curr Opin Neurol*. 2014;27(2):227-35.
52. Vespa PM, Ronne-Engstrom E, Smith M et al. Increase in extracellular glutamate caused by reduced cerebral perfusion pressure and seizures after human traumatic brain injury: a microdialysis study. *J. Neurosurg* 1998;89:971-982.
53. Vespa P, Martin NA, Nenov V et al. Delayed increase in extracellular glycerol with post-traumatic electrographic epileptic activity: Support for the theory that seizures induce secondary injury *Acta Neurochir, Suppl* 2002;81:355-357.
54. Vespa PM, Nuwer MR, Nenov V et al. Increased incidence and impact of nonconvulsive and convulsive seizures after traumatic brain injury as detected by continuous EEG in the intensive care unit. *J Neurosurg*. 1999;91:750-760.
55. Vespa PM, Miller C, McArthur D et al. Nonconvulsive electrographic seizures after traumatic brain injury result in a delayed, prolonged increase in intracranial pressure and metabolic crisis. *Crit Care Medicine*. 2007;35:1-7.
56. Vespa PM, McArthur DL, Xu Y et al. Nonconvulsive seizures after traumatic brain injury are associated with hippocampal atrophy. *Neurology*. 2010;75(9):792-8.
57. Vespa P, Tubi M, Claassen J et al. Metabolic Crisis occurs with Seizures and Periodic Discharges after Brain Trauma. *Ann Neurol*. 2016 Jan 27.
58. Wang H, Xin T, Sun X et al. Post-traumatic seizures--a prospective, multicenter, large case study after head injury in China. *Epilepsy Res*. 2013;107(3):272-8.
59. Wang B, Prastawa M, Irimia A et al. Analyzing imaging biomarkers for traumatic brain injury using 4D modeling of longitudinal MRI. *Proc IEEE Int Symp Biomed Imaging*. 2013;31:1392-1395.
60. Wang J, Qiu S, Xu Y et al. Graph theoretical analysis reveals disrupted topological properties of whole brain functional networks in temporal lobe epilepsy. *Clin Neurophysiol*. 2014;125(9):1744-56.

61. Wang J, Tan L, Tian Y et al. Circulating microRNAs are promising novel biomarkers for drug-resistant epilepsy. *Sci Rep*. 2015 May 18;5:10201.
62. Waziri A, Claassen J, Stuart RM et al. Intracortical electroencephalography in acute brain injury. *Ann Neurol*. 2009;66:366–377.
63. White A, Williams PA, Hellier JL et al. EEG spike activity precedes epilepsy after kainate-induced status epilepticus. *Epilepsia*. 2010;51(3):371-83
64. Worrell GA, Jerbi K, Kobayashi K et al. Recording and analysis techniques for high-frequency oscillations. *Prog Neurobiol*. 2012;98:265-278.
65. Wright MJ, McArthur DL, Alger JR et al. Early metabolic crisis-related brain atrophy and cognition in traumatic brain injury. *Brain Imaging Behav*. 2013;7(3):307-15.
66. Wulsin DF, Gupta JR, Mani R et al. Modeling electroencephalography waveforms with semi-supervised deep belief nets: fast classification and anomaly measurement. *J Neural Eng*. 2011;8(3):036015.
67. Xu Y, McArthur DL, Alger JR et al. Early nonischemic oxidative metabolic dysfunction leads to chronic brain atrophy in traumatic brain injury. *J Cereb Blood Flow Metab*. 2010;30(4):883-94.
68. Yuh EL, Mukherjee P, Lingsma HF et al. TRACK-TBI Investigators. Magnetic resonance imaging improves 3-month outcome prediction in mild traumatic brain injury. *Ann Neurol*. 2013;73(2):224-35.