

Hands-On: FukuiGrid - Fukui Functions $f^-(r)$ by Interpolation

Nicolás F. Barrera

November 6, 2025

1 Preparing the Input Files

Before executing the code, ensure that you have the required input files in the working directory. You should have four CHGCAR files corresponding to the system at different charge states:

- CHGCAR for the neutral system ($\delta N = 0.0$).
- CHGCAR files for the system with fraction charges ($\delta N = -0.05, -0.10, -0.15$).

2 Example: TiC (001) Surface

To run FukuiGrid, execute the following command in the terminal:

```
python FukuiGrid.py
```

Upon execution, the following main menu will be displayed:

```
***** Main Menu *****
1 — Fukui Function via Interpolation
2 — Fukui Potential via Electrodes
3 — Fukui Potential via SCPC
4 — Process Grid Data
5 — Perturbative Expansion
6 — Exit
```

Choose an option: 1

Select option 1 for Fukui Function via Interpolation:

```
Chose option 1: Fukui Function via Interpolation.
```

```
11 Electrophilic Fukui function  $f^-(r)$ .
```

```
12 Nucleophilic Fukui function  $f^+(r)$ .
```

Choose an option: 11

If you select option 11, you will be prompted to choose the δN values:

```
Do you want to use the default  $\delta N [-0.15, -0.10, -0.05, 0.0]$ ?
```

```
111 Yes, I would like to use the default array
```

```
112 No, I prefer to define my own values
```

Choose an option: 111

If you select option 111, you will be prompted to enter the CHGCAR files for the δN default values:

```
Name CHGCAR files with  $\delta N: -0.15, -0.10, -0.05$ , and  $0.0$ 
```

```
Enter name of file 1:
```

Once the calculation is complete, the output file “CHGCAR_FUKUI.vasp” will be generated. This file contains the computed Fukui function mapped onto the original charge density grid.

For the files available at <https://github.com/cacarden/FukuiGrid/tree/main/examples/Interpolation/fukui->, which correspond to the TiC (001) surface, the computed Fukui function $f^-(r)$ is obtained and is shown in Fig. 1.




Figure 1: Isosurface representation of $f^-(r)$ with iso value $0.002 a_0^{-3}$.