

3D Cuboid Labeling

Leo

November 18, 2016

1 Introduction

Given an image, we want to recover the inherent 3D structure behind it. For simplicity, we often treat the scene as a cuboid. Such approximation is quite enough for applications such as environment map generation. To label each face of the cuboid, we should firstly set up the 3 axes, as Figure 1 shows. p_0 is the intersection of the 3 axes and the vector $\overrightarrow{p_0p_1}$, $\overrightarrow{p_0p_2}$ and $\overrightarrow{p_0p_3}$ indicate the direction of x , y and z . So how do we get the 3D positions of these points based on those projected coordinates in 2D?




Figure 1: A image with labeled axes.

2 Inference

Assume the width and height of the image are w and h respectively. The focal length of the camera is f . Applying the projection transformation to a 3D point

$[X, Y, Z]$, we get

$$\begin{bmatrix} 2f/w & 0 & 0 & 0 \\ 0 & 2f/h & 0 & 0 \\ 0 & 0 & a & b \\ 0 & 0 & -1 & 0 \end{bmatrix} \begin{bmatrix} X \\ Y \\ Z \\ 1 \end{bmatrix} = \begin{bmatrix} 2fX/w \\ 2fY/h \\ aZ + b \\ -Z \end{bmatrix} \sim \begin{bmatrix} -2fX/(wZ) \\ -2fY/(hZ) \\ -(aZ + b)/Z \\ 1 \end{bmatrix}, \quad (1)$$

where $a = -\frac{F+N}{F-N}$, $b = -\frac{2FN}{F-N}$, F and N are the near and far plane. The 2D position $[u, v] = [-2fX/(wZ), -2fY/(hZ)]$ is the projected coordinate of $[X, Y, Z]$. The labeled 2D position $[u, v]$ is known, so we have

$$X = \left(-\frac{w}{2f}\right)uZ \quad (2)$$

$$Y = \left(-\frac{h}{2f}\right)vZ. \quad (3)$$

Back to Figure 1, assume the 2D position of the labeled point is $p_k = [u_k, v_k]$ and the corresponding 3D position is $P_K = [X_k, Y_k, Z_k]$. With $\overrightarrow{p_0 p_1} \perp \overrightarrow{p_0 p_2}$, we have

$$(X_1 - X_0)(X_2 - X_0) + (Y_1 - Y_0)(Y_2 - Y_0) + (Z_1 - Z_0)(Z_2 - Z_0) = 0. \quad (4)$$

Substituting Equation 2 and 3 into 4:

$$\begin{aligned} & \left(\frac{w}{2f}\right)^2 (u_1 Z_1 - u_0 Z_0)(u_2 Z_2 - u_0 Z_0) + \\ & \left(\frac{h}{2f}\right)^2 (v_1 Z_1 - v_0 Z_0)(v_2 Z_2 - v_0 Z_0) + \\ & (Z_1 - Z_0)(Z_2 - Z_0) = 0, \end{aligned} \quad (5)$$

or

$$\begin{aligned} & \left(\frac{w}{2f}\right)^2 (u_1 u_2 Z_1 Z_2 - u_0 u_2 Z_0 Z_2 - u_0 u_1 Z_0 Z_1 + u_0^2 Z_0^2) + \\ & \left(\frac{h}{2f}\right)^2 (v_1 v_2 Z_1 Z_2 - v_0 v_2 Z_0 Z_2 - v_0 v_1 Z_0 Z_1 + v_0^2 Z_0^2) + \\ & (Z_1 Z_2 - Z_0 Z_2 - Z_0 Z_1 + Z_0^2) = 0. \end{aligned} \quad (6)$$

Define $C_{mn} = \left(\frac{w}{2f}\right)^2 u_m u_n + \left(\frac{h}{2f}\right)^2 v_m v_n + 1$ and substitute it to Equation 6:

$$C_{12} Z_1 Z_2 - C_{02} Z_0 Z_2 - C_{01} Z_0 Z_1 + C_{00} Z_0^2 = 0, \quad (7)$$

i.e.

$$(C_{12} Z_2 - C_{01} Z_0) Z_1 = (C_{02} Z_2 - C_{00} Z_0) Z_0. \quad (8)$$

Symmetrically, we also have

$$(C_{23} Z_3 - C_{02} Z_0) Z_2 = (C_{03} Z_3 - C_{00} Z_0) Z_0 \quad (9)$$

$$(C_{31} Z_1 - C_{03} Z_0) Z_3 = (C_{01} Z_1 - C_{00} Z_0) Z_0. \quad (10)$$

According to Equation 10, $Z_3 = \frac{(C_{01} Z_1 - C_{00} Z_0) Z_0}{C_{31} Z_1 - C_{03} Z_0}$. Substituting it to Equation , we can express Z_2 using Z_0 and Z_1 . Then Substituting Z_2 into Equation 8 (I

use Matlab for such tedious work), we get

$$\begin{aligned}
& (C_{01}^2 C_{23} - C_{01} C_{03} C_{12} - C_{01} C_{02} C_{31} + C_{00} C_{12} C_{31}) Z_1^2 + \\
& (C_{01} C_{02} C_{03} - C_{00} C_{01} C_{23} + C_{01} C_{02} C_{03} - C_{00} C_{01} C_{23}) Z_1 Z_0 + \\
& C_{00}^2 C_{23} - C_{00} C_{02} C_{03}) Z_0^2 = 0. \quad (11)
\end{aligned}$$

Solving such a quadratic equation, we get Z_1 . Finally there are 2 sets of solution for Equation 8 - 10. We can choose either one.