KEY TO COMMON TREES IN PACIFIC SPIRIT PARK

Adapted from Straley, G.B. and R.P. Harrison. 1987. An Illustrated Flora of the University Endowment Lands. Botanical Garden, Technical Bulletin #12. University of British Columbia, Vancouver B.C.

A key is a tool for identifying plants. Characteristics of the plant are described in couplets (e.g. 1a and 1b). For each species, only one of the two statements is accurate. For each accurate description, follow the instructions indicated in the right column; continue to the next pair of couplets or note the identification of the plant. Ultimately you will determine the genus and species of each plant using this key or by referring to other field guides. Remember to follow protocol when using scientific names of plants. * indicates non-native species.

1a. 1b.		eaves needle or scale-like; seeds in cone or berry-like structure				
Divisi	on Pinop	hyta (C	Conifers)			
2a.	Leaves scale-like, smallest branchlets flattened; bark thin; reddish-brown, shredded; tree with upright stem, mature trees large and dominate the canopy; seeds in cones					
2b.	Leaves 4a. 4b.	Single bright small	Is seed in a fleshy red berry-like structure; needles sharp pointed, green on both sides of needle, with short stalk; tree with reddish to purplish-brown bark			

Division Magnoliophyta (Broadleaf trees)

8a	Broadleaved trees, may be deciduous or evergreen tree; leaves simple						
9a.	Leaves	Leaves spiny, simple, deciduous or evergreen; flowers white; fruits red;					
	bark sr	nooth, s	mall shrubby tree				
9b.	Leaves	s not spi	ny, deciduous in autumn	10			
	10a.		s lobed	11			
		11a.	Large leaves; 5 deep lobes beyond middle of leaf; bark greyish-brown, narrow scaly ridges, mature trees medium diameter, dominate the canopy	Acer macrophyllum (Bigleaf maple)			
		11b.	Leaves small, almost circular; 7-9 shallow lobes; bark smooth, thin, greenish to reddish-brown, small tree often with multiple stems, lower canopy				
	10b.	Leaves	s toothed				
		12a.	Coarsely toothed, leaves tapered from the middle to both ends; remain green until they fall in autumn; bark smooth, light grey; mature trees medium diameter, dominate the canopy				
		12b.	Leaves almost as wide as long, gradually tapered to tip; bright yellow in autumn; mature bark greyish-brown, deeply furrowed; mature trees medium diameter	Populus trichocarpa (Black cottonwood)			
8b.	Broadleaved tree, deciduous in autumn; leaves compound						
	13a.	Leaves	s in alternate pairs, 9-17 leaflets, coarsely toothed; bark smooth, brown;				
			small tree, lower canopy				
-	13b.	Leaves in opposite pairs, 5-7 leaflets, sharply toothed; bark smooth, greenish					
		small 1	ree often with multiple stems, lower canopy	Sambucus racemosa (Red elderberry)			