

DNA MARKER-ASSISTED BREEDING AND GENETIC TRANSFORMATION FOR PRODUCING SALT-TOLERANT RICE FOR BANGLADESH

ZEBA I. SERAJ, U.S. MAHZABIN AMIN AND M. SAZZADUR RAHMAN¹

Department of Biochemistry and Molecular Biology, University of Dhaka, Dhaka 1000, Bangladesh

¹*Plant Physiology Division, Bangladesh Rice Research Institute, Gazipur, Bangladesh*

ABSTRACT

Bangladesh is very small fertile country with a total land area of 14.8 Mha and cultivable area of 9.1 Mha (62%). The total irrigated land area is 52% with a cropping intensity of 181%. The share of agriculture in the GDP is 19.95%. The paddy production has so far been almost adequate for the growing population of rice eaters at 33.5 Mt (~150 million people), with the exception of import of ~1 Mt to mitigate floods or similar disasters. Another 100 million is likely to be added to the population, before stabilizing by the middle of this century. Therefore rice production has to nearly double. Abiotic stresses like salinity affects 1.2 Mha in the South. Drought affects 2.7 Mha in the North. In addition there is 1 Mha of flash flood-prone areas and 3.2 Mha of rainfed low-lying areas in Bangladesh.

Use of biotechnological tools like DNA marker-assisted breeding with the help of IRRI and currently funded by the Gates Foundation with the project STRASA (Stress tolerant Rice for Africa and South Asia) has resulted in the release of submergence tolerant rice like BRRIdhan 51 and 52. Some traditionally bred drought tolerant varieties as well as IR64 introgression lines with QTLs from drought tolerant donors have been produced and are in the pipeline. Our own work at Dhaka University helped incorporate salinity tolerance QTLs (*Saltol*) into farmer-popular BR11 and BRRIdhan 28, which are currently undergoing field trials in the South of Bangladesh. STRASA is also funding for pyramiding of multiple traits into farmer-popular rice through marker-assisted breeding. For example, the submergence tolerance trait, *Sub1* is being combined with each of stagnant flood tolerance (SF), anaerobic germination (AG), drought tolerant yield (DTY) and salinity tolerance (*Saltol*). Marker-assisted breeding for introgression of the kinase gene, *Pstol1*, which confers tolerance of phosphorous deficiency and drought and results in 20% increase in yield has just been initiated for Bangladesh.

Rice genetic transformation for incorporation of genes with multiple downstream effects has been shown to be effective in producing field level tolerance to drought, salinity and providing water-use efficiency. Moreover, tolerant traits coded for by different mechanisms can be pyramided providing more durable tolerance. Some of the target genes are those for antiporters, helicases and transcription factors like, HARDY and SNAC1. These genes are being cloned and incorporated into rice at Dhaka University. Some of these are collaborative efforts like those of Pea DNA Helicase with ICGEB, India.

INTRODUCTION

Tolerance to abiotic stresses such as drought, salinity or flooding is controlled by quantitative trait loci (QTLs) or multiple genes. In order to produce tolerant crops, for example, rice, these genes need to be combined with those responsible for good agronomic performance resulting in high yielding abiotic stress-tolerant genotypes suitable for cultivation. Breeding for high yield in rice has been in practice for several decades now and with help and inputs from the International Rice Research Institute (IRRI), the Bangladesh Rice Research Institute (BRRI) has produced at least 50 high yielding rice cultivars ⁽¹⁾. Climate change scenarios and scarcity of water resources has however now made introgression of stress tolerance genes necessary into these high-yielding cultivars.

One approach is the use of DNA markers associated with the tolerance genetic loci to guide introgression of the latter into the background of a high yielding cultivar. This is referred to as DNA-marker assisted backcrossing and involves the crossing of a donor genotype (having the tolerance genetic locus) with the high-yielder or the farmer-popular. This is followed by 2-3 backcrosses with the high-yielder, while selecting progenies having the tolerance loci aided by DNA markers. The backcross with the high-yielder ensures retention of its genetic background and the good agronomic traits associated with it. One very successful example of this approach was the introgression of the submergence tolerance locus or *Sub1A* from the rice landrace FR13A into the agronomically superior Swarna and BR11 ^(2, 3). These two genotypes were released as BRRIdhan 51 and 52 respectively in Bangladesh ⁽⁴⁾. One important factor in the success of this approach is that the targeted tolerant genetic locus has to be responsible for a high proportion of the phenotype of interest, since multiple loci may be responsible for the latter. For the *Sub1A* locus, this was precisely the case as it turned out to encode a transcription factor called ERF1 which in turn affected many downstream genes ^(5, 6). It is likely that the *Pstol1* locus encoding a protein kinase gene will also affect multiple downstream genes and be a good target for introgression of drought tolerance as well as efficient phosphorous absorption in a DNA marker-assisted backcross approach ⁽⁷⁾.

Knowledge of suitable DNA markers associated with abiotic traits of interest such as stagnant flooding (SF), anaerobic germination (AG) and drought tolerant yield (DTY) is essential for the success of the backcross breeding approach. Breeding programs for introgression and pyramiding of these traits are already underway ⁽⁸⁾. A major salinity tolerance locus, *Saltol*, was described in rice Chr 1 ^(9, 10) and has been the focus of salinity tolerance marker-assisted backcross breeding at IRRI, BRRI and Dhaka University (DU) and is briefly described in this paper.

Genetic transformation is another approach to producing abiotic stress tolerant crops such as rice. Initial efforts were made to overexpress single genes encoding rate limiting enzymes of osmolyte production. This resulted in limited improvement in tolerance at the vegetative state of several crops (reviewed by ⁽¹¹⁾). Overexpression of the Na/H antiporter which sequesters Na in the

vacuole provided good tolerance to dicots^(12, 13) but produced less dramatic effect in monocots like wheat⁽¹⁴⁾ and limited in rice⁽¹⁵⁾. Our own experience with the vacuolar Na/H antiporter transformation in rice has shown that its regulation is complex because it can produce 3 alternate transcripts from the single gene. We have also shown that inclusion of the 5' and 3' UTRs increases the level of tolerance compared to the cDNA only. Even so the vacuolar Na/H antiporter provides only moderate salt tolerance to rice (Seraj Z. I. unpublished results). Transformation with regulatory molecules like transcription factors and regulatory enzymes was more successful. These include the SNAC1⁽¹⁶⁾ and HARDY⁽¹⁷⁾ transcription factors as well as the RNA and DNA unwinding helicase enzyme^(18, 19). Work at Dhaka University, involving the cloning and transformation of these transcription factors and regulatory enzymes in rice, some of it in collaboration with the International Center for Genetic Engineering and Biotechnology (ICGEB, India) is described in this paper.

DNA MARKER-ASSISTED BACKCROSSING

Marker-assisted backcrossing (MABC) is now frequently used to introgress favorable alleles and major effect QTLs (Quantitative Trait Loci) in to mega varieties and elite genotypes for the improvement of complex abiotic and biotic stress tolerance traits specifically where conventional breeding fails to introgress stress tolerant traits to elite genotypes. *Saltol*, a major effect QTL for salinity tolerance was identified in rice chromosome 1, which was targeted for introgression to improve seedling stage salinity tolerance of two Bangladeshi mega rice variety BR11 (*Transplated Aman* or monsoon season) and BRRI dhan28 (BR28) (*Boro* or winter season). FL378, a tolerant F₈ Recombinant Inbred Line (RIL) was used as the donor to introgress *Saltol* alleles conferring seedling stage salinity tolerance into BR11 and BRRI dhan28 by MABC. Three-step marker aided selections i.e. Foreground, Recombinant and Background selection were employed to select progenies having precise QTL within a clean background of recurrent parent. In foreground selection 3 most tightly linked and robust SSR marker i.e. RM1287 (10.90 Mb), RM3412 (11.50 Mb) and RM493 (12.20 Mb) were used to locate the *Saltol* QTL in backcross progenies^(10, 20-22). Two to four SSR markers i.e. RM3627 (10.31 Mb) at the distal end of the QTL and RM10825 (13.30 Mb), RM10864 (14.20 Mb), RM562 (14.60 Mb), RM7075 (15.10 Mb) at the proximal end were used in recombinant selection to precisely limit the QTL segment (to reduce negative linkage drag). For the background selection 87-103 SSRs, InDel and gene-based markers were used to recover the recurrent genome⁽²³⁾. Three backcrosses and two selfs were done to transfer positive alleles of *Saltol* from FL378 into a clean and/or minimum background of BR11 and BRRI dhan28.

PROGENY SELECTION FOR TESTING AND RELEASE

Two and six Near Isogenic Lines (NILs) at BC₃F₃ stage were selected having 1.3-3.7 Mb introgression at the *Saltol* region with 97 to 99% recurrent parent genome for BR11 and BRRI dhan28 respectively. Fig 1 illustrates the MABC scheme for the development of introgression lines (BR11-*Saltol*) in a nutshell. Introgressing 1.3-3.7 Mb regions of target loci in to BR11 and

BRRI dhan28 slightly improved their overall tolerance at seedling stage (SES score) (Fig 2). All introgression lines however showed improved agronomic performance i.e. showed good stature, long panicles, high and dense grains, etc. The NILs looked very similar to the recurrent parents (BR11, BRRI dhan28), some of which however, showed greater yield potential in both saline and non-saline conditions. Tolerance to salinity is highly polygenic in nature; three different types of stresses (Osmotic, Ionic and Oxidative) are associated with the trait. So, in all likelihood a single QTL is not enough to significantly increase the tolerance level. Pyramiding of multiple QTL controlling different physiological mechanisms with different genetic background could help to achieve a higher level of salt tolerance.




Fig 1. Flow chart of 3-backcross step MABC scheme for the development of BR11-Saltol introgression lines with DNA markers used and the numbers of progenies genotyped in different backcross generation.



Fig 2. Photograph showing tolerance of introgression lines at seedling stage in hydroponics culture with stress @12 dSm⁻¹ in Net house condition. From L-R, Rows 1-3 are the negative controls without *Saltol*, rows 4-6 are lines with *Saltol* (show increased vigor). Rows 7, 8, 9 and 10 are respectively, the donor FL378, the sensitive parent BR28, the sensitive control IR29 and the tolerant control Pokkali

GENETIC TRANSFORMATION

The *Helicase* gene (*Pea DNA Helicase* or *PDH45*) construct was obtained from Dr. Narendra Tuteja of ICGEB under an MTA. This was transformed into *E.coli*, confirmed and subsequently transformed into *A. tumefaciens* for rice transformation as described in detail including testing of the transgenic rice, Binnatoa, for salt tolerance in Amin *et al.* ⁽¹⁹⁾. Since Binnatoa is a traditional cultivar with poor agronomic traits, the gene was backcrossed into three farmer-popular Bangladeshi rice cultivars, BRRIdhan 28, 29 and 47 as described below. Progenies showing good tolerance score at the seedling stage were advanced and stable inheritance of the transgene was shown upto the F₅ generation. These F₅ progenies are now being tested for reproductive stage stress tolerance.

The *SNAC1* RNA was isolated from the salt tolerant rice landrace Pokkali, converted to cDNA and cloned into the InVitrogen directional cloning ENTR vector, confirmed by sequencing and restriction digestion before recombining it into the Gateway *Agrobacterium*-compatible Binary Destination vector, pH7WG2 ⁽²⁴⁾. Standard rice transformation and

regeneration protocols were done, gene insertion and expression confirmed, plants advanced to T₂ and seedling salinity tolerance assays conducted as described in Amin et al. ⁽¹⁹⁾.

Similar protocols were used for cloning of the HARDY gene, except that the RNA was isolated from *Arabidopsis thaliana* Landsberg. All these transformations were done in the tissue culture responsive rice variety Binnatoa which is a traditional cultivar and hence not high-yielding. In order to transform the transcription factor genes into high yielding cultivars, we performed co-transformations with *Helicase* and *SNAC1* and *Helicase* and *HARDY* using an *in planta* transformation protocol according to Lin et al. ⁽²⁵⁾ (Please see below). The genes were co-transformed into BRRIdhan-27, BRRIdhan-29, BRRIdhan-43, BRRIdhan-49, BRRIdhan-52 and BRRIdhan-55. Transgenic T₂ progenies are being assessed for their seedling salinity tolerance as described in Amin et al. ⁽¹⁹⁾.

AGROBACTERIUM –MEDIATED TRANSFORMATION (TISSUE-CULTURE DEPENDENT)

Rice calli were transformed with *Agrobacterium tumefaciens* (strain LBA4404) containing the selected gene construct. The transformation method was carried out according to the method described by Khanna and Raina ⁽²⁶⁾ with some modifications ⁽²⁷⁾. The Bangladeshi tissue culture responsive cultivar Binnatoa, was used for transformation ⁽²⁸⁾.

AGROBACTERIUM-MEDIATED TRANSFORMATION (TISSUE-CULTURE INDEPENDENT OR IN PLANTA)

Mature rice seeds of individual variety were sterilized with 99% ethanol, 30% chlorox, Tween 20 and kept at 37°C. After 2 days the embryo region turned white. The embryonic apical meristem in the seed was pierced to a depth of 1-1.5 mm with a needle that had been dipped in the bacterial solution to inoculate gene construct in *Agrobacterium* into plumule where a shoot would later emerge as described in Lin et al. ⁽²⁵⁾ with the exception that 0.4% acetosyringone was added to the bacterial culture media. For co-transformation, a mixture of *Agrobacteria* solution, containing two gene constructs was used. The pierced seeds were then placed in a conical flask and soaked in the *Agrobacterium* inoculums and vacuum applied. The inoculated seeds were transferred onto petri dishes containing wet filter papers and incubated in the dark at 28°C for 6-7 days. The emerged seedlings were treated with 250mg/L carbenicillin solution, washed with ddH₂O and transferred to new petri dishes containing wet filter papers. Seedlings were then kept in light for 16 hours and in dark for 6 hours. When the seedlings turn green, they were transferred to hydroponic solution. After 2-3 days the hydroponic pots were transferred to net-house. When the seedlings were mature, they were transferred to soil. T₁ seeds only from those panicles whose flag leaf was resistant to hygromycin solution were germinated again in hygromycin. Seedlings which survived in hygromycin and were PCR-positive for the transgene were advanced to the T₂ generation and salinity tolerance tests performed.

SALT STRESS SCREENING FOR SELECTING TRANSGENICS WITH GOOD TOLERANCE SCORES

Transgenic plants were checked for tolerance level by seedling stage screening as described in detail in Amin *et al.*⁽¹⁹⁾. The germinated seeds of WT, transgenic lines and sensitive controls (IR29) were transferred to Hydroponics in netted, floating styrofoam in a completely randomized design, allowed to grow for 2 weeks and then salt stress up to 12 dS/m was applied gradually.⁽²⁹⁾ The EC of the solution was maintained at 12 dS/m until the end of the experiment. The pH of the non-aerated Yoshida⁽³⁰⁾ culture solution was adjusted to 5.0 every day and the culture solution was changed every 2 days. After 8–10 days, when the sensitive control IR29 in stress were dead, tolerance-related traits (Leaf Drying score, root length, shoot length) of all stressed as well as control plants were measured. Data for percent survival and total leaf area affected was recorded according to the standard evaluation system of rice at IRRI^(31; 32). The level of salinity tolerance was evaluated based mainly on the value of LDS, which is based on the percentage of leaf damage. The plants were scored according to the following scale: 1: highly tolerant (10%); 3: tolerant (10–30%); 5: moderately tolerant (30–50%); 7: moderately susceptible (50–70%) and 9: susceptible (70%)⁽³¹⁾. The chlorophyll content of the stressed and control transgenic shoots as well as WT was measured at this stage⁽³³⁾, as well as their dry weight after keeping them for 72 h at 70°C in a hot-air circulating oven (Honeywell, UK model DT200).

INTRODUCING THE HELICASE TRANSGENE INTO FARMER-POPULAR CULTIVARS BY BACKCROSSING AND SELECTION OF THE BEST TOLERANT LINES

The *Helicase* gene was backcrossed from the Binnatoa genetic background into three farmer-popular Bangladeshi rice cultivars, BRRIdhan 28, 29 and 47. The F₁ plantlets containing the *PDH45* transgene were selected by PCR and allowed to self up to the F₅ generation. The presence of the transgene was confirmed at each generation. Seedling stage screening was carried out at both F₃ and F₄ generations. Some transgenic progenies showed better tolerance than their sensitive parent (Fig 3). Significantly tolerant plants of each variety were selected and will be screened at reproductive stage to check the grain yield under salt stress.




Fig 3 : Transgenic rice with PDH45 performed better at physiological screening.(A) Seedlings of PDH45, PDH45×BR29 and BR29 after 12 dS/m salt stress in a hydroponic system. PDH45×BR29 seedling is more vigorous than the Wild type BR29. (B,C,D) SES Score of wild-type and transgenic rice seedlings after NaCl stress at 12 dS/m in hydroponics. Each bar represents the mean \pm SE (n = 18); $P < 0.05$

SELECTION OF THE BEST-PERFORMING LINES TRANSFORMED WITH THE TRANSCRIPTION FACTORS SNAC1 AND HARDY

Six transgenic rice lines containing the SNAC1 gene were selected by PCR and advanced to the T₃ generation. The presence of the transgene was confirmed at each generation. Depending on the expression levels of the transgene by RT-PCR, four lines have been selected for physiological screening. Among these four lines, progenies of one line showed better tolerance than the wild type (Fig 4).




Fig 4: Transgenic rice containing SNAC1 gene showed better tolerance. (A) SNAC1 gene containing transgenic seedlings, Wild Type (Binnatoa), Sensitive control IR29 after 12 dS/m salt stress in hydroponic system. (B)) SES Score of wild-type and transgenic rice seedlings after NaCl stress at 12 dS/m in hydroponics. Each bar represents the mean \pm SE (n = 18); P < 0.05

The presence of HARDY gene in the transgenic rice plants was confirmed by PCR and their expression checked by RT PCR. One of the three transgenic plants had higher expression of HARDY gene and was thus selected for further analysis. The co-transformed seedlings were screened by Hygromycin resistance assay for the presence of respective transgenes at the T_0 generation and further confirmed by PCR. T_1 seeds were collected from the hygromycin resistant PCR positive plants. The transgenic plants were further characterized by leaf disc senescence assay (Fig 5).



Fig 5: Leaf Senescence Assay at T_1 generation for the HARDY transformants

Acknowledgements: Funds for this work are gratefully acknowledged from the Generation Challenge Program, US Department of Agriculture (USDA) and BAS-USDA-PALS (Bangladesh Academy of Sciences-USDA program in Life Sciences).

REFERENCES

1. www.brri.gov.bd
2. Neeraja, C.N., R. Maghirang-Rodriguez, A. Pamplona, S. Heuer, B.C.Y. Collard, E.M. Septiningsih, G. Vergara, D. Sanchez, K. Xu, A.M. Ismail, and D.J. Mackill. 2007. A marker-assisted backcross approach for developing submergence-tolerant rice cultivars. *Theoretical Applied Genetics* 115:767–776.
3. Iftekharuddaula K. M., M. A. Newaz, M. A. Salam, H. U. Ahmed, M. A. A. Mahbub, E. M. Septiningsih, B. C. Y. Collard, D. L. Sanchez, A. M. Pamplona, D. J. Mackill. 2011. Rapid and high-precision marker assisted backcrossing to introgress the SUB1 QTL into BR11, the rainfed lowland rice mega variety of Bangladesh. *Euphytica* 178:83–97.
4. BRRI. 2011. *Adhunik dhaner chash* (a Bengali booklet for modern rice cultivation). 16th Edition. BRRI, Gazipur, Bangladesh.
5. Xu K, X. Xu, T. Fukao, P. Canlas, R. Maghirang-Rodriguez, S. Heuer, A. M. Ismail, J. Bailey-Serres, P. C. Ronald, D. J. Mackill(2006). Sub1A is an ethylene-response-factor-like gene that confers submergence tolerance to rice. *Nature* 442 (7103) :705-8.
6. Bailey-Serres J., T. Fukao, P. Ronald, A. Ismail, S. Heuer and D. Mackill (2010). Submergence Tolerant Rice: SUB1's Journey from Landrace to Modern Cultivar. *Rice* 3: 138-147.
7. Gamuyao R., J. H. Chin, J. Pariasca-Tanaka, P. Pesaresi, S. Catausan, C. Dalid, I. Slamet-Loedin I, E. M. Tecson-Mendoza, M. Wissuwa and S. Heuer (2012). The protein kinase Pstol1 from traditional rice confers tolerance of phosphorus deficiency. *Nature* 488 (7412): 535-9
8. Mackill D. J., A. M. Ismail, A. M. Pamplona, D. L. Sanchez, J. J. Carandang and E. M. Septiningsih (2010). Stress tolerant Rice varieties for adaptation to a changing climate. *Crop, Environment & Bioinformatics* 7:250-259
9. Bonilla, P.S., J. Dvorak, D. Mackill, K. Deal and G.B. Gregorio. 2002. RFLP and SSCP mapping of salinity tolerance genes in chromosome 1 of rice (*Oryza sativa* L.) using recombinant inbred lines. *The Philippines Agricultural Scientist* 85(1):64-74.
10. Islam, M.R. 2006. Validation and allelic diversity analysis of molecular markers to salinity tolerance in rice chromosome 1. Ph.D. Thesis, 2006, Department of Genetics and Plant Breeding, Bangladesh Agricultural University, Mymensingh, Bangladesh.
11. Agarwal P. K., P. S. Shukla, K. Gupta, B. Jha (2013). Bioengineering for salinity tolerance in plants: state of the art. *Mol Biotechnol* 54:102-23.
12. Zhang H-X. and E. Blumwald (2001). Transgenic salt-tolerant tomato plants accumulate salt in foliage but not in fruit. *Nature Biotech* 19:765-768
13. Zhang H-X, J. N. Hodson, J. P. Williams and E. Blumwald (2001). Engineering salt-tolerant Brassica plants: Characterization of yield and seed oil quality in transgenic plants with increased vacuolar sodium accumulation. *PNAS* 98:12832-12836

14. Xue Z-Y., D-Y. Zhi, G-P. Xue, H. Zhang, Y-X. Zhao and G-M. Xia (2004). Enhanced salt tolerance of transgenic wheat (*Triticum aestivum* L.) expressing a vacuolar Na⁺/H⁺ antiporter gene with improved grain yields in saline soils in the field and a reduced level of leaf Na⁺. *Plant Science* 167 (2004) 849–859
15. Fukuda A., A. Nakamura, A. Tagiri, H. Tanaka, A. Miyao, H. Hirochika and Y. Tanaka (2004). Function, Intracellular Localization and the Importance in Salt Tolerance of a Vacuolar Na⁺/H⁺ Antiporter from Rice. *Plant Cell Physiol* (2004) 45 (2):146-159.
16. Hu, H., M. Dai, J. Yao, B. Xiao, X. Li, Q. Zhang and L. Xiong (2006). "Overexpressing a NAM, ATAF, and CUC (NAC) transcription factor enhances drought resistance and salt tolerance in rice." *Proceedings of the National Academy of Sciences* 103(35): 12987-12992
17. Karaba, A., S. Dixit, R. Greco, A. Aharoni, K. R. Trijatmiko, N. Marsch-Martinez, A. Krishnan, K. N. Nataraja, M. Udayakumar and A. Pereira (2007). "Improvement of water use efficiency in rice by expression of HARDY, an Arabidopsis drought and salt tolerance gene." *Proceedings of the National Academy of Sciences* 104(39): 15270-15275.
18. Sanan-Mishra N., X. H. Pham, S. K. Sopory and N. Tuteja (2005). Pea DNA helicase 45 overexpression in tobacco confers high salinity tolerance without affecting yield. *Proc Natl Acad Sci USA* 102(2):509–514
19. Amin M., S. M. Elias, A. Hossain, A. Ferdousi, M. S. Rahman, N. Tuteja and Z. I. Seraj (2012). Over-expression of a DEAD-box helicase, PDH45, confers both seedling and reproductive stage salinity tolerance to rice (*Oryza sativa* L.). *Mol Breeding* (2012) 30:345–354
20. Ferdousi, A. 2008. Validation of DNA markers linked to salt tolerant traits in breeding populations and near isogenic lines of rice. MS thesis, Faculty of Biological Science, Department of Biochemistry and Molecular Biology, University of Dhaka, Dhaka, Bangladesh.
21. Thomson, M.J., M. de Ocampo, J. Egdane, M.A. Rahman, A.G. Sajise, D.L. Adorada, E. Tumimbang-Raiz, E. Blumwald, Z.I. Seraj, R.K. Singh, G.B. Gregorio and A.M. Ismail. 2010. Characterizing the Saltol Quantitative Trait Locus for Salinity Tolerance in Rice. *Rice* 3:148–160.
22. Alam, R., M.S. Rahman, Z. I. Seraj, M. J. Thomson, A. M. Ismail, E. Tumimbang-Raiz and G. B. Gregorio. 2011. Investigation of seedling-stage salinity tolerance QTLs using backcross lines derived from *Oryza sativa* L. Pokkali. *Plant Breeding* 130 (4):430-437.
23. Septiningsih, E.M., A. M. Pamplona, D. L. Sanchez, C. N. Neeraja, G. V. Vergara, S. Heuer, A. M. Ismail and D. J. Mackill. 2009. Development of submergence-tolerant rice cultivars: the Sub1 locus and beyond. *Annals of Botany* 103: 151–160.
24. Abdullah-Al-Emran, R. Malo, M. Amin, D. Chakraborty, A. Ferdousi and Z. I. Seraj (2010). Cloning and Transformation of the Transcription Factor SNAC1 from Rice (*Oryza sativa* L.) Landrace Pokkali. *Proceedings of the Sixth International Plant Tissue Culture and Biotechnology Conference*; Dec 3-5, 2010; BAPTC&B, Dhaka, Bangladesh.
25. Lin, J., B. Zhou, Y. Yang, J. Mei, X. Zhao, X. Guo, X. Huang, D. Tang and X. Liu (2009). "Piercing and vacuum infiltration of the mature embryo: a simplified method for

Agrobacterium-mediated transformation of indica rice." Plant cell reports 28(7): 1065-1074.

26. Khanna H. K., Raina SK (1999). Agrobacterium-mediated transformation of indica rice cultivars using binary and superbinary vectors. Aust J Plant Physiol 26:311–324
27. Rasul N (2005). Isolation and characterization of vacuolar *Oryza sativa* Na⁺/H⁺ antiporter and its expression in transgenic rice. PhD thesis, University of Dhaka
28. Rasul N, Ali K, Islam R, Seraj ZI (1997). Transformation of an indica rice cultivar Binnatoa with *Agrobacterium tumefaciens*. Plant Tissue Cult 7:71–80
29. Ponnamperna F (1977.) Screening rice for tolerance to mineral stress. IRRI research paper series no. 6
30. Yoshida S, Forno DA, Cock JH, Gomez KA (1976.) Laboratory manual for physiological studies of rice, 3rd edn. International Rice Research Institute, Los Banos, pp 61–66
31. IRRI (1996) Standard evaluation system manual, 4th edn. International Rice Research Institute, Los Banos
32. Gregorio G, Senadhira D, Mendoza R (1997) Screening rice for salinity tolerance. International Rice Research Institute, Los Banos. IRRI discussion paper series no. 22
33. Vernon L (1969) Spectrophotometric determination of chlorophylls and phaeophytins in plant extracts. Anal Chem 32:1144–1150

