





## Protocol and Pitfalls

### Laser Coagulation of Anal Fistula (LCAF)

3-D-Endosonografie	OP-Methode	Laser in der Fistel	Postoperativ
			

#### Before surgery:

Blood thinner: discontinue (controverse discussion)  
Bowel preparation: only evacuation with  
2 times Lecicarbon suppositories (CO2 gas) day of surgery

#### During surgery:

Antibiotics: single shot, cefuroxim + metronidazol  
Anaesthesia: general anaesthesia  
Local anaesthesia: 3 ml bupivacaine 0,5% with adrenaline 1:10.000 (not in recurrence = scar)  
Wavelength: 1470 nm (810 or 980 nm going deeper into tissue, damage of muscle possible)  
Probe: 600 µm fiber with special glas top with 360 spreading laser light  
Power: 10 Watts  
Shots: each 5 mm 3 sec, beginning at level of internal fistula opening, gently pulling until adherence loosed, remove gently pasting necrosis from glas top, than pushing probe into fistula tract again until you feel resistance  
In cases of fistula cavities you'll not feel resistance. Therefore try fan-shaped burning to reach every edge of cavity.

#### After surgery:

What normally happens: slight pain, purulent discharge until 6 weeks

Bandage: shower with water 3 times a day and dexpanthenol ointment on a gauze compress  
Sick note: 7 days, but man with own business will go for business next day  
Pain killer: Ibuprofen 600 if necessary  
Bowel movement: Psyllium (Plantago ovata) 5 gram after breakfast 3 weeks, not liquid bm  
Sports: no heavy weights and contact sports for 3 weeks  
no jogging, cycling for 1 week  
no swimming in pools until wound closure

#### Pitfalls:

- You can't see an internal fistula opening and if you push probe into fistula tract it stops beneath mucosal surface: Don't force and start with burning.
- Strong bleeding within first 3 weeks are very seldom possible.
- After 6 weeks external fistula opening heals quicker than fistula tract (in recurrences regularly happen): reopen in local anaesthesia, do Curettage if necessary (or LCAF once again)
- After doing LCAF two times fistula tract can changing from transsphincteric to subanodermal: Cut fistula tract.
- In cases of horse shoe fistula set a loop to get shortest straight path and burn side branches first.
- Crohn's disease is no contraindication if it is well treated with drugs simultaneously
- There will be a more or less synclinal scar in place of internal fistula opening.