
(a)
DCCM — BIN1 BAR (S1, 200 ns metadynamics)
PROA
PROB
PROB
PROA
Residue index
Residue index
Cross-correlation


(b)
BIN1 BAR — Site688/680 Pocket Opening vs Crescent Curvature
200 ns metadynamics
A BAR pocket opening vs crescent geometry
CV1 (cross-chain)
Tip-tip distance
CV1 (nm)
N-term tip-to-tip distance (nm)
Time (ns)
B Pearson r = -0.629 (p = 1.1e-44)
r = -0.629
N-term tip-to-tip distance (nm)
CV1 — cross-chain distance (nm)
Time (ns)


(c)
DCCM — PICALM ANTH (S3, 200 ns metadynamics)
PROA
PROA
Residue index
Residue index
Cross-correlation


(d)
PICALM ANTH — Site473 Pocket Opening vs PIP2-Binding Helix Displacement
200 ns metadynamics
A PICALM site473 opening vs PIP2-helix displacement
cv_site4 (pocket opening)
PIP2-helix RMSD
cv_site4 (nm)
PIP2-helix RMSD (Å)
Time (ns)
B Pearson r = 0.718 (p = 0.0e+00)
r = 0.718
PIP2-binding helix RMSD (Å)
cv_site4 — site473 opening (nm)
Time (ns)


(e)
PICALM ANTH — ΔDCCM: Apo vs Ibufenac Holo
Apo: 200 ns metadynamics | Holo: complex MD rep1+rep5 (100% engaged)
A Apo (metadynamics)
Cyan: PIP2-helix (19–40)
Yellow: site473 contacts
B Holo (Ibufenac, rep1+rep5)
C ΔDCCM (holo − apo)
Residue
Residue
Residue
C
C
ΔC