

GaviLyte™-C

generic of CoLyte®*

(PEG-3350, sodium sulfate, sodium bicarbonate, sodium chloride and potassium chloride)

GaviLyte-C, PEG-3350 and Electrolytes for Oral Solution, is an effective bowel prep kit used prior to colonoscopy or barium enema X-ray examination. [Click here to View Important Safety Information.](#)

Package contents:

- 4-liter container
- Flavor pack
- GaviLyte-C powder



Step by Step

GaviLyte™-C

(PEG-3350, sodium sulfate, sodium bicarbonate, sodium chloride and potassium chloride)



If choosing to add provided flavoring, add the lemon flavor pack to the GaviLyte-C powder in the 4L container before reconstituting and shake well to incorporate flavoring into the powder.



Add lukewarm drinking water up to the fill line mark on the bottle. Solution should be 4L upon reconstitution.

TIP: Make sure to fill all the way to the fill line.



Cap bottle securely and shake vigorously to dissolve ingredients.



Adults should drink one 8-ounce (240mL) cup of solution rapidly every 10 minutes until the watery stool is clear and free of solid matter. This is usually complete after drinking 3L to 4L. Rapid drinking of each portion is preferred to drinking small amounts continuously. The first bowel movement should commence in approximately one hour.

Tips for Effective Use



Patients should not have solid food for at least 2 hours before solution is given. Ideally, the patient should fast 3-4 hours prior to GaviLyte-C administration.



Lukewarm water facilitates dissolution.



The solution is more palatable if chilled before drinking. However, chilled solution is not recommended for infants.



GaviLyte-C should be used within 48 hours of reconstituting. Discard any solution that is not used within that time frame.