


CURRICULUM VITAE

Dr. dr. Dovy Djanas, Sp.OG (K-FM)



- Tempat / Tgl lahir : Padang / 19 April 1968
- S1 : Kedokteran Universitas Andalas.
- Spesialis 1 : Obgin FK Unand-RSUP Dr.M. Djamil Padang, 2005
- Spesialis 2 : Fetomaternal FKUI-RSCM Jakarta, 2011
- S 3 : Biomedik FK Unand, 2018
- Pendidikan S2 MARS
- Klinisi di Klinik Morula IVF RS Bunda BMC Padang
- Staff Obgin FK-Unand / RSUP Dr. Djamil Padang
- Kepala Instalasi Rawat Inap Ibu dan Anak RSUP Dr. M. Djamil Padang
- Ketua POGI Sumbar 2019 - 2021
- Pemakalah Tetap Media Koran Padang Ekspres 2012 sampai sekarang

DETEKSI DINI CACAT LAHIR PADA ANC



DOVY DJANAS

DIVISI FETOMATERNAL BAGIAN
OBSTETRI DAN GINEKOLOGI
FK UNAND/ RS INTERNASIONAL DR
M DJAMIL PADANG



PENDAHULUAN

Lebih dari 4 juta anak dilahirkan dengan cacat lahir setiap tahun


Merupakan tantangan kesehatan klinis dan publik yang cukup besar karena dampaknya di seluruh dunia terhadap kesehatan populasi

WHO :

Dari 2,68 juta kematian bayi, 11,3% disebabkan oleh kelainan bawaan (INFODATIN Kemenkes RI, 2014-2018)

Di Indonesia, hasil Riskesdas tahun 2007 menjelaskan kelainan bawaan menjadi salah satu penyebab kematian bayi

Gambar 1. Proporsi Penyebab Kematian Bayi di Dunia Tahun 2015



Gambar 2. Penyebab Kematian Bayi 0 - 6 Hari dan 7 - 28 Hari, Riskesdas 2007



Order	Birth defects	Cases	Prevalence(1/10000)
1	Neural tube defects	126	20.1
2	Congenital heart disease	107	17.1
3	Cleft lip and palate	101	16.2
4	Congenital hydrocephalus	62	9.9
5	Polydactylia	53	8.5
6	External ear malformation	27	4.3
7	Limb reduction defect	26	4.2
8	Inguinal hernia	25	4.0
9	Congenital hemangiomas	18	2.9
10	Talipes equinovarus	17	2.7

Organs or systems	Cases	Proportion(%)
Central nervous system	193	19.8
Face or eye	176	18.0
Extremities	129	13.2
Cardiovascular	118	12.1
Skeletal	76	7.8
Genitourinary tract	60	6.1
Thoracic or abdominal wall	55	5.7
Gastrointestinal tract	41	4.2
Tumors	10	1.0
Other	118	12.1
Total	976	100.0

SITUASI NASIONAL

March of Dimes Birth Defects Foundation pada tahun 2006 merilis *Global Report on Birth Defect* prevalensi bayi dengan kelainan bawaan di Indonesia adalah 59,3 per 1.000 kelahiran hidup

Gambar 3. Prevalensi Bayi dengan Kelainan Bawaan per 1000 Kelahiran Hidup di Asia Tenggara (Tahun 1980 - 2001)



Sumber : *Global Report on Birth Defects, March of Dimes Birth Defects Foundation, 2006*

CACAT LAHIR

NBBD

Newborn Birth Defects



Suatu kondisi pada saat bayi lahir yang dilihat tepat ketika bayi lahir

Kelainan struktur pada janin yang terjadi ketika janin masih berkembang didalam perut ibu

Cacat Lahir Struktural : Mayor atau Minor

Cacat lahir INTERNAL / EKSTERNAL




Sistem saraf atau masalah otak, masalah sensorik, gangguan metabolisme, gangguan degeneratif



@DOCTORS_OFFICIAL

APA PENYEBABNYA???

- Masalah genetik
- Masalah kromosom
- Infeksi
- Terpapar obat, bahan kimia, atau agen lain selama kehamilan
- **Gangguan Makronutrien dan Mikronutrien selama kehamilan**



Defisiensi VIT A

Defisiensi Asam Folat

Defisiensi Zn


Defisiensi vitamin E

TABEL KEBUTUHAN ASUPAN MAKRONUTRIEN DAN MIKRONUTRIEN PADA IBU HAMIL DAN TIDAK HAMIL


Nutrient	Recommendations for Interventions/Supplement Use ¹	Non Pregnant Adult Females (19–50 years) ²	Pregnant Adult Females (19–50 years) ²
Macronutrients			
Energy	Energy restriction reduces GWG but could adversely affect birthweight and is currently not recommended in pregnancy	EER (kcal/day) ³ = 354 – (6.91 × age [year]) + PA × [(9.36 × weight [kg]) + (726 × height [m])]	Non pregnant EER + 340 and 452 kcal/day in 2nd and 3rd trimesters
Protein	Balanced energy / protein supplements (≤ 25% total energy from protein) are recommended only in undernourished women to prevent stillbirth and SGA	0.8 g/kg/day (46 g/day)	0.8 increasing to 1.1 g/kg/day in 2nd half of pregnancy (71 g/day)
Total fibre ⁴	Fibre-rich diet may reduce preeclampsia and GDM but no specific recommendations are currently available; fibre supplements can be used to relieve constipation if diet modification is unsuccessful	14 g/1000 kcal (or 25 g/day)	14 g/1000 kcal or (or –28 g/day to account for GWG)
Carbohydrates (GI and GL)	Low GL or GI diets may be beneficial for women at risk of GDM or LGA but can increase risk of SGA. No specific recommendations are currently available	130 g/day of carbohydrates	175 g/day of carbohydrates
Essential fatty acids ⁴ (linoleic acid [n-6] and α-linoleic acid [n-3])	n-3 PUFAs may prevent preterm birth but can increase post-term birth and LGA. No specific recommendations are currently available	12 g/day (linoleic) 1.1 g/day (α-linoleic)	13 g/day (linoleic) 1.4 g/day (α-linoleic)
Micronutrients			
Folate/folic acid	Recommended (400 µg/day) from preconception until at least 12 weeks to prevent NTDs	400 µg/day	600 µg/day
Vitamin A	Not recommended except in areas with severe deficiency/night blindness	700 µg/day	770 µg/day
Thiamine (B ₁) Niacin (B ₂) Riboflavin (B ₂) Pyridoxine (B ₆) Cyanocobalamin (B ₁₂)	B-complex vitamins are not recommended to improve pregnancy outcomes until further evidence is available	1.1 mg/day 14 mg/day 1.1 mg/day 1.3 mg/day 2.4 µg/day	1.4 mg/day 18 mg/day 1.4 mg/day 1.9 mg/day 2.6 µg/day
Vitamin C Vitamin E	Not recommended until further evidence relating to safety and PROM is available	75 mg/day 15 mg/day	85 mg/day 15 mg/day
Vitamin D ⁴	Not recommended for improving pregnancy outcomes but should be given to women with deficiency (200 IU/day)	5 µg/day	5 µg/day
Calcium ⁴	Recommended (1.5–2.0 g/day) to prevent hypertensive disorders in women with low dietary calcium intake or who are at high risk of hypertension	1 g/day	1 g/day
Iodine	Recommended only in women at high risk to prevent IDD (i.e., in countries where < 20% of households have access to iodized salt)	150 µg/day	220–250 µg/day
Iron	Recommended (30–60 mg/day) to prevent maternal anaemia, puerperal sepsis, LBW and preterm birth	18 mg/day	27–60 mg/day
Zinc	Not recommended for improving pregnancy outcomes until more rigorous research is available	8 mg/day	11 mg/day
Alcohol	Not recommended during pregnancy until safe upper limits are established	NA	None
Caffeine	Reducing intake is recommended in women with high caffeine intake (> 300 mg/day) to prevent pregnancy loss and LBW infants	NA	<200 mg/day

TIMELINE KEHAMILAN

Human Development Timeline



Human Critical Periods of Development



WHO recommendations on antenatal care for a positive pregnancy experience



WHO GUIDELINES

- A. Intervensi nutrisi
- B. **Penilaian ibu dan janin**
- C. **Tindakan pencegahan**
- D. Intervensi untuk gejala fisiologis umum
- E. Intervensi sistem kesehatan untuk meningkatkan pemanfaatan dan kualitas ANC.

SKRINING CACAT LAHIR KETIKA ANTENATALCARE

TRIMESTER 1 pada 9–13 MINGGU KEHAMILAN

TRIMESTER 2 pada 14-18 MINGGU KEHAMILAN



Tes darah sederhana (Serum ibu)



Ultrasonografi Anomali

HASILNYA?

INCREASED RISK



NOT INCREASED RISK

TES DIAGNOSTIK

1 Tes prenatal noninvasif (NIPT)

2 Amniosentesis


3 Chorionic Villus Sampling (CVS)

4 Ultrasound resolusi tinggi


5 Alpha Fetoprotein (AFP)

6 Acetylcholinesterase (AChE)

Amniosentesis



Chorionic Villus Sampling (CVS)





APA SAJA TINDAKAN PREVENTIF???

INTERVENSI GIZI

- INTERVENSI DIET
- SUPLEMEN ZAT BESI DAN ASAM FOLAT
- SUPLEMEN KALSIMUM
- SUPLEMEN VITAMIN A
- SUPLEMEN SENG
- SUPLEMEN MULTIPLE MICRONUTRIENTS (MMN)
- SUPLEMEN VITAMIN B6 (PYRIDOXINE)
- SUPLEMEN VITAMIN E & C
- SUPLEMEN VITAMIN D
- MEMBATASI ASUPAN KAFEIN

Review

Macronutrient and Micronutrient Intake during Pregnancy: An Overview of Recent Evidence

Aya Mousa ^{1,*}, Amreen Naqash ² and Siew Lim ¹

¹ Monash Centre for Health Research and Implementation (MCHRI), School of Public Health and Preventive Medicine, Monash University, Melbourne VIC 3168, Australia; siew.lim1@monash.edu

² Department of Pharmaceutical Sciences, University of Kashmir, Srinagar, Jammu and Kashmir 190006, India; anaqash.scholar@kashmiruniversity.net

* Correspondence: aya.mousa@monash.edu; Tel.: +61-385-722-854

Received: 10 January 2019; Accepted: 14 February 2019; Published: 20 February 2019



Poor dietary intakes or deficiencies in key macronutrients and micronutrients can therefore have a substantial impact on pregnancy outcomes and neonatal health

FUNGSI MIKRONUTRIEN PADA PERIODE PERIKONSEPSI

Table 1 Main functions of micronutrients involved in the periconceptual period

Micronutrient	Function	References
Folate	Involvement in the DNA replication (cell cycle); methylation cycle (amino acids cysteine and methionine cycle)	FAO/WHO Report (2004a), Scholl and Johnson (2000)
Vitamin B ₁₂	Conversion of homocysteine to methionine as cofactor of the methionine synthase	Ryan-Harshman and Aldoori (2008)
Vitamin B ₆	Metabolism of amino acids, lipids, one-carbon units and glycogen as co-enzyme; pathways of gluconeogenesis, heme and neurotransmitter biosynthesis	Mackey <i>et al.</i> (2006)
Vitamin A	Growth and differentiation of a number of cells and tissues	Ross (2006)
Antioxidants	Defence systems against free radical species	Fang <i>et al.</i> (2002)
Iron	Haematopoiesis; nucleic acid metabolism; carrier of oxygen to the tissues by red blood cell haemoglobin; transport medium for electrons within cells; integrated part of important enzyme systems	Cook <i>et al.</i> (1992); Beard (2003), FAO/WHO Report (2004c), Wood and Ronnenberg (2006)
Zinc	Structural, regulatory and catalytic functions as cofactor for numerous metalloenzymes	McCall <i>et al.</i> (2000), Hambidge (2000)
Copper	Neurotransmission, neuropeptide maturation, oxidative phosphorylation, defence from free radical damage as cofactor for numerous cuproenzymes and copper-binding proteins	Turnlund (2006)

MIKRONUTRIEN DAN CACAT LAHIR

FOLAT

Larut dalam air, banyak dalam sayuran hijau berdaun, ekstrak ragi dan buah jeruk

Berfungsi sebagai koenzim dalam transfer satu karbon selama siklus metilasi dan integral untuk sintesis DNA dan neurotransmitter

Defisiensi folat menghasilkan akumulasi homocysteine, yang dapat meningkatkan risiko dan kehamilan yang buruk termasuk preeklampsia dan anomali janin

Suplementasi dengan asam folat selama prakonsepsi dan kehamilan dini sangat penting dan dapat mencegah 40-80% dari cacat tabung saraf seperti spina bifida

VITAMIN A

Vitamin yang larut dalam lemak berasal dari retinoid yang terbentuk sebelumnya atau provitamin karotenoid

Retinoid, seperti retinal dan asam retinoat, diperoleh dari sumber hewani termasuk telur, susu, hati, dan minyak hati ikan

Vitamin A tambahan diperlukan selama kehamilan untuk mendukung pertumbuhan dan pemeliharaan jaringan pada janin dan untuk menyediakan cadangan janin dan membantu metabolisme ibu

Vitamin B

B-complex termasuk didalamnya vitamin B1 (thiamine), B2 (riboflavin), B3 (niacin), B6 (pyridoxine) and B12 (cyanocobalamin)

Vitamin ini bertindak sebagai koenzim di beberapa jalur metabolisme perantara untuk pembangkit energi dan pembentukan sel darah

Kekurangan riboflavin dan niasin telah dikorelasikan dengan preeklampsia, kelainan jantung bawaan, dan bayi BBLR

ZINC

Kekurangan zinc pada awal kehamilan menyebabkan gangguan implantasi, aborsi dan malformasi janin, termasuk sumbing bibir dan langit-langit, malformasi otak dan mata, berbagai kelainan jantung, paru-paru dan urogenital sistem

COPPER

Teratogenisitas terkait defisiensi tembaga seperti swollen hidbrains, penyakit lepuh, penggumpalan darah dan distensi pembuluh darah besar

TAKE HOME MESSAGE

Cacat lahir adalah penyebab utama kematian bayi dan telah terjadi selama 25 tahun terakhir, menyebabkan 22% dari semua kematian bayi

Untuk memastikan awal yang sehat untuk bayi mereka, sebagian besar wanita sebaiknya menjalani beberapa bentuk skrining prenatal.

Skrining trimester pertama akurat dan efisien dalam praktik klinis

Kekurangan mikronutrien telah dikaitkan dengan risiko kehamilan yang sangat buruk, mulai dari infertilitas hingga cacat janin struktural dan penyakit jangka panjang

PELANTIKAN PENGURUS POGI
(Perkumpulan Obstetri dan Ginekologi Indonesia)
CABANG SUMATERA BARAT
Periode 2018-2021
PADANG, 20 MAREK 2019



TERIMA KASIH