

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS General Certificate of Education Ordinary Level

CANDIDATE NAME					
CENTRE NUMBER			CANDIDATE NUMBER		



PHYSICS 5054/04

Paper 4 Alternative to Practical

May/June 2008

1 hour

Candidates answer on the Question Paper.

No Additional Materials are required.

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use a soft pencil for any diagrams, graphs or rough working.

Do not use staples, paper-clips, highlighters, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer all questions.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

For Examiner's Use				
1	_			
2				
3				
4				
Total				

This document consists of $\bf 9$ printed pages and $\bf 3$ blank pages.



1 A student investigates the maximum height a ball reaches after bouncing on a hard surface. Fig. 1.1 shows the apparatus used.

For Examiner's Use




Fig. 1.1

(a) The ball is dropped from a height of 1.00 m above the hard surface.

State which part of the ball should be used when measuring its height above the surface.

.....[1]

(b) After the first bounce, the ball reaches a maximum height of 0.66 m.

On Fig. 1.1,

- (i) draw the ball at a height of 0.66 m,
- (ii) mark where you would position your eye to measure this height.

[2]

(c) Explain

© UCLES 2008

(i) why the maximum height h of the ball after the first bounce is difficult to measure,

(ii) how this height can be measured more accurately by two students working together.

5054/04/M/J/08

For Examiner's Use

				•		
(d)			can be estimate of the first bo		e <i>t</i> between the ball	being released and it
	The	students me	easure <i>t</i> five t	mes. The values	obtained are:	
().84	S	0.81 s	0.85 s	0.83 s	0.80 s
	(i)	Calculate th	ne average va	lue of t.		
		Give your a	nswer to a su	itable number of	significant figures.	
					<i>t</i> =	s [1]
	(ii)	An approxir	mate value of	h is given by		
				$h = (2.21t - 1)^2$		
		Calculate h	using this rel	ationship.		
					h =	m [1]

Question 1 continues on page 4

(e) The ball is allowed to bounce several times. The maximum height *h* after each bounce is measured and recorded in the table of Fig. 1.2.

For Examiner's Use

number of bounces N	<i>h </i> m
0	1.00
1	0.66
2	0.45
3	0.32
4	0.21

Fig. 1.2

On Fig. 1.3, plot the graph of *h* on the *y*-axis against the number of bounces *N* on the *x*-axis. Start your axes from the origin. Draw the curve of best fit. [4]




Fig. 1.3

© UCLES 2008 5054/04/M/J/08

(f)	Describe the relationship between N and h.							
		Examiner? Use						
	[1]							
(g)	Use your graph to estimate the total number of bounces N_T before h becomes zero.							
	$N_{\tau} = \dots [1]$							

2 Fig. 2.1 is a circuit containing a 1.5 V cell, a switch and a lamp labelled 1.5 V, 0.20 A.

For Examiner's Use




Fig. 2.1

(a) When the switch is closed the lamp does not light up.

Explain, with the aid of a diagram, how to use a voltmeter to find out whether the cell has run down.

												[2]
Suggest three	other	possible	faults	in	the	circuit	that	might	prevent	the	lamp	from


(b) Suggest three other possible faults in the circuit that might prevent the lamp from lighting.

١.	 	 	• • • • • • • •	 	 		 	 	 	
	 	 		 	 			 	 	[1]
	 	 		 	 		 	 	 	[.]
2.	 	 		 	 		 	 	 	
										[4]
•••	 	 		 	 	•••••	 	 	 	[1]
3.	 	 		 	 		 	 	 	

.....[1]

3 A group of students determine the approximate volume of air in their empty school laboratory.

For Examiner's Use



(a)	a) State						
	(i)	the measuring instrument used					
		[1]					
	(ii)	the measurements taken					
		[1]					
((iii)	how the volume of the air is calculated					

(b)	State two possible sources of error in their answer.
	1
	2

			8							
4	Fig.	4.1	on page 9 shows four thermometers used in a science laboratory.							
	(a)	Stat	State the temperature reading on thermometer A[1]							
	(b)		cm ³ of boiling water is poured into a beaker as shown in Fig. 4.2. The temperature is asured every 30 s for 10 minutes.							
			thermometer							
			boiling water beaker							
			Fig. 4.2							
		(i)	(i) State and explain which thermometer from Fig. 4.1 is the most suitable for this experiment.							
			[3]							
		(ii)	Describe how the thermometer is used in this experiment to obtain accurate readings.							
	(c)	The	rmometer D in Fig. 4.1 is used to measure the temperature of a person.							
	(0)	Fig.	4.3 shows a modern forehead thermometer. It is a thin flexible plastic strip that is sed on the forehead. The colour of the numbers changes to show the temperature.							
		piac								
			C 35 36 37 38 39 40 F 95 96! 98! 100! 102! 104							
			Fig. 4.3							
		Stat	te one advantage of this thermometer when taking the temperature of a young child.							

For Examiner's Use

© UCLES 2008 5054/04/M/J/08




Fig. 4.1

© UCLES 2008

BLANK PAGE

BLANK PAGE

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.