

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS International General Certificate of Secondary Education

| CANDIDATE NAME | | | |
|-------------------|---------------------|-----|-------|
| CENTRE NUMBER | CANDIDATE NUMBER | | |
| CHEMISTRY | | 062 | 20/31 |

Paper 3 (Extended)

October/November 2010

1 hour 15 minutes

Candidates answer on the Question Paper.

No Additional Materials are required.

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use a pencil for any diagrams, graphs or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer all questions.

A copy of the Periodic Table is printed on page 16.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

| For Examiner's Use | | | | |
|--------------------|--|--|--|--|
| 1 | | | | |
| 2 | | | | |
| 3 | | | | |
| 4 | | | | |
| 5 | | | | |
| 6 | | | | |
| 7 | | | | |
| 8 | | | | |
| Total | | | | |

This document consists of 15 printed pages and 1 blank page.



1 The table gives the composition of three particles.

| particle | number of protons | number of electrons | number of neutrons | | |
|----------|-------------------|---------------------|--------------------|--|--|
| Α | 15 | 15 | 16 | | |
| В | 15 | 18 | 16 | | |
| С | 15 | 15 | 17 | | |

| (a) Wh | at is the evidence in the table for each of the following? | |
|----------------|---|------------|
| (i) | Particle A is an atom. | |
| | | |
| (ii) | They are all particles of the same element. | [1] |
| | | [1] |
| (iii) | Particle B is a negative ion. | |
| | | [2] |
| (iv) | Particles A and C are isotopes. | [-] |
| | | [2] |
| (b) (i) | What is the electronic structure of particle A ? | |
| | | [1] |
| (ii) | What is the valency of the element? | |
| (iii) | Is the element a metal or a non-metal? Give a reason for your choice. | [1] |
| | | [1] |
| | | [Total: 9] |


| 2 | About 4 tin. | 000 years ago the Bronze Age started in Britain. Bronze is an alloy of copper and |
|---|----------------|--|
| | (a) (i) | Suggest a reason why a bronze axe was better than a copper axe. [1] |
| | (ii) | Brass is another copper alloy. Name the other metal in brass. |
| | | [1] |
| | (b) The | e diagram below shows the arrangement of particles in a pure metal. |
| | | |
| | (i) | What is the name given to a regular arrangement of particles in a crystalline solid? |
| | | [1] |
| | (ii) | Draw a diagram which shows the arrangement of particles in an alloy. |
| | | |
| | | |
| | | |
| | | |
| | | [2] |
| | (iii) | Explain the term <i>malleable</i> . |
| | | [1] |
| | (iv) | Why are metals malleable? |
| | | |
| | | [2] |

| (c) | The | common | ore | of | tin | is | tin(IV) | oxide | and | an | ore | of | copper | is | malachite, |
|-----|-----|------------------------|------------------|----|-----|----|---------|-------|-----|----|-----|----|--------|----|------------|
| | CuC | O ₃ .Cu(OH) |) ₂ . | | | | | | | | | | | | |

| (i) | Write a word equation for the reduction of tin(IV) oxide by carbon. | |
|------|--|-----|
| | | [1] |
| (ii) | Malachite is heated to form copper oxide and two other chemicals. Name these chemicals. | |

......[2]

(iii) Copper oxide is reduced to copper which is then refined by electrolysis. Label the diagram of the apparatus which could be used to refine copper.




(iv) Give **one** use of copper, other than making alloys.

[Total: 15]

[3]

3 The diagram shows a cell. This is a device which produces electrical energy. The reaction in a cell is a redox reaction and involves electron transfer.



A cell will change energy into electrical energy. [1]

- (ii) Draw an arrow on the diagram to show the direction of the electron flow. [1]
- (iii) In the left hand beaker, the colour changes from brown to colourless. Complete the equation for the reaction.

$$Br_2 + \dots \rightarrow \dots$$
 [2]

(iv) Is the change in (iii) oxidation or reduction? Give a reason for your choice.

| | [41] |
|------|------|
| | |

(v) Complete the following description of the reaction in the right hand beaker.

$$Fe^{2+}$$
 changes into [1]

(vi) When a solution of bromine is replaced by a solution of chlorine, the voltage increases. When a solution of bromine is replaced by a solution of iodine, the voltage decreases.

Suggest an explanation for this difference.

| | |
|------|---------|
| | [1] |

[Total: 7]

| Ammor | nia is | an important industrial ch | emical. | | | | |
|---------------|--------|--|------------|-------------|-------------|------------|---------------|
| (a) (i) | Giv | ve the electron structure of | f an atom | of nitroge | en. | | |
| | | | | | | | |
| (ii) | | e this electronic structure, mula of ammonia is NH ₃ n | | an the val | ency of ni | trogen, to | explain why t |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| (b) Am | nmor | nia is made by the Haber F | Process. | | | | |
| | | | | | | | |
| N_2 | (g) + | $+3H_2(g) \rightleftharpoons 2NH_3(g)$ for | ward reac | tion is exc | othermic | | |
| - | | 2 - | | | | | |
| Th | е ре | rcentage of ammonia in th | e equilibr | ium mixtu | re varies v | with condi | itions. |
| | | | | | | | |
| | | pressure / atmospheres | 100 | 200 | 300 | 400 | |
| | | % ammonia at 300 °C | 45 | 65 | 72 | 78 | |
| | | % ammonia at 500 °C | 9 | 18 | 25 | 31 | |
| | | | 000 1 | | 450.00 | | |
| In | e co | nditions actually used are | 200 atmo | spheres, | 450°C an | d an iron | catalyst. |
| (i) | Th | e original catalyst was pla | tinum. Su | agest a re | eason why | it was ch | anged to iron |
| () | | 5 · · · · · · · · · · · · · · · · · · · | | 55 | , | | 3 |
| | | | | | | | |
| | | | | | | | |
| (ii) | | plain why the highest presuilibrium mixture. | ssure give | es the hig | hest perce | entage of | ammonia in t |
| | | | | | | | |
| | •••• | | | | | | |
| | | | | | | | |
| (iii) | Wł | nat happens to the unreac | ted nitrog | en and hy | drogen? | | |
| | | | | | | | |
| | | | | | | | |
| | - | | | | | | |
| | | | | | | | |


| (iv) | State one advantage and one disadvantage of using a lower temperature. | E |
|------|--|---|
| | advantage | |
| | [1] | |
| | disadvantage | |
| | [1] | |
| | [Total: 9] | |

- **5** Monomers polymerise to form polymers or macromolecules.
 - (a) (i) Explain the term polymerise.

....

(ii) There are two types of polymerisation - addition and condensation. What is the difference between them?

(b) An important monomer is chloroethene which has the structural formula shown below.



It is made by the following method.

$$C_2H_4 + Cl_2 \rightarrow C_2H_4Cl_2$$
 dichloroethane

This is heated to make chloroethene.

$$C_2H_4Cl_2 \rightarrow C_2H_3Cl + HCl$$

(i) Ethene is made by cracking alkanes. Complete the equation for cracking dodecane.

$$C_{12}H_{26} \rightarrow \dots + 2C_2H_4$$
 [1]

Another method of making dichloroethane is from ethane.

$$C_2H_6 + 2Cl_2 \rightarrow C_2H_4Cl_2 + 2HCl$$

(ii) Suggest a reason why the method using ethene is preferred.

(iii) Describe an industrial method of making chlorine.

(iv) Draw the structural formula of poly(chloroethene).

Include three monomer units.

[2]

[Total: 9]


6 The table below shows the elements in the second period of the Periodic Table and some of their oxidation states in their most common compounds.

| element | Li | Ве | В | С | N | 0 | F | Ne |
|---------------------------|----|----|----|----|----|----|----|----|
| number of outer electrons | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 |
| oxidation state | +1 | +2 | +3 | +4 | -3 | -2 | -1 | 0 |

| (a) (i) | What does it mean when the only oxidation state of an element is zero? |
|---------------|--|
| (ii) | Explain why some elements have positive oxidation states but others have negative |
| | ones. [2] |
| (iii) | Select two elements in the table which exist as diatomic molecules of the type X ₂ . [1] |
| (b) Be | ryllium hydroxide, a white solid, is an amphoteric hydroxide. |
| (i) | Name another metal which has an amphoteric hydroxide. |
| (ii) | Suggest what you would observe when an excess of aqueous sodium hydroxide is added gradually to aqueous beryllium sulfate. |
| | [2] |
| (c) (i) | Give the formulae of lithium fluoride and nitrogen fluoride. |
| | lithium fluoride |
| | nitrogen fluoride[2] |

| (ii) | Predict two differences in their properties. | |
|-------|--|-----|
| | | |
| (iii) | Explain why these two fluorides have different properties. | [4] |
| | | [2] |
| | [Total: | 13] |

7 The diagram shows part of the carbon cycle. This includes some of the processes which determine the percentage of carbon dioxide in the atmosphere.



| Carbon dioxide is one greenhouse gas. Name another one. |
|--|
| [1] |
| Explain the term <i>respiration</i> and how this process increases the percentage of carbon dioxide in the atmosphere. |
| |
| |
| [3] |
| Explain why the combustion of waste crop material should not alter the percentage of carbon dioxide in the atmosphere. |
| |
| [2] |
| In 1960 the percentage of carbon dioxide in the atmosphere was 0.032% and in 2008 it was 0.038%. Suggest an explanation for this increase. |
| |
| [2] |
| [Total: 8] |

| 8 | Soluble sal | ts can b | he made | using a | hase and | an acid |
|---|-------------|----------|---------|---------|----------|----------|
| U | COIUDIC Sai | lo carri | oc maac | using a | base and | an acid. |

(a) Complete this method of preparing dry crystals of the soluble salt cobalt(II) chloride-6-water from the insoluble base cobalt(II) carbonate.

| Step 1 Add an excess of cobalt(II) carbonate to hot dilute hydrochloric acid. |
|---|
| Step 2 |
| |
| |
| |
| Step 3 |
| |
| |
| Step 4 |
| |
| [4] |

(b) 6.0 g of cobalt(II) carbonate was added to 40 cm³ of hydrochloric acid, concentration 2.0 mol/dm³. Calculate the maximum yield of cobalt(II) chloride-6-water and show that the cobalt(II) carbonate was in excess.

$$\begin{aligned} \mathsf{CoCO_3} \ + 2\mathsf{HC}l \ \to \ \mathsf{CoC}l_2 \ + \ \mathsf{CO_2} \ + \ \mathsf{H_2O} \\ \\ \mathsf{CoC}l_2 \ + \ \mathsf{6H_2O} \ \to \ \mathsf{CoC}l_2.\mathsf{6H_2O} \end{aligned}$$

Maximum yield

| Number of moles of HCl used = | |
|---|-------|
| Number of moles of $CoCl_2$ formed = | |
| Number of moles of $CoCl_2$.6H ₂ O formed = | |
| Mass of one mole of $CoCl_2$.6H ₂ O = 238 g | |
| Maximum yield of $CoCl_2.6H_2O = \dots g$ | [4] |
| To show that cobalt(II) carbonate is in excess | |
| Number of moles of HCl used = (use value from above) | |
| Mass of one mole of $CoCO_3$ = 119 g | |
| Number of moles of CoCO ₃ in 6.0 g of cobalt(II) carbonate = | [1] |
| Explain why cobalt(II) carbonate is in excess | |
| | [1] |
| [Total: | : 101 |

BLANK PAGE

DATA SHEET
The Periodic Table of the Elements

| | 0 | 4 He Helium 2 | Neon Neon 40 Ar | Argon 18 | 8 X | Krypton 36 | 131 Xe Xenon Xenon 54 | Radon 86 | | 175 Lu Lutetium 71 | Lr Lawrendur 103 |
|-------|-----|----------------------|---------------------------------------|------------------|-----------------|-----------------|-------------------------------------|----------------------------------|----------------------------------|---|---|
| | II/ | | 19 Fluorine 9 35.5 C1 | Chlorine 17 | 8 Q | ø. | | At | | 173 Yb Ytterbium 70 | Nobelium |
| | | | 16 Oxygen 8 | Sulfur 16 | 79 Se | Selenium 34 | 128 Te Tellurium | | | 169 Tm Thulium 69 | Md Mendelevium 101 |
| | > | | 14 Nitrogen 7 31 | Phosphorus 15 | 75 As | Arsenic 33 | 122 Sb Antimony 51 | | | 167 Er Erbium 68 | E min |
| | > | | Carbon 6 Carbon 8 S | _ | 73 Ge | Ε | Sn In 50 | 207 Pb Lead | | 165 Ho Holmium | Es n Einsteinium 99 |
| | = | | 11 B Boron 5 27 A1 | Aluminium 13 | o G | Gallium 31 | 115 In Indium | 204 T 1 Thallium 81 | | 162 Dy Dysprosium 66 | Cf Californium 98 |
| | | | | | 65 Zn | Zinc 30 | 112 Cd Cadmium 48 | | | 159 Tb Terbium 65 | BK Berkelium 97 |
| | | | | | ⁶⁴ C | Copper 29 | 108 Ag Silver 47 | 197 Au Gold | | 157 Gd Gadolinium 64 | |
| Group | | | | | 26 26 | Nickel 28 | 106 Pd Palladium 46 | 195 Pt Platinum 78 | | 152 Eu Europium 63 | |
| Gre | | | | | _{စ္} | Cobalt 27 | TO3 Rhodium 45 | 192 I.r Iridium | | Sm Samarium 62 | |
| | | Hydrogen | | | ₅₆ | Iron 26 | 101 Ru Ruthenium 44 | 190 Os Osmium 76 | | Pm Promethium 61 | Neptunium |
| | | | | | 55 N | /anganese | Tc Tc | 186 Re Rhenium | | Neodymium 60 | 238 U Uranium 92 |
| | | | | | C | Chromium 24 | 96 Mo Molybdenum 7 | 184 W Tungsten 74 | | 141 Pr Praseodymium 59 | Pa Protactinium 91 |
| | | | | | 51 | Ē | 93 Nb Niobium 41 | 181 Ta Tantalum | | 140 Ce Cerium | 232 Th Thorium |
| | | | | | 48 | Ę | 91 Zr Zirconium 40 | 178 # Hafnium | | | nic mass bol nic) number |
| | | | | | گو د | Scandium 21 | 89 × | 139 La Lanthanum * | 227 Ac Actinium 89 | series eries | a = relative atomic mass X = atomic symbol b = proton (atomic) number |
| | = | | Beryllium 4 24 Mg | Magnesium 12 | ⁶ 0 | Calcium 20 | Strontium | 137 Ba Barium 56 | 226 Ra Radium 88 | *58-71 Lanthanoid series 190-103 Actinoid series | « X |
| | _ | | 7 Li Lithium 3 23 Na | Sodium 11 | ® × | Potassium 19 | Rb Rubidium | 133 Cae sium | Fr Francium 87 | *58-71 L | Key |

The volume of one mole of any gas is 24 dm³ at room temperature and pressure (r.t.p.).

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.