

Cambridge International Examinations

Cambridge Ordinary Level

| CANDIDATE NAME | | | | | |
|-------------------|--|--|---------------------|--|--|
| CENTRE NUMBER | | | CANDIDATE NUMBER | | |



CHEMISTRY 5070/22

Paper 2 Theory

May/June 2016

1 hour 30 minutes

Candidates answer on the Question Paper.

No Additional Materials are required.

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Section A

Answer all questions.

Write your answers in the spaces provided in the Question Paper.

Section B

Answer any three questions.

Write your answers in the spaces provided in the Question Paper.

Electronic calculators may be used.

You may lose marks if you do not show your working or if you do not use appropriate units.

A copy of the Periodic Table is printed on page 20.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.




Section A

Answer **all** the questions in this section in the spaces provided.


The total mark for this section is 45.

A1 Choose from the following polymers to answer the questions.


polymer A




polymer B



polymer C



polymer D




polymer E

polymer F


5070/22/M/J/16

© UCLES 2016


polymer G



polymer H



polymer I



Each polymer can be used once, more than once or not at all.

| (a) | Which two polymers are polyesters? | |
|-----|--|-----|
| | and | [1] |
| (b) | Which polymer is used to make both clingfilm and plastic bags? | |
| | | [1] |
| (c) | Give the letter of an addition polymer | |
| | Give the letter of a condensation polymer | [1] |
| (d) | Give the letter of a polymer that is a saturated hydrocarbon. | |
| | | [1] |
| (e) | Which polymer could be part of a protein? | |
| | | [1] |

[Total: 5]

| Нус | iroge | n fluoride, HF, has a simple molecular structure. It is soluble in water. |
|-----|------------|---|
| (a) | Sug | gest one other physical property of hydrogen fluoride. |
| | | [1] |
| (b) | Hyd | rogen fluoride dissociates in water to form dilute hydrofluoric acid. |
| | (i) | Write an equation to show the dissociation of hydrogen fluoride. |
| | | [4] |
| | (ii) | Explain why an acidic solution is formed when hydrogen fluoride dissociates in water. |
| | | [1] |
| (c) | Dilu | te hydrofluoric acid reacts with aqueous calcium hydroxide. |
| | | $2HF(aq) + Ca(OH)_2(aq) \rightarrow CaF_2(aq) + 2H_2O(I)$ |
| | | at is the minimum volume, in cm 3 , of 0.150 mol/dm 3 Ca(OH) $_2$ required to react completely a solution containing 0.200 g of HF? |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | volume of Ca(OH) ₂ (aq) =cm ³ [3] |
| | (a) (b) | (a) Sug (b) Hyd (i) (ii) (c) Dilu Wha |

| (d) | Mag | gnesium reacts with fluorine to make the ionic compound magnesium fluoride. |
|-----|------|--|
| | (i) | Predict two physical properties of magnesium fluoride. |
| | | 1 |
| | | 2[2] |
| | (ii) | Explain, in terms of electrons, how a magnesium atom reacts with a fluorine molecule, F_2 , to make a magnesium ion and two fluoride ions. |
| | | |
| | | |
| | | [2] |
| | | [Total: 10] |

| Δ3 | Fsters | are | used | as | food | flavouring | s and | solvents |
|----|--------|-----|------|----|------|------------|-------|------------|
| ~~ | LSICIS | aic | useu | as | ioou | navounny | o anu | SUIVEIIIS. |

| (0) | Drow the structure | of othyl mothonog | to chowing all | of the atoms | and all of the bonds. |
|-----|-----------------------|-------------------|------------------|--------------|-----------------------|
| (a) | Draw the structure of | ot ethvi methanoa | ate. snowing ali | of the atoms | and all of the bonds. |

| | | | [1] |
|-----|------|---|------|
| (b) | Eth | yl ethanoate evaporates at room temperature. | |
| | (i) | What is meant by the term evaporation? | |
| | | | |
| | | | |
| | | | [1] |
| | (ii) | A sample of ethyl ethanoate in a beaker is moved into a colder room. | |
| | | Explain, in terms of the kinetic particle theory, why this results in a decrease in the rate evaporation. | ∍ of |
| | | | |
| | | | |
| | | | |
| | | | |

(iii) The table shows some information about different esters.

| name | structure | relative molecular mass (M_r) |
|------------------|--|---------------------------------|
| methyl ethanoate | CH ₃ CO ₂ CH ₃ | 74 |
| ethyl ethanoate | CH ₃ CO ₂ C ₂ H ₅ | 88 |
| propyl ethanoate | CH ₃ CO ₂ C ₃ H ₇ | 102 |
| butyl ethanoate | CH ₃ CO ₂ C ₄ H ₉ | 116 |
| pentyl ethanoate | CH ₃ CO ₂ C ₅ H ₁₁ | 130 |

| Which ester has the lowest rate of evaporation at room temperature and pressure? |
|---|
| |
| Explain your answer. |
| |
| |
| [2] |
| [Total: 6] |

| A 4 | Sulf | furic acid is manufactured by the contact process. | |
|------------|------|---|-----|
| | (a) | State the conditions used in the contact process. | |
| | | temperature | |
| | | pressure | |
| | | catalyst | |
| | (b) | In the contact process, sulfur dioxide reacts with oxygen. | [2] |
| | (5) | | |
| | | $2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g)$ | |
| | | Describe and explain the effect of increasing the concentration of oxygen on the rate of t reaction. | his |
| | | | |
| | | | |
| | | | |
| | | | [2] |
| | (c) | The catalyst used in the contact process increases the rate of the reaction. Describe one other advantage of using a catalyst in an industrial process. | |
| | | | |
| | (d) | Sulfuric acid is used to make the fertiliser potassium sulfate, $\rm K_2SO_4$. | |
| | | Calculate the percentage by mass of potassium in this fertiliser. | |
| | | | |

[2]

[Total: 7]

A5 The statements give some of the chemical properties of cobalt and its compounds.

- Cobalt does not react with cold water.
- Cobalt fizzes slowly with dilute hydrochloric acid.
- Cobalt does not react with aqueous zinc nitrate.
- Cobalt reacts with aqueous silver nitrate.
- Cobalt(II) oxide reacts with magnesium to form cobalt.

| (a) | Use the information | on to help arrange the following metals in order of reactivity. | |
|------|---------------------|---|------------|
| | | cobalt, magnesium, silver, sodium and zinc | |
| | most reactive | | |
| | | | |
| | | | |
| | | | |
| | least reactive | | [2] |
| (b) | Construct the equ | nation for the reaction between cobalt(II) oxide, CoO, and magnesium. | |
| | | | [1] |
| (c) | | pens when cobalt(II) carbonate is heated strongly. | |
| (-) | • | [| [1] |
| (al\ | | | ני. |
| (d) | | ing point of 1495°C. | |
| | | of structure and bonding, why a metal such as cobalt has a high meltir e a labelled diagram in your answer. | ng |
| | | | |
| | | | |
| | | | |
| | | | ••• |
| | | , | |
| | | | [2] |
| (e) | The symbol for or | ne isotope of cobalt is ${}^{57}_{27}$ Co. | |
| | Another isotope o | f cobalt has a nucleon number of 59. | |
| | Write its symbol. | | |
| | | , | [4] |
| | | | [1] |

[Total: 7]

| ۸6 | Div. | River water contains dissolved minerals and gases. | | | | | | |
|----|------|--|--|--|--|--|--|--|
| AU | LIVE | er wate | er contains dissolved militerals and gases. | | | | | |
| | (a) | Carb | on dioxide is one of the gases dissolved in river water. | | | | | |
| | | | the 'dot-and-cross' diagram to show the bonding in a molecule of carbon dioxide. Only the outer-shell electrons. | | | | | |
| | | | | | | | | |
| | | | | | | | | |
| | | | | | | | | |
| | | | | | | | | |
| | | | | | | | | |
| | | | | | | | | |
| | | | [1] | | | | | |
| | (b) | | water often contains dissolved compounds such as ammonium nitrate and calcium phate. | | | | | |
| | | (i) : | State one source of both of these compounds. | | | | | |
| | | | [1] | | | | | |
| | | | Describe and explain the environmental effect of the presence of these dissolved compounds in river water. | | | | | |
| | | | | | | | | |
| | | | | | | | | |
| | | | | | | | | |
| | | | | | | | | |
| | | | | | | | | |
| | | | | | | | | |
| | | | | | | | | |

| (c) | Rive | er water is often purified for use as drinking water. | |
|-----|------|--|-------------|
| | Des | scribe three processes involved in the purification of river water. | |
| | prod | cess 1 | |
| | | | |
| | | | |
| | prod | cess 2 | |
| | | | |
| | | | |
| | prod | cess 3 | |
| | | | |
| | | | [3] |
| | | | [0] |
| (d) | Wat | ter has a low melting point and is neutral ($pH = 7$). | |
| | (i) | Explain why water has a low melting point. | |
| | | | |
| | | | [1] |
| | (ii) | A pH meter can be used to confirm that water is neutral. | |
| | | Describe another way in which a student can confirm that water is neutral. | |
| | | | |
| | | | |
| | | | [1] |
| | | | [Total: 10] |

BLANK PAGE

Section B

Answer three questions from this section in the spaces provided.

The total mark for this section is 30.

| B7 | The formula of lead(II) nitrate is Pb(NO ₃) ₂ . | | | | | | |
|----|--|------|--|--|--|--|--|
| | (a) | | scribe how a pure sample of lead(II) nitrate crystals can be prepared from lead(II) oxide, ch is insoluble in water. | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | [4] | | | | |
| | (b) | | | | | | |
| | | Cor | nstruct the ionic equation, with state symbols, for this reaction. | | | | |
| | | | [2] | | | | |
| | (c) | | blueous lead(II) nitrate is electrolysed using graphite electrodes. Bubbles of colourless gas formed at both electrodes. | | | | |
| | | (i) | Identify the gas formed at each electrode. | | | | |
| | | | negative electrode (cathode) | | | | |
| | | | positive electrode (anode) | | | | |
| | | | [2] | | | | |
| | | (ii) | Construct the equation for the reaction at the cathode. | | | | |
| | | | [1] | | | | |
| | (d) | On | heating, lead(II) nitrate decomposes to form PbO, NO_2 and O_2 . | | | | |
| | | Cor | nstruct the equation for this reaction. | | | | |
| | | | [1] | | | | |

 $\textbf{B8} \quad \text{Cyclohexene, C}_{6}\textbf{H}_{10}, \text{ is a cycloalkene.}$

Cycloalkenes react in a similar way to alkenes.

cyclohexene

| (a) | Cyc | clohexene is an unsaturated hydrocarbon. |
|-----|------|--|
| | (i) | What is meant by the term unsaturated? |
| | | |
| | | [1] |
| | (ii) | What is meant by the term <i>hydrocarbon</i> ? |
| | | |
| | | [1] |
| (b) | Cor | nstruct the equation for the complete combustion of cyclohexene. |
| | | [1] |
| (c) | Сус | clohexene reacts with bromine. |
| | This | s is an addition reaction. |
| | (i) | Write the molecular formula of the product of this reaction. |
| | | [1] |
| | (ii) | What would be observed in this reaction? |
| | | [1] |

| (d) | Сус | lohexene can be manufactured from hexane as shown in the equation. | |
|-----|------|---|------|
| | | $C_6H_{14} \rightarrow C_6H_{10} + 2H_2$ | |
| | | culate the mass of cyclohexene that can be made from 258 g of hexane. of cyclohexene = 82] | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | mass of cyclohexene = g | [2] |
| (e) | Ano | ther cycloalkene has the following percentage composition by mass. | |
| | | C, 88.2%; H, 11.8% | |
| | (i) | Use the percentage composition by mass to show that the empirical formula of cycloalkene is ${\rm C_5H_8}.$ | this |
| | | | |
| | | | |
| | | | |
| | | | [2] |
| | (ii) | The cycloalkene has a relative molecular mass, $M_{\rm r}$, of 68. | |
| | | Draw the structure of the cycloalkene, showing all of the atoms and all of the bonds. | |
| | | | |
| | | | |

[1]

[Total: 10]

| RQ | Carbon | monoxide | reacts | with | hydrogen | in a | reversible | reaction |
|----|--------|-------------|--------|-------|----------|------|----------------|-----------|
| DJ | Carbon | IIIOIIOXIUE | reacis | willi | nyuruqen | ша | . I evel Sible | reaction. |

$$CO(g) + 2H_2(g) \rightleftharpoons CH_3OH(g) \Delta H = -91 \text{ kJ/mol}$$

The reaction reaches an equilibrium if carried out in a closed container.

| (a) | Explain, in terms of bond breaking and bond forming, why this reaction is exothermic. | | | |
|-----|---|---|-----|--|
| | | | | |
| | | | | |
| (b) | Whe | en one mole of methanol, CH ₃ OH, is formed, 91 kJ of energy is released. | [2] | |
| | Cald | culate the amount of energy released when 160 g of methanol is formed. of methanol = 32] | | |
| | | | | |
| | | | | |
| | | energy released =kJ | [2] | |
| (c) | Pre | dict, with a reason, how the position of equilibrium of this reaction changes as the | | |
| | (i) | pressure is increased at constant temperature, | | |
| | | | | |
| | | | | |
| | (ii) | temperature is increased at constant pressure. | [~] | |
| | | | | |
| | | | | |

| (d) | (d) Methanol and compound X react together to form methyl butanoate. | | | | | |
|-----|--|--|--|--|--|--|
| | (i) | Name X. | | | | |
| | | [1 | | | | |
| | (ii) | The reaction is normally carried out using a catalyst. | | | | |
| | | Name a suitable catalyst for this reaction. | | | | |
| | | [1 | | | | |
| | | [Total: 10 | | | | |

 ${\bf B10}~{\rm Manganese}({\rm IV})$ oxide, ${\rm MnO_2},$ can be used in the preparation of both chlorine and oxygen.

Reaction 1
$$MnO_2(s) + 4HCl(aq) \rightarrow Cl_2(g) + 2H_2O(l) + MnCl_2(aq)$$

Reaction 2
$$2H_2O_2(aq) \rightarrow O_2(g) + 2H_2O(l)$$

In reaction 2 manganese(IV) oxide acts as a catalyst.

(a) Reaction 1 converts chloride ions into chlorine molecules.

| Explain why this is an example of oxidation. | | | | | | |
|--|-----|--|--|--|--|--|
| | | | | | | |
| | | | | | | |
| | 171 | | | | | |

(b) Reaction 1 is investigated using different masses of MnO₂. The results are shown in the table.

| volume of HC <i>l</i> /cm ³ | concentration of HC <i>I</i> (aq) in mol/dm ³ | mass of MnO ₂ used /g | volume of Cl_2 formed at room temperature and pressure /dm 3 |
|--|--|--|---|
| 100 | 1.0 | 1.74 | 0.48 |
| 100 | 1.0 | 0.87 | 0.24 |

| xplain the difference in the volume of chlorine formed. | | | | | | |
|---|-----|--|--|--|--|--|
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | [O. | | | | | |

(c) Reaction 2 is investigated using different masses of MnO₂. The results are shown in the table.

| volume of H ₂ O ₂ (aq) /cm ³ | concentration of H ₂ O ₂ in mol/dm ³ | mass of MnO ₂ used /g | volume of O ₂ formed at room temperature and pressure /dm ³ |
|---|---|--|---|
| 100 | 1.0 | 1.74 | 1.20 |
| 100 | 1.0 | 0.87 | |

Predict the volume of oxygen, measured at room temperature and pressure, when 0.87g of MnO₂ is used. Write your answer in the table. [1]

(d) Chlorine is bubbled through aqueous iron(II) chloride to form iron(III) chloride.

| | Explain, with the aid of equations, how aqueous sodium hydroxide can be used to distinguish between aqueous iron(II) chloride and aqueous iron(III) chloride. | ish |
|-----|---|------------------|
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | [4 . |
| (e) | Describe the chemical test for chlorine. | |
| | test | |
| | | |
| | observation | |
| | | |
| | | [2] |

[Total: 10]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

The Periodic Table of Elements

| | ■/ | 2 He | helium 4 | 10 | Ne | neon 20 | 18 | Ā | argon 40 | 36 | 궃 | krypton 84 | 54 | Xe | xenon 131 | 98 | R | radon | | | |
|-------|----|---------|---------------|---------------|---------------|------------------------------|----|----|------------------|----|----|-----------------|----|----------|------------------|-------|-------------|-----------------|--------|-----------|--------------------|
| | => | | | 6 | ш | fluorine 19 | 17 | Cl | chlorine 35.5 | 35 | B | bromine 80 | 53 | Н | iodine 127 | 85 | Αŧ | astatine - | | | |
| | > | | | 8 | 0 | oxygen 16 | 16 | ഗ | sulfur 32 | 34 | Se | selenium 79 | 52 | <u>e</u> | tellurium 128 | 84 | Ро | polonium – | 116 | _ | livermorium - |
| | > | | | 7 | Z | nitrogen 14 | 15 | ₾ | phosphorus 31 | 33 | As | arsenic 75 | 51 | Sb | antimony 122 | 83 | Ξ | bismuth 209 | | | |
| | 2 | | | 9 | ပ | carbon 12 | 14 | S | silicon 28 | 32 | Ge | germanium 73 | 50 | Sn | tin 119 | 82 | Ъ | lead 207 | 114 | Εl | flerovium — |
| | ≡ | | | 2 | В | boron 11 | 13 | Αl | aluminium 27 | 31 | Ga | gallium 70 | 49 | I | indium 115 | 81 | 11 | thallium 204 | | | |
| | | | | | | | | | | 30 | Zu | zinc 65 | 48 | g | cadmium 112 | 80 | Нg | mercury 201 | 112 | ပ် | copemicium |
| | | | | | | | | | | 29 | Cn | copper 64 | 47 | Ag | silver 108 | 62 | Au | gold 197 | 111 | Rg | roentgenium - |
| Group | | | | | | | | | | 28 | Ë | nickel 59 | 46 | Pq | palladium 106 | 78 | 五 | platinum 195 | 110 | Ds | darmstadtium - |
| Ğ | | | | | | | | | | 27 | ဝိ | cobalt 59 | 45 | 몬 | rhodium 103 | 77 | 'n | iridium 192 | 109 | Ĭ | meitnerium - |
| | | - I | hydrogen 1 | | | | | | | 26 | Ь | iron 56 | 44 | Ru | ruthenium 101 | 92 | Os | osmium 190 | 108 | Hs | hassium – |
| | | | | | | | | | | 25 | Mn | manganese 55 | 43 | ပ | technetium - | 75 | Re | rhenium 186 | 107 | Bh | bohrium – |
| | | | | | pol | ass | | | | 24 | ပ် | chromium 52 | 42 | Mo | molybdenum 96 | 74 | ≯ | tungsten 184 | 106 | Sg | seaborgium - |
| | | | Key | atomic number | atomic symbol | name relative atomic mass | | | | 23 | > | vanadium 51 | 41 | qN | niobium 93 | 73 | Та | tantalum 181 | 105 | Dp | dubnium – |
| | | | | | atc | rek | | | | 22 | j | titanium 48 | 40 | Zr | zirconium 91 | 72 | 茔 | hafnium 178 | 104 | 꿆 | rutherfordium - |
| | | | | | | | | | | 21 | Sc | scandium 45 | 39 | > | yttrium 89 | 57–71 | lanthanoids | | 89–103 | actinoids | |
| | = | | | 4 | Be | beryllium 9 | 12 | Mg | magnesium 24 | 20 | Ca | calcium 40 | 38 | Š | strontium 88 | 56 | Ba | barium 137 | 88 | Ra | radium |
| | _ | | | 3 | := | lithium 7 | 1 | Na | sodium 23 | 19 | ¥ | potassium 39 | 37 | & | rubidium 85 | 22 | S | caesium 133 | 87 | Ā | francium – |

| | _ | E | _ | | inm |
|----|----|---------------------|-----|-----------|---------------------|
| 71 | | lutetium 175 | 103 | ت | lawrenc |
| 70 | Υp | ytterbium 173 | 102 | ٥ N | nobelium – |
| 69 | Tm | thulium 169 | 101 | Md | mendelevium – |
| 89 | щ | erbium 167 | 100 | Fm | fermium – |
| 29 | 운 | holmium 165 | 66 | Es | einsteinium – |
| 99 | ò | dysprosium 163 | 86 | ŭ | californium - |
| 65 | Д | terbium 159 | 26 | Ř | berkelium – |
| 64 | gq | gadolinium 157 | 96 | Cm | curium — |
| 63 | Ш | europium 152 | 95 | Am | americium - |
| 62 | Sm | samarium 150 | 94 | Pn | plutonium – |
| 61 | Pm | promethium - | 93 | ď | neptunium — |
| 09 | PΝ | neodymium 144 | 92 | \supset | uranium 238 |
| 59 | Ą | praseodymium 141 | 91 | Ра | protactinium 231 |
| 58 | Ce | cerium 140 | 06 | Ч | thorium 232 |
| 22 | Га | lanthanum 139 | 89 | Ac | actinium _ |

lanthanoids

actinoids

The volume of one mole of any gas is $24\,\text{dm}^3$ at room temperature and pressure (r.t.p.)