



UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS International General Certificate of Secondary Education

CANDIDATE NAME					
CENTRE NUMBER			CANDIDATE NUMBER		

CAMBRIDGE INTERNATIONAL MATHEMATICS

0607/06

Paper 6 (Extended)

October/November 2010

1 hour 30 minutes

Candidates answer on the Question Paper.

Additional Materials: Graphics Calculator

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

Do not use staples, paper clips, highlighters, glue or correction fluid.

You may use a pencil for any diagrams or graphs.

Answer both parts **A** and **B**.

You must show all relevant working to gain full marks for correct methods, including sketches.

In this paper you will also be assessed on your ability to provide full reasons and communicate your mathematics clearly and precisely.

At the end of the examination, fasten all your work securely together.

The total number of marks for this paper is 40.



Answer both parts A and B.

For Examiner's Use

A INVESTIGATION

THE FIBONACCI SEQUENCE

24 marks

You are advised to spend no more than 55 minutes on this part.

The Fibonacci sequence is a sequence of numbers that is found in many real-life situations.

The Fibonacci sequence begins

1 1 2 3 5

where, apart from the first two terms, each term is the sum of the previous two terms.

For example

1+1=2, 1+2=3, 2+3=5 and so on.

1 Complete the table for the first 15 Fibonacci numbers. You must show your working.

Term position	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
Fibonacci number	1	1	2	3	5	8	13	21	34	55	89	144	233		

2 (a) The table shows Fibonacci numbers that are multiples of 2.

For Examiner's Use

Complete the table.

Term position	3		9	
Fibonacci number	2	8		

Notice that:

2 is the third term in the Fibonacci sequence, 8 is the sixth term in the Fibonacci sequence, and so on.

Every third term in the Fibonacci sequence is a multiple of 2.

(b) The next two tables show other patterns.

Complete the tables and the statements that follow.

(i)

Term position	4	8	12	
Fibonacci number	3			

3 is the _____ term in the Fibonacci sequence.

Every term in the Fibonacci sequence is a multiple of 3.

(ii)

Term position			20
Fibonacci number	5	55	6765


5 is the _____ term in the Fibonacci sequence.

Every term in the Fibonacci sequence is a multiple of


(c) Complete the following statement.

Every term in the Fibonacci sequence is a multiple of 8.

3 A Golden Rectangle is a rectangle with width and length that are **consecutive** Fibonacci numbers.




When a Golden Rectangle is divided into the **least number** of squares, the length of the side of each square is a Fibonacci number.



The diagram above shows the 2 by 3 Golden Rectangle.

The least number of squares it can be divided into is three.

These squares have sides 1, 1 and 2.



The diagram above shows the 3 by 5 Golden Rectangle.


The least number of squares it can be divided into is four.

These squares have sides 1, 1, 2 and 3.


© UCLES 2010 0607/06/O/N/10

For Examiner's Use (a) On the grid below, draw the 5 by 8 Golden Rectangle. Show how this can be divided into the least number of squares. These squares have sides 1, 1, 2, 3 and 5.

Examiner's Use



(b) On the grid below, draw the 8 by 13 Golden Rectangle. Show how this can be divided into the least number of squares.



(c) (i) Complete the table to show the least number of squares in each of the first six Golden Rectangles.

For Examiner's Use

Size of rectangle	1 by 1	1 by 2	2 by 3	3 by 5	5 by 8	8 by 13
Least number of squares	1			4		


	Least number of squares	1			4			
	(ii) Write down the least	number of	squares the	ere are in th	e 21 by 34	Golden Re	ctangle.	
	(::) When the least num	han af agu	uamaa ia 11	vrmita dav	von the voic	1th and the	lamath of	thia
	(iii) When the least num Golden Rectangle.	iber of squ	iares is 11	, write dov	wn the wic	un and the	e length of	unis
					and			
(d)	When the width and the l Fibonacci sequence, write	_		_			nth terms o	f the

(e) In this part, numbers that are **two** positions apart in the Fibonacci sequence are used for the width and length of a rectangle. For example, 1 by 2, 1 by 3, 2 by 5, 3 by 8 and so on.

For Examiner's Use

Term position	1	2	3	4	5	6	
Fibonacci number	1	1	2	3	5	8	
						ı	

Write down, in words, the connection between the term positions of the two Fibonacci numbers used for the width and length of a rectangle and the least number of squares in the rectangle.



B MODELLING

THE SOLAR SYSTEM

16 marks

Examiner's Use

You are advised to spend no more than 35 minutes on this part.

Logarithms to base 10 are written as log.

1 The table below shows information about seven planets in the Solar System.

Planet	Distance from the Sun (S km)	Time to orbit the Sun (<i>T</i> days)	$\log S$	log T
Mercury	5.79×10^7	88	7.8	1.9
Venus	1.08 x 10 ⁸	225	8.0	2.4
Earth	1.50 x 10 ⁸	365	8.2	2.6
Mars	2.28 x 10 ⁸	687		
Jupiter	7.78 x 10 ⁸	4330		
Saturn	1.43 x 10 ⁹	10800		
Pluto	5.91 x 10 ⁹	90 800	9.8	5.0

Complete the table of values for $\log S$ and $\log T$. Give each value correct to 2 significant figures.

- 2 (a) On the grid opposite, plot the seven points ($\log S$, $\log T$).
 - **(b)** Plot the mean point (8.6, 3.2) and use this to draw a line of best fit. (Do this by eye. Do not use your calculator.)
- 3 The time taken for the planet Uranus to orbit the Sun is 30 685 days.

Use your graph to estimate the distance of Uranus from the Sun. Give your answer correct to 2 significant figures.

|--|

Let $x = \log S$ and $y = \log T$. The equation of the line of best fit is y = mx + c.

Use your calculator to find the values of m and c. Give each answer correct to 2 significant figures.


<i>m</i> =	
<i>c</i> =	

5 A model for this is $\log T = m \log S + c$. The distance of the planet Neptune from the Sun is 4.50×10^9 km.

Use the model to find the time taken for Neptune to orbit the Sun. Give your answer in standard form correct to 2 significant figures.

For

Use



For Examiner's Use

6	Writing c as $\log k$, the model can be written as $\log T = m \log S + \log k$.			
	(a)	Show that $T = kS^m$.		
	(b)	Find the value of k .		
			<i>k</i> =	
	(c)	Write the model $T = kS^m$ using your values of k and m .		

Use the data for Earth to test this model.

BLANK PAGE

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.