

DRAFT VERSION JULY 6, 2018

Typeset using **L^AT_EX modern** style in AASTeX62

The Mass of Stirring Bodies in the AU Mic Debris Disk Inferred from Resolved Vertical Structure

CAIL DALEY,¹ A. MEREDITH HUGHES,¹ EVAN S. CARTER,¹ KEVIN FLAHERTY,¹
ZACHARY LAMBROS,¹ MARGARET PAN,² HILKE SCHLICHTING,^{3,2}
EUGENE CHIANG,^{4,5} MARK WYATT,⁶ DAVID WILNER,⁷ SEAN ANDREWS,⁷ AND
JOHN CARPENTER⁸

¹*Department of Astronomy, Van Vleck Observatory, Wesleyan University, 96 Foss Hill Drive, Middletown, CT 06459, USA*

²*Department of Earth, Atmospheric and Planetary Sciences, Massachusetts Institute of Technology, Cambridge, MA 02139, USA*

³*Department of Earth and Space Sciences, University of California, Los Angeles, CA 90095, USA*

⁴*Department of Astronomy, University of California at Berkeley, Campbell Hall, Berkeley, CA 94720-3411*

⁵*Department of Earth and Planetary Science, University of California at Berkeley, McCone Hall, Berkeley, CA 94720-4767*

⁶*Institute of Astronomy, University of Cambridge, Madingley Road, Cambridge CB3 0HA, United Kingdom*

⁷*Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA, 02138, USA*

⁸*Joint ALMA Observatory (JAO), Alonso de Cordova 3107 Vitacura-Santiago de Chile, Chile*

ABSTRACT

The vertical distribution of dust in debris disks is particularly sensitive to the total mass of planetary bodies stirring the disk. Inferences of dynamical mass from debris disk vertical structure have previously been applied to infrared and optical observations of several systems, but theoretical works predict that the small particles traced by short-wavelength observations will be ‘puffed up’ by radiation pressure, yielding only upper limits on the total dynamical mass. The large grains that dominate the emission at millimeter wavelengths are much less sensitive to the effects of stellar radiation or stellar winds, and therefore trace the underlying mass distribution more directly. Here we present ALMA 1.3 mm dust continuum observations of the debris disk around the nearby M star AU Mic. The 3 au spatial resolution of the observations, combined with the favorable edge-on geometry of the system, allows us to measure the vertical thickness of the disk; we report a scale height-to-radius aspect ratio of $h = 0.032 \pm 0.005$ between ~ 23 au and ~ 42 au. Comparing the observed

aspect ratio to a theoretical model of steady-state size-dependent velocity distributions in the collisional cascade, we find that the total mass of bodies embedded in the local disk is less than $1.5 M_{\oplus}$, and that these bodies have individual sizes greater than about 400 km. These measurements rule out the presence of a gas giant or Neptune analog at radii near the ~ 40 au extent of the debris ring, but are suggestive of the presence of large planetesimals (or an Earth-sized planet) stirring the dust distribution.

1. INTRODUCTION

Planets form during a relatively short and early stage in the lifetime of stellar systems, when the host star is still encircled by a protoplanetary disk. Planet formation, as well as processes including accretion, photoevaporation, and winds, causes first-generation protoplanetary material to dissipate over time (Williams & Cieza 2011; Ercolano & Pascucci 2017). The first-generation material is replaced by second-generation ‘debris,’ produced by collisional grinding of larger planetesimals into small dust grains in a process known as a collisional cascade (Wyatt 2008). The resulting debris disks, optically thin and significantly less luminous than their protoplanetary counterparts, are currently observable around at least 25% of Solar-type stars and are likely to be at least as common as the exoplanetary systems with which they are thought to be associated (Montesinos et al. 2016).

Analysis of the morphological and emissive properties of debris disks sheds light on the final stages of planetary system evolution and can reveal the presence of planets hidden within. Planets can imprint features such as rings, gaps, clumps, or other asymmetries on their parent disks, although it is rarely straightforward to infer the properties of planets directly from the disk morphology (see the review by Hughes et al. 2018, and references therein). In gas-poor systems, the vertical structure of a debris disk can serve as a probe of the total mass of large bodies stirring the collisional cascade (Thébault 2009). The presence of massive bodies increases the inclination dispersion of the dust particle orbits and thus the scale height H of the observed dust distribution.

The dynamical excitation of a disk can therefore be measured from its aspect ratio H/r , which in turn allows inferences about the presence and size of the bodies responsible for the dynamical stirring. Such work has been undertaken by several authors using visible and infrared observations (Artymowicz 1997; Thébault & Augereau

2007; Quillen et al. 2007). However, Thébault (2009) demonstrates that radiation pressure from the host star should preferentially excite the smallest dust grains in a disk, imparting a ‘natural’ scale height to the system even in the absence of large stirring bodies. Thus longer-wavelength ($\lambda \geq 50 \mu\text{m}$ for typical grain blow-out sizes of ~ 2 to $10 \mu\text{m}$) observations are required to measure the disk scale height as determined by dynamical stirring alone, as the large grains dominating the emission at these wavelengths are much less sensitive to the effects of radiation pressure.

The M3IVe star AU Mic presents a particularly favorable target for such observations because of its proximity (9.725 ± 0.005 pc; Gaia Collaboration et al. 2016, 2018), edge-on inclination, and apparently symmetric morphology at millimeter wavelengths. The first M star detected to have a far-infrared excess, AU Mic hosts one of the best-studied debris disks (Moshir et al. 1990). As a member of the β Pic Moving Group, it is thought to be relatively young: 23 ± 2 Myr (Binks & Jeffries 2014; Mamajek & Bell 2014; Malo et al. 2014). A wide range of masses ($0.3 M_{\odot}$ to $0.6 M_{\odot}$; Plavchan et al. 2009; Houdebine & Doyle 1994) are reported in the literature; Schüppler et al. (2015) assume a stellar mass of $0.5 M_{\odot}$ based on the mean of these values. The disk around AU Mic was first resolved by Kalas et al. (2004) in scattered light, and a host of observations spanning the optical to the submillimeter have followed (Augereau & Beust 2006; MacGregor et al. 2013; Schneider et al. 2014; Matthews et al. 2015; Wang et al. 2015).

Notably, the radial structure of the debris around AU Mic exhibits a so-far-unique time variability at scattered light wavelengths. Boccaletti et al. (2015, 2018) identify several local intensity maxima offset from the disk midplane on the southeast side of the disk. These features are moving away from the star along the disk midplane at projected velocities that are not consistent with Keplerian rotation; in fact, the outermost features appear to be unbound from the star. Sezestre et al. (2017) provide kinematic fits and invoke a dust source of unspecified nature to explain the features. Chiang & Fung (2017) propose that these fast-moving features are made up of dust particles repelled by a time-variable stellar wind that triggers dust avalanches when the wind blows strongest. These avalanches would be seeded by the debris remaining from the recent disruption of a ~ 400 km sized progenitor. In this paper we will provide an independent constraint on the presence of comparably sized planetesimals.

We present here $0.3''$ Atacama Large Millimeter/submillimeter Array (ALMA) 1.3 mm observations of the AU Mic debris disk. These observations represent a

factor of ~ 2 improvement in both spatial resolution and rms noise relative to previous ALMA observations of the system by MacGregor et al. (2013), and our analysis indicates that the vertical structure of the disk is resolved with 4σ confidence. In §2 we present the new observations and describe the data reduction. Section 3 documents the basic observational results regarding the disk flux, morphology, and gas content. In §4 we conduct a parametric exploration of an axisymmetric disk model in order to investigate the degeneracy between vertical structure, radial structure, and viewing geometry. In §5 we discuss our results, particularly the constraints on the dynamical excitation of the disk imposed by our measurement of the scale height, and compare them to previous observations. The results of our measurement of the scale height and its implications for the total disk mass are summarized in §6.

2. OBSERVATIONS

AU Mic was observed with ALMA on three dates: 26 March 2014, 18 August 2014, and 24 June 2015 (see Table 1). All observations employed ALMA’s 12m antennas and Band 6 receivers, including four independently tunable spectral windows each with a bandwidth of 1.875 GHz. One spectral window was centered around the CO $J = 2 - 1$ transition at a rest frequency of 230.538001 GHz and a channel spacing of 0.6 km/s. The remaining three spectral windows were configured to detect continuum emission with channel spacings of 21.7 km/s and central frequencies of 228.5, 213.5, and 216.0 GHz. The mean of the four central wavelengths is 1.35 mm. The baseline lengths range between 12 m and 1320 m; the longest baseline among the three observations traces an angular scale of 0.22'' and a spatial scale of 2.1 au.

Calibration, reduction, and imaging were carried out using the CASA software package. Standard ALMA reduction scripts were applied to the datasets: phase calibration was accomplished via gain calibration and water vapor radiometry tables, while system temperature calibrations were performed to account for variations in instrument and weather conditions. Flux and bandpass calibrations were subsequently applied; the flux calibration is subject to a 10% systematic uncertainty. In addition to these standard procedures, the weights of the visibilities were recalculated using the variance around each baseline as in Flaherty et al. (2017).

During the last segment of the June observation (04:23:38-04:29:58 UT), the host star flared. To determine the flux density of the flare as a function of time, we first binned the data into one-minute intervals using the task `split`. Fitting a point source to the long baselines in each bin tended to overestimate the stellar flux when

Observational parameters	26 March 2014	18 August 2014	24 June 2015
Antennas:	32	35	37
Baseline length (m):	12–406	19–1160	30–1320
On-source time (min):	35	35	33
Flux calibrator:	Titan	J2056-472	Titan
Bandpass calibrator:	J1924-2914	J2056-4714	J1924-2914
Gain calibrator:	J2101-2933	J2101-2933	J2056-3208
pwv range (mm):	[0.63, 0.66]	[1.58, 1.69]	[0.67, 0.74]
<hr/>			
Imaging parameters			
Beam size (arcsec):	1.27×0.74	0.33×0.30	0.47×0.31
Peak intensity ($\mu\text{Jy}/\text{beam}$):	630	240	320
rms noise ($\mu\text{Jy}/\text{beam}$):	30	30	20

Table 1. Observational and imaging parameters for the three datasets used in this work. Images were created using the task `tclean` with natural weighting.

compared with the results of our parametric modeling presented in §4, as even the longest baselines are sensitive to disk emission. Instead, we used the task `tclean` to image the emission with natural weighting and determined the flux density of the peak pixel in each image. To recover the flare flux density, we subsequently subtracted the disk-only flux density in the corresponding pixel of our best-fit disk model image convolved with the ALMA visibilities (see 4 for details of the model fitting process and Table 3 for a list of best-fit parameters). The resulting flare flux densities can be found in Table 2. We exclude from our analysis of the disk emission the six minutes during which the flare occurred, as it proved difficult to separate the stellar emission from that of the disk while it was changing so rapidly.

Since AU Mic is a high proper motion system, its equatorial coordinates changed significantly over the 1.5 years between the first and last observations. We were able to obtain a more precise alignment of the datasets from the bright chromospheric emission of the central star than from the measured proper motion. We fit an image-domain 2-D Gaussian to a small region around the star on each date with the task `imfit`, and used the centroid of the Gaussian fit to define the star position. Each dataset was then phase-shifted using the task `fixvis` so that the pointing center of the data was the same as the fitted star position. We note that the centroid of the Gaussian fit for the June observations changes by 1.13 ± 0.18 au (30% of the

Time (UTC)	Peak Flux Density (mJy/beam)
03:45:0–04:20:0 (no flare)	0.26 ± 0.02
4:23:38–4:24:00	0.84 ± 0.15
4:24:00–4:25:00	11.10 ± 0.12
4:25:00–4:26:00	3.46 ± 0.10
4:26:00–4:27:00	1.40 ± 0.10
4:27:00–4:28:00	0.48 ± 0.11
4:28:00–4:29:00	0.40 ± 0.10
4:29:00–4:29:58	0.33 ± 0.10

Table 2. The peak flux densities of the central source before and during the June 24 flare, determined from naturally-weighted clean images of one-minute time bins.

synthesized beam) between the flare and non-flare segments of the data; this change in the fit stellar location could be explained if the flare were not symmetric with respect to the star.

Imaging was performed using standard Fourier inversion methods as implemented in the **CASA** task **tclean**. A natural weighting scheme was used to trace small-scale disk structure, resulting in an rms noise of $15\,\mu\text{Jy}/\text{beam}$ and a $0.52'' \times 0.39''$ restoring beam with a position angle (PA) of $77^\circ.9$. Because the **CASA** task **tclean** preserves pointing center offsets when converting several visibility datasets into an image, it was necessary to combine the data into a single file before cleaning in order to account for the offset in phase center between datasets. This was done using the task **concat**, which combines datasets with pointing centers aligned so long as their pointing centers do not differ by a value greater than the parameter **dirtol**.

3. RESULTS

Figure 1 shows the combined dust continuum emission from all three observations at 1.3 mm; chromospheric emission from the M star is visible as a point source at the center of the image (Cranmer et al. 2013). The peak signal-to-noise ratio of the dust emission is ~ 23 . Using the **MIRIAD** task **cgcurs**, we measure an integrated flux density of $4.97 \pm 0.08\,\text{mJy}$ enclosed within the 3σ contours of the naturally weighted image. We note that this value represents the combined emission from the disk *and* the star. Faithfully disentangling the two components proved difficult, both because the stellar flux varied significantly across the three nights of observation (Table 3) and because emission from the star and edge-on disk overlap in the sky




Figure 1. The AU Mic system imaged by ALMA at a wavelength of 1.3 mm using natural weighting. The rms noise is $15 \mu\text{Jy}/\text{beam}$, and the restoring beam has dimensions $0.52'' \times 0.39''$ with a PA of 77.9° . Contours are integer multiples of the three times the rms noise. The hatched ellipse in the bottom left of each pane designates the size and shape of the restoring beam.

plane. Consequently, the most accurate way to isolate the disk flux from the stellar contribution is through parametric modeling (see §4) where the two components can be specified separately; our modeling yields a disk flux density of $4.81_{-0.05}^{+0.04} \text{ mJy}$. This value is significantly smaller than the best-fit 1.28 mm disk flux density of $7.14_{-0.25}^{+0.12} \text{ mJy}$ reported by MacGregor et al. (2013) (equivalent to $6.46_{-0.23}^{+0.11} \text{ mJy}$ when

scaled to our observing wavelength of 1.35 mm assuming a blackbody temperature of 25 K).

The ansa to the NW exhibits a maximum flux density of $330 \pm 15 \mu\text{Jy}/\text{beam}$ at a separation of $24.1 \pm 0.2 \text{ au}$ and PA of $128^\circ 2 \pm 0.4$, while the ansa to the SE exhibits a maximum flux density of $340 \pm 15 \mu\text{Jy}$ at a separation of $29.0 \pm 0.2 \text{ au}$ and PA of $128^\circ 7 \pm 0.4$. Here PA is measured counterclockwise with respect to the north celestial pole. The discrepancy in position angle between the two peaks falls within the estimated uncertainties and so we are not able to confirm the scattered light PA offset observed by [Boccaletti et al. \(2015\)](#). The peak flux densities of the ansae differ by less than the rms noise, indicating that there is no significant difference in brightness between the two sides of the disk. Indeed, the apparent flux asymmetry in these data is in the opposite direction of the apparent flux asymmetry in [MacGregor et al. \(2013\)](#), providing further circumstantial evidence that there is no significant flux asymmetry at millimeter wavelengths.

While comparison of the ansae peak flux densities suggests that the AU Mic disk does not exhibit severe asymmetry, a more detailed analysis can be conducted by extracting surface brightness and vertical structure profiles from the naturally weighted image in Figure 1. These profiles, shown in Figure 2, were created by fitting a one-dimensional Gaussian to the disk vertical surface brightness profile at a series of stellocentric separations along the disk major axis. The top, center, and bottom panes correspond respectively to the amplitude, centroid, and beam-subtracted FWHM of the Gaussian fits. Broadening effects of the synthesized beam in the vertical direction have been removed by subtracting in quadrature the Gaussian beam FWHM from the Gaussian fit FWHM. In more physical terms, the three panes represent the intensity at the location of the disk ‘spine,’ the spine’s elevation from the midplane, and the disk FWHM.

As seen in the top pane of Figure 2, the surface brightness profiles of the two ansae generally mirror each other. However, we note the presence of slightly asymmetric local intensity maxima in the inner regions of the disk, one $\sim 10.5 \text{ au}$ to the SE of the star and one $\sim 7.3 \text{ au}$ to the NW. It is unclear whether these are real features of the disk or artifacts of the rms noise or cleaning process; we examine the significance of these features in §4. Three-sigma emission as determined from the surface brightness profile extends to a radial distance of $\sim 42 \text{ au}$ on the NW side and $\sim 44 \text{ au}$ on the SE side. Thus, the disk is resolved across ~ 17 beams along the major axis.




Figure 2. The AU Mic debris disk’s radial and vertical structure, extracted from the naturally weighted image in Figure 1. The solid blue line shows the southeast limb of the disk, while the dashed red line shows the northeast limb; the shaded bands designate 1σ uncertainties. From top to bottom, the three panes show i) the disk spine surface brightness, ii) the disk spine deviation from the midplane, and iii) the disk FWHM as a function of projected separation from the star after subtraction of the beam FWHM in the vertical direction. Because the disk is viewed edge-on and a wide range of stellocentric separations contribute to the FWHM at a given projected separation, the apparent vertical FWHM has no radial dependence. The Gaussian profile inset in the first pane shows the size of the combined dataset synthesized beam projected onto the *radial* axis of the disk, while the profile inset in the bottom pane shows its projection onto the *vertical* axis. The bottom inset also shows 40 disk vertical profiles (dotted lines) sampled at random from $r \leq 35$ au; the profiles have all been scaled to the same amplitude and centroid, showing that the FWHMs of the disk profiles are larger than that of the beam. Although the last pane indicates a beam-subtracted disk FWHM that is technically smaller than the vertical FWHM of the beam, the fact that the image-domain vertical height of the disk is consistently in excess of the beam contribution implies that our data spatially resolve the vertical structure of the disk.

We were unable to detect (in either the combined dataset or the three individual epochs) the intensity variations or excursions of the disk spine from the midplane that characterize the fast-moving features observed by [Boccaletti et al. \(2015, 2018\)](#). This is not entirely surprising, as both [Sezestre et al. \(2017\)](#) and [Chiang & Fung \(2017\)](#) suggest that the features are composed of sub-micron-sized grains which do not emit efficiently in the millimeter.

Cursory analysis indicates the disk is marginally resolved perpendicular to the major axis as well, exhibiting a vertical FWHM of $2.8 \text{ au} \approx 2/3 \text{ beam}$ after taking into account the broadening effects of the beam (Figure 2, bottom pane). Our ability to resolve the vertical structure of the AU Mic debris disk is discussed in greater detail in §5.5.

3.1. ^{18}CO Content

There is no evidence that the AU Mic system harbors a significant reservoir of molecular gas. We set a 3σ upper limit of $0.07 \text{ Jy km s}^{-1}$ on the CO $J = 2 - 1$ integrated flux, obtained by integrating the flux density in an $8'' \times 8''$ box around the star between V_{LSRK} velocities of -10 and 10 km s^{-1} . For a given excitation temperature, an upper limit on the total gas mass can be inferred from the upper limit on integrated flux. Assuming local thermal equilibrium (LTE) and that the gas is cospatial with the dust disk, we find a total gas mass upper limit between $1.7 \times 10^{-7} M_\oplus$ and $8.7 \times 10^{-7} M_\oplus$ for excitation temperatures between 10 K and 250 K.

4. ANALYSIS

Previous studies of the scale height of debris disks have demonstrated a degeneracy between vertical structure, radial structure, and viewing geometry (e.g. [Milli et al. 2014](#)). For example, it can be difficult to distinguish a disk that is vertically thin but slightly inclined from one that is vertically broad but perfectly edge-on. In light of this, we adopt a modeling approach that combines a simple ray-tracing code to properly project the radial and vertical flux distribution of the optically thin emission onto the sky plane with an MCMC fitting algorithm that allows us to explore the degree to which these known degeneracies impact our ability to measure the vertical structure of the disk.

4.1. Modeling Formalism

We use the parametric structure and ray tracing disk code described in [Flaherty et al. \(2015\)](#), itself an adaptation of earlier work by [Rosenfeld et al. \(2013\)](#). Synthetic

sky-projected images are generated from a given temperature and density structure and are subsequently Fourier transformed to create model visibilities that can be directly compared to the interferometric data.

We assume the disk to be azimuthally symmetric and vertically isothermal. At a given radius r , the vertical density profile is assumed to be Gaussian with a standard deviation equal to the scale height $H(r)$. The scale height is given by $H(r) = hr$, where the aspect ratio h is a constant. It is common for debris disk models to assume that scale height is a linear function of radius (Sai et al. 2015; Olofsson et al. 2016), and the theoretical work of Thébault (2009) on vertical structure in debris disks also assumes a ‘global’ aspect ratio. We cannot justify choosing a more complex parameterization of the scale height given the resolution of the data compared to the FWHM of the disk. A Gaussian vertical structure is consistent with Brown (2001), who finds that the ecliptic inclination distribution of Kuiper Belt Objects (KBOs) is well described by the sum of two Gaussians. Furthermore, the author notes that a Gaussian appears to be a “natural functional form” for the distribution of ecliptic inclinations in the Kuiper Belt; Monte Carlo simulations of dynamical interactions in a disk with initial inclinations of zero produce an ecliptic inclination distribution that is perfectly fit by a Gaussian.

The dust opacity is set to $2.3 \text{ cm}^2 \text{ g}^{-1}$ (Beckwith et al. 1990), placing the model disk in the optically thin regime for the range of dust masses explored. For an optically thin disk, the observed thermal emission is determined by both the surface density and temperature of the dust; to break the degeneracy between these two parameters, we assume that the dust grains are in blackbody equilibrium with the central star. Thus the dust temperature at a distance r from the host star is given by

$$T_{dust}(r) = \left(\frac{L_\star}{16\pi r^2 \sigma} \right)^{1/4} \quad (1)$$

where L_\star is the bolometric luminosity of the star and σ is the Stefan-Boltzmann constant. We assume that the radial surface density takes the form of a power law:

$$\Sigma(r) = \begin{cases} \Sigma_c r^p & r_{in} \leq r \leq r_{out} \\ 0 & \text{otherwise} \end{cases} \quad (2)$$

where p is the power law exponent, and r_{in} and r_{out} are the disk inner and outer radius. The critical surface density Σ_c normalizes the surface density structure for a

given total dust mass M_{dust} :

$$\Sigma_c = \frac{M_{dust} (p+2)}{2\pi \left[r_{out}^{(p+2)} - r_{in}^{(p+2)} \right]}. \quad (3)$$

The observed disk PA and inclination i are free parameters. We adopt a stellar luminosity of $0.09 L_\odot$ (Plavchan et al. 2009) and a distance to the star of 9.725 ± 0.005 pc (Gaia Collaboration et al. 2018); the observed 1.3 mm stellar flux density F_\star is left as a free parameter for each observation date. We note that the uncertainty in the stellar distance could affect the modeled disk mass and disk radial extent, while the 10% systematic flux uncertainty could affect the modeled disk mass and stellar flux density. The disk center is set by the measured position of the central point source, which were obtained in §2. The spatial resolution of the resulting model sky image is set to 0.15 au per pixel, $\sim 7\%$ of the spatial scale sampled by the longest baseline in the data. After the model image is generated by the ray tracing code, it is Fourier transformed into the visibility domain and sampled at the same spatial frequencies as the ALMA data with the MIRIAD task `uvmodel`. This allows the model to be compared directly to the visibilities in the Fourier domain, where uncertainties are better characterized than in the image domain.

We explore the parameter space of the model using the affine-invariant formulation of the MCMC algorithm described by Goodman & Weare (2010) and implemented in Python as `emcee` (Foreman-Mackey et al. 2013). MCMC routines sample parameter space such that the density of samples in a given region is proportional to the local probability density, allowing estimation of the posterior probability functions themselves. The process therefore not only identifies regions of high probability in parameter space, but also allows uncertainties and degeneracies between parameters to be determined from the correlations between the posteriors of each parameter. A log-likelihood metric $\ln \mathcal{L} = -\chi^2/2$ is used to assess the quality of fit between the synthetic and observed visibilities.

We assume uniform priors for all parameters. The dust mass was sampled in logarithmic space, formally equivalent to assuming a log uniform prior. Bounds placed on the logarithm of the dust mass were chosen to be wide enough to encompass all currently detectable optically thin dust masses. Priors placed on the stellar flux density and disk inner radius, width, power law exponent, and aspect ratio ensured that these parameters were always greater than zero. The position angle was confined




Figure 3. Kernel density estimates of the marginalized posterior probability distributions for the fiducial run. The central dashed line designates the median of each distribution while the outer lines mark the 16th and 84th percentiles (1σ confidence intervals).

to the range $1^\circ < \text{PA} < 360^\circ$, and the inclination to $0^\circ < i < 90^\circ$. Because AU Mic is so close to edge-on, the preferred inclination falls very close to the 90° prior upper bound. To ensure that the proximity of the solution to the edge of parameter space does not affect the posterior distribution, we investigated the effect of allowing the inclination to vary above 90° . Doing so produced a inclination distribution with two symmetric modes on either side of 90° . When the resulting unbounded inclination distribution was reparameterized such that all values fell between 0° and 90° , the original bounded inclination distribution was recovered, indicating that the placing a 90° prior upper bound has no significant effect on the inclination posterior.

We performed several MCMC runs in order to investigate a variety of model formalisms. All runs used 50 walkers, and 10^5 samples were drawn from each run to allow accurate statistical comparison between runs. Initially we varied ten parameters: the logarithm of the disk dust mass ($\log M_{dust}$), the disk inner radius (r_{in}), width (Δr), power law exponent (p), scale height aspect ratio (h), inclination (i), position angle (PA), and finally a separate stellar flux density (F_{\star}) for each of the three observation dates. After the fact, the posterior distribution for Δr was replaced by the outer radius posterior $r_{out} = r_{in} + \Delta r$ to allow for easier interpretation. This model formalism resulted in a best-fit χ^2 value of 626157.84 (reduced $\chi^2 = 1.027$), and we treat this parameterization as our fiducial model.

4.2. Investigating Radial Structure

Marginalized posterior probability distributions for the fiducial model parameters are shown in Figure 3. While the majority of the distributions are Gaussian, the three parameters that determine the radial structure of the disk (p , r_{in} , and r_{out}) exhibit slight bimodality. All three parameters are degenerate as can be seen from the ‘corner’ plot in Figure 4; in fact, the bimodality of the three parameters is a result of the existence of two distinct families of solutions in parameter space. While the fiducial best-fit model has $p = 0.9$, $r_{in} = 23.3$ au, and $r_{out} = 41.5$ au, a lower-likelihood family of solutions can clearly be seen in Figure 4. The highest-likelihood model associated with this family has $p = 2.8$, $r_{in} = 6.1$ au, and $r_{out} = 40.4$ au. More generally, mild degeneracies between the parameters determining vertical structure, radial structure, and viewing geometry are visible, but the aspect ratio h is correlated to a significant degree only with the inclination i .

The fiducial best-fit model image and residuals found in Figure 5a provide further information as to the cause of the bimodal posterior distribution. As can be seen from the residual map, the outer regions of the disk are reproduced well by the model; however semi-symmetric 3σ residuals, one ‘above’ the disk midplane and one ‘below,’ remain at projected stellocentric separations of ~ 10 au. We note these residuals share the locations of the local intensity maxima described in §3. The convergence of these features leads us to consider the possibility of either a dust density enhancement (a ring) or reduction (a gap) in the inner regions of the disk. As a gap/ring would cause the radial surface brightness profile of the disk to deviate from that given by the simple power law used in our modeling, it could explain the bimodality in the posterior distributions of the parameters governing disk radial structure.

We first explored the effects of adding a gap to the inner regions of the disk. The gap inner and outer radii were left as free parameters and the dust density within the gap was set to zero. The gap consistently converged to regions where the dust density was already zero (interior to the disk inner radius or exterior to the disk outer radius); we take this as evidence that the data are not well characterized by a gap. As a next step, we experimented with adding a ring to the disk. The ring inner radius r_{ring} was once again left as a free parameter and the ring width was fixed to 0.1 au. The ring was also characterized by a dust mass M_{ring} , which was evenly distributed across the radial extent of the ring. As can be seen in Figure 5b






Figure 4. ‘Corner’ plot showing degeneracies between a subset of the free parameters for the fiducial model. Two families of solutions are visible, and are most prominent in the p , r_{in} , and r_{out} distributions. In addition, the two-dimensional slices through parameter space show limited degeneracies between radial structure, vertical structure, and viewing geometry. Of particular interest is the scale height h , which exhibits a noticeable degeneracy with the inclination i .

(a) Fiducial run best-fit model and residuals.



(b) Ring run best-fit model and residuals.



(c) ‘Skinny’ disk run best-fit model and residuals.




Figure 5. Best-fit model image and residuals, sampled at the spatial frequencies of the ALMA data and cleaned using natural weighting. Contours are integer multiples of the ALMA data 3σ confidence level. In the residual maps the stellar location is marked with a star, and the disk extent and position angle are given by the dotted black line. For the fiducial model (a), 3σ residuals can clearly be seen at the semi-symmetric positions ~ 10 au from the star. Adding an additional ring of dust to the model (b) reduces these residuals by $\sim 25\%$, but statistical analysis indicates that best-fit ring model does not conclusively describe the data better than the fiducial model. Fixing the scale height well below ALMA’s resolution limits (c) results in a less statistically significant model, especially in the outer regions of the disk where ALMA is sensitive to the vertical structure of the disk.

Note: The new Gaia ring run hasn’t finished yet, which is why the second subplot hasn’t been updated..

this parameterization was better able to reproduce the ‘bump’ in the radial surface brightness profile at $r \sim 10$ au, reducing the best-fit model residuals by $\sim 25\%$. However, 2σ residuals are still present at the location of the bumps; the ring model’s failure to accurately reproduce these features could be explained by the fact that the two local intensity maxima are not perfectly symmetric about the star, and together exhibit a position angle that deviates from that of the main disk. Image domain analysis indicates that the bump to the SE has a separation of ~ 10.5 au, while the NW bump has a separation of ~ 7.3 au (Figure 2). This discrepancy could be explained if the hypothetical ring were eccentric or if the two bumps were instead localized, asymmetric density enhancements within the outer ring.

The median values and best-fit model parameters for the fiducial and ring parameterizations can be found in Table 3. We use both the AICc, a form of the Akaike Information Criterion (AIC) corrected for finite datasets, and the Bayesian Information Criterion (BIC) to compare goodness of fit between models with different numbers of free parameters. The best-fit ring model is preferred to the fiducial model with 3.7σ confidence on the AICc; conversely, the fiducial model is preferred to the ring model on the BIC with $\Delta\text{BIC} = 4.3$. As such, we are not able to conclusively confirm the presence of an additional ring of mm dust grains in the AU Mic disk.

A model with a single F_\star across all three dates was also investigated, but did not reproduce the data to the same degree of accuracy as the fiducial model. Significant residuals were visible at the location of the star, and a stellar flux density varying by more than a factor of 2 over a period of months to years is preferred with 7.3σ confidence by the AICc and with $\Delta\text{BIC} = 35.0$.

4.3. Investigating Vertical Structure

The posterior distribution for the aspect ratio h suggests that the data are capable of measuring AU Mic’s scale height despite the mild degeneracy between the aspect ratio and other parameters like the radial structure and viewing geometry. Even when marginalized over these other parameters, the posterior distribution indicates a measured value of $h = 0.032 \pm 0.005$, which translates to a $\sim 6\sigma$ measurement of the scale height rather than an upper limit. At the ~ 40 au outer edge of the disk, this aspect ratio implies a vertical scale height of 1.3 ± 0.2 au. The corresponding disk FWHM is 3.0 ± 0.5 au, which is consistent with the mean image-domain FWHM of ~ 2.8 au for projected separations ≤ 35 au (Figure 2; due to the edge-on inclination

Table 3. MCMC Fitting Results

Parameter	Fiducial		Disk + Ring ^a	
	Median	Best Fit	Median	Best Fit
$\log M$ (M_\odot)	$-7.561^{+0.006}_{-0.006}$	-7.561	$-7.548^{+0.007}_{-0.007}$	-7.545
p	$0.9^{+0.5}_{-0.4}$	0.8	$0.8^{+0.6}_{-0.4}$	0.9
h	$0.032^{+0.005}_{-0.005}$	0.029	$0.027^{+0.004}_{-0.005}$	0.028
r_{in} (au)	$23.3^{+0.6}_{-1.0}$	23.4	$24.1^{+0.6}_{-0.9}$	23.9
r_{out} (au)	$41.5^{+0.4}_{-0.6}$	41.5	$42.3^{+0.4}_{-0.5}$	42.4
i ($^\circ$)	$88.6^{+0.4}_{-0.3}$	88.5	$88.27^{+0.22}_{-0.16}$	88.3
PA ($^\circ$)	$128.50^{+0.07}_{-0.07}$	128.50	$128.50^{+0.07}_{-0.07}$	128.48
March F_* (μJy)	390^{+20}_{-20}	390	390^{+20}_{-20}	370
August F_* (μJy)	150^{+30}_{-30}	150	150^{+20}_{-30}	140
June F_* (μJy)	230^{+20}_{-20}	230	220^{+20}_{-20}	210
r_{ring} (au)			$11.9^{+1.7}_{-1.8}$	10.8
M_{ring} ($M_\oplus \times 10^{-4}$)			$1.8^{+0.5}_{-0.7}$	1.7
$\ln \mathcal{L}$	-313083^{+2}_{-4}	-313078	-313077^{+2}_{-5}	-313072

^aThe new Gaia ring run hasn't finished yet, so these values aren't currently comparable to the fiducial values.

of the disk, the outer edge of the disk sets the apparent disk thickness at all projected separations).

To verify that we in fact measured the scale height, we investigate a model parameterization in which the scale height is set to a value well below ALMA's resolution limits. The aspect ratio is fixed at a value of 0.003, so that even at the outer edge of the disk the scale height is $\sim 3\%$ of the beam size along the disk's vertical axis. If the disk is in fact resolved by the observations, such a ‘skinny’ disk model should perform significantly worse than the fiducial model. Hence, we can quantify the statistical significance of our detection of the scale height by comparing the best-fit skinny model to the best-fit fiducial model. The best-fit model image and residuals are shown in Figure 5c. The skinny model results in a significantly poorer fit to the data than the fiducial model with variable aspect ratio, with the best fits differing at the 3.7σ level according to the AICc and by a value of 5.8 on the BIC.

5. DISCUSSION

Parametric modeling suggests that AU Mic’s debris disk is nearly edge on, exhibits an increasing surface density with radius until ~ 42 au, and reaches a maximum scale height of ~ 1.2 au. There is also marginal evidence for an additional annulus of dust at $r \sim 10$ au. Here we compare the results of our analysis with previous studies of AU Mic’s debris disk. Relevant quantities in the literature have been scaled by the new Gaia distance, although in most cases the scaling has no effect on the first significant figure. For cases where scaling by the Gaia distance changes any of the significant digits reported, the resulting value is referred to as “Gaia-corrected.”

5.1. Stellar Activity

Our analysis indicates with high confidence that AU Mic exhibited long-term millimeter variability over the ~ 1.5 yr observation period, with stellar flux densities preferred by our modeling varying by more than a factor of two (150_{-30}^{+20} μ Jy to 390 ± 20 μ Jy) from one observation to the next. Our values are comparable to the best-fit 1.28 mm flux density of 320 ± 60 μ Jy from MacGregor et al. (2013), which is equivalent to 290 ± 50 μ Jy when scaled to our observing wavelength of 1.35 mm assuming an effective stellar temperature of 3500 ± 100 K from Plavchan et al. (2009). It seems likely that this variability is distinct from the 6 minute flare spanning two orders of magnitude that occurred during the June observations. That being said, we cannot rule out the possibility that the observations caught the star in varying states of flare decay, especially as White (1996) note that shorter-wavelength radio flares tend to have slower decay timescales. Simulations of radio emission from low-mass stars allow for variability of more than a factor of two over the entire longitudinal extent of a star (Llama et al. 2018); it is also possible that different longitudinal regions of AU Mic were facing Earth during the three observations. Cox & Gibson (1985) detect small-scale ($\sim 50\%$) variations over a two-week period in 2 cm to 20 cm observations of AU Mic with the VLA, possibly due to ‘several independent mini-flares’ or rotational modulation of the star. In fact, factor of ~ 2 variability over months to years is typical of ‘quiescent’ microwave emission from cool stars (Guedel 1994). Regardless of the origin of AU Mic’s short- and long-term variability, we emphasize the importance of accounting for stellar emission/variability in observations of circumstellar disks around low-mass stars, especially in light of the Proxima Centauri flare discovered by MacGregor et al. (2018).

5.2. Dust & Gas Mass

MCMC fitting prefers a median dust mass of $9.08 \pm 0.13 \times 10^{-3} M_{\oplus}$ for an opacity of $2.3 \text{ cm}^2 \text{ g}^{-1}$. Our preferred dust mass tends to be slightly smaller than previous estimates; we note that the reported uncertainties take into account neither the 10% flux uncertainty nor assumptions about the dust opacity. Assuming an opacity of $2.7 \text{ cm}^2 \text{ g}^{-1}$ and a dust temperature of 25 K, MacGregor et al. (2013) estimate a dust mass of $\sim 0.01 M_{\oplus}$ from a 1.3 mm disk flux density of $7.14^{+0.12}_{-0.25} \text{ mJy}$. Matthews et al. (2015) report an identical mass with 20% uncertainty from a $850 \mu\text{m}$ disk flux density of $12.5 \pm 1.5 \text{ mJy}$ and an opacity of $1.7 \text{ cm}^2 \text{ g}^{-1}$. Liu et al. (2004) also take the opacity to be $1.7 \text{ cm}^2 \text{ g}^{-1}$, and report $0.011 M_{\oplus}$ from a disk flux density of $14.4 \pm 1.8 \text{ mJy}$ at $850 \mu\text{m}$. Finally, Strubbe & Chiang (2006) calculate $0.01 M_{\oplus}$ by deriving a steady-state collisional cascade grain size distribution and fitting the disk's surface brightness and thermal spectrum to *V*- and *H*-band HST observations.

Detections of gas in debris disks are becoming increasingly common, and are changing our understanding of dust morphology in gas-bearing systems (see §4 of Hughes et al. 2018, and references therein). Theoretical works such as that of Takeuchi & Artymowicz (2001) predict that small dust grain dynamics will be affected by the presence of gas for gas-to-dust ratios $\gtrsim 1$; Section 5.2 of Hughes et al. (2017) provides a discussion of the ways in which gas can interact with dust in debris disks. As discussed in §3, even a large gas excitation temperature of 250 K implies a 3σ upper limit on the ^{12}CO mass of only $8.7 \times 10^{-7} M_{\oplus}$. Combined with our preferred dust mass of $9.08 \times 10^{-3} M_{\oplus}$ and assuming a conservative CO/H₂ ratio of 10^{-4} , this CO mass corresponds to an upper limit of ~ 1 on the gas-to-dust ratio. Such a low gas-to-dust ratio excludes the possibility of gas influencing the millimeter grain kinematics, and thus our measurement of the scale height.

5.3. Disk Geometry

The geometric properties of the disk inferred from our modeling agree well with the literature. Our median inclination of $88.6^{+0.4}_{-0.3}$ is consistent with previous estimates of the disk's inclination within uncertainties (Metchev et al. 2005; Krist et al. 2005). The median PA of $128^\circ 49 \pm 0.07$ falls within uncertainties of measurements by MacGregor et al. (2013) and Krist et al. (2005); however, the limb-averaged PA interior to 50 au from Metchev et al. (2005) is $129^\circ 8 \pm 0.2$, and Schneider et al. (2014) report an optical-wavelength PA of $127^\circ 8 \pm 0.2$ between 50 au to 100 au.




Figure 6. A plot of the probability that the AU Mic disk has a given uncertainty. There is a 50% chance that the eccentricity of is less than 0.15, a 65% chance that it is less than 0.28, and a 95% chance that it is less than 0.78.

The eccentricity of the AU Mic debris disk can be constrained from the SE-NW ansa flux density ratio $f_{\text{SE-NW}} = \frac{340 \pm 15 \text{ mJy/beam}}{330 \pm 15 \text{ mJy beam}} = 1.035 \pm 0.065$ reported in §3. For a perfectly circular disk, the flux ratio for all pairs of points should be 1. For a disk with a nonzero eccentricity, the flux ratio will be largest between periapse and apoapse. When the major axis of the disk falls on the sky plane (perpendicular to the line of sight), $f_{\text{SE-NW}}$ is simply the apoapse-periapse ratio. If the major axis of the disk has any other orientation, $f_{\text{SE-NW}}$ is closer to unity than the apoapse-periapse ratio, and $f_{\text{SE-NW}}$ should equal one when the major axis points exactly along our line of sight.

For a disk with a small eccentricity, the fluxes at apoapse and periapse are still fairly close to one another and so the ratios for all orientations of the major axis disk fall within the measured 1σ confidence interval. The disk may also exhibit

a larger eccentricity, such that the apoapse-periapse flux ratio is greater than the observed upper limit of $f_{\text{SE-NW}} = 1.095$; then the disk must have its major axis oriented at an angle to the sky plane in order to satisfy the constraint on $f_{\text{SE-NW}}$. As the disk eccentricity increases further, the disk major axis must lie closer and closer to our line of sight for $f_{\text{SE-NW}}$ to fall within the observed range. Assuming all edge-on disk orientations are equally likely, higher eccentricities are thus less likely given the observed fluxes. On the other hand, all eccentricities below 0.13—the value at which the apoapse-periapse flux ratio is equal to the observed upper limit of $f_{\text{SE-NW}} = 1.095$ —are equally likely.

Figure 6 shows the probability distribution of the disk’s eccentricity given the observed value of $f_{\text{SE-NW}}$. We assume a disk radius of 40 au, and a stellar mass, effective temperature, and luminosity of $0.5 M_\odot$, 3500 K, and $0.09 L_\odot$ respectively. The grain size distribution is determined by a power law index $q = 3.5$, and grain absorptivity scales as λ^{-1} for wavelengths larger than the grain size. We find that there is a 50% probability that the eccentricity of the AU Mic debris disk is less than 0.15, a 65% probability that it is less than 0.28, and a 95% probability that it is less than 0.78.

5.4. Radial Structure

Modeling of the radial structure discussed in §4.2 is broadly consistent with the previous analysis of millimeter wavelength emission from the disk performed by MacGregor et al. (2013) as can be seen in Figure 7. However, the factor of ~ 2 increase in spatial resolution introduces additional interpretive complexities when comparing the results. Our outer radius ($41.5^{+0.4}_{-0.6}$ au) is significantly larger than the previously reported Gaia-corrected value of 38.9 ± 0.4 au. In contrast to our inner radius of $r_{in} = 23.8^{+0.6}_{-0.9}$ au, MacGregor et al. (2013) report a Gaia-corrected $r_{in} = 8.5^{+11.0}_{-1.0}$ au. Neither value of the inner radius is well constrained: the authors report a Gaia-corrected 3σ upper limit of ~ 20 au, whereas conversely we find a 2σ lower limit of ~ 1 au. Similar discrepancies arise in the posterior distribution for p , which is degenerate with r_{in} . While MacGregor et al. (2013) finds $p = 2.32^{+0.21}_{-0.31}$, we recover a median value of $p = 0.9^{+0.5}_{-0.4}$. In our analysis, the uncertainties in these two parameters arise from the existence of two families of solutions: one at $r_{in} \sim 23$ au, $r_{out} \sim 41$ au, $p \sim 0.9$ and another at $r_{in} \sim 6$ au, $r_{out} \sim 40$ au, $p \sim 2.8$ (Figure 7). The latter family, which has a lower likelihood, is probably associated with the local intensity maxima located at stellocentric separations of ~ 7.3 au to the




Figure 7. Comparison between the best-fit surface density profiles obtained in MacGregor et al. (2013) (dashed red line) and this work (solid blue line). Also included is the best-fit surface density profile associated with the lower-likelihood family of solutions (dotted black line). The vertical extent of the shaded regions marks 1σ confidence intervals on the surface density. The points designate the best-fit inner and outer radii for each model, and the horizontal extent of the more transparent shaded regions show 1σ confidence intervals on the inner and outer radii. Not included in the uncertainties are the 10% flux uncertainties of each observation.

NW and ~ 10.5 au to the SE. Our exploratory modeling of these features in §4 raises the possibility of an annulus of dust interior to the main disk—if such an annulus does exist, the ~ 2 times lower resolution observations from MacGregor et al. (2013) would probably not have been able to distinguish the annulus emission from that of the main disk. In fact, an unresolved annulus would likely bias the inner radius preferred by the authors’ modeling towards smaller values and could account for the differences in our characterization of the radial structure.

While the annulus model provides only a marginally significantly improved fit to the data, previous observations spanning a wide range of wavelengths have recovered

surface brightness enhancements at the same projected stellocentric separation (~ 10 au) as the hypothetical annulus. A local maximum is present to the SE of the star at this separation in lower-resolution ALMA observations by MacGregor et al. (2013), and there is a suggestive peak in the noise on the opposite side of the star as well. Although these surface brightness enhancements do not attain 3σ significance after subtraction of an axisymmetric model, the independent detection of these features in both data sets suggests that they may be real. Schneider et al. (2014) also observe an optical-wavelength ‘bump’ ~ 13 au SE of the star and slightly elevated from the disk midplane (i.e. to the NE). No matching surface brightness enhancement is observed on the NW side of the disk, which is obscured by the STIS occulting wedge for $r \lesssim 12$ au. On the other hand, the scattered light emission on the NW side is not symmetric about the midplane and the authors tentatively identify a warp below the midplane extending to a Gaia-corrected distance of ~ 43 au from the star. They note that these features may share a common cause, and posit that dust orbits in the inner disk are non-coplanar with those found at larger separations.

Near-infrared GPI observations presented in Wang et al. (2015) further corroborate the presence of a ‘bump’ to the SE, characterized by a FWHM roughly triple the FWHM at an equivalent separation on the NW side of the disk. No features are detected at a corresponding separation to the NW. Figure 4 of Wang et al. (2015) shows a composite map of the bump as seen by GPI, STIS, and ALMA; the common location of the bump in both scattered light and mm observations would indicate that the mm- and micron-sized grains are cospatial. The micron-sized grains traced by scattered light, and all larger bodies ranging up to the cm-to-m sizes characterizing the top of the collisional cascade, are thought to originate from a narrow ‘birth ring’ at ~ 40 au (Strubbe & Chiang 2006). Thus it is possible that all features are located at a true stellocentric separation of ~ 40 au, with the apparent stellocentric separation of ~ 10 au being due projection effects. In principle, the true stellocentric separation could be constrained by the range of temperatures allowed by AU Mic’s SED. However such an analysis would be complicated by the relatively small mass contribution of the feature, the need to incorporate constraints on the small grains from scattered light imaging, and assumptions about the grain size distribution, which is beyond the scope of this work.

Planets are often invoked to explain rings in debris disks; P. Plavchan et al. (in preparation) propose a Jovian-mass exoplanet candidate interior to 1 au, but AU

Mic's stellar activity make it difficult to confirm the radial velocity detection. It is unlikely that a planet so close to the star would be responsible for a ring at ~ 10 au. On the other hand, a planet might be related to AU Mic's fast-moving features in the point-source parent body picture of [Sezestre et al. \(2017\)](#). The < 1 au separation of the potential planet detection by Plavchan et al. is far interior to the preferred 8 au separation that [Sezestre et al. \(2017\)](#) derive to explain the origin of the fast-moving features, but separations < 5 au are not ruled out by the latter's analysis. The exoplanet proposed by Plavchan et al. (in preparation) seems an unlikely candidate to resolve the unexplained aspects of AU Mic's radial structure, although planetary interactions should not be wholly discounted.

5.5. Vertical Structure


As discussed in §4.3, our MCMC analysis yields a median aspect ratio of $h = 0.032 \pm 0.005$; at a reference radius of 40 au, this translates to a vertical scale height $H = 1.3 \pm 0.2$ au and a FWHM of 3.0 ± 0.5 au. The FWHM is thus only $\sim 2/3$ the size of the combined-data beam projected onto the vertical axis of the disk. While it may seem improbable that we measure a vertical thickness below the image-domain spatial resolution of the combined data, our results can be explained by the following observations. First, the beam size of the naturally weighted combined image ($0.52'' \times 0.39''$) is not wholly representative of the resolution of the data. The smallest naturally weighted beam FWHM from the three individual observations is $0.30'' = 2.9$ au (Table 1), while the spatial scale traced by the longest baseline is $0.22'' = 2.1$ au. Second, the scale height H refers to the disk height as measured from the midplane; the observable quantity that must be resolved in order to measure the scale height is actually the total vertical thickness $2H$. Third, the scale height represents the standard deviation of a Gaussian distribution of dust particles that in fact reaches well beyond the extent of the scale height. For example, considering that the peak signal-to-noise (SNR) of the data is ~ 23 and that the vertical distribution of dust is assumed to be Gaussian, the SNR should remain above 3 over a total vertical extent of $\sim 0.35''$ (3.4 au). The combination of these factors indicates that it is plausible that we are able to detect a scale height smaller than the resolution of the image-domain data.

Our ability to measure the disk's vertical thickness could also be affected by phase noise, which could cause the true angular resolution of the data to be worse than that given by the beam. [Boehler et al. \(2013\)](#) find that introducing Gaussian-distributed

phase noise to synthetic ALMA observations of highly-inclined proplanetary disks has little effect on the measured scale height value, although the corresponding uncertainties do grow. The authors also find that introducing phase noise dramatically increases model reduced χ^2 values; our low reduced χ^2 value of 1.027 provides an indication that phase noise does not affect our results. To further investigate the effects of phase noise on the resolution, we imaged the test quasars included in the ALMA data with the same clean parameters that were used to image the disk. We find that the image-domain FWHMs of the quasars observed during the two long-baseline observations are slightly larger than the corresponding beam FWHMs; however the beam-subtracted quasar FWHMs ($\sim 0.23'' \times 0.07''$) are still smaller than the mean beam-subtracted vertical disk FWHM ($0.28''$), indicating that phase noise does not prevent us from resolving the vertical structure of AU Mic's disk.

Our measured HWHM of 1.5 ± 0.2 au at $r = 40$ au is marginally consistent with values cited in the literature derived from observations spanning the optical to the sub-millimeter, ranging from roughly 0.2 au to 2 au. Schüppler et al. (2015) place an upper limit of 0.05 rad on the 1.3 mm opening angle (equivalent to the aspect ratio for small angles) by extracting the image-domain vertical profile from vertically unresolved ALMA observations presented in MacGregor et al. (2013), and estimate a visible-wavelength opening angle of 0.03 rad by reading off the vertical scale height from STIS observations of the disk in Schneider et al. (2014). Metchev et al. (2005) approximate a *H*-band Keck FWHM of ~ 4 au at a separation of 40 au; Krist et al. (2005) fit a vertical Lorentzian profile to multicolor HST observations of the disk and find the Gaia-corrected FWHM interior to 50 au to fall between 2.4 au and 3.4 au.

Because the measurements quoted above are determined from the observed vertical thickness of AU Mic's disk, they can be affected by the radial structure and viewing geometry of the disk as well as scattering effects. As such, parametric modeling provides a more reliable way to assess AU Mic's vertical structure, and must be taken into account in any comparison of our results with values in the literature. For example, Krist et al. (2005) report a FWHM between 1.73 au and 1.74 au at a separation of 20 au from three-dimensional scattering models. Metchev et al. (2005) find that a constant scale height of 0.8 au adequately reproduces the observed mean disk thickness. Collisional modeling can also be used to learn about AU Mic's vertical structure: collisional velocities—and thus dust production—are affected by the maximum eccentricity e_{max} of planetesimal orbits, which in turn can be related to

Figure 8.

Label	Reference	Wavelength	HWHM at 40 au	Methodology
1a	Schüppler et al. (2015)	< 70 μ m	2 au	Collisional modeling
1b	—	0.445 μ m–1.3 mm	0.6 au	Collisional modeling
1c	—	> 100 μ m	0.2 au	Collisional modeling
2a	Metchev et al. (2005)	1.63 μ m	2 au	Apparent
2b	—	0.0647 μ m–1.63 μ m	0.8 au	Model
3a	Krist et al. (2005)	0.0430 μ m–0.0833 μ m	1.5 au	Apparent
3b	—	—	0.9 au	Modeled
4	Schneider et al. (2014)	0.2 μ m–1.05 μ m	1.2 au	Apparent
5	MacGregor et al. (2013)	1.28 mm	\leq 2 au	Apparent
6	This work	1.35 mm	1.5 ± 0.2 au	Model

Table 4. Measurements of the vertical structure of AU Mic’s debris disk.

the disk vertical structure (see §5.6 below). Schüppler et al. (2015) perform such modeling constrained by photometric observations spanning visible to millimeter wavelengths and quote a reference model opening angle of 0.015 rad. The authors go on to note that an opening angle of 0.005 rad better reproduces the disk spectral energy distribution (SED) in the long-wavelength regime beyond 100 μ m, but is unable to reproduce flux measurements for $\lambda \leq 70 \mu$ m.

In sum, estimates of AU Mic’s vertical structure vary depending on the wavelength of observation, the techniques used, and the assumptions made. Rigorous comparison

between measurements in the literature is difficult—for example, Krist et al. (2005) assume a flared vertical profile while other authors use a linear parameterization for the scale height. Nevertheless, some general statements may be made. HWHM values determined from optical and near-infrared observations range from roughly 0.8 au to 2 au at a radius of 40 au, while the HWHMs determined at least in part from mm observations range from 0.2 au to 2 au, the latter being an upper limit. Estimates based on millimeter observations (including this work) tend to be smaller than those derived from shorter-wavelength observations, although the two wavelength regimes do not provide radically different values. We might explain this result by reasoning that shorter-wavelength data probe smaller particles, and smaller particles experience higher collision rates and therefore stronger damping of random velocities.

In recent years, the vertical structure of several other debris disks has been tentatively resolved with ALMA. Observations of CO J = 3 – 2 and 2 – 1 line emission from the edge-on β Pic debris disk at a spatial resolution of ~ 5.5 au presented by Matrà et al. (2017) suggest that the disk is resolved in the vertical direction. The beam-subtracted apparent FWHM, determined in a similar manner as the bottom pane of Figure 2 in this work, ranges between ~ 7 au and 12 au over the ~ 100 au radial extent of the disk. Assuming Keplerian rotation and an edge-on inclination, the authors report the following scale height-radius relation:

$$H = 7.0 \pm 0.6 \times \left(\frac{R}{85 \text{ au}} \right)^{0.75 \pm 0.02} \text{ au} \quad (4)$$

where the scale height H is the standard deviation of the Gaussian vertical density distribution. For reference to our work, this corresponds to a scale height of 4 au at a separation of 40 au.

As pointed out in §4.9 of Marino et al. (2016), the scale height of a narrow ring may be resolved even if the ring is not close to edge-on. The authors estimate that for ~ 0.25 arcsec resolution observations of the low-inclination ($i \approx 30^\circ$) HD 181327 debris disk at a SNR of ~ 50 , an aspect ratio of $h = 0.083$ could be constrained to within ± 0.005 . Along the same lines, Kennedy et al. (2018) analyze 11.6 au \times 13.1 au spatial resolution observations of the moderately-inclined ($i \approx 77^\circ$) HR 4796A debris disk, reporting a marginal detection of the disk scale height. The authors find that a vertically resolved ring (FWHM = 7 ± 1 au) is favored over a vertically unresolved ring (FWHM < 4 au) with $\Delta BIC = 6.8$, meeting the condition for ‘strong’ evidence of a statistically higher-quality fit ($\Delta BIC > 6$). This FWHM corresponds to a

Gaussian standard deviation $H \approx 3.0 \pm 0.4$ au and, at the ~ 80 au radial location of the ring, a scale factor of $h \approx 0.038 \pm 0.005$. In sum, the vertical structure of debris disks is just beginning to be resolved in the millimeter thanks to the improvements in sensitivity and resolution provided by ALMA; to our knowledge, AU Mic exhibits the narrowest millimeter-wavelength scale height of any known debris disk.

Though dust dynamics in protoplanetary disks are governed by different physics than in gas-poor debris disks, observations of protoplanetary disk scale heights provide an interesting point of comparison to our results. Dust settling in protoplanetary disks should lead to a wavelength-dependence in protoplanetary dust scale height measurements similar to that predicted by Thébault (2009), although for different reasons. Boehler et al. (2013) find that at a reference radius of 100 au around a $1 M_\odot$ star, micron-sized grains have the same scale height as the gas—14.6 au. On the other hand, 1 mm grains exhibit a scale height of only ~ 0.9 au, corresponding to an aspect ratio $h \sim 0.009$. Observations of protoplanetary disks often indicate a flared disk geometry, with typical scale heights of ~ 10 au for small grains and ~ 1 to 6 au for large grains at a stellocentric separation of 100 au (Pinte et al. 2008, 2016; Gräfe et al. 2013). As such, our preferred 1.3 mm aspect ratio $h \sim 0.032$ is comparable to the aspect ratios for corresponding grain sizes in protoplanetary disks.

5.6. Inferring the Mass of Stirring Bodies

Information regarding the bodies dynamically stirring AU Mic’s disk can be recovered by relating the scale height to the dynamical excitation of the disk’s dust grains. As discussed by Thébault (2009) and Quillen et al. (2007), the planetesimals responsible for stirring the disk impart kinetic energy to the dust, perturbing them from a Keplerian orbit and thus increasing their orbital eccentricity dispersion $\langle e^2 \rangle$. Here we define $\bar{e} = \sqrt{\langle e^2 \rangle}$ and $\bar{i} = \sqrt{\langle i^2 \rangle}$, where $\langle i^2 \rangle$ is the inclination dispersion of the dust grain orbits. In equilibrium there is an equipartition between the vertical and in-plane components of the velocities imparted to the grains, so $\bar{i} = \bar{e}/2$. Inclination can be related to the observed FWHM using $\bar{i} = \sqrt{2}\text{FWHM}$ for small angles; thus inclination is related to our Gaussian-standard-deviation aspect ratio by $\bar{i} \approx 2.355h/\sqrt{2}$. The interparticle relative velocity $\langle v_{rel} \rangle$ can then be determined directly from observables using the following relation (Wetherill & Stewart 1993; Wyatt & Dent 2002):

$$\langle v_{rel} \rangle \approx v_{Kep}(r)\sqrt{\bar{i}^2 + 1.25\bar{e}^2} \approx v_{Kep}(r)2.355\sqrt{3}h \quad (5)$$

where $v_{Kep}(r)$ is the Keplerian velocity at radius r . Adopting a stellar mass of $0.5 M_\odot$ and taking $r = 40$ au, $h = 0.032 \pm 0.005$ yields $v_{rel} = 430 \pm 70$ m/s.

The velocity dispersion of the dust grains will be excited to about the escape velocity of the largest bodies governing the disk dynamics in the absence of any significant damping of their velocity dispersion. This result arises because viscous stirring has a larger cross section (i.e. shorter characteristic timescale) than collisions as long as the velocity dispersion is less than the escape velocity of the largest bodies that dominate the stirring. The two cross sections (and timescales) become comparable as v_{rel} approaches v_{esc} and collisions start to dominate, limiting the growth of v_{rel} to about v_{esc} (e.g., [Schlichting 2014](#)). As such, we can use our estimate for v_{rel} to place a lower limit on the escape velocity v_{esc} and thus size a_{big} of bodies stirring the disk. Assuming a density of 2 g cm^{-3} , we find $a_{big} \sim 410 \pm 60$ km and $m_{big} \sim 5.8 \pm 2.7 \times 10^{20}$ kg, i.e. $\sim 4\%$ the mass of Pluto.

On the other hand, the disk may be in a steady state in which velocity damping is balanced with excitation (rather than damping being inefficient). Under this condition, the scale height provides an estimate of the total mass of stirring bodies rather than their size. To convert the observed disk scale height into an upper limit on the total mass of bodies stirring the disk, we refer to theoretical models of steady state size-dependent velocity distributions in the collisional cascade described in [Pan & Schlichting \(2012\)](#). We assume a disk in steady state in which velocity excitation (“stirring”) occurs via close encounters with the planet(s) and velocity damping occurs via direct collisions with other (small) bodies. Because the disk is assumed to be in steady state, we can equate the expressions for stirring and damping rates and solve for the velocity of dust grains. The computation equating the stirring and damping is analogous to that of Equations 15 & 16 in [Pan & Schlichting \(2012\)](#); however, while Equations 15 & 16 assume that catastrophic collisional destruction dominates the velocity damping, we assume that *all* collisions with smaller bodies contribute to damping. We then relate the dust grain velocity to the disk scale height as described above. Assuming a scale height of 1.2 au at 40 au, a stellar mass of $0.5 M_\odot$, and a dust mass of $0.01 M_\oplus$, these models indicate a total dynamical mass of $\sim 1.5 M_\oplus$. If all of the $1.5 M_\oplus$ were locked up in a single body, it would have a radius of $\sim 1.1 R_\oplus$ assuming a mean density of 5.5 g cm^{-3} characteristic of Earth.

In sum, the largest bodies in the AU Mic disk cannot be smaller than 410 ± 60 km or they would not be able to stir the dust grains to the velocity dispersion inferred

from the measured scale height. Conversely, the largest body residing within the disk cannot be larger than $\sim 1.5 M_{\oplus}$ or the scale height would exceed the value measured in this work. It should also be noted that because the scale height measured at the outer edge of the disk is comparable to the spatial resolution of the data, and since scale height decreases with decreasing radius, the observations are likely only sensitive to the scale height near the ~ 40 au outer disk radius. If the disk is stirred by an ensemble of 410 km planetesimals, they would necessarily be located within this outer region of the disk. Bodies external to the disk are also capable of stirring the disk at a distance, through secular and/or mean-motion resonant perturbations; consideration of external perturbers is deferred to future work. If external stirring is significant, then the above constraints on the properties of embedded (non-external) perturbers are all upper limits.

The properties of the perturbers estimated in this work are consistent with those inferred by Chiang & Fung (2017) on independent grounds using the time-variable infrared observations by Boccaletti et al. (2015, 2018). Our lower limit on the perturbing body size of ~ 410 km fits with the ~ 400 km radius for their Varuna-like progenitor (for reference, Varuna is among the larger objects in the Kuiper belt). Moreover, there should be on the order of 10^3 such objects according to their scenario [the argument is as follows: since the lifetime of the debris left from the catastrophic disruption of a Varuna-sized progenitor is $\sim 3 \times 10^4$ yr (Chiang & Fung 2017, their equation 21), and since the age of the AU Mic system is ~ 20 Myr, there must have been $20 \text{ Myr} / 3 \times 10^4 \text{ yr} \sim 700$ such progenitors over the system lifetime. A population on the order of 10^3 bodies then yields an order-unity probability for observing the aftermath of a single catastrophic collision today.] An ensemble of $\sim 10^3$ bodies each with a mass of $4\pi/3 \cdot 2.5 \text{ g cm}^{-3} \cdot (400 \text{ km})^3 \sim 1 \times 10^{-3} M_{\oplus}$ amounts to a total mass of $\sim 0.1 M_{\oplus}$, safely below the upper limit of $1.5 M_{\oplus}$ established by our present work.

6. CONCLUSION

We have presented new 1.3 mm ALMA observations of thermal dust emission from the debris disk around AU Mic at nearly double the angular resolution of previous observations. Both the vertical and radial structure of the disk are resolved. We see no indication of a millimeter complement to the fast-moving features detected in the optical by Boccaletti et al. (2015), but the data are suggestive of an additional ring of dust at ~ 10 au. Models with a ring interior to the main disk provide a better fit

to the data, but measures of statistical significance remain equivocal as to whether such models minimize the information lost in the modeling process—possibly due to eccentricity or non-coplanarity of the feature. MCMC analysis prefers an aspect ratio $h = 0.032$, corresponding to a vertical scale height $H = 1.3$ au in the outer regions of the disk. Our analysis suggests that this is not an upper limit; a model with vertically resolved structure provides a statistically improved fit to the data over a model with unresolved vertical structure at a 4σ confidence level. Furthermore, the disk vertical FWHM derived from parametric modeling corresponds well with image-domain estimates of the beam-subtracted FWHM of the emission perpendicular to the disk plane.

By comparing our measurement of the scale height to the steady-state collisional modeling of Pan & Schlichting (2012) we are able to place constraints on the mass and sizes of bodies stirring AU Mic’s disk. In the lower-limit case where collisional velocity damping is inefficient, the stirring bodies would have a radius of 410 ± 60 km, corresponding to a characteristic mass ~ 20 times less than that of Pluto. On the other hand, velocity damping may balance stirring; this condition allows us to place an upper limit of $\sim 1.5 M_{\oplus}$ on the mass of stirring bodies. This is a dynamical mass, distinct from the lunar mass of dust grains inferred from the millimeter flux density of the disk. These results rule out the presence of a gas giant or Neptune analog embedded within the outer disk, but are suggestive of a significant population of large asteroids having a combined mass of less than one Earth (this is also an upper limit on the mass of embedded objects because it neglects the possibility of perturbations by external bodies). Such a population has been inferred on independent grounds using time-variable infrared observations (Chiang & Fung 2017).

Looking forward, the scale height measurement presented in this work could be combined with other measurements of AU Mic’s scale height at widely-separated (sub)millimeter wavelengths. This would allow the size-dependent velocity dispersion and internal strengths of bodies in AU Mic’s collisional cascade to be constrained, testing the assumption of collisional cascade theory that velocity dispersion is constant with grain size. The $450 \mu\text{m}$ ALMA Band 9 observations of AU Mic presented in the forthcoming work of Carter et al. (in preparation) may be able to provide such constraints. Our measurements of the AU Mic system provide a proof of concept that spatially resolved observations of the vertical structure at millimeter wavelengths can constrain the presence of Uranus and Neptune analogs and even large Kuiper belt

object analogues, which are undetectable by standard planet-detection techniques. Applying this technique to other high-inclination debris disks with a range of central stellar masses will provide unique constraints on the prevalence of large perturbing bodies throughout the Galaxy.

ACKNOWLEDGEMENTS

C.D. is sponsored by a NASA CT Space Grant Undergraduate Research Fellowship and Wesleyan University’s Research in the Sciences Fellowship. C.D., A.M.H., E.C., and K.F. gratefully acknowledge support from NSF grant AST-1412647. M.P. gratefully acknowledges support from NASA grants NNX15AM35G and NNX15AK23G. H.E.S gratefully acknowledges support from NASA grant NNX15AK23G. E.I.C. acknowledges support from the National Science Foundation. J.M.C. acknowledges support from the National Aeronautics and Space Administration under grant No. 15XRP15_20140 issued through the Exoplanets Research Program. This paper makes use of the following ALMA data: ADS/JAO.ALMA#2012.1.00198.S. ALMA is a partnership of ESO (representing its member states), NSF (USA) and NINS (Japan), together with NRC (Canada), MOST and ASIAA (Taiwan), and KASI (Republic of Korea), in cooperation with the Republic of Chile. The Joint ALMA Observatory is operated by ESO, AUI/NRAO and NAOJ. The National Radio Astronomy Observatory is a facility of the National Science Foundation operated under cooperative agreement by Associated Universities, Inc. This work has made use of data from the European Space Agency (ESA) mission *Gaia* (<https://www.cosmos.esa.int/gaia>), processed by the *Gaia* Data Processing and Analysis Consortium (DPAC, <https://www.cosmos.esa.int/web/gaia/dpac/consortium>). Funding for the DPAC has been provided by national institutions, in particular the institutions participating in the *Gaia* Multilateral Agreement.

Software: CASA (McMullin et al. 2007), MIRIAD (Sault et al. 1995), NumPy (Van Der Walt et al. 2011), Astropy (The Astropy Collaboration et al. 2018), Pandas (McKinney 2010), emcee (Foreman-Mackey et al. 2013), Uncertainties, <http://pythonhosted.org/uncertainties/>

REFERENCES

- Artymowicz, P. 1997, Annual Review of Earth and Planetary Sciences, 25, 175, doi: [10.1146/annurev.earth.25.1.175](https://doi.org/10.1146/annurev.earth.25.1.175)
- Augereau, J.-C., & Beust, H. 2006, A&A, 455, 987, doi: [10.1051/0004-6361:20054250](https://doi.org/10.1051/0004-6361:20054250)
- Beckwith, S. V. W., Sargent, A. I., Chini, R. S., & Gusten, R. 1990, AJ, 99, 924, doi: [10.1086/115385](https://doi.org/10.1086/115385)
- Binks, A. S., & Jeffries, R. D. 2014, MNRAS, 438, L11, doi: [10.1093/mnrasl/slt141](https://doi.org/10.1093/mnrasl/slt141)
- Boccaletti, A., Thalmann, C., Lagrange, A.-M., et al. 2015, Nature, 526, 230, doi: [10.1038/nature15705](https://doi.org/10.1038/nature15705)
- Boccaletti, A., Sezestre, E., Lagrange, A. M., et al. 2018, ArXiv e-prints, arXiv:1803.05354. <https://arxiv.org/abs/1803.05354>
- Boehler, Y., Dutrey, A., Guilloteau, S., & Piétu, V. 2013, MNRAS, 431, 1573, doi: [10.1093/mnras/stt278](https://doi.org/10.1093/mnras/stt278)
- Brown, M. E. 2001, AJ, 121, 2804, doi: [10.1086/320391](https://doi.org/10.1086/320391)
- Chiang, E., & Fung, J. 2017, ApJ, 848, 4, doi: [10.3847/1538-4357/aa89e6](https://doi.org/10.3847/1538-4357/aa89e6)
- Cox, J. J., & Gibson, D. M. 1985, in IN: Radio stars; Proceedings of the Workshop on Stellar Continuum Radio Astronomy, Boulder, CO, August 8-10, 1984 (A86-38003 17-90). Dordrecht, D. Reidel Publishing Co., 1985, p. 233-236., Vol. 116, 233–236
- Cranmer, S. R., Wilner, D. J., & MacGregor, M. A. 2013, ApJ, 772, 149, doi: [10.1088/0004-637X/772/2/149](https://doi.org/10.1088/0004-637X/772/2/149)
- Ercolano, B., & Pascucci, I. 2017, Royal Society Open Science, 4, 170114, doi: [10.1098/rsos.170114](https://doi.org/10.1098/rsos.170114)
- Flaherty, K. M., Hughes, A. M., Rosenfeld, K. A., et al. 2015, ApJ, 813, 99, doi: [10.1088/0004-637X/813/2/99](https://doi.org/10.1088/0004-637X/813/2/99)
- Flaherty, K. M., Hughes, A. M., Rose, S. C., et al. 2017, ApJ, 843, 150, doi: [10.3847/1538-4357/aa79f9](https://doi.org/10.3847/1538-4357/aa79f9)
- Foreman-Mackey, D., Hogg, D. W., Lang, D., & Goodman, J. 2013, PASP, 125, 306, doi: [10.1086/670067](https://doi.org/10.1086/670067)
- Gaia Collaboration, Brown, A. G. A., Vallenari, A., et al. 2018, ArXiv e-prints, arXiv:1804.09365. <https://arxiv.org/abs/1804.09365>
- Gaia Collaboration, Prusti, T., de Bruijne, J. H. J., et al. 2016, A&A, 595, A1, doi: [10.1051/0004-6361/201629272](https://doi.org/10.1051/0004-6361/201629272)
- Goodman, J., & Weare, J. 2010, Communications in Applied Mathematics and Computational Science, Vol. 5, No. 1, p. 65-80, 2010, 5, 65, doi: [10.2140/camcos.2010.5.65](https://doi.org/10.2140/camcos.2010.5.65)
- Gräfe, C., Wolf, S., Guilloteau, S., et al. 2013, A&A, 553, A69, doi: [10.1051/0004-6361/201220720](https://doi.org/10.1051/0004-6361/201220720)
- Guedel, M. 1994, The Astrophysical Journal Supplement Series, 90, 743, doi: [10.1086/191899](https://doi.org/10.1086/191899)
- Houdebine, E. R., & Doyle, J. G. 1994, A&A, 289, 185
- Hughes, A. M., Duchene, G., & Matthews, B. 2018, ArXiv e-prints. <https://arxiv.org/abs/1802.04313>
- Hughes, A. M., Lieman-Sifry, J., Flaherty, K. M., et al. 2017, ApJ, 839, 86, doi: [10.3847/1538-4357/aa6b04](https://doi.org/10.3847/1538-4357/aa6b04)
- Kalas, P., Liu, M. C., & Matthews, B. C. 2004, Science, 303, 1990, doi: [10.1126/science.1093420](https://doi.org/10.1126/science.1093420)

- Kennedy, G. M., Marino, S., Matrà, L., et al. 2018, MNRAS, 475, 4924, doi: [10.1093/mnras/sty135](https://doi.org/10.1093/mnras/sty135)
- Krist, J. E., Ardila, D. R., Golimowski, D. A., et al. 2005, AJ, 129, 1008, doi: [10.1086/426755](https://doi.org/10.1086/426755)
- Liu, M. C., Matthews, B. C., Williams, J. P., & Kalas, P. G. 2004, ApJ, 608, 526, doi: [10.1086/392531](https://doi.org/10.1086/392531)
- Llama, J., Jardine, M. M., Wood, K., Hallinan, G., & Morin, J. 2018, ApJ, 854, 7, doi: [10.3847/1538-4357/aaa59f](https://doi.org/10.3847/1538-4357/aaa59f)
- MacGregor, M. A., Weinberger, A. J., Wilner, D. J., Kowalski, A. F., & Cranmer, S. R. 2018, ApJ, 855, doi: [10.3847/2041-8213/aaad6b](https://doi.org/10.3847/2041-8213/aaad6b)
- MacGregor, M. A., Wilner, D. J., Rosenfeld, K. A., et al. 2013, ApJL, 762, L21, doi: [10.1088/2041-8205/762/2/L21](https://doi.org/10.1088/2041-8205/762/2/L21)
- Malo, L., Doyon, R., Feiden, G. A., et al. 2014, ApJ, 792, 37, doi: [10.1088/0004-637X/792/1/37](https://doi.org/10.1088/0004-637X/792/1/37)
- Mamajek, E. E., & Bell, C. P. M. 2014, MNRAS, 445, 2169, doi: [10.1093/mnras/stu1894](https://doi.org/10.1093/mnras/stu1894)
- Marino, S., Matrà, L., Stark, C., et al. 2016, MNRAS, 460, 2933, doi: [10.1093/mnras/stw1216](https://doi.org/10.1093/mnras/stw1216)
- Matrà, L., Dent, W. R. F., Wyatt, M. C., et al. 2017, MNRAS, 464, 1415, doi: [10.1093/mnras/stw2415](https://doi.org/10.1093/mnras/stw2415)
- Matthews, B. C., Kennedy, G., Sibthorpe, B., et al. 2015, ApJ, 811, 100, doi: [10.1088/0004-637X/811/2/100](https://doi.org/10.1088/0004-637X/811/2/100)
- McKinney, W. 2010, in Proceedings of the 9th Python in Science Conference, ed. S. van der Walt & J. Millman, 51 – 56
- McMullin, J. P., Waters, B., Schiebel, D., Young, W., & Golap, K. 2007, in Astronomical Data Analysis Software and Systems XVI ASP Conference Series, Vol. 376, proceedings of the conference held 15-18 October 2006 in Tucson, Arizona, USA. Edited by Richard A. Shaw, Frank Hill and David J. Bell., p.127, Vol. 376, 127
- Metchev, S. A., Eisner, J. A., Hillenbrand, L. A., & Wolf, S. 2005, ApJ, 622, 451, doi: [10.1086/427869](https://doi.org/10.1086/427869)
- Milli, J., Lagrange, A.-M., Mawet, D., et al. 2014, A&A, 566, A91, doi: [10.1051/0004-6361/201323130](https://doi.org/10.1051/0004-6361/201323130)
- Montesinos, B., Eiroa, C., Krivov, A. V., et al. 2016, A&A, 593, A51, doi: [10.1051/0004-6361/201628329](https://doi.org/10.1051/0004-6361/201628329)
- Moshir et al. 1990, in IRAS Faint Source Catalogue, version 2.0 (1990)
- Olofsson, J., Samland, M., Avenhaus, H., et al. 2016, A&A, 591, A108, doi: [10.1051/0004-6361/201628196](https://doi.org/10.1051/0004-6361/201628196)
- Pan, M., & Schlichting, H. E. 2012, ApJ, 747, 113, doi: [10.1088/0004-637X/747/2/113](https://doi.org/10.1088/0004-637X/747/2/113)
- Pinte, C., Dent, W. R. F., Ménard, F., et al. 2016, ApJ, 816, 25, doi: [10.3847/0004-637X/816/1/25](https://doi.org/10.3847/0004-637X/816/1/25)
- Pinte, C., Padgett, D. L., Ménard, F., et al. 2008, A&A, 489, 633, doi: [10.1051/0004-6361:200810121](https://doi.org/10.1051/0004-6361:200810121)
- Plavchan, P., Werner, M. W., Chen, C. H., et al. 2009, ApJ, 698, 1068, doi: [10.1088/0004-637X/698/2/1068](https://doi.org/10.1088/0004-637X/698/2/1068)
- Quillen, A. C., Morbidelli, A., & Moore, A. 2007, MNRAS, 380, 1642, doi: [10.1111/j.1365-2966.2007.12217.x](https://doi.org/10.1111/j.1365-2966.2007.12217.x)
- Rosenfeld, K. A., Andrews, S. M., Hughes, A. M., Wilner, D. J., & Qi, C. 2013, ApJ, 774, 16, doi: [10.1088/0004-637X/774/1/16](https://doi.org/10.1088/0004-637X/774/1/16)

- Sai, S., Itoh, Y., Fukagawa, M., Shibai, H., & Sumi, T. 2015, PASJ, 67, 20, doi: [10.1093/pasj/psu152](https://doi.org/10.1093/pasj/psu152)
- Sault, R. J., Teuben, P. J., & Wright, M. C. H. 1995, in Astronomical Society of the Pacific Conference Series, Vol. 77, Astronomical Data Analysis Software and Systems IV, ed. R. A. Shaw, H. E. Payne, & J. J. E. Hayes, 433
- Schlichting, H. E. 2014, ApJ, 795, L15, doi: [10.1088/2041-8205/795/1/L15](https://doi.org/10.1088/2041-8205/795/1/L15)
- Schneider, G., Grady, C. A., Hines, D. C., et al. 2014, AJ, 148, 59, doi: [10.1088/0004-6256/148/4/59](https://doi.org/10.1088/0004-6256/148/4/59)
- Schüppler, C., Löhne, T., Krivov, A. V., et al. 2015, A&A, 581, A97, doi: [10.1051/0004-6361/201525664](https://doi.org/10.1051/0004-6361/201525664)
- Sezestre, É., Augereau, J.-C., Boccaletti, A., & Thébault, P. 2017, ArXiv e-prints. <https://arxiv.org/abs/1707.09761>
- Strubbe, L. E., & Chiang, E. I. 2006, ApJ, 648, 652, doi: [10.1086/505736](https://doi.org/10.1086/505736)
- Takeuchi, T., & Artymowicz, P. 2001, ApJ, 557, 990, doi: [10.1086/322252](https://doi.org/10.1086/322252)
- The Astropy Collaboration, Price-Whelan, A. M., Sipócz, B. M., et al. 2018, ArXiv e-prints. <https://arxiv.org/abs/1801.02634>
- Thébault, P. 2009, A&A, 505, 1269, doi: [10.1051/0004-6361/200912396](https://doi.org/10.1051/0004-6361/200912396)
- Thébault, P., & Augereau, J.-C. 2007, A&A, 472, 169, doi: [10.1051/0004-6361:20077709](https://doi.org/10.1051/0004-6361:20077709)
- Van Der Walt, S., Colbert, S. C., & Varoquaux, G. 2011, Computing in Science & Engineering, 13, 22
- Wang, J. J., Graham, J. R., Pueyo, L., et al. 2015, ApJL, 811, L19, doi: [10.1088/2041-8205/811/2/L19](https://doi.org/10.1088/2041-8205/811/2/L19)
- Wetherill, G. W., & Stewart, G. R. 1993, Icarus, 106, 190, doi: [10.1006/icar.1993.1166](https://doi.org/10.1006/icar.1993.1166)
- White, S. M. 1996, in Cool stars; stellar systems; and the sun : 9 : Astronomical Society of the Pacific Conference Series, volume 109; Proceedings of the 9th Cambridge workshop; held 3-6 October 1995 in Florence; Italy; San Francisco: Astronomical Society of the Pacific (ASP); —c1996; edited by Roberto Pallavicini and Andrea K. Dupree, p.21, Vol. 109, 21
- Williams, J. P., & Cieza, L. A. 2011, ARA&A, 49, 67, doi: [10.1146/annurev-astro-081710-102548](https://doi.org/10.1146/annurev-astro-081710-102548)
- Wyatt, M. C. 2008, ARA&A, 46, 339, doi: [10.1146/annurev.astro.45.051806.110525](https://doi.org/10.1146/annurev.astro.45.051806.110525)
- Wyatt, M. C., & Dent, W. R. F. 2002, MNRAS, 334, 589, doi: [10.1046/j.1365-8711.2002.05533.x](https://doi.org/10.1046/j.1365-8711.2002.05533.x)