

NON-KEPLERIAN MOLECULAR GAS MOTIONS IN THE INNER ($\lesssim 100$ AU) DISK AROUND HD 100546

CATHERINE WALSH^{1,2}, CAIL DALEY³, STEFANO FACCHINI⁴, & ATTILA JÚHASZ⁵

¹School of Physics and Astronomy, University of Leeds, Leeds, LS2 9JT, UK

²Leiden Observatory, Leiden University, P. O. Box 9531, 2300 RA Leiden, The Netherlands

³Astronomy Department, Wesleyan University, 96 Foss Hill Drive, Middletown, CT 06459, USA

⁴Max-Planck-Institut für Extraterrestrische Physik, Giessenbackstrasse 1, 85748 Garching, Germany

⁵Institute of Astronomy, University of Cambridge, Madingley Road, Cambridge CB3 0HA, UK

ABSTRACT

We present spatially resolved Atacama Large Millimeter/submillimeter Array (ALMA) images of ^{12}CO $J = 3 - 2$ emission from the protoplanetary disk around the Herbig Ae star, HD 100546. We expand upon earlier analyses of this data and model the spatially-resolved kinematic structure (i.e., the first moment map) of the CO emission. Assuming a velocity profile which prescribes a flat or flared emitting surface in Keplerian rotation, we uncover significant residuals with a peak of $\approx 7\delta v$, where $\delta v = 0.21 \text{ km s}^{-1}$ is the width of a single spectral resolution element. The shape and extent of the residuals reveal the possible presence of a severely warped and twisted inner disk extending to at most $90 - 100 \text{ au}$ (**it strikes me as a little awkward to give a range for a maximum value.. what if we just say ‘at most 100’?**). Adapting the model to include a misaligned inner disk with (i) an inclination almost edge-on to the line of sight and (ii) a position angle almost orthogonal to that of the outer disk reduces the residuals to $< 3\delta v$. HD 100546 joins several other protoplanetary disks that exhibit non-Keplerian kinematics as evidenced by molecular emission in the inner region. However, our findings are contrasted by recent VLT/SPHERE, MagAO/GPI, and VLTI/PIONIER observations of HD 100546 that show no evidence of a severely misaligned inner *dust* disk down to spatial scales of $\sim 1 \text{ au}$. An alternative explanation for the observed kinematics are fast radial flows across the inner dust cavity. Regardless of the process, the most likely cause is the presence of an unseen planetary companion.

1. INTRODUCTION

Observations of protoplanetary disks around nearby young stars offer unique insight into the initial conditions of planetary system formation. Resolved continuum observations spanning optical to cm wavelengths reveal the spatial distribution of dust across a range of grain sizes, which in turn, can highlight signposts of ongoing planet formation and/or as yet unseen massive companions/planets (e.g., cavities, gaps, rings, and spirals; see the recent reviews by [Espaillat et al. 2014](#), [Andrews 2015](#), and [Grady et al. 2015](#)). Likewise, spectrally and spatially resolved observations of molecular line emission disclose the spatial distribution and excitation of various gas species, from which information on disk gas properties can be extracted (e.g., [Dutrey et al. 2014](#); [Sicilia-Aguilar et al. 2016](#)).

Second only to H_2 in gas-phase molecular abundance, CO is a powerful diagnostic of various properties including the disk gas mass, radial surface density, and temperature. The primary isotopologue ^{12}CO is optically thick, and thus emits from the warm disk atmosphere; this allows the gas temperature in this region to be derived. (e.g., [Williams & Cieza 2011](#); [Dutrey et al. 2014](#)). The rarer isotopologues (^{13}CO , C^{18}O , C^{17}O and $^{13}\text{C}^{18}\text{O}$) have progressively lower opacities and so enable penetration towards and into the disk midplane (see, e.g., recent theoretical studies by [Bruderer 2013](#), [Miotello et al. 2016](#), and [Yu et al. 2016](#)). In observations with sufficiently high spatial resolution, now routine with ALMA, this allows a direct determination of the location of the CO snowline with high precision (see, e.g., [Nomura et al. 2016](#); [Schwarz et al. 2016](#); [Zhang et al. 2017](#)). However, it has been demonstrated recently that chemistry, in particular isotope-selective photodissociation ([Visser et al. 2009](#)), can complicate the extraction of disk gas masses from CO isotopologue emission ([Miotello et al. 2014, 2016](#)). Chemical conversion of CO to a less volatile

form, e.g., CO_2 , complex organic molecules, or hydrocarbons, is an alternative explanation for apparently low disk masses derived from CO observations (Helling et al. 2014; Furuya & Aikawa 2014; Eistrup et al. 2016; Yu et al. 2017).

Because emission from ^{12}CO (and often ^{13}CO) at (sub)mm wavelengths is bright, it has historically been used as a tracer of disk kinematics allowing a dynamical determination of the mass of the central star (e.g., Simon et al. 2000). However, gas motion can deviate from that expected due to Keplerian rotation alone because of a variety of different physical effects that can be inferred from spatially-resolved observations. These include spiral density waves, a substantial (and thus measurable) gas pressure gradient, radial flows mediated by accreting planets across cavities, or a disk warp (see, e.g., Rosenfeld et al. 2012; Tang et al. 2012; Casassus et al. 2013; Rosenfeld et al. 2014; Christiaens et al. 2014; Casassus et al. 2015). Spirals, radial flows, and warps can all signify the presence of (potentially massive) planetary companions; hence, perturbations from Keplerian motion traced in bright and spectrally- and spatially-resolved CO emission may expose unseen planets.

Here, we present high signal-to-noise and spectrally-resolved ALMA Cycle 0 images of ^{12}CO $J = 3 - 2$ emission from the protoplanetary disk around the nearby Herbig Ae star, HD 100546. The HD 100546 disk has been proposed to host (at least) two massive companions (see, e.g., Ake & van den Ancker 2006; Quanz et al. 2013; Walsh et al. 2014). However, recent MagAO/GPI observations presented in Rameau et al. (2017) have raised doubt on the previous identification of the point source at 50 au as a (proto)planet by two previous and independent groups (Currie et al. 2015; Quanz et al. 2015). In Walsh et al. (2014), henceforth referred to as Paper I, we presented the ^{12}CO ($J = 3 - 2$, $\nu = 345.796$ GHz) first moment map and dust continuum emission (at 302 and 346 GHz). These data spatially resolved the CO emission and allowed direct determination of the radial extent of the molecular disk (≈ 390 au, see also Pineda et al. 2014). The continuum data analysed in Paper I showed that the (sub)mm-sized dust grains had been sculpted into two rings. Pinilla et al. (2015) showed that this dust morphology is consistent with dust trapping by two massive companions: one with mass $\approx 20M_J$ at 10 au, and one with mass $\approx 15M_J$ at 70 au. Hence, the ALMA data support the presence of an outer (proto)planet. Emission from ^{12}CO ($J = 3 - 2$, $6 - 5$, and $7 - 6$) from HD 100546 had been detected previously in single-dish observations with APEX (Panić et al. 2010). The APEX data revealed an asymmetry in the red and blue peaks in the double-peaked line profiles most apparent in the $J = 3 - 2$ and $6 - 5$ transitions. Panić et al. (2010) hypothesised that the asymmetry may arise due to shadowing of the outer disk by a warp in the inner disk. Using the same data set as here, Pineda et al. (2014) showed that the position-velocity ($p - v$) diagram across the major axis of the disk is better described by a disk inclination of $\approx 30^\circ$, rather than an inclination of 44° that best reproduces the aspect ratio of the disk as seen in continuum emission (Paper I).

In this work we revisit the HD 100546 ALMA Cycle 0 data and conduct a deeper analysis of the spatially and spectrally resolved ^{12}CO $J = 3 - 2$ emission. The focus of this work is the search for evidence of a warp in the inner regions of the disk, as suggested by the single dish data presented in Panić et al. (2010). In Section 2, we outline the imaging presented in the paper, and in Section 3 we describe the modelling techniques used and present the results. Sections 4 and 5 discuss the implications and state the conclusions, respectively.

2. ALMA IMAGING OF HD 100546

HD 100546 was observed with ALMA on 2012 November 24 with 24 antennas in a compact configuration, with baselines ranging from 21 to 375 m. The self-calibrated and phase-corrected measurement set, produced as described in Paper I, is used in these analyses. In this work, we adopt the revised distance to HD 100546 determined by Gaia (109 ± 4 pc, Gaia Collaboration 2016a,b), and a stellar mass of $2.4M_\odot$, (van den Ancker et al. 1998)

In Paper I, the integrated intensity and first moment maps from the ^{12}CO $J = 3 - 2$ rotational transition at 345.795 GHz ($E_{\text{up}} = 33.19$ K and $A_{\text{ul}} = 2.497 \times 10^{-6} \text{ s}^{-1}$) were presented. The data cube from which those maps were produced was itself produced using the CASA task `clean` with Briggs weighting (robust=0.5) at a spectral resolution of 0.15 km s^{-1} . The resulting channel maps had an rms noise of $19 \text{ mJy beam}^{-1} \text{ channel}^{-1}$ and a synthesised beam of $0''.95 \times 0''.42$ (38°). The ^{12}CO was strongly detected with a signal-to-noise ratio (S/N) of 163 in the channel maps.

Because of the high S/N the imaging is redone here using uniform weighting which results in a smaller beam (and improved spatial resolution) at the expense of sensitivity. The resulting channel maps have an rms noise of $26 \text{ mJy beam}^{-1} \text{ channel}^{-1}$, a S/N of 106, and a synthesised beam of $0''.92 \times 0''.38$ (37°). Figure ?? presents the channel maps. Emission is detected ($\geq 3\sigma$) across 111 channels: the central channel is centred at the source velocity of 5.7 km s^{-1} as constrained previously by these data (see Paper I). The highest velocity emission detected is $\pm 8 \text{ km s}^{-1}$ relative to the source velocity. Given that the disk inclination (as constrained by the outer disk) is 44° and that the stellar mass is $2.4M_\odot$, emission is detected down to a radius of 15 au from the central star. Using the estimate of $\approx 30^\circ$ for the inclination of the inner disk from Pineda et al. (2014), reduces this radius to 8 au.

The channel maps in Figure ?? reveal the classic “butterfly” morphology of spectrally- and spatially-resolved line

emission from an inclined and rotating protoplanetary disk (see, e.g., [Semenov et al. 2008](#)). Compared with the resolved ^{12}CO emission from the disk around the Herbig Ae star HD 163296 which has a similar inclination, there is no evidence of emission from the back side of disk that is a signature of CO freezeout in the disk midplane coupled with emission from a flared surface ([de Gregorio Monsalvo et al. 2013](#); [Rosenfeld et al. 2013](#)). The blue-shifted emission also appears mostly symmetric about the disk major axes (especially that from the south-east of the disk) indicating that it arises from a relatively “flat” surface. This is in contrast with emission from the disk around the Group I Herbig Ae disk HD 97048 ([Walsh et al. 2016](#); [van der Plas et al. 2017](#)). However, the emission is not wholly symmetric about the disk *minor* axis, with the red-shifted emission from the north-west quadrant appearing both fainter, and with a positional offset, relative to blue-shifted emission at the same velocity. In Figure ?? the channel maps from ± 0.45 to $\pm 1.5 \text{ km s}^{-1}$ are shown, now rotated counter-clockwise by 34° ($180^\circ - \text{P.A.}$) to align the disk major axis in the vertical direction, and mirrored across the disk minor axis. Exhibiting the data in this velocity range and in this manner highlights the described asymmetry in brightness across the disk minor axis, the flatness of the emission, and the positional offset of the red-shifted north-west lobe relative to its blue-shifted counterpart. The brightest lobe to the north east, an apparent CO “hot spot”, is consistent in position angle with a proposed (proto)planetary companion seen in direct imaging ([Quanz et al. 2013](#); [Currie et al. 2015](#); [Quanz et al. 2015](#); [Rameau et al. 2017](#)).

Figure ?? presents the moment maps (zeroth, first, second, and eighth). The zeroth moment map (integrated intensity) was produced using a 3σ rms noise clip. The first (intensity-weighted velocity), second (intensity-weighted velocity dispersion), and eighth (peak flux density) moment maps were produced using a more conservative clip of 6σ .

The integrated intensity appears relatively symmetric about the disk minor axis; however, the ^{12}CO integrated emission extends further to the south-west than it does to the north-east. This asymmetry is also evident in the eighth-moment map with the north-east side of the disk appearing brighter than the south-west side with a dark lane in the east-west direction also seen in scattered light images with VLT/SPHERE ([Garufi et al. 2016](#)). Both maps hint at emission from a flared disk which would lead to an asymmetry in integrated emission across the disk major axis (i.e., the axis of inclination). The first and second moment maps also hint at asymmetric emission, in particular, the emission at the source velocity through the inner disk is twisted relative to the disk minor axis determined from the continuum emission. The velocity dispersion in the inner disk is also not wholly symmetric across the disk minor axis. A “by-eye” inspection of the first- and eighth-moment maps, in particular, suggest the possible presence of a warp in the inner disk.

3. MODELLING THE KINEMATICS

Modelling of the kinematics as traced by the ^{12}CO emission is conducted using analytical models which describe the line-of-sight projected velocity, i.e. the first moment map (**isn’t the first moment map technically the intensity-weighted projected velocity?**). The model moment maps are convolved with the synthesised beam of the observations. The residuals (data – model) are in units of a single spectral resolution element ($\delta v = 0.21 \text{ km s}^{-1}$). This discretisation is necessary because features smaller than the native spectral resolution of the data cannot be fit. Note that this analysis will not address the asymmetry in CO brightness across the disk seen in the channel maps and zeroth and eighth moment maps. We leave such an analysis to future work when data for multiple CO transitions and isotopologues allow a robust extraction of the gas surface density and gas temperature.

We explore several metrics of “best fit”: (i) the total number of pixels for which the analytical and smoothed projected velocity reproduces the data within one spectral resolution element, (ii) the sum of the square of the residuals normalised by the total number of unmasked pixels, and (iii) the magnitude of the peak residual. The total number of unmasked pixels in the observed first moment map is 2562. Given the relatively small number of parameters for each model considered, the modelling approach is grid based, i.e. all possible grid combinations are explored.

3.1. A flat emitting surface

The simplest prescription for describing the first moment map of spectrally-resolved line emission from a disk is axisymmetric emission arising from a geometrically flat surface inclined to the line of sight. Assuming that the position angle of the disk is aligned with the y axis, the projected velocity on the sky relative to the observer is described by

$$v(x', y') = \sqrt{\frac{GM_\star}{\rho}} \sin i \sin \theta, \quad (1)$$

where G is the gravitational constant, M_\star is the mass of the central star, $\rho = \sqrt{x^2 + y^2}$ is the radius, i is the inclination, and $\theta = \arctan(y/x)$ (e.g., [Rosenfeld et al. 2013](#)). In this projection and for this particular orientation, $x = x' / \cos i$,

$y = y'$, and $z = 0$. Model first moment maps for a flat disk with the same P.A. as HD 100546 and inclinations of 30° , 45° , and 60° are shown in Figure ?? in the Appendix.

The wide range of disk inclinations ($[20^\circ, 60^\circ]$) and disk position angles ($[120^\circ, 170^\circ]$) explored are motivated by previous analyses of the continuum data which suggested a P.A. of $146^\circ \pm 4^\circ$ and an inclination of $44^\circ \pm 3^\circ$ (see Paper I and Pineda et al. 2014). Using the CO data (I'm a little confused—which CO data is being referred to here? The data used in this paper?), Pineda et al. (2014) suggest that the inner disk may be better described with an inclination of $\approx 30^\circ$; as such, we extend our explored range accordingly to ensure good coverage over the parameter space. First, a coarse grid with a resolution of 5° is run over the full parameter space, followed by a zoomed in region with a resolution of 1° .

The top-left panel of Figure ?? presents a 3D plot showing the total number of pixels which fit the data velocity field within one spectral resolution element, δv , as a function of disk inclination and position angle. The distribution is strongly peaked: the best-fit flat disk model using this metric has an inclination of 36° and a P.A. of 145° with 62.1% of model pixels lying within one spectral resolution element of the data. These data are also listed in Table 1. The P.A. is in excellent agreement with that derived from the continuum observations. The inclination, on the other hand, is lower and closer to the suggested inclination from Pineda et al. (2014).

The left hand plots of Figure ?? show the distribution of residuals summed over the entire disk (top panel) and the residual first moment map (bottom panel). The histogram of residuals shows small dispersion of about 0 with 96.0% of pixels matching the data within $\pm 0.315 \text{ km s}^{-1}$. The residual map shows that a flat disk well reproduces the large-scale velocity field: the largest deviations from this model occur in the innermost disk where the model velocity field over-predicts (by up to $\approx 7\delta v$) the magnitude of the projected line-of-sight velocity along the minor axis of the disk. This leads to negative residuals in the north-east and positive residuals in the south-west. The morphology of the residuals suggests that the inner disk has an additional inclination along the minor axis of the outer disk, i.e., close to orthogonal to that of the outer disk.

3.2. A flared emitting surface

Although a geometrically flat disk well reproduces much of the velocity field, particularly for the outer disk, we test next whether emission from a flared surface can improve upon the flat disk fit. This is important to check because HD 100546 is classified as a Group I (i.e., flared) Herbig Ae star (Meeus et al. 2001), so one might expect the ^{12}CO emission to arise from a layer higher up in the disk atmosphere. Indeed, thermo-chemical modelling of the disk around HD 100546 by Bruderer et al. (2012) suggests that the ^{12}CO line emission arises from a layer $z/\rho \approx 0.2$.

Rosenfeld et al. (2013) modelled the emission from the disk around the Herbig Ae star HD 163296 by assuming that the emission arises from an inclined and flared surface with some opening angle, α , relative to the (x, y) plane (the disk midplane), i.e., a “double-cone” morphology. In this way, the front and back sides of the disk with the same projected line-of-sight velocity are spatially offset (see also de Gregorio Monsalvo et al. 2013). Here, a similar toy model is used; however, to determine the line-of-sight velocity, the radius is defined using spherical coordinates ($r = \sqrt{x^2 + y^2 + z^2}$) rather than cylindrical coordinates ($\rho = \sqrt{x^2 + y^2}$; Rosenfeld et al. 2013). For small opening angles the two methods give similar results: the radii differ by no more than 10% for $\alpha \leq 25^\circ$. A flared disk with this emission morphology has two possible orientations, with either the lower or the upper face of the cone visible to the observer (see e.g., figure 3 in Rosenfeld et al. 2013). Model first moment maps for a flared disk with the same P.A. and inclination as HD 100546 but with different opening angles (20° , 45° , and 60°) are shown in Figure ?? in the Appendix.

The range of surface opening angles ($[0^\circ, 20^\circ]$) is motivated by previous thermo-chemical modelling of CO emission from HD 100546 which suggest an opening angle $\alpha \approx 11^\circ$ (Bruderer et al. 2012). The symmetry in the channel maps (Figure ??) also suggests that the emitting layer lies relatively close to the midplane. As before, a coarse grid with a resolution of 5° is initially run over the full parameter space, followed by a zoomed in grid with a resolution of 1° .

Figures ?? and ?? present the statistics and residuals for the best-fit lower cone and upper cone of a flared disk. Using metric (i), the best-fit upper cone model fits the data marginally better (reproducing 65.0% of the velocity field) than both the flat disk and the best-fit lower cone model (62.1% and 62.6%, respectively, see Table 1). The best-fit inclination, P.A., and opening angle are 36° , 145° , and 9° , respectively (see Table 1). The opening angle of the ^{12}CO -emitting surface agrees well with that suggested by thermo-chemical models of HD 100546 (Bruderer et al. 2012).

The best-fit lower cone model has an inclination of 38° , a P.A. of 142° , and an opening angle of 13° . The inclination of this model lies closest to that derived from the continuum observations ($44^\circ \pm 3^\circ$). Despite resulting in a marginally worse fit to the data than the upper cone model (see Table 1), a “by-eye” examination of the residual map (bottom left panel of Figure ??) shows that this morphology best reproduces the velocity field in all quadrants of the outer

disk excepting the north-west quadrant for which the magnitude of the velocity field is over-estimated. Comparing this residual map to both the channel map (Figure ??) and the eighth moment map (bottom right panel of Figure ??) shows that emission from this quadrant appears less bright and exhibits a positional offset relative to that mirrored across the minor axis of the disk. However, that the upper cone model fits the data best using this metric is in agreement with recent VLT/SHERE images that confirm that the the far side of the flared disk surface lies towards the north (Garufi et al. 2016).

The global best-fit across all three models as determined by metric (ii), i.e., sum of the squares of the residuals normalised by the total number of pixels, is a flat disk with an inclination of 37° and a P.A. of 142° . The best-fit model selected by the smallest peak residual, i.e., metric (iii), is also shared by all three models and is a flat disk with an inclination of 39° (again in good agreement with the other two metrics); however, the disk P.A. which gives the smallest peak residual is 126° . The residual histograms and maps for both of these models are shown in Figure ?. That the inner disk velocity structure is better fit with a shallower P.A. than the outer disk, highlights the presence of a twisted warp: this is investigated in the subsequent section.

Table 1. Best-fit parameters for the flat and flared kinematic models.

Model	Metric of best-fit	Inclination	P.A.	Opening angle	Pixel number	Percentage	Sum of residual squares	Peak residual
Flat	(i)	36°	145°	0°	1590	62.1%	0.737	7.23
	(ii)	37°	142°	0°	1550	60.5%	0.665	6.71
	(iii)	39°	126°	0°	224	8.8%	3.610	3.63
Flared (lower cone)	(i)	38°	142°	13°	1605	62.6%	0.718	6.64
Flared (upper cone)	(i)	36°	145°	9°	1665	65.0%	0.764	7.28

3.3. A warped disk

The residual maps displayed in Figures ?? and ?? reveal two features: (i) a rotating disk within $\approx 1''0$ of the source position with an inclination angle approximately orthogonal to the line of sight, and (ii) a shallower position angle on small scales ($\lesssim 1''0$) than on larger scales. Both results point towards a twisted warp in the inner disk (see, e.g, Juhász & Facchini 2017, and references therein).

Because the residuals are of the order of the size of the synthesised beam, a simple toy prescription for the warp is used. The inner disk is modelled as a planar disk within a fixed radius which possesses its own inclination and P.A., i.e., the inner disk is misaligned relative to the outer disk. This is similar to the approach used by Rosenfeld et al. (2014) to model the kinematics of HD 142527. Figure ?? presents model first moment maps for a warped disk for a range of inclinations and position angles; the transition radius is fixed at 100 au, and the outer disk parameters are given values appropriate for the HD 100546 disk.

The outer disk velocity structure is fixed to that of the best-fit upper cone model. As mentioned in the previous section, recent VLT/SPHERE images of scattered light from HD 100546 suggest that the far side of the (flared) disk lies towards the north-east (Garufi et al. 2016). This results in three additional fitting parameters only: the inner disk inclination ($[40^\circ, 90^\circ]$), the inner disk P.A. ($[40^\circ, 100^\circ]$), and a transition radius marking the boundary between the inner and outer disks ($[40, 120]$ au). Note that, for simplicity, we assume that the inner disk velocity structure is described using the flat disk prescription (i.e., Equation 1). A coarse grid with a resolution of 10° and 10 au is first run to identify the parameter space containing the global best-fit, followed by a finer grid over this zoomed-in region (with a resolution of 2° and 2 au).

Figures ?? and ?? present the statistics and residuals for the best-fit warped disk, respectively. Metric (i) favours a model with an inner disk that is almost “edge-on” ($i = 80^\circ$) to the line of sight, almost orthogonal to the outer disk major axis (P.A. = 60°), and with a transition radius of 90 au (see Figure ?? and Table 2). These values are consistent with the morphology of the residuals of both the flat and flared models (see Figures ?? and ??). The magnitude of the peak residual of this model is significantly smaller than the previous models selected using metric (i), $4\delta v$ versus $7\delta v$. Metrics (ii) and (iii) select the same model (see Table 2) with parameters similar to those using metric (i); an inclination of 84° , a P.A. of 64° , and a transition radius of 100 au. Comparing the residual histograms and maps

for these two models (shown in Figure ??), highlights how a small change in inclination and/or position angle can significantly reduce the magnitude of the peak residual. This latter model results in a peak residual of only $2.4\delta v$ and has the smallest dispersion of residuals: 98% of pixels match the data within $\pm 0.315 \text{ km s}^{-1}$ and 100% of pixels match within $\pm 0.525 \text{ km s}^{-1}$.

Table 2. Best-fit parameters for the warped kinematic models.

Model	Metric of best-fit	Inclination	P.A.	Transition radius (au)	Pixel number	Percentage	Sum of residual squares	Peak residual
Warped	(i)	80°	60°	90	1722	67.2%	0.387	4.064
	(ii) & (iii)	84°	64°	100	1710	66.4%	0.350	2.441

4. DISCUSSION

The analysis of the kinematics of the HD 100546 protoplanetary disk presented here has indicated the possible presence of a severely warped (misaligned) inner disk within 100 au of the central star. The inner gas disk exhibits a position angle that is almost orthogonal to that of the outer disk, and is inclined almost edge-on to the line-of-sight. Figure ?? shows an idealised model of the HD 100546 protoplanetary disk. The morphology of the intermediate region between the inner and outer disks is not yet known, and so is intentionally left blank in the cartoon. Higher spatial resolution data are needed to reveal the intermediate region morphology and determine if the proposed warp lies within a smaller radial region than suggested by these data. Scattered light images taken with VLT/SPHERE and MagAO/GPI with a spatial resolution of $0''.02$ and $0''.01$, respectively, reveal no evidence of a severely misaligned dust disk beyond $\approx 10 \text{ au}$ (Garufi et al. 2016; Follette et al. 2017). However, both datasets do suggest the presence of spirals in the inner disk within 50 au, and resolve the inner edge of the outer dust disk traced in small grains (11-15 au, depending on wavelength). Further, recent VLTI/PIONIER interferometric observations of HD 100546 reported in the survey by Lazareff et al. (2017), and which have a spatial resolution of order $\sim 1 \text{ au}$, suggest that the very innermost dust disk has a similar position angle (152°) and inclination (46°) as the outer disk traced in sub-mm emission (146° and 44° , respectively; Walsh et al. 2014). Hence, if the non-Keplerian motion of the CO gas $\lesssim 100 \text{ au}$ is indeed caused by an extreme warp, in light of these new observations it appears that the small dust grains may have a different morphology than the gas in the inner disk. This is discussed further below.

The presence of a disk warp in HD 100546 was originally proposed by Panić et al. (2010) and based on asymmetries seen in the red and blue peaks of single-dish spectra observed with APEX. The ALMA data presented here show no such asymmetries in the peaks of the line profile (see Figure ??); however, the mirrored line profile does highlight that the red and blue lobes of emission have different shapes when integrated over the disk. This is consistent with the morphology seen in the channel maps and described in Section 2.

A warp in the inner disk could shadow the outer disk and lead to an asymmetric temperature structure, because one side of the disk is more illuminated by the central star than the opposite side (see figure 3 in Panić et al. 2010). Because the ^{12}CO emission is optically thick, it traces the gas temperature, and thus any perturbations, in the emitting layer, usually the disk atmosphere. The APEX ^{12}CO $J = 3 - 2$ line profile from Panić et al. (2010) shows a stronger blue peak. The dates of the observations taken with APEX and ALMA are November 2008 (Panić et al. 2010) and November 2012 (Walsh et al. 2014), respectively. There also exist multiple APEX observations of the ^{12}CO $J = 6 - 5$ transition. The line profile presented in Panić et al. (2010) shows an even higher-contrast asymmetry in the blue and red peaks than in the $J = 3 - 2$ line. More recent observations taken between April and July 2014 and presented in Kama et al. (2016) show a similar asymmetry; however, the ratio between the peaks is within the flux density uncertainties of the observations (M. Kama, priv. commun.). Thus, it is currently difficult to draw concrete conclusions on any periodicity in the shapes of the line profiles. Conservatively assuming that the warp has rotated $\approx 90^\circ$ between acquirement of the original APEX data (with the stronger blue peak) and the ALMA data (with no asymmetry in the line profile) suggests a warp precession period of the order of 16 yr.

The presence of a warp indicates the presence of a perturbing companion. A massive ($\approx 20M_J$) planetary companion has been proposed to reside at $\approx 10 \text{ au}$ in HD 100546. This is thought to be responsible for clearing the inner cavity in both the dust and gas (Mulders et al. 2013; Panić et al. 2014; Walsh et al. 2014; Pinilla et al. 2015; Wright et al. 2015;

Garufi et al. 2016), as well as inducing dynamical perturbations in [OI], OH, and CO line emission from the inner region (Acke & van den Ancker 2006; van der Plas et al. 2009; Brittain et al. 2014). In addition, a second companion, which may still be in the act of formation, has been proposed to reside at ≈ 50 au with a mass estimate of $\approx 15M_J$ (Quanz et al. 2013; Walsh et al. 2014; Quanz et al. 2015; Currie et al. 2015; Pinilla et al. 2015). However, as mentioned in the introduction, the identification of this point source as a (proto)planet has been disputed in a recent analysis of MagAO/GPI observations showing close-to-zero proper motion of the point source on a 4.6 yr timescale at the 2σ level (Rameau et al. 2017). Given that the presence of the outer ring of (sub)mm dust emission is difficult to explain in the absence of an outer companion (Walsh et al. 2014; Pinilla et al. 2015), we proceed in remainder of the discussion with the assumption that this disk “feature” indeed arises due to a (proto)planet.

The precession period of a warp induced by a perturbing companion can be estimated using equation (2) in Debes et al. (2017), used for the case of TW Hya, and which was originally derived by Lai (2014),

$$P \propto (\mu_c \cos i_c)^{-1} (M_*)^{-1/2} (r_{\text{disk}})^{-3/2} (a_c)^3 \quad (2)$$

where μ_c is the mass ratio of the companion to the central star (here $2.4M_\odot$), i_c is the angle between the star-planet orbital plane and disk plane (here 80°), r_{disk} is the radius of the outer edge of the inner disk, and a_c is the orbital radius of the planet. Assuming that the protoplanet at 50 au is the perturbing body, and setting $\mu_c = 0.006$, $r_{\text{disk}} = 40$ au, and $a_c = 50$ au results in a precession period of 1940 yr. Hence, it is unlikely that the outer companion has perturbed the inner disk creating a warp that explains the change in shape in line profile seen in the single-dish data and these ALMA data on a 4-year baseline. Assuming instead that the inner companion is triggering the warp, and that the inner disk cavity is not completely devoid of gas ($\mu_c = 0.008$, $r_{\text{disk}} = 10$ au, $i_c = 80^\circ$, and $a_c = 10$ au), leads to a precession period of 111 yr. Keeping all other parameters constant, a companion mass of $\approx 140M_J$ ($\approx 0.1M_\odot$) is required for a warp precession period of ≈ 16 yr. There is no evidence that HD 100546 is in a binary system with an M-dwarf star. Alternatively, a closer-in companion of only $\approx 4.5M_J$ at 1 au and assuming an inner disk radius of 1 au, leads to a similar precession period. However, these ALMA data suggest that the non-Keplerian gas motion lies significantly beyond 1 au since the highest velocity gas detected only probes down to a radius of 8-15 au depending upon the assumed disk inclination. Longer baseline (in time) observations are required to determine if any clear periodicity in the line profile shape exists, and to then relate this more concretely to the precession period of an inner disk warp.

Similar analyses have revealed warps in disks around other stars. Rosenfeld et al. (2012) showed that the kinematic structure of TW Hya could be reproduced using a parametric model of a disk warp with a moderate inclination ($\approx 8^\circ$) at a radius of 5 au. Facchini et al. (2014) demonstrated the observed warp amplitude could be induced by a misaligned close-in companion as massive as $\approx 14M_J$ orbiting within 4 au of the central star. New scattered light data from HST/STIS, coupled with archival data over a 17-year baseline, reveal an orbiting azimuthal brightness asymmetry, with a period of ≈ 16 yr (Debes et al. 2017). The authors argue that this is consistent with partial shadowing of the outer disk by a misaligned inner disk interior to 1 au and that is precessing due to the presence of a roughly Jupiter-mass planetary companion.

Given that the ALMA data presented here suggest non-Keplerian motion of the gas on spatial scales $\gtrsim 10$ au, it is worth considering alternative explanations to a warp. HD 142527 is another protoplanetary disk in which non-Keplerian motion traced by HCO^+ ($J = 4 - 3$) and ^{12}CO ($J = 6 - 5$) emission has been observed (Casassus et al. 2013, 2015). HD 142527 is a transition disk which possesses the largest known dust cavity in sub(mm) emission (≈ 140 au; Casassus et al. 2013; Fukagawa et al. 2013). The observed gas motions were postulated to arise from fast (near free-fall) radial flows across the cavity and accretion onto the central star via a severely misaligned inner disk (Casassus et al. 2013, 2015; Marino et al. 2015). Marino et al. (2015) propose that that inner and outer disk are misaligned by 70° . Compared with the misalignment parameters derived for HD 142527, those derived here for HD 100546 appear not so extreme. Also, “hot-off-the-press” ALMA observations of HCO^+ ($J = 3 - 2$) line emission from AA Tau at an angular resolution of $0''.2$ (≈ 30 au), also show evidence of non-Keplerian motions due to either a warp or fast radial inflow in the inner regions (Loomis et al. 2017). Similar to here, this was revealed by a clear twist in the first moment map of the molecular line emission. However, Rosenfeld et al. (2014) conducted a theoretical study of fast radial flows in transition disks and showed that the derived velocity profile can also mimic that characteristic of an inner disk warp. To determine which physical mechanism is responsible for the observed gas motions in HD 100546 will require data at sufficient spatial and spectral resolution to resolve molecular emission within the gas cavity i.e., within a diameter of ≈ 20 au.

5. CONCLUSION

The data and analyses shown here clearly demonstrate that the kinematic structure of the disk around HD 100546 cannot be described by a purely Keplerian velocity profile with a universal inclination and position angle. Given the current evidence for the presence of (at least) one massive planetary companion orbiting within 10 au of the central star, the presence of a disk warp, albeit with extreme parameters, has been our favoured explanation. Higher spatial resolution data are vital for corroborating this warp hypothesis. However, recent near-IR interferometric observations suggest that the very inner dust disk is not misaligned relative to the outer dust disk traced in (sub)mm emission. This suggests that the gas and dust disks may have different morphologies on radial scales of ~ 1 to 10's of au. To determine whether the alternative explanation of fast radial flows could be responsible for the observed kinematic structure of HD 100546, higher spatial *and* spectral resolution data are required to resolve gas emission across the inner dust ($\lesssim 25$ au) and gas ($\lesssim 10$ au) cavities and explore any spatial association with the proposed planetary candidate(s).

REFERENCES

- Acke, B. & van de Ancker, M. E. 2006, *A&A*, 449, 267
- van den Ancker, M. E., de Winter, D., Tijn A Djie, H. R. E., *A&A*, 330, 145
- Andrews, S. M. 2015, *PASP*, 127, 961
- Brittain, S. D., Najita, J. R., & Carr, J. S. 2009, *ApJ*, 702, 85
- Brittain, S. D., Carr, J. S., Najita, J. R., Quanz, S. P., & Meyer, M. R. *ApJ*, 791, 136
- Bruderer, S., van Dishoeck, E. F., Doty, S. D., & Herczeg, G. 2012, *A&A*, 541, 91
- Bruderer, S. 2013, *A&A*, 559, A46
- Casassus, S., van der Plas, G., Pérez, S. M., et al. 2013, *Nature*, 493, 191
- Casassus, S., Marino, S., Pérex, S., et al. 2015, *ApJ*, 811, 92
- Christiaens, V., Casassus, S., Perez, S., van der Plas, G., & Ménard, F. 2014, *ApJL*, 785, L12
- Currie, T., Cloutier, R., Brittain, S., et al. 2015, *ApJL*, 814, L27
- Debes, J. H., Poteet, C. A., Jang-Condell, H., et al. 2017, *ApJ*, 835, 205
- Dutrey, A., Semenov, D., Chapillon, E., et al. 2014, in *Protostars and Planets VI*, ed. H. Beuther et al. (Tucson, AZ: Univ. Arizona Press), 317
- Eistrup, C., Walsh, C., & van Dishoeck, E. F. 2016, *A&A*, 595, A83
- Españolat, C., Muzurrolle, J., Najita, J., et al. 2014, in *Protostars and Planets VI*, ed. H. Beuther et al. (Tucson, AZ: Univ. Arizona Press), 497
- Facchini, S., Ricci, L., & Lodato, G. 2014, *MNRAS*, 442, 3700
- Follette, K. B., Rameau, J., Dong, R., et al. 2017, *AJ*, in press (arXiv:1704.06260)
- Fukagawa, M., Tsukagoshi, T., Momose, M., et al. 2013, *PASJ*, 65, L14
- Furuya, K. & Aikawa, Y. 2014, *ApJ*, 790, 97
- Gaia Collaboration et al. 2016a, *A&A*, 595, A1
- Gaia Collaboration et al. 2016b, *A&A*, 595, A2
- Garufi, A., Quanz, S. P., Schmid, H. M., et al. 2016, *A&A*, 588, A8
- Grady, C., Fukagawa, M., Maruta, Y., et al. 2015, *Ap&SS*, 355, 253
- de Gregorio Monsalvo, I., Ménard, F., Dent, B., et al. 2013, *A&A*, 557, 133
- Helling, C., Woitke, P., Kamp, I., et al. 2014, *Life*, 4, 142
- Juhász, A. & Facchini, S. 2017, *MNRAS*, in press
- Kama, M., Bruderer, S., van Dishoeck, E. F., et al. 2016, *A&A*, 592, A83
- Lai, D. 2014, *MNRAS*, 440, 3532
- Lazareff, B., Berger, J.-P., Kluska, J., et al. 2017, *A&A*, 599, A85
- Loomis, R. A., Öberg, K. I., Andrews, S. M., & MacGregor, M. A. 2017, *ApJ*, in press (arXiv : 1704.02006)
- Marino, S., Pérez, S., & Casassus, S. 2015, *ApJL*, 798, L44
- Meeus, G., Waters, L. B. F. M., Bouwman, J., et al. 2001, *A&A*, 365, 476
- Miotello, A., Bruderer, S., & van Dishoeck, E. D. 2014, *A&A*, 572, A96
- Miotello, A., van Dishoeck, E. D., Kama, M., & Bruderer, S. 2016, *A&A*, 594, A85
- Mulders, G. D., Paardekooper, S.-J., Panić, O., et al. 2013, 557, 68
- Nomura, H., Tsukagoshi, T., Kawabe, R., et al. 2016, *ApJL*, 819, L7
- Panić, O., van Dishoeck, E. F., Hogerheijde, M., et al. 2010, *A&A*, 519, 110
- Panić, O., Ratzka, Th., Mulders, G. D., et al. 2014, 562, A101
- Perez, S., Dunhill, A., Casassus, S., et al. 2015, *ApJ*, 811, L5
- Pineda, J., Quanz, S. P., Meru, F., et al. 2014, *ApJL*, 788, L34
- Pinilla, P., Birnstiel, T., & Walsh, C. 2015, *A&A*, 580, A105
- van der Plas, G., van de Ancker, M. E., Acke, B., et al. 2009, *A&A*, 500, 1137
- van der Plas, G., Wright, C. M., Ménard, F., et al. 2017, *A&A*, 597, A32
- Quanz, S., Amara, A., Meyer, M. R., Kenworthy, M. A., Kasper, M., & Girard, J. A. 2013, *ApJL*, 766, L1
- Quanz, S. P., Amara, A., Meyer, M., et al. 2015, *ApJ*, 807, 64
- Rameau, J., Follette, K. B., Pueyo, L., et al. 2017, *AJ*, 153, 244
- Rosenfeld, K. A., Qi, C., Andrews, S. M., et al. 2012, *ApJ*, 757, 129
- Rosenfeld, K. A., Andrews, S. M., Hughes, A. M., Wilner, D. J., & Qi, C. 2012, *ApJ*, 757, 129
- Rosenfeld, K. A., Chiang, E., & Andrews, S. M. 2014, *ApJ*, 782, 62
- Schwarz, K. R., Bergin, E. A., Cleeves, L. I., et al. *ApJ*, 823, 91
- Semenov, D., Pavlyuchenkov, Ya., Henning, Th., Wolf, S., & Launhardt, R. 2008, *ApJL*, 673, L195
- Sicilia-Aguilar, A., Banzatti, A., Carmona, A., et al. 2016, *PASA*, 33, 59
- Simon, M., Dutrey, A., & Guilloteau, S. 2000, *ApJ*, 545, 1034
- Tang, Y.-W., Guilloteau, S., Piétu, V., et al. 2012, *A&A*, 547, A84
- Visser, R., van Dishoeck, E. F., & Black, J. H. 2009, *A&A*, 503, 323
- Williams, J. P. & Cieza, L. A. 2011, *ARA&A*, 49, 67
- Walsh, C., Juhász, A., Pinilla, P., et al. 2014, *ApJL*, 791, L6
- Walsh, C., Juhász, A., Meeus, G., et al. 2016, *ApJ*, 831, 200
- Wright, C. M., Maddison, S. T., Wilner, D. J., et al. 2015, *MNRAS*, 453, 414
- Yu, M., Willacy, K., Dodson-Robinson, S. E., Turner, N. J., & Evans II, N. J. 2016, *ApJ*, 822, 53
- Yu, M., Evans II, N. J., Dodson-Robinson, S. E., Willacy, K., & Turner, N. J. 2017, *ApJ*, in press (arXiv : 1704.05508)

Zhang, K., et al. 2017, Nature Astronomy,

APPENDIX

A. MODEL FIRST MOMENT MAPS

Figure ?? shows the first moment maps for a geometrically flat protoplanetary disk for a range of disk inclinations. Figure ?? shows model first moment maps for a flared disk with a fixed inclination for a range opening angles, α , of the emitting surface. Figure ?? shows model first moment maps for a warped inner disk with a fixed transition radius of 100 au, and a range of warp inclinations and position angles. The outer disk parameters are those from the best-fit upper cone model (i.e., an inclination of 36° , a P.A. of 145° , and an opening angle of 9°).