

# Bayesian hierarchical models for NHPP using **rstan**

*Miao Cai miao.cai@slu.edu*

*2019-06-20*

## Contents

### 1 Model setting

1

## 1 Model setting

Let  $T_{d,s,i}$  denote the time to the  $d$ -th driver's  $s$ -th shift's  $i$ -th critical event. The total number critical events of  $d$ -th driver's  $s$ -th shift is  $n_{d,s}$ . The ranges of these notations are:

- $i = 1, 2, \dots, n_{d,S_d}$ ,
- $s = 1, 2, \dots, S_d$ ,
- $d = 1, 2, \dots, D$ .

We assume the times of critical events within the  $d$ -th driver's  $s$ -th shift were generated from a non-homogeneous Poisson process (NHPP) with a power law process (PLP), with a fix rate parameter  $\beta$  and varying scale parameters  $\theta_{d,s}$  across drivers. The data generating process is then:

$$T_{d,s,1}, T_{d,s,2}, \dots, T_{d,s,n_{d,s}} \sim \text{PLP}(\beta, \theta_{d,s})$$

$$\beta \sim \text{Gamma}(1, 1)$$

$$\log \theta_{d,s} = \gamma_{0d} + \gamma_1 x_{d,s,1} + \gamma_2 x_{d,s,2} + \dots + \gamma_k x_{d,s,k}$$

$$\gamma_{0i}, \gamma_{0i}, \dots, \gamma_{0D} \sim \text{i.i.d. } N(\mu_0, \sigma_0^2)$$

$$\gamma_1, \gamma_2, \dots, \gamma_k \sim \text{i.i.d. } N(0, 10)$$

$$\mu_0 \sim N(0, 10)$$

$$\sigma_0 \sim \text{Gamma}(1, 1)$$