

Does A Medical Consortium Influence Health Outcomes of Hospitalized Cancer Patients?

An Integrated Care Model in Shanxi, China


Miao Cai, MS, Echu Liu, PhD, Hongbing Tao, PhD,
Zhengmin Qian, PhD, John Fu, PhD, Xiaojun Lin, MS,
Manli Wang, PhD, Chang Xu, PhD



SAINT LOUIS UNIVERSITY
—
COLLEGE FOR PUBLIC HEALTH
AND SOCIAL JUSTICE

1. Background

- **A Medical Consortium** is a vertical integrated care that involves one widely recognized tertiary hospital and several secondary hospitals or community health centers.
- **Hypothesis:** The medical consortium policy has a significant positive effect on outcomes of cancer patients admitted to secondary hospitals in Shanxi, China.



2. Methods

- **Data:** Electronic medical records of lung cancer (n = 8193), stomach cancer (n = 5693) and esophagus cancer (n = 2802) patients hospitalized in secondary hospitals were used.
- **Sample matching:** Propensity scores were used to match each patient enrolled in medical consortium hospitals with a counterpart admitted in non-medical consortium hospitals.
- **Statistical models:** Cox proportional hazard models were used to estimate the hazard ratio of patients enrolled different categories of hospitals. Controlling variables include gender, age, comorbidities, urgency of disease, and surgery.
- **Test of Assumptions:** with the proportional hazards assumption evaluated by the Empirical Score Process with cumulative sums of martingale-based residuals



3. Plots

Figure 1. Product-Limit Survival Estimates of Matched Full Sample Patients







Figure 2. Product-Limit Survival Estimates of 3 Matched Cancer Patients
Lung Cancer Patients



Stomach Cancer Patients



Esophageal Cancer



Note: strata = 0 denotes patients enrolled in non-Medical Consortium Hospitals,
strata = 1 denotes patients enrolled in Medical Consortium Hospitals



4. Results, conclusion, and implications

- **Results:** significantly lower hazard ratios were consistently associated with lung (hazard ratio (HR) = 0.405, $p < 0.001$), stomach (HR = 0.406, $p < 0.001$), and esophagus (HR= 0.439, $p < 0.001$) cancer patients in medical consortium hospitals, compared with those in non-medical consortium hospitals.
- **Conclusion:** The medical consortium provides an effective strategy to improve the outcomes of cancer patients in Shanxi, China.
- **Implications for policies:** Policy makers can learn from the positive outcomes of the medical consortium and improve the outcomes of cancer patients through the vertical integration of medical providers.

