

UNICAMP

UNIVERSIDADE ESTADUAL DE CAMPINAS (UNICAMP)

INSTITUTO DE MATEMÁTICA, ESTATÍSTICA E COMPUTAÇÃO CIENTÍFICA (IMECC)

PROGRAMA DE PÓS-GRADUAÇÃO EM ESTATÍSTICA

CAIO GOMES ALVES

Title of the dissertation/thesis

Título da dissertação/tese (em português)

Campinas

Ano

Caio Gomes Alves

Title of the dissertation/thesis

Título da dissertação/tese (em português)

Dissertação apresentada ao Instituto de Matemática, Estatística e Computação Científica da Universidade Estadual de Campinas (UNICAMP) como parte dos requisitos exigidos para a obtenção do título de Mestre em Estatística.

Dissertation presented to the Institute of Mathematics, Statistics and Scientific Computing of the University of Campinas (UNICAMP) as part of the requirements for the degree of Master in Statistics.

Orientador: Prof. Dr. Guilherme Vieira Nunes Ludwig

Coorientador: Prof. Dr. Maurício Enrique Zevallos Herencia

Este trabalho corresponde à versão final da Dissertação defendida.

A ficha catalográfica deverá ser solicitada online via
http://www.sbu.unicamp.br/sbu/elaboracao-de-ficha-catalografica/

A folha de aprovação será fornecida pela Secretaria de Pós-Graduação

Este trabalho é dedicado às crianças adultas que,
quando pequenas, sonharam em se tornar cientistas.

Acknowledgements

Inserir aqui os agradecimentos. ATENÇÃO ALUNO BOLSISTA: você precisa mencionar agradecimento ao órgão de fomento de sua bolsa, incluindo explicitamente o código do processo. Caso seja bolsista CAPES, especial atenção ao Art. 3º da Portaria CAPES 206/2018, que prevê texto padrão de agradecimento.

"If you can solve your problem, then what is the need of worrying?
If you cannot solve it, then what is the use of worrying"
Shantideva

Resumo

Segundo a ABNT (2003, 3.1-3.2), o resumo deve ressaltar o objetivo, o método, os resultados e as conclusões do documento. A ordem e a extensão destes itens dependem do tipo de resumo (informativo ou indicativo) e do tratamento que cada item recebe no documento original. O resumo deve ser precedido da referência do documento, com exceção do resumo inserido no próprio documento. Neste trabalho, devem ser utilizadas até 500 palavras. (. . .) As palavras-chave devem figurar logo abaixo do resumo, antecedidas da expressão Palavras-chave:, separadas entre si por ponto e finalizadas também por ponto.

Palavras-chave: Latex. Abntex. Editoração de texto.

Abstract

This is the english abstract.

Keywords: Latex. Abntex. Text editoration.

List of Figures

Figure 1 – Branching of an self-exciting point process. 21

List of Tables

Table 1 – Example of a earthquake catalog. 19

List of abbreviations and acronyms

UNICAMP	Universidade Estadual de Campinas
IMECC	Instituto de Matemática, Estatística e Computação Científica
LabCSD	Laboratório de Controle e Sistemas Dinâmicos
EPIFISMA	Laboratório de Epidemiologia e Fisiologia Matemática
LCP	Laboratório de Computação Paralela
LGC	Laboratório de Geofísica Computacional
LMDC	Laboratório de Matemática Discreta e Códigos
MiLAB	Laboratório de Tratamento Matemático de Imagens e Inteligência Computacional
LPOO	Laboratório de Pesquisa Operacional e Otimização
LEM	Laboratório de Ensino de Matemática
PAPMEM	Programa de Aperfeiçoamento para Professores de Matemática do Ensino Médio
OMU	Olimpíada de Matemática da Unicamp
CCPG	Comissão Central de Pós-Graduação
ABNT	Associação Brasileira de Normas Técnicas
abnTeX	ABsurdas Normas para TeX

List of symbols

$\mathbb{M}_{m\times n}(\mathbb{R})$	Conjunto das matrizes de ordem $m \times n$ com entradas reais
$\mathbb{R}^n_+$	Conjunto dos vetores x pertencentes a $\mathbb{R}^n$ que satisfazem $x \geqslant 0$
$\mathbb{R}^n_{++}$	Conjunto dos vetores x pertencentes a $\mathbb{R}^n$ que satisfazem $x > 0$
$\|\cdot\|$	Norma-p vetorial
$\|\|\|\cdot\|\|\|_p$	Norma-p matricial
$\mathbf{I}_n$	Matriz identidade de ordem $n \times n$
∇f	Gradiente da função f
∞	Infinito

List of Algorithms

List of Source Codes

Contents

Introdução 17

I Theoretical foundations 18

1 Point Processes 19

1.1 Marked Spatio-Temporal Point Processes 19

1.2 Hawkes Processes and the ETAS Model 20

1.3 Likelihood and Stochastic Declustering 23

II Materials and Methods 26

2 Methods 27

2.1 Parameter Estimation 27

2.2 Goodness of fit tests 27

3 Considerações Finais 28

BIBLIOGRAPHY 29

Appendix 31

APPENDIX A Cumulative distributions of $g(t)$ and $f(x, y)$ 32

APPENDIX B Quisque libero justo 34

APPENDIX C Nullam elementum urna vel imperdiet sodales elit ipsum pharetra ligula ac pretium ante justo a nulla curabitur tristique arcu eu metus 35

Annex 36

ANNEX A Morbi ultrices rutrum lorem. 37

ANNEX B Cras non urna sed feugiat cum sociis natoque penatibus et magnis dis parturient montes nascetur ridiculus mus 38

ANNEX C Fusce facilisis lacinia dui 39

Introdução

Este documento e seu código-fonte são exemplos de referência de uso da classe abntex2 e do pacote abntex2cite. O documento exemplifica a elaboração de trabalho acadêmico (teses e dissertações) produzido inicialmente conforme a **Informação CCPG/001/2015** (que trata das *Normas para impressão de teses/dissertações* da UNICAMP). Posteriormente, foi adequada considerando a Instrução Normativa CCPG/06/2025 (CCPG–UNICAMP, 2025). Encorajamos o leitor a consultar estas instruções antes de iniciar as alterações neste documento e seu código-fonte.

A elaboração deste modelo teve como base uma customização do "Modelo Canônico de Trabalho Acadêmico com abnTEX2" (ARAUJO, 2015b) para que as normas presentes na Informação CCPG/001/2015 fossem respeitadas. O modelo original produzido pela equipe abnTEX2 cumpre as seguintes normas ABNT:

1. **ABNT NBR 14724:2011**: Informação e documentação - Trabalhos acadêmicos - Apresentação;
2. **ABNT NBR 10520:2002**: Informação e documentação - Citações;
3. **ABNT NBR 6034:2004**: Informação e documentação - Índice - Apresentação;
4. **ABNT NBR 6028:2003**: Informação e documentação - Resumo - Apresentação;
5. **ABNT NBR 6027:2012**: Informação e documentação - Sumário - Apresentação;
6. **ABNT NBR 6024:2012**: Informação e documentação - Numeração progressiva das seções de um documento - Apresentação
7. **ABNT NBR 6023:2002**: Informação e documentação - Referência - Elaboração.

Este documento deve ser utilizado como complemento dos manuais do abnTEX2 (ARAUJO, 2015a; ARAUJO, 2015c; ARAUJO, 2015d) e da classe memoir (WILSON; MADSEN, 2010).

A leitura do teor desde documento (tanto o PDF quando os arquivos que compõem seu código-fonte), bem como do arquivo LEIAME.txt é altamente recomendada para melhor entendimento da dinâmica de funcionamento da classe abntex2 e do pacote abntex2cite. Seus principais comandos e usos estão exemplificados no decorrer do texto, bem como outras informações relevantes para a escrita de seu trabalho acadêmico.

Part I

Theoretical foundations

1 Point Processes

1.1 Marked Spatio-Temporal Point Processes

A spatio-temporal point process $\mathcal{X}$ is a sequence of random variables on $\{S \subset \mathbb{R}^2 \times [0,T]\}$ in which the number of points in $S \times T$ has a counting measure $N(S \times T)$ who is almost surely finite, when $S \times T$ is bounded. Usually $\mathcal{X}$ is observed in a spatio-temporal window $W = S \times T$, and a realization of $\mathcal{X}$ in W is a set $\mathbf{x} = \{(s_i, t_i), i = 1, 2, \ldots, n\}$, where $s_i = (x_i, y_i)$ and t_i represent the spatial coordinates and times of occurrence of the i-th event, respectively.

A marked spatio-temporal point process incorporates a uniquely associated component m_i with each event of $\mathcal{X}$. In earthquake's catalogs, the marks m_i usually denote the magnitude of each event, while (s_i, t_i) represents the longitude/latitude of the earthquake's epicenter and the ordered times of occurrence.

Table 1 – Example of a earthquake catalog.

Event	**Time**	**Longitude**	**Latitude**	**Magnitude**
1	t_1	x_1	y_1	m_1
2	t_2	x_2	y_2	m_2
$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$
n	t_n	x_n	y_n	m_n

Source: The author.

Consider the event history $\mathcal{H}_t = \{(x_i, y_i, t_i, m_i) \in \mathcal{X} : t_i < t\}$. A spatio-temporal point process can be fully characterized by it's conditional intensity function $\lambda(s, t|\mathcal{H}_t)$ (DALEY; VERE-JONES, 2003)

$$\lambda(s, t|\mathcal{H}_t) = \lim_{\delta s, \delta t \to 0} \frac{\mathbb{E}\left[N(\delta s \times \delta t)|\mathcal{H}_t\right]}{\delta s \, \delta t}, \tag{1.1}$$

that represents the expected number of events observed in an infinitesimal region of volume $(\delta s \times \delta t)$ centered around (s, t), conditional in the observed history of the process up to time t. With it, we can define the process's associated compensator $\Lambda(t)$

$$\Lambda(t|\mathcal{H}_u) = \int_0^t \int_S \lambda(s, u|\mathcal{H}_u) ds \, du, \tag{1.2}$$

in which $N(s, t|\mathcal{H}_t) - \Lambda(t|\mathcal{H}_t)$ is an local $\mathcal{H}_t$-martingale. The simplest and most common kind of spatio-temporal point process is the Poisson Process, defined as such:

Definition 1 (Poisson Process)**.** *A spatio-temporal counting process* $\{N(s,t) : s \in \mathbb{R}^2, t \in \mathbb{R}_+\}$ *is a Inhomogeneous Poisson Process with intensity function* $\lambda(s,t)$ *if:*

- *For any bounded window* $W = (S \times T)$, $N(W) \sim \text{Pois}\left(\int_T \int_S \lambda(s,t) ds\, dt\right)$;
- *For any* n *disjoint windows* $W_1, W_2, \ldots, W_n$, *the random variables* $N(W_1), N(W_2), \ldots, N(W_n)$ *are mutually independent.*

The Poisson Process have two properties that makes it very versatile for modeling point processes in general: **Superposition** and **Thinning**:

Theorem 1 (Superposition of Poisson Processes)**.** *Given* k *independent Poisson Processes* $\mathcal{X}_1, \mathcal{X}_2, \ldots, \mathcal{X}_k$ *with intensity functions* $\lambda_1, \lambda_2, \ldots, \lambda_k$, *the new process* $\mathcal{X} = \sum_{i=1}^{k} \mathcal{X}_i$ *is a Poisson Process, with intensity function* $\lambda = \sum_{i=1}^{k} \lambda_i$.

Theorem 2 (Thinning of Poisson Processes)**.** *Let* $\mathcal{X}$ *be a Poisson Process with intensity function* λ. *For each occurrence of* $\mathcal{X}$, *assign it to process* $\mathcal{X}_1$ *with probability* p_1, *to process* $\mathcal{X}_2$ *with probability* p_2, *and so on up to process* $\mathcal{X}_k$ *with probability* p_k, *such that* $\sum_{i=1}^{k} p_i = 1$. *The new processes* $\{\mathcal{X}_1, \mathcal{X}_2, \ldots, \mathcal{X}_k\}$ *are Poisson Processes with intensity functions* $\{p_1\lambda, p_2\lambda, \ldots, p_k\lambda\}$, *and are mutually independent.*

With these two results, many point processes can have their intensity function modelled using Poisson Processes. However, one condition necessary for that is that $\lambda(s,t)$ must remain constant along time (i.e. $\lambda(s,t) = \lambda(s)$, $\forall t \in [0,T]$), and in many cases that is not necessarily true.

1.2 Hawkes Processes and the ETAS Model

In many pontual phenomena, the ocurrence of events increases the probability of further events occuring, such as the spread of diseases, urban violence (Mohler et al. (2011), Ludwig et al. (2025)) and earthquakes. For these kinds of processes, the intensity function must change dynamically, in order to accommodate the increase in probability triggered by a previous event.

A self-exciting point process (occuring in time) was proposed by Hawkes (1971a), having intensity function given by


$$\lambda(t|\mathcal{H}_t) = \mu + \sum_{i:t_i<t} g(t - t_i), \tag{1.3}$$

where $\mu > 0$ is the baseline intensity of the process, and $\sum_{i:t_i<t} g(t-t_i)$ represents the contribution of past events to the increase in the conditional intensity up to time t, with

$g(v) \geqslant 0$ for nonnegative v and 0 otherwise. To guarantee stationarity for this process (Hawkes (1971b)), is necessary that $\int_0^\infty g(v)dv < 1$, otherwise the process can have infinite events in finite time.

The **ETAS** (*Epidemic-Type Aftershock Sequence*) Model was proposed by Ogata (1988), and is the most used to describe earthquake occurrences. The initial model was only concerned with the event times and was a particular case of a marked point process, for it's inclusion of the earthquake's magnitudes. This model assumes that any earthquake can come from one of two sources: the **background** events come from a Inhomogeneous Poisson Process, independent of time, and the **offspring** were triggered by a previous event (either a background, or another offspring). Another way to call these events are **aftershocks**, and have a more geological interpretation. Figure 1 shows the branching structure of a temporal **ETAS**, with events beign parents if they come from generation 0, and offsprings if they come from any other generation.

Figure 1 – Branching of an self-exciting point process.



In a later work, Ogata (1998) extended the ETAS model to accommodate the spatio-temporal distribution of earthquakes, in the form

$$\lambda_\theta(x, y, t|\mathcal{H}_t) = \mu(x, y) + \sum_{i:t_i<t} \kappa(m_i)g(t - t_i; m_i)f(x - x_i, y - y_i; m_i), \tag{1.4}$$

with it's components being:

- $\mu(x, y)$ is the background rate, representing a smooth function on the region S, with $\mu(x, y) > 0$. The parent events occur following this background rate, and it is assumed to be independent of time.

- $\kappa(m_i)$ is the productivity of the process, representing the expected number of events triggered from a event of magnitude m_i, taking form

$$\kappa(m) = A \exp[\alpha(m - m_0)],$$

with $A > 0$ and $\alpha > 0$ being unknown parameters, and m_0 the cutoff magnitude being considered in the study (i.e., all the n events in the catalog have magnitudes $m_i \geqslant m_0, i = 1, 2, \ldots, n$).

- $g(t - t_i)$ is the probability density function of the time (in days) of a triggered event occuring at time t_i, and is assumed to be independent of the event magnitude m_i. A power law density is considered, based in the modified Omori's law (Utsu et al. (1995))

$$g(t - t_i) = \begin{cases} (p-1)c^{p-1}\left((t - t_i) + c\right)^{-p}, & t - t_i > 0 \\ 0, & t - t_i \leqslant 0 \end{cases},$$

with $c > 0$ and $p > 1$ being unknown parameters.

- $f(x - x_i, y - y_i; m_i)$ is the probability density function of the location in space of a event triggered from a event occurred in (x_i, y_i) and magnitude m_i. The distances are considered to be in degrees[2] (x is the longitude and y is the latitude), and this can cause some considerable distortion in areas with big latitudes. An isotropic density function is considered

$$f(x - x_i, y - y_i; m_i) = \frac{q-1}{\pi D \exp[\gamma(m_i - m_0)]} \left(1 + \frac{(x - x_i)^2 + (y - y_i)^2}{D \exp[\gamma(m_i - m_0)]}\right)^{-q},$$

with $D > 0, \gamma > 0$ and $q > 1$ are unknown parameters.

Other parametrizations for $\lambda_\theta(x, y, t|\mathcal{H}_t)$ were proposed by Ogata, but this one offers great flexibility by allowing the times of triggered events to be independent of the magnitudes, and by having separated parameters that control different aspects of the earthquake's intensity function behavior (temporal decay rate, influence of magnitudes in the number of expected offspring, spatial clustering, etc). The model can also incorporate the distribution of the marks/magnitudes

$$\lambda_{\theta,\beta}(x, y, t, m|\mathcal{H}_t) = \nu(m)\lambda_\theta(x, y, t|\mathcal{H}_t), \tag{1.5}$$

where $\nu(m)$ follows the Gutenberg-Richter distribution (Gutenberg e Richter (1944))

$$\nu(m) = \beta \exp[-\beta(m - m_0)], \ \beta > 0.$$

The expected number of offsprings generated from any event, regardless of its magnitude is given by

$$e_{\text{off}} = \int_{m_0}^{\infty} \kappa(m)\nu(m)dm = \frac{A\beta}{\beta - \alpha}. \tag{1.6}$$

If $e_{\text{off}} < 1$, the model is stable (Daley e Vere-Jones (2003)), and has agglomerate size of $\frac{1}{(1-e_{\text{off}})} = \frac{\beta-\alpha}{\beta(1-A)-\alpha}$. Because the marks distribution is independent of the other parameters (as the complete intensity $\lambda_{\theta,\beta}(x,y,t,m|\mathcal{H}_t)$ is the product of $\nu(v)$ and $\lambda_\theta(x,y,t|\mathcal{H}_t)$), the estimation of β can be done without the need to estimate the other parameters.

1.3 Likelihood and Stochastic Declustering

Estimating the model parameters β and $\theta = (A, \alpha, c, p, D, \gamma, q)$ and the background intensity function $\mu(x,y)$ is crucial to understand the seismological dynamic of a certain region, as it dictates when and where the aftershocks of an earthquake have more probability of happening. Moreover, early-warning systems depend on this estimates to make predictions in feasible time, as the time needed for these systems to mitigate further damages are between 1 and 10 minutes (Allen e Melgar (2019)).

For this, maximum likelihood methods for estimating the vector parameters have many desirable properties, such as being consistent, asymptotically unbiased and efficient (Ogata (1978)). The ETAS model complete log-likelihood is given by

$$l(\theta,\beta|\mathcal{H}_t) = \sum_{i=1}^{n} \log(\lambda_{\theta,\beta}(x_i,y_i,t_i,m_i|\mathcal{H}_t)) - \int_{m_0}^{\infty}\int_0^T\int\int_S \lambda_{\theta,\beta}(x,y,t,m|\mathcal{H}_t)dx\,dy\,dt\,dm, \tag{1.7}$$

where T is the time of the last event in the catalog (the time integral starts at 0 because we can re-scale the event times $(t_1,t_2,\ldots,t_n)$ such that we have the fist event occur at time 0 with $(t_1-t_1, t_2-t_1,\ldots,t_n-t_1) = (0,t_2^*,\ldots,T)$). As seen in **??**, β and θ are separable, and $\int_{m_0}^{\infty}\nu(m)dm = 1$ (because it is a probability density function), we can decompose the previous log-likelihood as

$$l(\theta,\beta|\mathcal{H}_t) = l_1(\beta|\mathcal{H}_t) + l_2(\theta|\mathcal{H}_t)$$

where

$$l_1(\beta|\mathcal{H}_t) = \sum_{i=1}^{n}\log(\nu(m_i)) = n\log(\beta) - \beta\sum_{i=1}^{n}(m_i - m_0) \tag{1.8}$$

and

$$l_2(\theta|\mathcal{H}_t) = \sum_{i=1}^{n}\log(\lambda_\theta(x_i,y_i,t_i|\mathcal{H}_t)) - \int_0^T\int\int_S \lambda_\theta(x,y,t|\mathcal{H}_t)dx\,dy\,dt. \tag{1.9}$$

The maximum likelihood estimator of β can be obtained analytically using Equation (1.8)

$$\hat{\beta} = \frac{n}{\sum_{i=1}^{n}(m_i - m_0)},$$

whereas the maximization of Equation (1.9) requires some form of numerical optimization method, to approximate the integral part. When we evaluate the integral term, we have

$$\int_0^T \int \int_S \lambda_\theta(x, y, t|\mathcal{H}_t) = T \int \int_S \mu(x, y)dx\, dy \; + \sum_{i:t_i<t} \kappa(m_i)G(T - t_i)F(S - s_i; m_i)$$

where $G(t)$ and $F(s)$ are the cumulative probability functions of $g(t)$ and $f(s)$, respectively, and are defined in Appendix A. For any event not occurring near the spatio-temporal border (so that all the probability of being a offspring comes from the events inside the observed window, and not from any external event), we have that $G(T - t_i) \approx 1$ and $F(S - s_i) \approx 1$. This approximation simplifies the integral, making the log-likelihood easier to evaluate, when the assumptions are met.

However, the direct optimization of the complete log-likelihood remains complicated, due to two main factors: the $\log(\lambda_\theta(x_i, y_i, t_i|\mathcal{H}_t))$ term is the logarithm of a sum that has all the parameters and is not separable, and the computational cost for each evaluation is in the order of $\mathcal{O}(n^2)$. If the underlying branching structure were known, the log-likelihood would be separable because the terms for the contribution of the background events and the offsprings would be either 1 or 0 for any occurrence.

Consider that each event j has a parent $u_j \in \{0, 1, \ldots, j-1\}$, in which $u_j = 0$ indicates a event that has come from the background rate function. If u_j were observed, we would have

$$\begin{aligned} l_2(\theta|\mathcal{H}_t) = \sum_{j=1}^{n} \left[\mathbb{1}(u_j = 0) \log(\mu(x_j, y_j)) + \sum_{i<j} \mathbb{1}(u_j \neq 0) \log(\kappa(m_i)g(t_j - t_i)f(s_j - s_i; m_i)) \right] - \\ T \int \int_S \mu(x, y)dx\, dy + \sum_{i:t_i<t} \kappa(m_i)G(T - t_i)F(S - s_i; m_i), \end{aligned} \tag{1.10}$$

making the log-likelihood separable, and with direct evaluation of the parameters. Zhuang, Ogata e Vere-Jones (2002) proposed a stochastic declustering method based on the decomposition of the conditional intensity in the ratio between the total intensity and the local intensity. Denote ρ_j as the probability that the event j is an background event, and ρ_{ij} the probability that event j was triggered by event i. Then, we have

$$\rho_{ij} = \frac{\kappa(m_i)g(t_j - t_i)f(s_j - s_i; m_i)}{\lambda_\theta(x_j, y_j, t_j|\mathcal{H}_{t_j})}, \tag{1.11}$$

and

$$\rho_j = \frac{\mu(x_j, y_j)}{\lambda_\theta(x_j, y_j, t_j|\mathcal{H}_{t_j})} = 1 - \sum_{i<j} \rho_{ij}. \tag{1.12}$$

Thus, the correct estimation of θ should in turn help estimate wheter an earthquake was triggerred by the background rate, or if it is an aftershock of another

previous event in the catalog. When all the events in the catalog are far from the spatio-temporal border of the observed window W, we can consider the approximation $G(T-t_i) = F(S-s_i;m_i) \approx 1$, and with the results from Equations (1.11) and (1.12), we can substitute the indicator function for the unobserved u_j's in Equation (1.10) for their estimated probabilities as such

$$l_2(\theta|\mathcal{H}_t) = \sum_{j=1}^{n} \rho_j \log\Big(\mu(x_j,y_j)\Big) + \sum_{i=1}^{n}\sum_{i<j}(1-\rho_j)\log\Big(\kappa(m_i)g(t_j-t_i)f(s_j-s_i;m_i)\Big) - T\int\int_S \mu(x,y)dx\,dy + \sum_{i:t_i<t}\kappa(m_i). \tag{1.13}$$

This form of the log-likelihood is easier to maximize given it's additive terms, but requires the calculated probabilities ρ_j

Part II

Materials and Methods

2 Methods

2.1 Parameter Estimation

2.2 Goodness of fit tests

Theorem 3 (Random time change)**.** *Let* $0 < t_1 < t_2 < \ldots < t_n$ *be the realization of a Point Process with conditional intensity function* $\lambda(t|\mathcal{H}_t)$*, where* $\lambda(t|\mathcal{H}_t) > 0$ *for all* $t \in [0, T]$*. Then, the transformed times:*

$$\{\Lambda(t_1), \Lambda(t_2), \ldots, \Lambda(t_n)\} = \{\tau_1, \tau_2, \ldots, \tau_n\},$$

are the points of a unidimensional unit rate Poisson Process, if $\Lambda(t|\mathcal{H}_t)$ *is almost surely finite for all* $t \in [0, T]$*.*

This theorem holds for spatio-temporal point processes, and it is used for goodness of fit tests.

3 Considerações Finais

Precisa de ajuda?

Consulte os manuais abnTEX2 (ARAUJO, 2015a; ARAUJO, 2015c; ARAUJO, 2015d) e da classe memoir (WILSON; MADSEN, 2010).

Envie um e-mail para `gfabinhomat@gmail.com` (Fábio Rodrigues Silva), que é o responsável pela manutenção desta versão de adaptação do modelo canônico abnTEX2.

Consulte a FAQ com perguntas frequentes e comuns no portal do abnTEX2: https://github.com/abntex/abntex2/wiki/FAQ.

Inscreva-se no grupo de usuários LATEX: http://groups.google.com/group/latex-br, tire suas dúvidas e ajude outros usuários.

Participe também do grupo de desenvolvedores do abnTEX2: http://groups.google.com/group/abntex2 e faça sua contribuição à ferramenta.

Bibliography

ALLEN, R. M.; MELGAR, D. Earthquake Early Warning: Advances, Scientific Challenges, and Societal Needs. *Annual Review of Earth and Planetary Sciences*, v. 47, n. 1, p. 361–388, maio 2019. ISSN 0084-6597, 1545-4495. Disponível em: https://www.annualreviews.org/doi/10.1146/annurev-earth-053018-060457. Citado na página 23.

ARAUJO, L. C. *A classe abntex2: Modelo canônico de trabalhos acadêmicos brasileiros compatível com as normas ABNT NBR 14724:2011, ABNT NBR 6024:2012 e outras*. [S.l.], 2015. Disponível em: http://www.abntex.net.br/. Citado 2 vezes nas páginas 17 and 28.

_____. *Modelo Canônico de Trabalho Acadêmico com abnTeX2*. [S.l.], 2015. Disponível em: http://www.abntex.net.br/. Citado na página 17.

_____. *O pacote abntex2cite: Estilos bibliográficos compatíveis com a ABNT NBR 6023*. [S.l.], 2015. Disponível em: http://www.abntex.net.br/. Citado 2 vezes nas páginas 17 and 28.

_____. *O pacote abntex2cite: tópicos específicos da ABNT NBR 10520:2002 e o estilo bibliográfico alfabético (sistema autor-data)*. [S.l.], 2015. Disponível em: http://www.abntex.net.br/. Citado 2 vezes nas páginas 17 and 28.

ASSOCIAÇÃO BRASILEIRA DE NORMAS TÉCNICAS. *NBR 6028*: Resumo - apresentação. Rio de Janeiro, 2003. 2 p. Citado na página 8.

COMISSÃO CENTRAL DE PÓS-GRADUAÇÃO – UNIVERSIDADE ESTADUAL DE CAMPINAS. *Informação CCPG/06/2025*. Campinas, 2025. 27 p. Disponível em: https://prpg.unicamp.br/documentos-e-normas/normas/instrucoes-normativas/. Acesso em: 06 fev. 2026. Citado na página 17.

DALEY, D. J.; VERE-JONES, D. *An introduction to the theory of point processes*. 2nd ed. ed. New York: Springer, 2003. ISBN 9780387955414 9780387213378 9780387498355. Citado 2 vezes nas páginas 19 and 23.

GUTENBERG, B.; RICHTER, C. F. Frequency of earthquakes in California*. *Bulletin of the Seismological Society of America*, v. 34, n. 4, p. 185–188, out. 1944. ISSN 1943-3573, 0037-1106. Disponível em: https://pubs.geoscienceworld.org/ssa/bssa/article/34/4/185/101140/Frequency-of-earthquakes-in-California. Citado na página 22.

HAWKES, A. G. Point Spectra of Some Mutually Exciting Point Processes. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, v. 33, n. 3, p. 438–443, out. 1971. ISSN 1369-7412, 1467-9868. Disponível em: https://academic.oup.com/jrsssb/article/33/3/438/7027167. Citado na página 20.

_____. Spectra of some self-exciting and mutually exciting point processes. *Biometrika*, v. 58, n. 1, p. 83–90, 1971. ISSN 0006-3444, 1464-3510. Disponível em: https://academic.oup.com/biomet/article-lookup/doi/10.1093/biomet/58.1.83. Citado na página 21.

LUDWIG, G.; WANG, Y.; CHU, T.; WANG, H.; ZHU, J. Bayesian analysis and variable selection for spatial count data with an application to Rio de Janeiro gun violence. *Spatial Statistics*, v. 67, p. 100890, jun. 2025. ISSN 22116753. Disponível em: https://linkinghub.elsevier.com/retrieve/pii/S2211675325000120. Citado na página 20.

MOHLER, G. O.; SHORT, M. B.; BRANTINGHAM, P. J.; SCHOENBERG, F. P.; TITA, G. E. Self-Exciting Point Process Modeling of Crime. *Journal of the American Statistical Association*, v. 106, n. 493, p. 100–108, mar. 2011. ISSN 0162-1459, 1537-274X. Disponível em: http://www.tandfonline.com/doi/abs/10.1198/jasa.2011.ap09546. Citado na página 20.

OGATA, Y. The asymptotic behaviour of maximum likelihood estimators for stationary point processes. *Annals of the Institute of Statistical Mathematics*, v. 30, n. 2, p. 243–261, dez. 1978. ISSN 0020-3157, 1572-9052. Disponível em: http://link.springer.com/10.1007/BF02480216. Citado na página 23.

_____. Statistical Models for Earthquake Occurrences and Residual Analysis for Point Processes. *Journal of the American Statistical Association*, v. 83, n. 401, p. 9–27, mar. 1988. ISSN 0162-1459, 1537-274X. Disponível em: http://www.tandfonline.com/doi/abs/10.1080/01621459.1988.10478560. Citado na página 21.

_____. Space-Time Point-Process Models for Earthquake Occurrences. *Annals of the Institute of Statistical Mathematics*, v. 50, n. 2, p. 379–402, jun. 1998. ISSN 0020-3157, 1572-9052. Disponível em: https://link.springer.com/10.1023/A:1003403601725. Citado na página 21.

UTSU, T.; OGATA, Y.; S, R.; Matsu'ura. The Centenary of the Omori Formula for a Decay Law of Aftershock Activity. *Journal of Physics of the Earth*, v. 43, n. 1, p. 1–33, 1995. ISSN 1884-2305, 0022-3743. Disponível em: http://www.jstage.jst.go.jp/article/jpe1952/43/1/43_1_1/_article. Citado na página 22.

WILSON, P.; MADSEN, L. *The Memoir Class for Configurable Typesetting - User Guide*. Normandy Park, WA, 2010. Disponível em: http://mirrors.ctan.org/macros/latex/contrib/memoir/memman.pdf. Acesso em: 19 dez. 2012. Citado 2 vezes nas páginas 17 and 28.

ZHUANG, J.; OGATA, Y.; VERE-JONES, D. Stochastic Declustering of Space-Time Earthquake Occurrences. *Journal of the American Statistical Association*, v. 97, n. 458, p. 369–380, jun. 2002. ISSN 0162-1459, 1537-274X. Disponível em: http://www.tandfonline.com/doi/abs/10.1198/016214502760046925. Citado na página 24.

Appendix

APPENDIX A – Cumulative distributions of $g(t)$ and $f(x, y)$

Considering the formulation of the probability density functions $g(t)$ and $f(x, y)$, in section 1.2, we can calculate their cumulative distribution function as such

$$\begin{aligned}
G(T) = \int_0^T g(t)dt &= \int_0^T (p-1)c^{p-1}(t+c)^{-p}dt \\
&= (p-1)c^{p-1}\int_0^T (t+c)^{-p}dt \\
&= (p-1)c^{p-1}\left[\frac{(t+c)^{1-p}}{1-p}\Big|_0^T\right] \\
&= (p-1)c^{p-1}\left[\frac{(T+c)^{1-p}}{1-p} + \frac{c^{1-p}}{p-1}\right] \\
&= 1 - \frac{(p-1)c^{p-1}}{(p-1)}(T+c)^{1-p} \\
&= 1 - \frac{c^{p-1}}{(T+c)^{p-1}} \\
&= 1 - \left(\frac{c}{T+c}\right)^{p-1}.
\end{aligned}$$

With this result, we can see that this is indeed a cumulative probability function, as $\lim_{T\to 0} G(T) = 0$ and $\lim_{T\to\infty} G(T) = 1$. For $f(x, y)$, consider the spherical coordinates $x = r\cos(\theta)$ and $y = r\text{sen}(\theta)$ and $D[\gamma(m - m_0)] = Q$. With those, we have

$$\begin{aligned}
F(R) = \int_0^{2\pi}\int_0^R f(r)dr\,d\theta &= \int_0^{2\pi}\int_0^R \frac{(q-1)}{\pi Q}\left(1+\frac{r^2}{Q}\right)^{-q} r\,dr\,d\theta \\
&= \frac{(q-1)}{\pi Q}\left(\int_0^{2\pi} d\theta\right)\left(\int_0^R r\left(1+\frac{r^2}{Q}\right)dr\right) \\
&= \frac{2(q-1)}{Q}\int_0^R r\left(1+\frac{r^2}{Q}\right)dr.
\end{aligned}$$

Let $u = 1 + \frac{r^2}{Q}$, so that $du = \frac{2r}{Q}dr$ and $r\,dr = \frac{Q}{2}du$. Using substitution, we have

$$
\begin{aligned}
F(R) &= \frac{2(q-1)}{Q} \int_1^{1+\frac{R^2}{Q}} u^{-q} \left(\frac{Q}{2}\right) du \\
&= (q-1) \int_1^{1+\frac{R^2}{Q}} u^{-q} du \\
&= (q-1) \left(\frac{u^{1-q}}{1-q} \Big|_1^{1+\frac{R^2}{Q}} \right) \\
&= -u^{1-q} \Big|_1^{1+\frac{R^2}{Q}} \\
&= 1 - \left(1 + \frac{R^2}{Q}\right)^{1-q},
\end{aligned}
$$

that is a valid cumulative probability function, as $\lim_{R^2 \to 0} = 0$ and $\lim_{R^2 \to \infty} = 1$.

APPENDIX B – Quisque libero justo

Quisque facilisis auctor sapien. Pellentesque gravida hendrerit lectus. Mauris rutrum sodales sapien. Fusce hendrerit sem vel lorem. Integer pellentesque massa vel augue. Integer elit tortor, feugiat quis, sagittis et, ornare non, lacus. Vestibulum posuere pellentesque eros. Quisque venenatis ipsum dictum nulla. Aliquam quis quam non metus eleifend interdum. Nam eget sapien ac mauris malesuada adipiscing. Etiam eleifend neque sed quam. Nulla facilisi. Proin a ligula. Sed id dui eu nibh egestas tincidunt. Suspendisse arcu.

APPENDIX C – Nullam elementum urna vel imperdiet sodales elit ipsum pharetra ligula ac pretium ante justo a nulla curabitur tristique arcu eu metus

Nunc velit. Nullam elit sapien, eleifend eu, commodo nec, semper sit amet, elit. Nulla lectus risus, condimentum ut, laoreet eget, viverra nec, odio. Proin lobortis. Curabitur dictum arcu vel wisi. Cras id nulla venenatis tortor congue ultrices. Pellentesque eget pede. Sed eleifend sagittis elit. Nam sed tellus sit amet lectus ullamcorper tristique. Mauris enim sem, tristique eu, accumsan at, scelerisque vulputate, neque. Quisque lacus. Donec et ipsum sit amet elit nonummy aliquet. Sed viverra nisl at sem. Nam diam. Mauris ut dolor. Curabitur ornare tortor cursus velit.

Morbi tincidunt posuere arcu. Cras venenatis est vitae dolor. Vivamus scelerisque semper mi. Donec ipsum arcu, consequat scelerisque, viverra id, dictum at, metus. Lorem ipsum dolor sit amet, consectetuer adipiscing elit. Ut pede sem, tempus ut, porttitor bibendum, molestie eu, elit. Suspendisse potenti. Sed id lectus sit amet purus faucibus vehicula. Praesent sed sem non dui pharetra interdum. Nam viverra ultrices magna.

Aenean laoreet aliquam orci. Nunc interdum elementum urna. Quisque erat. Nullam tempor neque. Maecenas velit nibh, scelerisque a, consequat ut, viverra in, enim. Duis magna. Donec odio neque, tristique et, tincidunt eu, rhoncus ac, nunc. Mauris malesuada malesuada elit. Etiam lacus mauris, pretium vel, blandit in, ultricies id, libero. Phasellus bibendum erat ut diam. In congue imperdiet lectus.

Annex

ANNEX A – Morbi ultrices rutrum lorem.

Sed mattis, erat sit amet gravida malesuada, elit augue egestas diam, tempus scelerisque nunc nisl vitae libero. Sed consequat feugiat massa. Nunc porta, eros in eleifend varius, erat leo rutrum dui, non convallis lectus orci ut nibh. Sed lorem massa, nonummy quis, egestas id, condimentum at, nisl. Maecenas at nibh. Aliquam et augue at nunc pellentesque ullamcorper. Duis nisl nibh, laoreet suscipit, convallis ut, rutrum id, enim. Phasellus odio. Nulla nulla elit, molestie non, scelerisque at, vestibulum eu, nulla. Ut odio nisl, facilisis id, mollis et, scelerisque nec, enim. Aenean sem leo, pellentesque sit amet, scelerisque sit amet, vehicula pellentesque, sapien.

ANNEX B – Cras non urna sed feugiat cum sociis natoque penatibus et magnis dis parturient montes nascetur ridiculus mus

Sed consequat tellus et tortor. Ut tempor laoreet quam. Nullam id wisi a libero tristique semper. Nullam nisl massa, rutrum ut, egestas semper, mollis id, leo. Nulla ac massa eu risus blandit mattis. Mauris ut nunc. In hac habitasse platea dictumst. Aliquam eget tortor. Quisque dapibus pede in erat. Nunc enim. In dui nulla, commodo at, consectetuer nec, malesuada nec, elit. Aliquam ornare tellus eu urna. Sed nec metus. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas.

ANNEX C – Fusce facilisis lacinia dui

Phasellus id magna. Duis malesuada interdum arcu. Integer metus. Morbi pulvinar pellentesque mi. Suspendisse sed est eu magna molestie egestas. Quisque mi lorem, pulvinar eget, egestas quis, luctus at, ante. Proin auctor vehicula purus. Fusce ac nisl aliquam ante hendrerit pellentesque. Class aptent taciti sociosqu ad litora torquent per conubia nostra, per inceptos hymenaeos. Morbi wisi. Etiam arcu mauris, facilisis sed, eleifend non, nonummy ut, pede. Cras ut lacus tempor metus mollis placerat. Vivamus eu tortor vel metus interdum malesuada.