

# Dietary fiber intervention on gut microbiota composition in healthy adults: a systematic review and meta-analysis

Daniel So,<sup>1</sup> Kevin Whelan,<sup>2</sup> Megan Rossi,<sup>2</sup> Mark Morrison,<sup>3,4</sup> Gerald Holtmann,<sup>4,5</sup> Jaimon T Kelly,<sup>1</sup> Erin R Shanahan,<sup>3,5</sup> Heidi M Staudacher,<sup>4</sup> and Katrina L Campbell<sup>1,6</sup>

<sup>1</sup>Faculty of Health Sciences and Medicine, Bond University, Gold Coast, Australia; <sup>2</sup>Department of Nutritional Sciences, King's College, London, United Kingdom; <sup>3</sup>The University of Queensland Diamantina Institute, Translational Research Institute; <sup>4</sup>Faculty of Medicine, University of Queensland, Brisbane, Australia; <sup>5</sup>Department of Gastroenterology & Hepatology; and <sup>6</sup>Department of Nutrition and Dietetics, Princess Alexandra Hospital, Brisbane, Australia

## ABSTRACT

**Background:** Dysfunction of the gut microbiota is frequently reported as a manifestation of chronic diseases, and therefore presents as a modifiable risk factor in their development. Diet is a major regulator of the gut microbiota, and certain types of dietary fiber may modify bacterial numbers and metabolism, including short-chain fatty acid (SCFA) generation.

**Objective:** A systematic review and meta-analysis were undertaken to assess the effect of dietary fiber interventions on gut microbiota composition in healthy adults.

**Design:** A systematic search was conducted across MEDLINE, EMBASE, CENTRAL, and CINAHL for randomized controlled trials using culture and/or molecular microbiological techniques evaluating the effect of fiber intervention on gut microbiota composition in healthy adults. Meta-analyses via a random-effects model were performed on alpha diversity, prespecified bacterial abundances including *Bifidobacterium* and *Lactobacillus* spp., and fecal SCFA concentrations comparing dietary fiber interventions with placebo/low-fiber comparators.

**Results:** A total of 64 studies involving 2099 participants were included. Dietary fiber intervention resulted in higher abundance of *Bifidobacterium* spp. (standardized mean difference (SMD): 0.64; 95% CI: 0.42, 0.86;  $P < 0.00001$ ) and *Lactobacillus* spp. (SMD: 0.22; 0.03, 0.41;  $P = 0.02$ ) as well as fecal butyrate concentration (SMD: 0.24; 0.00, 0.47;  $P = 0.05$ ) compared with placebo/low-fiber comparators. Subgroup analysis revealed that fructans and galacto-oligosaccharides led to significantly greater abundance of both *Bifidobacterium* spp. and *Lactobacillus* spp. compared with comparators ( $P < 0.00001$  and  $P = 0.002$ , respectively). No differences in effect were found between fiber intervention and comparators for  $\alpha$ -diversity, abundances of other prespecified bacteria, or other SCFA concentrations.

**Conclusions:** Dietary fiber intervention, particularly involving fructans and galacto-oligosaccharides, leads to higher fecal abundance of *Bifidobacterium* and *Lactobacillus* spp. but does not affect  $\alpha$ -diversity. Further research is required to better understand the role of individual fiber types on the growth of microbes and the overall gut microbial community. This review was registered at PROSPERO as CRD42016053101. *Am J Clin Nutr* 2018;107:965–983.

**Keywords:** diet, dietary fiber, gastrointestinal microbiome, gastrointestinal microbiota, gut microbiota, prebiotic

## INTRODUCTION

The gut microbiota is a highly diverse and metabolically active community, consisting of  $\sim 3.9 \times 10^{13}$  microbial cells (1). These microbes participate in several functions beneficial to the host, including the fermentation of undigested nutrients (2, 3), synthesis of vitamins (4), and interaction with the immune system (5, 6). A number of disorders, including irritable bowel syndrome and type 2 diabetes mellitus, have been linked with disturbances in gut microbiota composition (2, 7–9). Such an association presents the gut microbiota as a potentially modifiable risk factor in the etiology of these conditions.

The gut microbiota can be detected and enumerated via different methods ranging from culture to next-generation sequencing (6, 10, 11), and can be characterized by measures of diversity and bacterial abundances (12, 13). Alpha diversity of the gut microbiota describes the richness (number of taxonomically distinct organisms present) and evenness (relative abundances of organisms) of its composition (12, 13), with cross-sectional studies demonstrating inverse associations between  $\alpha$ -diversity and disease states (7–9). Specific bacteria shown to be more abundant in

The authors reported no funding received for this study.

Supplemental Tables 1–7 and Supplemental Figures 1–7 are available from the “Supplementary data” link in the online posting of the article and from the same link in the online table of contents at <https://academic.oup.com/ajcn>.

Address correspondence to KLC (e-mail: [kcampbel@bond.edu.au](mailto:kcampbel@bond.edu.au)).

Abbreviations used: FISH, fluorescence in situ hybridization; GI, gastrointestinal; HMO, human milk oligosaccharide; MD, mean difference; OTU, operational taxonomic unit; qPCR, quantitative polymerase chain reaction; RCT, randomized controlled trial; SCFA, short-chain fatty acid; SMD, standardized mean difference.

Received August 15, 2017. Accepted for publication February 14, 2018.

First published online May 11, 2018; doi: <https://doi.org/10.1093/ajcn/nqy041>.

health compared with disease states include *Bifidobacterium* and *Lactobacillus* spp. (2, 7, 14), whose functions include carbohydrate fermentation and vitamin synthesis (15–18). Furthermore, increasing evidence supports the importance of “keystone” bacterial species, whose absence may have profound consequences for the host, as well as other members of the microbial community and their metabolic outputs, including the short-chain fatty acid (SCFA) butyrate (19–23). Butyrate is of particular relevance to health owing to its beneficial properties such as its immunomodulatory effects (24, 25).

Dietary fiber is defined as nondigestible carbohydrates of  $\geq 3$  monomeric units found inherently in foods, and also includes isolated or synthetic fibers with demonstrated physiologic benefits (26–28). It is a key candidate in facilitating changes in the gut microbiota, as it escapes digestion by the host in the small intestine to pass into the colon where it is available to the microbial community. Dietary fiber encompasses an array of heterogeneous compounds whose physicochemical properties vary based on their particle size, chemical structure, solubility, viscosity, and fermentability (29, 30). Fibers with fermentable characteristics are substrates for the microbial population in the colon, stimulating growth of specific organisms and leading to production of various metabolites including SCFA (19, 29, 31). Indeed, some fibers can be further classified as “prebiotic” (e.g., fructans) if they have been shown to be selectively utilized by host microorganisms conferring a health benefit (32).

The current body of evidence regarding the effect of dietary fiber on the gut microbiota is informed via specific prebiotic fiber interventions (33, 34), whole-diet interventions (35–37), and cross-sectional associations (38, 39). However, these investigations are limited in that prebiotic fibers represent only a subset of total dietary fiber, and confounding factors such as disease states and intake of other fermentable substrates are unaccounted for in whole-diet studies and cross-sectional studies (40). Therefore, there is a gap in knowledge regarding the precise impact of dietary fiber intervention on the gut microbiota in healthy subjects, and this is the focus of this systematic review.

## METHODS

This systematic review was conducted in line with the guidelines of the Preferred Reporting Items for Systematic Reviews and Meta-Analysis (PRISMA) statement (41), and the guidelines of the *Cochrane Handbook for Systematic Reviews and Interventions* (42). The methods including the eligibility criteria, search strategy, extraction process, and analysis were prespecified and documented in a protocol that was published in the PROSPERO as CRD42016053101: [http://www.crd.york.ac.uk/PROSPERO/display\\_record.asp?ID=CRD42016053101](http://www.crd.york.ac.uk/PROSPERO/display_record.asp?ID=CRD42016053101).

## Literature search

A literature search was performed in the electronic databases MEDLINE, EMBASE, CENTRAL, and CINAHL (from inception to 4 October 2017), through the use of a combination of subject headings, free text terms, and synonyms relevant to this review, in consultation with an experienced systematic review search librarian (Supplemental Tables 1–4). There was no date

or language restriction in the search strategy. A multistep search approach was taken to retrieve relevant studies through additional hand-searching; contacting field experts; searching conference abstracts; theses and dissertations (ProQuest); and the International Clinical Trials Register Search Portal and clinicaltrials.gov to identify ongoing trials. Two review authors (DS and HMS) screened articles in a blinded, standardized manner, with disagreements in judgment resolved by consensus or a third reviewer (KLC).

## Study selection

Search results were merged into the reference management software Endnote (X7; Thomson Reuters) and de-duplicated before screening with the use of Rayyan (Qatar Computing Research Institute) (43). Full-text articles of potentially relevant studies were sought and reviewed. Attempts were made to contact the corresponding author when the full-text article provided inadequate information to assess eligibility or extract relevant data. Studies were included if they met all of the following criteria: 1) randomized controlled trial (RCT), cluster RCT, or quasi-RCT; 2) inclusion of healthy adult participants ( $\geq 18$  y of age); 3) intervention aimed at increasing fiber intake; 4) inclusion of a placebo for supplement interventions (e.g., maltodextrin), and either low-fiber control (e.g., white bread) or habitual diet group for food interventions as comparators; 5) measured fecal microbiota related outcomes at the end of intervention.

Studies that were solely investigating enteral nutrition and those that included participants with an acute or chronic disease, including gastrointestinal (GI) conditions such as celiac disease, inflammatory bowel disease, irritable bowel syndrome, and other functional GI disorders, were excluded. Studies including mixed population groups in which the healthy subgroup was not reported separately were also excluded. Studies that included overweight and obese participants who were otherwise healthy and without any abnormal clinical parameters (e.g., elevated blood pressure) were included. Interventions eligible for inclusion provided an increase in fiber intake achieved through 1) dietary counselling to increase dietary fiber intake from food; 2) food intervention (e.g., added cereals); or 3) fiber supplementation. Dietary counselling studies or food interventions were only included if fiber modification was the primary aim of the intervention.

The primary outcome was between-group differences in  $\alpha$ -diversity of fecal microbiota at the end of the intervention. Measures of  $\alpha$ -diversity included the total number of observed operational taxonomic units (OTUs) (the number of taxonomically related groups of bacteria, evaluating richness); Chao1 index (a nonparametric richness estimator); Shannon diversity index (a metric combining richness and evenness, with equal weighting given to abundant and rare species); and Simpson diversity index (a metric of richness and evenness, in which more weighting is given to abundant species). Secondary outcomes were between-group differences in abundances of the following commonly measured bacterial groups: *Bifidobacterium* spp.; *Lactobacillus* spp.; *Roseburia* spp.; *Akkermansia muciniphila*; *Eubacterium hallii*; *Eubacterium rectale*; *Faecalibacterium prausnitzii*; and *Ruminococcus bromii*. Studies were included if they reported on either primary or secondary outcomes. Between-group differences in fecal SCFAs (individual and total) were included as an exploratory outcome.

## Data extraction and management

Two reviewers (DS and HMS) independently extracted the data from eligible studies. Data extracted included: study design (duration, location, details of “run-in” and “washout” periods); participant characteristics; intervention details (fiber type, fiber dose, intervention delivery, compliance, assessment and control of dietary intake); and other information including antibiotic or probiotic use.

For all prespecified primary, secondary, and exploratory outcome data, the mean, SD, SE, or 95% CIs that were reported at end of intervention were extracted for analysis. When studies used multiple intervention groups of different fiber doses, data for the highest intervention dose were extracted. When studies used multiple intervention groups of different fibers at the same dose compared with a single control group, data were extracted from each intervention group and pooled together. A weighted average of the intervention groups and the study variance were then calculated (44).

Risk of bias was independently assessed by 2 reviewers (DS and HMS) using Cochrane methodology (45). The review assessed “other bias” regarding the control of dietary intake during the study. This included examining whether dietary advice (e.g., to maintain dietary intake or avoid probiotic food sources) was provided, whether dietary compliance and/or intake were measured and reported, and whether adjustments in statistical analysis were made if differences in dietary intake were found.

## Statistical analysis

The overall treatment effect of fiber on primary and secondary outcomes was calculated as the difference between the end of intervention values for the intervention and comparator groups. Data reported as median and IQR were converted to mean and SD as previously described (46). Variance was calculated from the SD and SE of end of intervention values, or from the CIs when these values were not available (46). In crossover studies, the mean and SD, SE, or CI of intervention and control periods were extracted and analyzed separately (47). When end of intervention endpoint data were unable to be obtained, the results were described in text only.

Meta-analysis was performed when outcomes were reported in  $\geq 2$  studies with the use of Revman (version 5.3; Cochrane Collaboration). The mean difference (MD) was used to calculate effect sizes when outcome data were presented in the same units (Shannon diversity index, total number of observed OTUs). When outcome data were reported in different units, effect sizes were calculated with the use of the standardized mean difference (SMD) (bacterial abundances, fecal SCFA concentration).

A random-effects model was used to produce a pooled estimate of the MD or SMD, and the fixed-effects model was used to check for robustness and potential outliers. Inconsistencies between studies were assessed with the use of the  $I^2$  statistic, in which significant heterogeneity was defined as  $I^2 \geq 50\%$ .

Predefined subgroup analyses were undertaken for primary and secondary outcomes that were reported in  $\geq 2$  studies in each subgroup. Predefined subgroup analyses included intervention types (supplements and dietary interventions), fiber types (accepted and candidate prebiotic fibers defined by Roberfroid et al., and general fibers defined by the review) (34), dose-response

(comparing difference in fiber intake between intervention and control groups of  $<5$  g/d, 5–10 g/d, and  $>10$  g/d), trial design (parallel and crossover), and microbial analysis method (e.g., culture, sequencing). Fructans and galacto-oligosaccharides were classified as “accepted prebiotic” fibers, whereas “candidate prebiotic” fibers included a broader range of fibers including polydextrose and resistant starch (34). The term “general fiber” was used by the review to describe fibers not classified as either accepted or candidate prebiotics, and is not a formal term used to describe fibers in the literature. Post hoc subgroup analyses were undertaken for exploratory outcomes based on reporting method of fecal SCFA concentrations (dry weight of feces and wet weight of feces).

For the fiber type subgroup analysis only, the fiber arm with the highest prebiotic classification (e.g., accepted prebiotic as opposed to a general fiber) was selected if multiple intervention groups were reported. When multiple arms of the same prebiotic classification were presented, the interventions were pooled together and a weighted average of the intervention arms and study variance were calculated (44). Significant outliers were determined by visual inspection as well as through a study-by-study sensitivity analysis, in which each study was sequentially omitted and the remaining data reassessed. If a study contributed to over 30% heterogeneity (based on changes to the  $I^2$  statistic) then it was removed from the analysis in the sensitivity analysis. Funnel plots were generated for outcomes in which  $\geq 10$  studies were included in meta-analysis (48) and reporting bias was detected by assessment of funnel plot asymmetry by visual inspection.


## RESULTS

### Study characteristics

Study identification and selection are detailed in the PRISMA flow chart (Figure 1). The initial electronic and manual search generated 3829 records. After review of full texts (Supplemental Table 5), 64 publications, along with 3 secondary studies (49–51) reporting additional outcomes from the primary publications, fulfilled the inclusion criteria and were included in the review.

The 64 included primary studies analyzed a total of 2099 participants. Of these 64 studies, 29 were parallel RCTs (52–80) and 35 were crossover RCTs (81–115). Five crossover trials did not include a washout period (84, 93, 95, 105, 108). The majority of studies (52 studies) used fiber supplementation, including: accepted prebiotic fiber (26 studies) (52, 54–58, 61, 62, 65, 67, 70, 74, 86, 90, 92, 95, 97, 100, 102, 103, 105, 107, 109–111, 115); candidate prebiotic fiber (18 studies) (53, 63, 64, 66, 68, 69, 73, 77, 81, 83, 84, 87, 88, 91, 99, 101, 112, 113); general fiber (7 studies) (59, 60, 72, 76, 80, 93, 94); and a fiber mix (108). The remaining 12 studies used food intervention by providing key food items (e.g., whole-grain cereal) to supplement the diet (71, 78, 82, 85, 89, 96, 98) or provided all food and fluid to participants (75, 79, 104, 106, 114). Intervention doses ranged from 1.2 g/d to 50 g/d and treatment periods ranged from 5 d to 3 mo, with a median length of 3 wk.

Analysis techniques used to characterize fecal microbiota included: culture (15 studies) (52, 54–58, 65, 66, 69, 71, 73, 96, 98, 105, 114); fluorescence in situ hybridization (FISH) (20 studies) (53, 70, 74, 76, 82, 85, 89–92, 94, 99, 100, 103, 106, 108–110, 112, 113); quantitative polymerase chain reaction (qPCR) (11



**FIGURE 1** Flow diagram of studies evaluated in the systematic review. RCT, randomized controlled trial.

studies) (60, 63, 68, 81, 86, 87, 95, 102, 104, 107, 111); and next-generation sequencing (including 454 pyrosequencing and Illumina sequencing) (12 studies) (59, 62, 64, 72, 75, 77–80, 97, 101, 115). A combination of techniques were used in 6 primary studies (61, 67, 83, 84, 88, 93) and one secondary publication (49).

The outcomes of each meta-analysis are reported in **Table 1**. Results from subgroup analyses performed are included in **Supplemental Table 6**. Overall, outcome data from 58 studies were suitable for meta-analysis; results from the following studies were unable to be statistically pooled and are presented

**TABLE 1**  
Statistical analysis for the outcomes reported in  $\geq 2$  randomized controlled trials and included in the meta-analysis<sup>1</sup>

| Outcomes                               | No. of studies in meta-analysis (references)  | $n^2$ | Results                                 |          | Heterogeneity   |          |           |
|--|---|-------|---|----------|-----------------|----------|-----------|
|  |   |       | Meta-analysis overall estimate (95% CI) | $P$      | Chi-square test | $P$      | $I^2$ (%) |
| Shannon diversity index                | 6 (64, 72, 75, 80, 84, 88)  | 127   | MD: -0.06 (-0.25, 0.12)                 | 0.48     | 10.73           | 0.06     | 53        |
| Total number of observed OTUs          | 3 (72, 75, 84)  | 53    | MD: -4.37 (-42.92, 34.19)               | 0.82     | 0.07            | 0.97     | 0         |
| <i>Bifidobacterium</i> spp.            | 51 (52-58, 60, 61, 63-68, 70, 71, 73-76, 82, 84-94, 96-112, 114)                                | 1629  | SMD: 0.64 (0.42, 0.86)                  | <0.00001 | 327.93          | <0.00001 | 85        |
| <i>Lactobacillus</i> spp. <sup>3</sup> | 23 (52, 55, 56, 60, 63-65, 67, 68, 73, 75, 76, 84, 87, 93, 96, 97, 99, 104, 105, 107, 111, 114) | 670   | SMD: 0.22 (0.03, 0.41)                  | 0.02     | 42.8            | 0.005    | 49        |
| <i>Faecalibacterium prausnitzii</i>    | 13 (53, 61, 67, 68, 74, 84, 88, 94, 99-101, 110, 112)   | 519   | SMD: 0.14 (-0.12, 0.39)                 | 0.29     | 37.53           | 0.0002   | 68        |
| <i>Roseburia</i> spp.                  | 4 (68, 79, 84, 97)  | 189   | SMD: 0.33 (-0.14, 0.80)                 | 0.17     | 10.16           | 0.02     | 70        |
| <i>Eubacterium rectale</i>             | 2 (84, 101)   | 30    | SMD: -0.26 (-1.20, 0.67)                | 0.58     | 3.94            | 0.05     | 75        |
| <i>Ruminococcus bromii</i>             | 3 (81, 84, 101)   | 76    | SMD: 0.15 (-0.15, 0.45)                 | 0.33     | 1.1             | 0.58     | 0         |
| Total SCFA                             | 13 (52, 55, 59, 63, 64, 67, 73, 80, 82, 84, 86, 91, 94)   | 406   | SMD: 0.11 (-0.05, 0.27)                 | 0.19     | 6.46            | 0.89     | 0         |
| Acetate                                | 18 (52, 53, 63, 66, 71, 74, 77, 80, 82, 84, 86, 90, 91, 93, 94, 96, 103, 112)                   | 657   | SMD: 0.28 (-0.08, 0.63)                 | 0.13     | 119.36          | <0.00001 | 86        |
| Propionate                             | 19 (52, 53, 63, 66, 71, 74, 77, 80, 82, 84, 86, 90, 91, 93, 94, 96, 103, 112, 115)              | 677   | SMD: 0.01 (-0.20, 0.22)                 | 0.95     | 46.23           | 0.0003   | 61        |
| Butyrate                               | 20 (52, 53, 59, 63, 66, 71, 74, 77, 80, 82, 84, 86, 90, 91, 93, 94, 96, 103, 112, 115)          | 712   | SMD: 0.24 (0.00, 0.47)                  | 0.05     | 64.21           | <0.00001 | 70        |

<sup>1</sup>Data were meta-analyzed through the use of a random-effects model and presented as MDs or SMDs as appropriate. Statistical heterogeneity was assessed via the chi-square test and quantified with the use of the  $I^2$  statistic. MD, mean difference; OTU, operational taxonomic unit; SCFA, short-chain fatty acid; SMD, standardized mean difference.

<sup>2</sup>Number of participants in meta-analysis.

<sup>3</sup>Results from outlier study excluded from this meta-analysis.



narratively under their respective subheadings (59, 62, 69, 77–79, 83, 93, 95, 97, 101, 113, 115). The characteristics of included studies, separated into fiber supplementation studies and food interventions, are presented in **Tables 2** and **3**, respectively.

### Dietary fiber and gut microbiota diversity ( $\alpha$ -diversity)

Alpha-diversity was measured in 13 studies involving 393 participants (49, 59, 64, 72, 75, 77, 79, 80, 83, 88, 93, 97, 101).

Ten studies reported  $\alpha$ -diversity through the use of Shannon diversity index. Of these, 6 reported the metric in a form suitable for inclusion in the meta-analysis (49, 64, 72, 75, 80, 88). Dietary fiber intervention had no effect on  $\alpha$ -diversity compared with placebo/low-fiber comparators (MD:  $-0.06$  Shannon diversity index; 95% CI:  $-0.25, 0.12$ ;  $P = 0.48$ ), albeit with substantial heterogeneity ( $I^2 = 53\%$ ). In 2 of the studies not included in the meta-analysis, raffinose and resistant starch interventions did not lead to significant difference in  $\alpha$ -diversity compared with placebo (93, 101). A significant reduction in the  $\alpha$ -diversity of fecal microbiota from baseline was detected in a trial involving flaxseed mucilage, measured by both the exponential of Shannon diversity index ( $-38,010$ ; 95% CI:  $-64,473, -11,546$ ;  $P = 0.007$ ) as well as Simpson's inverse index ( $-17,515$ ; 95% CI:  $-30,992, -4038$ ;  $P = 0.014$ ), although a between-group comparison was not reported (59). Conversely, significant end of intervention differences in  $\alpha$ -diversity measured by Shannon diversity index ( $P = 0.013$ ) and inverse Simpson index ( $P = 0.004$ ) were detected between intervention and comparator groups in a supplementation trial involving resistant starch type 2 (77).

A study evaluating  $\alpha$ -diversity through Simpson's index found that it was significantly higher in the intervention group receiving polydextrose compared with placebo after 21 d ( $P = 0.014$ ) (88). A trial involving 15 g/d arabinoxylan supplementation reported variable intervention effects when  $\alpha$ -diversity was evaluated with different metrics:  $\alpha$ -diversity was significantly lower compared with placebo when measured through observed species ( $P = 0.029$ ), but there were no significant differences when assessed by Simpson's evenness ( $P = 0.063$ ) (80).

A separate meta-analysis was performed for the 3 studies reporting  $\alpha$ -diversity measured by total number of observed OTUs (49, 72, 75). Dietary fiber had no effect on  $\alpha$ -diversity compared with placebo/low-fiber comparators (MD:  $-4.37$  OTUs; 95% CI:  $-42.92, 34.19$ ;  $P = 0.82$ ), with no heterogeneity ( $I^2 = 0\%$ ). The Chao1 index was used to report  $\alpha$ -diversity in 2 studies, although there were insufficient data available, precluding meta-analysis. Neither trial reported significant differences between fiber intervention and placebo or low-fiber control (49, 83). A feeding trial comparing whole-grain and refined-grain diets found no difference in  $\alpha$ -diversity at end of intervention between the 2 groups, although the metric used to measure  $\alpha$ -diversity was not reported (79).

### Dietary fiber and bacterial abundances

Reporting of bacterial abundances differed across studies. Of the taxa of interest in this review, abundances of *Bifidobacterium* spp. (59 studies) and *Lactobacillus* spp. (28 studies) were most commonly reported. No studies reported on the abundance of *A. muciniphila*.

A total of 59 studies including 1896 participants reported the effect of dietary fiber on *Bifidobacterium* spp. abundance and, of these, 51 trials (1629 participants) reported data in a form suitable for meta-analysis (53–58, 60, 61, 63–68, 70, 71, 73–76, 81, 82, 84–94, 96–112, 114). Dietary fiber led to a significantly greater *Bifidobacterium* spp. abundance compared with placebo/low-fiber comparators (SMD: 0.64; 95% CI: 0.42, 0.86;  $P < 0.00001$ ), albeit with considerable heterogeneity ( $I^2 = 85\%$ ) (**Figure 2**).

However, subgroup analysis showed fiber interventions delivered through supplements resulted in a significantly higher *Bifidobacterium* spp. abundance compared with placebo/low-fiber controls (SMD: 0.75; 95% CI: 0.52, 0.98;  $P < 0.00001$ ,  $I^2 = 83\%$ ), whereas no differences were found between food interventions and comparators (SMD: 0.20; 95% CI:  $-0.36, 0.76$ ;  $P = 0.49$ ,  $I^2 = 88\%$ ), although considerable heterogeneity persisted in both analyses.

Subgroup analysis demonstrated interventions investigating fibers classified as accepted prebiotics and candidate prebiotics resulted in a significantly higher *Bifidobacterium* spp. abundance compared with placebo/low-fiber controls (accepted prebiotic fiber SMD: 0.68; 95% CI: 0.38, 0.98;  $P < 0.00001$ ,  $I^2 = 81\%$ ; candidate prebiotic fiber SMD: 0.77; 95% CI: 0.30, 1.24;  $P < 0.00001$ ,  $I^2 = 86\%$ ) (**Figure 2**). However, there was no difference in effect between the general fiber subgroup and comparators (SMD: 0.25; 95% CI:  $-0.16, 0.65$ ;  $P = 0.24$ ,  $I^2 = 86\%$ ). This subgroup analysis did not reduce the considerable heterogeneity across each subgroup.

Subgroup analysis of dose-response showed dietary fiber led to significantly higher *Bifidobacterium* spp. abundance compared with placebo/low-fiber comparators at all predefined dosages ( $\leq 5$  g/d fiber SMD: 0.51; 95% CI: 0.18, 0.84;  $P = 0.003$ ,  $I^2 = 70\%$ ; 5–10 g/d SMD: 0.48; 95% CI: 0.13, 0.83;  $P = 0.007$ ,  $I^2 = 87\%$ ;  $> 10$  g/d SMD: 0.85; 95% CI: 0.45, 1.25;  $P < 0.00001$ ,  $I^2 = 85\%$ ). No differences were found in subgroup analyses of trial design or microbiota analysis method (Supplemental Table 6).

Eight trials were not included in the meta-analysis. In the supplement trials of accepted prebiotics, a significantly higher *Bifidobacterium* spp. abundance was reported after supplementation involving inulin (115) and human milk oligosaccharides (HMO) (62) compared with placebo at the end of intervention, whereas a significant within-group increase from baseline was detected after 10 g/d inulin supplementation (95). In the candidate prebiotic trial of resistant starch supplementation, *Bifidobacterium* spp. abundance was significantly higher in the intervention group compared with placebo at end of intervention (77). In the supplement studies of general fiber, *Bifidobacterium* spp. abundance was higher after xylo-oligosaccharide supplementation compared with placebo (69) whereas manno-oligosaccharides had no effect on *Bifidobacterium* spp. compared with placebo (113). The third supplement trial of general fiber (resistant maltodextrin) reported no change in *Bifidobacterium* spp. abundance within groups using FISH, although a significant increase from baseline was reported for the intervention group on qPCR analysis (83). Finally, a food study comparing intakes of whole grains with refined-grain products found no significant difference in *Bifidobacterium* spp. abundance at the end of the intervention period (78).

*Lactobacillus* spp. abundance was measured in 28 studies involving 867 participants. Data from 24 studies (730 participants)

TABLE 2

Characteristics of randomized controlled trials of fiber supplementation comparing dietary fiber with placebo or low-fiber comparators in healthy adults<sup>1</sup>

| Study (reference)                             | Participants<br><i>n</i> ; age; <sup>2</sup> % F | Interventions   |           |                                | RCT design              |           |                  |        |         |                                      |
|---|--|---|-----------|--------------------------------|-------------------------|-----------|------------------|--------|---------|--------------------------------------|
|   |  | Fiber, daily dose, g  | Prebiotic | Comparator; daily dose, g      | Compliance <sup>3</sup> | Design    | Duration, d      | Run in | Washout | Analysis                             |
| Abell 2008 (81)                               | 46; 25–66; 65                                    | RS, 22  | C         | RS, 1                          | Y                       | Crossover | 28               | Y      | Y       | qPCR                                 |
| Alfa 2017 (77)                                | 84; 32–96; 42                                    | RS2, 21   | C         | Corn starch, 21                | Y                       | Parallel  | 72               | Y      | N       | Illumina                             |
| Alles 1999 (52)                               | 27.4; 40.4; 45                                   | TOS, 15   | A         | Glucose & lactose mix, 15      | Y                       | Parallel  | 21               | Y      | N       | Culture                              |
| Baer 2014 (83)                                | 14; 47; 9  | Resistant maltodextrin, 50  | C         | Maltodextrin, 50               | Y                       | Crossover | 21               | N      | Y       | 454 pyrosequencing; DGGE; FISH; qPCR |
| Beards 2010 (53)                              | 30; 33; <sup>4</sup> 66 <sup>4</sup>             | PDX; RS, 45.6   | C         | Maltitol, 45.6                 | N                       | Parallel  | 44               | N      | N       | FISH                                 |
| Blädel 2016 (115)                             | 21; 23–45; 100                                   | Inulin, 15  | A         | Placebo                        | Y                       | Crossover | 21               | N      | Y       | Illumina                             |
| Boler 2011 (84); Hooda 2012 (49) <sup>5</sup> | 21; 21–28; 0                                     | PDX; <sup>6</sup> Soluble maize fiber, 21   | C         | Placebo                        | N                       | Crossover | 21               | N      | N       | qPCR; pyrosequencing <sup>4</sup>    |
| Bouhnik 1996 (54)                             | 10; 22–39; 50                                    | SC-FOS, 12.5  | A         | Saccharose, 10                 | N                       | Parallel  | 12               | Y      | Y       | Culture                              |
| Bouhnik 1999 (58)                             | 8; 29.6; 55                                      | SC-FOS, 20  | A         | Saccharose, 20                 | N                       | Parallel  | 7                | N      | N       | Culture                              |
| Bouhnik 2004 (57)                             | 64; 30; <sup>4</sup> 55 <sup>4</sup>             | SC-FOS; <sup>6</sup> GOS; <sup>6</sup> Isomalto-OS; Inulin; <sup>6</sup> RS; Soybean-OS, 10 | A         | Sucrose & maltodextrin mix, 10 | N                       | Parallel  | 7                | Y      | N       | Culture                              |
| Bouhnik 2006 (56)                             | 40; 29; 55                                       | SC-FOS (Actilight), 10  | A         | Sucrose & maltodextrin mix, 10 | N                       | Parallel  | 7                | Y      | N       | Culture                              |
| Bouhnik 2007 (55)                             | 39; 33.9; NR                                     | Inulin, 5   | A         | Sucrose & maltodextrin mix, 5  | N                       | Parallel  | 28               | Y      | Y       | Culture                              |
| Brahe 2015 (59)                               | 35; 59.6; <sup>4</sup> 100                       | Flaxseed mucilage, 10   | G         | Placebo                        | Y                       | Parallel  | 42               | N      | N       | Quantitative metagenomics            |
| Calame 2008 (60)                              | 16; 30.9; NR                                     | Arabic gum, 40  | G         | Placebo                        | Y                       | Parallel  | 28               | N      | N       | qPCR                                 |
| Clarke 2016 (86)                              | 30; 27; 57                                       | Beta 2-1 fructan, 15  | A         | Maltodextrin, 15               | Y                       | Crossover | 28               | N      | Y       | qPCR                                 |
| Cloetens 2010 (87)                            | 20; 24; 70                                       | AXOS, 10  | C         | Maltodextrin, 20               | N                       | Crossover | 21               | N      | Y       | qPCR                                 |
| Costabile 2010 (90)                           | 31; 25; 56                                       | Very long chain inulin, 10  | A         | Maltodextrin, 10               | N                       | Crossover | 21               | N      | Y       | FISH                                 |
| Costabile 2012 (88)                           | 31; 33; 52                                       | PDX, 8  | C         | Maltodextrin, 8                | N                       | Crossover | 21               | N      | Y       | DGGE; FISH                           |
| Damen 2012 (91)                               | 27; 25; 63                                       | AXOS, 2.14  | C         | Placebo                        | Y                       | Crossover | 21               | Y      | Y       | FISH                                 |
| Depeint 2008 (92)                             | 30; 36.3; 60                                     | Beta-GOS, 7   | A         | Sucrose, 7                     | N                       | Crossover | 7                | Y      | Y       | FISH                                 |
| Dewulf 2013 (61)                              | 30; 47.5; 100                                    | Inulin-type fructan (Synergy 1), 16   | A         | Maltodextrin, 16               | N                       | Parallel  | Reported as 3 mo | N      | N       | qPCR; phylogenetic microarray        |
| Elison 2016 (62)                              | 40; 22–57; 52                                    | HMO; <sup>7</sup> 2'FL; LNnT; mixture (2:1 mixture of 2'FL + LNnT), 20                      | A         | Glucose, 2                     | Y                       | Parallel  | 14               | Y      | N       | Illumina                             |
| Fasting 2008 (63)                             | 25; 26.7; 50                                     | Resistant maltodextrin, 15  | C         | Maltodextrin, 15               | N                       | Parallel  | 21               | Y      | Y       | qPCR                                 |
| Fernando 2010 (93)                            | 12; 25.6; 42                                     | Raffinose, 5  | G         | Placebo                        | N                       | Crossover | 21               | N      | N       | qPCR; T-RFLP                         |
| Finegold 2014 (64)                            | 16; 21–49; <sup>4</sup> 66 <sup>4</sup>          | XOS, 2.8  | C         | Maltodextrin, 2.8              | N                       | Parallel  | 56               | Y      | Y       | Pyrosequencing                       |
| François 2012 (94)                            | 52; 42; 48                                       | Wheat bran extract, 10  | G         | Placebo                        | N                       | Crossover | 21               | Y      | Y       | FISH                                 |

(Continued)

(Continued)

TABLE 2 (Continued)

| Study (reference)                                       | Participants<br><i>n</i> ; age; <sup>2</sup> % F | Interventions   |           |                           | Compliance <sup>3</sup> | RCT design |             |        |         |                |
|---|--|---|-----------|---------------------------|-------------------------|------------|-------------|--------|---------|----------------|
|   |  | Fiber, daily dose, g  | Prebiotic | Comparator, daily dose, g |                         | Design     | Duration, d | Run in | Washout | Analysis       |
| Fuller 2007 (95); Ramirez-Farias 2009 (50) <sup>5</sup> | 12; 38.1; 75                                     | Inulin, 10  | A         | Nil                       | Y                       | Crossover  | 16          | N      | N       | qPCR           |
| Gopal 2003 (65)   | 19; 20–60; <sup>4</sup> 44 <sup>4</sup>          | GOS, 2.4  | A         | Placebo                   | Y                       | Parallel   | 28          | Y      | Y       | Culture        |
| Holscher 2015 (97)                                      | 29; 27; 52                                       | Agave inulin, 7.5   | A         | Placebo                   | N                       | Crossover  | 21          | Y      | Y       | Illumina       |
| Jie 2000 (66)   | 30; 29.9; 45                                     | PDX, 12   | C         | Nil                       | N                       | Parallel   | 28          | Y      | N       | Culture        |
| Kleessen 2007 (67)                                      | 45; 23.5; 55                                     | Inulin; <sup>7</sup> Chicory inulin; Jerusalem artichoke inulin, 15.4 | A         | Placebo                   | N                       | Parallel   | 21          | Y      | N       | Culture; FISH  |
| Lecerf 2012 (68)  | 59; 20.1; 57                                     | XOS; <sup>6</sup> Inulin-XOS mix, 6.64                                | C         | Wheat dextrin, 6.64       | N                       | Parallel   | 28          | N      | N       | qPCR           |
| Lin 2016 (69)   | 20; 24.2; 80                                     | XOS, 1.2  | C         | Placebo                   | N                       | Parallel   | 42          | Y      | Y       | Culture        |
| Lomax 2012 (70)   | 43; 55; 74                                       | Beta 2-1 fructan, 8   | A         | Maltodextrin, 8           | Y                       | Parallel   | 28          | Y      | N       | FISH           |
| Maki 2012 (99)  | 55; 35.1; <sup>4</sup> 54 <sup>4</sup>           | AXOS, 2.4   | C         | Placebo                   | N                       | Crossover  | 21          | N      | Y       | FISH           |
| Maneerat 2013 (100)                                     | 35; 67.4; <sup>4</sup> 53 <sup>4</sup>           | GOS, 8  | A         | Maltodextrin, 8           | N                       | Crossover  | 21          | N      | Y       | FISH           |
| Martínez 2010 (101)                                     | 10; 23–38; 50                                    | RS; <sup>7</sup> RS2; RS4, 33.2                                       | C         | Native wheat starch, 33.2 | N                       | Crossover  | 21          | Y      | Y       | Pyrosequencing |
| Pallav 2014 (72)  | 14; 31.4; <sup>4</sup> 65                        | Polysaccharidepeptide (I'm-Yunity), 3.6                               | G         | Nil                       | N                       | Parallel   | 14          | N      | N       | Pyrosequencing |
| Pasman 2006 (73)  | 29; 34.1; 0                                      | Nutrisse FB (dextrin), 45   | A         | Maltodextrin, 22.5        | Y                       | Parallel   | 35          | Y      | N       | Culture        |
| Petry 2012 (102)  | 32; 18–40; 100                                   | Inulin, 20  | A         | Maltodextrin, 20          | N                       | Crossover  | 28          | N      | Y       | qPCR           |
| Rammani 2010 (74)                                       | 66; 32.9; 50                                     | Inulin, 5   | A         | Placebo                   | Y                       | Parallel   | 21          | Y      | Y       | FISH           |
| Rammani 2015 (103)                                      | 38; 35.1; <sup>4</sup> 50                        | Agave inulin, 5   | A         | Maltodextrin, 5           | Y                       | Crossover  | 21          | Y      | Y       | FISH           |
| Salden 2017 (80)  | 27; 48; 48                                       | Arabinoxylans, 15   | G         | Maltodextrin, 15          | Y                       | Parallel   | 42          | N      | N       | Illumina       |
| Slavin 2011 (105)                                       | 10; 27–49; <sup>4</sup> 0                        | Chicory inulin, 20  | A         | Placebo                   | Y                       | Crossover  | 21          | N      | N       | Culture        |
| Ten Bruggencate 2006 (107)                              | 29; 22.7; 0                                      | FOS, 20   | A         | Sucrose, 6                | Y                       | Crossover  | 14          | N      | Y       | qPCR           |
| Tuohy 2001 (108)  | NR; NR; 55                                       | Mix (FOS & PHGG), 10  | Mix       | Placebo                   | Y                       | Crossover  | 21          | N      | N       | FISH           |
| Vulevic 2008 (109)                                      | 41; 69.3; <sup>4</sup> 64 <sup>4</sup>           | GOS (Bimuno), 5.5   | A         | Maltodextrin, 5.5         | Y                       | Crossover  | 70          | N      | Y       | FISH           |
| Vulevic 2015 (110)                                      | 40; 70.4; 62                                     | GOS (Bimuno), 5.5   | A         | Maltodextrin, 5.5         | Y                       | Crossover  | 70          | N      | Y       | FISH           |
| Walton 2010 (113)                                       | 31; 21; 58                                       | MOS, 5  | C         | Placebo                   | Y                       | Crossover  | 21          | N      | Y       | FISH           |
| Walton 2012 (111)                                       | 37; 58.9; <sup>4</sup> 57 <sup>4</sup>           | GOS, 8  | A         | Placebo                   | N                       | Crossover  | 21          | Y      | Y       | qPCR           |
| Walton 2012 (112)                                       | 40; 31.4; <sup>4</sup> 60 <sup>4</sup>           | AXOS, 2.2   | C         | Placebo                   | Y                       | Crossover  | 21          | Y      | Y       | FISH           |
| Wu 2011 (76)  | 15; 40.6; 93                                     | Konjac glucomannan, 4.5   | G         | Nil                       | N                       | Parallel   | 28          | N      | N       | FISH           |

<sup>1</sup>A, accepted prebiotic fiber; AXOS, arabinoxylan-oligosaccharide; C, candidate prebiotic fiber; DGGE, denaturing gradient gel electrophoresis; FISH, fluorescent in situ hybridization; FOS, fructo-oligosaccharide; G, general fiber; GOS, galacto-oligosaccharide; HMO, human milk oligosaccharide; LNNt, lacto-N-neotetraose; MOS, manno-oligosaccharide; NR, not reported by study; OS, oligosaccharide; PDX, polydextrose; PHGG, partially hydrolyzed guar gum; qPCR, quantitative polymerase chain reaction; RS, resistant starch; RS2, resistant starch 2; RS4, resistant starch 4; SC-FOS, short-chain fructo-oligosaccharide; TOS, trans-galacto-oligosaccharide; T-RFLP, Terminal restriction fragment length polymorphism; XOS, xylo-oligosaccharide; 2FL, 2'-O-fucosyllactose.

<sup>2</sup>Age expressed as mean y; age range provided when means were not obtainable.

<sup>3</sup>Compliance to intervention; assessed by primary study.

<sup>4</sup>Refers to randomized population rather than actual population.

<sup>5</sup>Secondary publication reporting additional outcomes from the primary study.

<sup>6</sup>Refers to analyzed intervention arm with the highest prebiotic classification (accepted prebiotic fiber > candidate prebiotic fiber > general fiber) selected for fiber type subgroup analysis.

<sup>7</sup>Refers to intervention fibers that have been pooled together for meta-analyses.



**TABLE 3**  
 Characteristics of randomized controlled trials of food interventions comparing dietary fiber with low-fiber comparators in healthy adults<sup>1</sup>

| Interventions                                     |  |                             |                       |                              |                            |                         |            |             |        |         |                    |
|---|--|-----------------------------|-----------------------|------------------------------|----------------------------|-------------------------|------------|-------------|--------|---------|--------------------|
| Study (reference)                                 | Participants<br><i>n</i> ; age; <sup>2</sup> % F | Intervention                | Comparator            | Daily fiber<br>difference, g | Study<br>diet <sup>3</sup> | Compliance <sup>4</sup> | RCT design |             |        |         |                    |
|   |  |                             |                       |                              |                            |                         | Design     | Duration, d | Run in | Washout | Analysis           |
| Ampatzoglou 2015 (82)                             | 33; 48.8; 64                                     | WG diet                     | RG diet               | 10                           | N                          | Y                       | Crossover  | 14          | Y      | Y       | FISH               |
| Carvalho-Wells 2010 (85)                          | 32; 31.6; 66                                     | WG cereal                   | Non-WG cereal         | 6.5                          | N                          | N                       | Crossover  | 21          | Y      | Y       | FISH               |
| Cooper 2017 (78)                                  | 46; 25.8; 46                                     | WG market basket            | RG market basket      | 5                            | N                          | Y                       | Parallel   | 42          | N      | N       | Illumina           |
| Costabile 2008 (89)                               | 31; 25; 52                                       | WG cereal                   | Wheat bran cereal     | 7.4                          | N                          | N                       | Crossover  | 21          | Y      | Y       | FISH               |
| Gråsten 2007 (96)                                 | 14; 59.7; <sup>5</sup> 100                       | Rye bread                   | White wheat bread     | 19                           | N                          | Y                       | Crossover  | 56          | Y      | Y       | Culture            |
| Jenkins 1999 (98)                                 | 24; 33; 50                                       | Wheat bran                  | Wheat flour           | 19                           | N                          | Y                       | Crossover  | 14          | N      | Y       | Culture            |
| Karl 2017 (79);<br>Vanegas 2017 (51) <sup>6</sup> | 81; 40–65; <sup>5</sup> 60                       | WG diet                     | RG diet               | 8                            | Y                          | Y                       | Parallel   | 42          | Y      | N       | Illumina           |
| Nemoto 2011 (71)                                  | 36; 22–67; 63                                    | Fermented brown rice        | “Non-functional food” | 4.62                         | N                          | Y                       | Parallel   | 14          | N      | N       | Culture            |
| Ross 2011 (104)                                   | 17; 35; 65                                       | WG diet                     | RG diet               | 13                           | Y                          | Y                       | Crossover  | 14          | Y      | Y       | qPCR               |
| Smith 2006 (106)                                  | 18; 42.8; 0                                      | Lupin kernel fiber diet     | Control diet          | 22                           | Y                          | N                       | Crossover  | 28          | N      | Y       | FISH               |
| Tap 2015 (75)                                     | 19; 19–25; 53                                    | High-fiber diet             | Low-fiber diet        | 30                           | Y                          | Y                       | Crossover  | 5           | N      | Y       | 454 pyrosequencing |
| Zeng 2015 (114)                                   | 77; 63.4; 70                                     | Whole cereal legume<br>diet | Control diet          | 14.5                         | Y                          | Y                       | Parallel   | 90          | N      | N       | Culture            |

<sup>1</sup>FISH, fluorescent in situ hybridization; qPCR, quantitative polymerase chain reaction; RG, refined grain; WG, whole grain.


<sup>2</sup>Age expressed as mean y; age range provided when means were not obtainable.

<sup>3</sup>Whether the participant's entire diet was provided by the study.


<sup>4</sup>Compliance to intervention; assessed by primary study.

<sup>5</sup>Refers to randomized population rather than actual population.

<sup>6</sup>Secondary publication reporting additional outcomes from the primary study.



**FIGURE 2** Forest plot of randomized controlled trials in healthy adults comparing dietary fiber with placebo/low-fiber comparators. Studies are subgrouped by fiber type, with the overall effect included at the bottom. Data are presented as means and SDs of *Bifidobacterium* spp. abundance at end of intervention. Effects of trials are presented as weights (percentages) and standardized mean differences (95% CIs). IV, inverse variance; Std., standardized.



**FIGURE 3** Forest plot of randomized controlled trials in healthy adults comparing dietary fiber with placebo/low-fiber comparators. Studies are subgrouped by fiber type, with the overall effect included at the bottom. Data are presented as means and SDs of *Lactobacillus* spp. abundance at end of intervention. Effects of trials are presented as weights (percentages) and standardized mean differences (95% CIs). IV, inverse variance; Std., standardized.

were reported in a form suitable for meta-analysis (52, 55, 56, 60, 63–68, 73, 75, 76, 84, 87, 93, 96, 97, 99, 104, 105, 107, 111, 114). Dietary fiber led to a significantly greater *Lactobacillus* spp. abundance compared with placebo/low-fiber comparators (SMD: 0.37; 95% CI: 0.07, 0.68;  $P = 0.02$ ). However, heterogeneity was considerable ( $I^2 = 80\%$ ), and was skewed by results from a single outlier study (66) (4.70; 95% CI: 3.69, 5.70). A sensitivity analysis excluding this study produced a more homogeneous study population ( $I^2 = 49\%$ ), with a modest impact on the result (SMD: 0.22; 95% CI: 0.03, 0.41;  $P = 0.02$ ) (Figure 3). The outlier study (66) was excluded from subsequent subgroup analyses.


Subgroup analysis demonstrated interventions involving fiber supplements resulted in a significantly higher *Lactobacillus* spp. abundance compared with placebo/low-fiber controls while substantially reducing study heterogeneity (SMD: 0.16; 95% CI: 0.01, 0.31;  $P = 0.04$ ,  $I^2 = 7\%$ ). No significant differences in effect were found between food interventions and comparators (SMD: 0.35; 95% CI: -0.46, 1.16;  $P = 0.40$ ,  $I^2 = 84\%$ ).

Subgroup analysis of fiber types showed accepted prebiotic fiber interventions led to a significantly greater *Lactobacillus*

spp. abundance compared with placebo/low-fiber controls and further reduced heterogeneity (SMD: 0.34; 95% CI: 0.13, 0.55;  $P = 0.002$ ,  $I^2 = 0\%$ ) (Figure 3). There were no differences in effect in the candidate prebiotic (SMD: -0.06; 95% CI: -0.29, 0.16;  $P = 0.58$ ,  $I^2 = 0\%$ ) and general fiber (SMD: 0.22; 95% CI: -0.31, 0.75;  $P = 0.42$ ,  $I^2 = 74\%$ ) subgroups when compared with comparators.

Subgroup analysis of analysis method demonstrated dietary fiber led to significantly higher *Lactobacillus* spp. abundance compared with placebo/low-fiber comparators when enumerated via culture (SMD: 0.61; 95% CI: 0.13, 1.08;  $P = 0.01$ ). There were no significant differences between intervention and comparator when *Lactobacillus* spp. was detected with the use of FISH, qPCR, or sequencing (Supplemental Table 6). There were no differences in effect when sub-analyzing by intervention type or dose-response (Supplemental Table 6).

There were 4 studies that could not be pooled into the meta-analysis. A prebiotic supplementation trial of HMOs reported no difference in *Lactobacillus* spp. abundance between intervention and control groups (62). There was also no significant



**FIGURE 4** Forest plot of randomized controlled trials in healthy adults comparing dietary fiber with placebo/low-fiber comparators. Data are presented as means and SDs of *Faecalibacterium prausnitzii*, *Roseburia* spp., *Eubacterium rectale*, and *Ruminococcus bromii* abundance at end of intervention. Effects of trials are presented as weights (percentages) and standardized mean differences (95% CIs). IV, inverse variance; Std., standardized.

difference in *Lactobacillus* spp. reported in a whole-grain food intervention study compared with controls (78). Of the 2 remaining studies, there was higher *Lactobacillus* spp. abundance after xylo-oligosaccharide supplementation compared with placebo (69), and significant within-group increases in *Lactobacillus* spp. abundance were demonstrated after manno-oligosaccharide supplementation (113).

Abundance of *F. prausnitzii* was measured in 15 studies investigating 566 participants. Thirteen studies (519 participants) were able to be meta-analyzed (53, 61, 67, 68, 74, 84, 88, 94,

99–101, 110, 112). There was no difference between dietary fiber and placebo/low-fiber comparators for *F. prausnitzii* abundance (SMD: 0.14; 95% CI: -0.12, 0.39;  $P = 0.29$ ), with substantial heterogeneity between studies ( $I^2 = 68\%$ ) (Figure 4). Aside from trial design, no differences with respect to the prespecified subgroups were found (Supplemental Table 6). Two studies reporting abundances of *F. prausnitzii* were unable to be pooled into the meta-analysis. Both studies measured the relative abundance of *F. prausnitzii* and reported only within-group changes, with 1 study reporting a decrease in abundance after supplementation

of flaxseed mucilage (59) and the other reporting an increase in abundance after inulin supplementation (50).

Seven studies including 261 participants measured *Roseburia* spp. abundance. Four studies (189 participants) were included in the meta-analysis (49, 68, 79, 97). Dietary fiber had no effect on *Roseburia* spp. abundance compared with placebo/low-fiber comparators (SMD: 0.33; 95% CI: -0.14, 0.80;  $P = 0.17$ ) although substantial heterogeneity was detected ( $I^2 = 70\%$ ) (Figure 4). Similar results were reported in the studies excluded from meta-analysis. No between- or within-group differences were detected between intervention and placebo groups in 2 prebiotic fiber supplement trials (50, 62). A third trial found the relative abundance of *Roseburia* spp. was lower after inulin supplementation compared with control at end of intervention, although significance was not reported (115).

Two studies of 32 participants measured *E. hallii* abundance. These results could not be statistically pooled because 1 study did not report data in a suitable form. One study reported no within-group difference in *E. hallii* abundance (50, 62), the other reported a significant decrease in *E. hallii* abundance compared with placebo (49).

*E. rectale* was measured in 3 studies including 42 participants. Two studies (30 participants) were suitable for meta-analysis (84, 101). Dietary fiber did not affect *E. rectale* abundance compared with placebo/low-fiber comparators (SMD: -0.26; 95% CI: -1.20, 0.67;  $P = 0.58$ ) and substantial heterogeneity was detected ( $I^2 = 75\%$ ) (Figure 4). The study not eligible for meta-analysis was an inulin supplementation trial which reported no difference for within-group effects for *E. rectale* abundance (50).

*R. bromii* abundance was measured in 3 studies encompassing 76 participants, of which all were suitable for meta-analysis (49, 81, 101). Dietary fiber had no effect on *R. bromii* abundance compared with placebo/low-fiber comparators (SMD: 0.15; 95% CI: -0.15, 0.45;  $P = 0.33$ ), with no heterogeneity detected ( $I^2 = 0\%$ ) (Figure 4).

### Dietary fiber and SCFAs

A total of 25 studies involving 870 participants reported between-group differences in fecal SCFA concentrations after fiber intervention (52, 53, 55, 59, 63, 64, 66–68, 71, 73, 74, 77, 80, 82, 84, 86, 90, 91, 93, 94, 96, 103, 112, 115). Fecal SCFA concentrations were determined through gas-liquid chromatography in all but 1 study (90) in which HPLC was used.

Total fecal SCFA concentration was measured in 13 studies encompassing 406 participants (52, 55, 59, 63, 64, 67, 73, 80, 82, 84, 86, 91, 94). Dietary fiber had no effect on total SCFA concentration compared with placebo/low-fiber comparators (SMD: 0.11; 95% CI: -0.05, 0.27;  $P = 0.19$ ), with similar intervention effects across studies ( $I^2 = 0\%$ ).

Fecal acetate concentration was reported in 18 studies involving 657 participants (52, 53, 63, 66, 71, 74, 77, 80, 82, 84, 86, 90, 91, 93, 94, 96, 103, 112). There was no difference in fecal acetate after fiber intervention compared with placebo/low-fiber comparators (SMD: 0.28; 95% CI: -0.08, 0.63;  $P = 0.13$ ), with substantial heterogeneity between studies ( $I^2 = 86\%$ ).

The effect of fiber intervention on fecal propionate concentration was reported in 19 studies of 677 participants (52, 53, 63, 66, 71, 74, 77, 80, 82, 84, 86, 90, 91, 93, 94, 96, 103, 112, 115). No differences were found between fecal propionate and

comparators (SMD: -0.01; 95% CI: -0.20, 0.22;  $P = 0.95$ ), with moderate heterogeneity detected ( $I^2 = 61\%$ ).

The effect of fiber intervention on fecal butyrate concentration was reported in 20 studies of 712 participants (52, 53, 59, 63, 66, 71, 74, 77, 80, 82, 84, 86, 90, 91, 93, 94, 96, 103, 112, 115). Fecal butyrate was significantly higher after fiber intervention compared with placebo/low-fiber comparators (SMD: 0.24; 95% CI: 0.00, 0.47;  $P = 0.05$ ), although considerable heterogeneity was present ( $I^2 = 70\%$ ).

Of the studies evaluating differences in fecal SCFA concentrations after fiber intervention compared with placebo/low-fiber comparators, 13 studies expressed mean SCFA concentrations per wet weight of feces (52, 53, 66, 67, 71, 73, 74, 77, 82, 90, 91, 96, 115), 10 studies as dry weight of feces (55, 59, 63, 64, 68, 80, 93, 94, 103, 112), 1 study as molar ratio (84), and 1 study as a combination of wet weight of feces and molar ratio (86). Additional subgroup analyses were performed to compare differences in fecal SCFA concentrations when expressed as wet weight compared with dry weight (Supplemental Table 7). Fiber intervention led to significantly higher fecal concentrations of total SCFA, acetate, and butyrate compared with comparators when expressed per wet weight of feces. However, there were no significant differences when mean SCFA concentrations were expressed per dry weight of feces. Study heterogeneity was considerably greater for fecal acetate and butyrate, but not total fecal SCFA concentration when expressed as wet compared with dry weight of feces. There were no differences in effect based on analysis method for fecal propionate concentration, although heterogeneity was greater when results were expressed per wet weight of feces (Supplemental Table 7).

### Differences in intervention effects based on trial design

There were differences in intervention effects in subgroup analyses depending upon trial design. Dietary fiber led to significantly lower  $\alpha$ -diversity compared with placebo/low-fiber comparators in crossover design trials, in which  $\alpha$ -diversity was reported with the use of Shannon diversity index (MD: -0.10; 95% CI: -0.19, -0.01;  $P = 0.03$ ), whereas there was no difference in  $\alpha$ -diversity in parallel design trials (MD: -0.03; 95% CI: -0.57, 0.51;  $P = 0.91$ ) (Supplemental Table 6). The presence and duration of washout periods were inconsistent across the 3 crossover trials included in this analysis. One study did not include a washout period (84), and washout periods lasted 14 (75) and 21 d (88) in the other 2. Regarding bacterial abundances however, intervention effects were significant in parallel trials but not in crossover trials for *Lactobacillus* and *Roseburia* spp. and *F. prausnitzii*, but not for *Bifidobacterium* spp. (Supplemental Table 6). Statistical heterogeneity was lower in crossover trials compared with parallel trials for  $\alpha$ -diversity reported with the use of Shannon diversity index, *Bifidobacterium* and *Lactobacillus* spp., as well as *F. prausnitzii*, but there was no difference in statistical heterogeneity for *Roseburia* spp. (Supplemental Table 6).

### Risk of bias

The risk of bias was low-to-moderate across the 64 included studies (Supplemental Figure 1). Selection bias was unclear in most studies. Random sequence generation and allocation



concealment were adequately described by 26% (59–62, 70–72, 77, 79, 80, 84, 86, 94, 103, 113–115) and 16% (59, 61, 62, 70, 77, 79, 80, 86, 94, 115) of studies, respectively. There was low risk of bias across included studies regarding performance and detection bias, as most trials investigated objective outcomes and incorporated a double-blind design. Attrition bias was adequately addressed by only 41% (54–58, 62, 67, 69, 71, 74–76, 79, 82, 86–89, 92, 93, 98, 99, 105, 107, 108, 110) of the included studies. Selective reporting was unclear in the majority of studies. Published protocols or clinical registrations were reported by only 26% (59, 61, 68–70, 75, 77–80, 86, 97, 100–102, 110, 115) of included studies. Bias related to control of dietary intake was unclear in half of included studies (55%) (54, 56–60, 62, 64–67, 71, 72, 74, 78, 80, 81, 83, 85–93, 96, 98, 102, 103, 105, 108, 110, 115); even fewer studies were judged to have a low risk of bias regarding dietary advice and assessment of dietary compliance (33%) (52, 55, 63, 68, 69, 73, 75, 76, 79, 82, 84, 94, 97, 99, 104, 106, 107, 111–114). Furthermore, 13% (53, 61, 70, 77, 95, 100, 101, 109) of studies did not provide dietary advice or assess intake, and were judged to have a high risk of bias relating to the potential influence of background dietary intake.

### Reporting bias

Funnel plots were generated for abundances of *Bifidobacterium* spp., *Lactobacillus* spp., *F. prausnitzii*, and total SCFA; and for acetate, propionate, and butyrate concentrations. Visual inspection found no evidence of funnel plot asymmetry, indicating reporting bias was unlikely (Supplemental Figures 2–7).

### DISCUSSION

This systematic review and meta-analysis found dietary fiber intervention had no effect on the diversity of the gut microbiota but did increase abundance of *Bifidobacterium* and *Lactobacillus* spp. as well as fecal butyrate concentration in healthy adults.

The lack of effect on  $\alpha$ -diversity of the gut microbiota found in this review is similar to other dietary interventions documented in the literature. For instance, controlled feeding studies lasting 4 d to 3 wk found that despite significant changes to fiber intake, there was no effect on microbial diversity (35–37). These findings suggest that short-term dietary interventions are unlikely to facilitate changes in the  $\alpha$ -diversity of the gut microbiota. Indeed, study design is likely important, as subgroup analysis demonstrated different effects between crossover and parallel trials. The lower  $\alpha$ -diversity between fiber and control groups in crossover trials may be related to a lack of or insufficient washout between interventions, as well as potential differences in the microbiota and habitual diet of individuals at baseline.

These null findings are in contrast to the findings from observational studies that report a correlation between fiber intakes in habitual diet and diversity of the gut microbiota, for example in studies comparing agrarian dietary habits with Western populations (38, 39). Interestingly, a positive correlation has also been reported between dietary diversity and microbiota diversity (116). Taken together, long-term dietary diversity as opposed to changes in isolated nutrients or foods over a short period of time may be a stronger driver of microbial diversity. It must also be noted that the stability of the gut microbiota, as well as the abundances and metabolites of the individual members of the microbial

community, also contribute to maintaining an ecosystem that promotes health (117, 118). Therefore, the totality of findings here, including that microbial diversity was not compromised, support the favorable effects of dietary fiber on the gut microbiota.

In regard to particular bacterial groups, this review demonstrated that dietary fiber interventions involving accepted prebiotic fibers led to higher abundance of *Bifidobacterium* and *Lactobacillus* spp. These results support the selectivity criteria of the prebiotic concept, in which the host microorganisms selectively utilize the prebiotic fibers as substrates, which may confer health benefits to the host (32). However, candidate prebiotic interventions produced different effects on the abundance of these 2 genera, with significant effects demonstrated for *Bifidobacterium* but not *Lactobacillus* spp. This may represent differences in substrate preferences between the 2 genera, in which *Bifidobacterium* spp. may be less discriminating than *Lactobacillus* spp. regarding fermentation substrates (119, 120). Conversely, fibers not classified as accepted or candidate prebiotics, here termed general fibers, did not affect the abundance of these taxa. This may be due to the heterogeneity of the general fibers, including their degree of polymerization, viscosity, and fermentability, whereas accepted and candidate prebiotic fibers are mostly highly fermentable oligosaccharides (29, 30).

Subgroup analysis separating the effects of food and supplement interventions showed food interventions had no effect on *Bifidobacterium* and *Lactobacillus* spp. This result may be attributed to a lack of statistical power, due to the food interventions comprising a relatively small number of low sample size studies (10 studies, 301 participants; and 4 studies, 127 participants, respectively). It must also be noted that most of the trials using food interventions supplemented with grain and cereal foods to increase fiber intake (71, 78, 79, 82, 85, 89, 96, 98, 104). Therefore, the food interventions evaluated may be more representative of grains and cereals per se rather than a diverse range of fibrous foods.

Interestingly, there were no differences in the effect of dietary fiber interventions on *Bifidobacterium* spp. abundance with varying doses of fiber. Dietary fiber intervention led to an effect at all levels of consumption in subgroup analysis (<5 g, 5–10 g, >10 g) with no discernible gradient in effectiveness, suggesting <5 g of dietary fiber is sufficient. This may represent a potential limit to the amount of fiber that can be fermented by *Bifidobacterium* spp. The lack of a dose-response effect may also be attributed to the percentage increase in fiber intake from baseline rather than the intervention dose, which was unable to be accounted for in this review due to the inconsistent reporting of baseline values across included studies. This requires further clarification but lower-dose supplementation may be advantageous in patients who experience GI symptoms with higher fiber loads.

There was more variability in intervention effects for abundances of *Bifidobacterium* spp. ( $I^2 = 85\%$ ) compared with *Lactobacillus* spp. ( $I^2 = 49\%$ ). Although this may be related to differences in the accuracy of techniques used to determine specific bacterial abundances (121, 122), there were no differences in effect based on analysis method for *Bifidobacterium* spp. Another plausible explanation is the differences in nutrient requirements of these taxa as discussed previously. Furthermore, “responder and nonresponder” effects for *Bifidobacterium* spp. abundance, which have been shown previously (97, 123, 124), may be affected by individual host factors, such as differences in baseline

abundances (124), or the presence/absence of specific strains of *Bifidobacterium* able to utilize the particular fiber under investigation.

There were differences in intervention effects based on trial design, with parallel design studies demonstrating stronger intervention effects and greater statistical heterogeneity compared with crossover design studies for several outcomes. This may in part be due to inter-individual differences in microbiota composition as well as carry-over effects from a lack of or insufficient washout periods in the crossover studies as discussed previously.

There was no effect of dietary fiber interventions on abundance of other commonly measured bacterial groups (e.g., *F. prausnitzii*), suggesting that these species may be stimulated by dietary components other than fiber, such as polyols and polyphenols (125). However, the number of studies evaluating species of other bacterial groups was small, and therefore further studies are needed to investigate the effect of fiber and other dietary components on these groups.

The higher fecal concentration of butyrate after fiber intervention highlights the ability of dietary fiber to beneficially modulate the metabolic outputs of the gut microbiota. This is likely due to cross-feeding interactions between butyrate producers and *Bifidobacterium* or *Lactobacillus* species, which are noted lactate and acetate producers (25, 120, 126). As the preferred energy source for colonic epithelial cells, butyrate is a microbial by-product that is of particular relevance to host health, exhibiting a wide spectrum of positive effects, such as inhibiting colonic carcinogenesis and ameliorating mucosal inflammation (31, 127, 128). However, it is acknowledged that the variability in the reporting of SCFA results may limit the applicability of these findings, particularly when considering the variance in results when expressed as wet compared with dry weight of feces.

This study is the first systematic review and meta-analysis to assess the effect of dietary fiber intervention on gut microbiota composition. Major strengths of this study include its robust design, comprehensive search strategies, and the use of 2 independent reviewers.

It is acknowledged this study has some limitations. Firstly, there were only a limited number of studies reporting the primary outcome of  $\alpha$ -diversity, and a small proportion presenting data via the same diversity indexes. Secondly, baseline fiber intake was not able to be accounted for due to the paucity of reporting by included studies. Furthermore, included studies sampled feces as a surrogate for gut microbiota profile, and although feces are a common sampling route, the microbial composition of feces differs from the mucosal microbiota (10, 11), which is in closer contact with the host and may be more important when considering the relation between microbiota and disease pathophysiology or outcomes. Finally, the limited number of taxa assessed in the review may not convey the overall effect elicited by dietary fiber intervention on gut microbiota composition and metabolic outputs, although the selection of taxa was guided by the available literature. Thus, the taxa selected may be more representative of the scope of research in the field to date, rather than a limitation of the review.

Dietary fiber intervention leads to a higher abundance of fecal *Bifidobacterium* and *Lactobacillus* spp., as well as higher fecal concentration of butyrate compared with placebo/low-fiber comparators. Accepted prebiotic fibers had an effect on the abundances of both *Bifidobacterium* and *Lactobacillus* spp. whereas

candidate prebiotic fibers had an effect on *Bifidobacterium* spp. abundance but not *Lactobacillus* spp. General fibers appear to have a limited effect on gut microbiota composition. Although the diversity of the gut microbiota, abundances of other commonly measured bacterial groups, and concentrations of other fecal SCFAs were not significantly different compared with controls after dietary fiber intervention, it is worth noting that a short-term increase in fiber intake does not appear to be rate-limiting to these outcomes. These results further support the favorable effects of dietary fiber and contribute to our understanding of its effect on the gut microbiota.

Future RCTs investigating the effect of fiber on the gut microbiota should adjust for participants' baseline microbiota composition and dietary characteristics as well as controlling for dietary intake in order to determine the precise effect of dietary fiber. Scope may also need to be broadened to evaluate more taxa than those considered here, including the eukaryote (e.g., fungi) members of the gut microbiota. In addition, longer-duration studies are needed to better assess the chronic effect of fiber on microbiota diversity.

We thank David Honeyman for assisting with the development of the search strategy. Many thanks to the authors of included studies who provided outcome data necessary for the extraction of data of the variables included in the meta-analyses.

The authors' responsibilities were as follows—HMS and KLC: initiated the study; DS, KW, HMS, MR, and KLC: developed the protocol; DS and HMS: performed eligibility screening and data extraction; DS and JTK: analyzed the data and performed the statistical analysis; DS, KW, MR, MM, JTK, ERS, HMS, and KLC: interpreted the data; DS: wrote the initial manuscript; KW, MR, MM, GH, JTK, ERS, HMS, and KLC: critically revised the manuscript; and authors: read and approved the final manuscript. None of the authors reported a conflict of interest related to the study.

## REFERENCES

1. Sender R, Fuchs S, Milo R. Revised estimates for the number of human and bacteria cells in the body. *PLoS Biol* 2016;14:e1002533.
2. Marchesi JR, Adams DH, Fava F, Hermes GDA, Hirschfield GM, Hold G, Quraishi MN, Kinross J, Smidt H, Tuohy KM et al. The gut microbiota and host health: a new clinical frontier. *Gut* 2016;65:330–9.
3. Scott KP, Duncan SH, Flint HJ. Dietary fibre and the gut microbiota. *Nutr Bull* 2008;33:201–11.
4. LeBlanc JG, Milani C, de Giori GS, Sesma F, van Sinderen D, Ventura M. Bacteria as vitamin suppliers to their host: a gut microbiota perspective. *Curr Opin Biotechnol* 2013;24:160–8.
5. Round JL, Mazmanian SK. The gut microbiota shapes intestinal immune responses during health and disease. *Nat Rev Immunol* 2009;9:313–23.
6. Sekirov I, Russell SL, Antunes LC, Finlay BB. Gut microbiota in health and disease. *Physiol Rev* 2010;90:859–904.
7. Clemente JC, Ursell LK, Parfrey LW, Knight R. The impact of the gut microbiota on human health: an integrative view. *Cell* 2012;148:1258–70.
8. de Vos WM, de Vos EA. Role of the intestinal microbiome in health and disease: from correlation to causation. *Nutr Rev* 2012;70(Suppl 1):S45–56.
9. Zhernakova A, Kurilshikov A, Bonder MJ, Tigchelaar EF, Schirmer M, Vatanen T, Mujagic Z, Vila AV, Falony G, Vieira-Silva S et al. Population-based metagenomics analysis reveals markers for gut microbiome composition and diversity. *Science (New York, NY)* 2016;352:565–9.
10. Fraher MH, O'Toole PW, Quigley EM. Techniques used to characterize the gut microbiota: a guide for the clinician. *Nat Rev Gastroenterol Hepatol* 2012;9:312–22.
11. Zoetendal EG, von Wright A, Vilpponen-Salmela T, Ben-Amor K, Akkermans AD, de Vos WM. Mucosa-associated bacteria in the

- human gastrointestinal tract are uniformly distributed along the colon and differ from the community recovered from feces. *Appl Environ Microbiol* 2002;68:3401–7.
12. Lozupone CA, Knight R. Species divergence and the measurement of microbial diversity. *FEMS Microbiol Rev* 2008;32:557–78.
  13. Morgan XC, Huttenhower C. Chapter 12: human microbiome analysis. *PLoS Comput Biol* 2012;8:e1002808.
  14. Tojo R, Suarez A, Clemente MG, de los Reyes-Gavilan CG, Margolles A, Gueimonde M, Ruas-Madiedo P. Intestinal microbiota in health and disease: role of bifidobacteria in gut homeostasis. *World J Gastroenterol* 2014;20:15163–76.
  15. Bottacini F, Ventura M, van Sinderen D, O'Connell Motherway M. Diversity, ecology and intestinal function of bifidobacteria. *Microbial Cell Factories* 2014;13(Suppl 1):S4.
  16. Rossi M, Amaretti A, Raimondi S. Folate production by probiotic bacteria. *Nutrients* 2011;3:118–34.
  17. Satokari RM, Vaughan EE, Smidt H, Saarela M, Matto J, de Vos WM. Molecular approaches for the detection and identification of bifidobacteria and lactobacilli in the human gastrointestinal tract. *Syst Appl Microbiol* 2003;26:572–84.
  18. Wells JM. Immunomodulatory mechanisms of lactobacilli. *Microbial Cell Factories* 2011;10(Suppl 1):S17.
  19. Flint HJ, Duncan SH, Louis P. The impact of nutrition on intestinal bacterial communities. *Curr Opin Microbiol* 2017;38:59–65.
  20. Flint HJ, Duncan SH, Scott KP, Louis P. Links between diet, gut microbiota composition and gut metabolism. *Proc Nutr Soc* 2015;74:13–22.
  21. Ze X, Duncan SH, Louis P, Flint HJ. *Ruminococcus bromii* is a keystone species for the degradation of resistant starch in the human colon. *ISME J* 2012;6:1535–43.
  22. Ze X, Le Mougen F, Duncan SH, Louis P, Flint HJ. Some are more equal than others: the role of “keystone” species in the degradation of recalcitrant substrates. *Gut Microbes* 2013;4:236–40.
  23. Scott KP, Antoine J-M, Midtved T, van Hemert S. Manipulating the gut microbiota to maintain health and treat disease. *Microb Ecol Health Dis* 2015;26:25877.
  24. O'Keefe SJ. Diet, microorganisms and their metabolites, and colon cancer. *Nat Rev Gastroenterol Hepatol* 2016;13:691–706.
  25. Rivi re A, Selak M, Lantin D, Leroy F, De Vuyst L. Bifidobacteria and butyrate-producing colon bacteria: importance and strategies for their stimulation in the human gut. *Front Microbiol* 2016;7:979.
  26. FAO/WHO. CODEX Alimentarius (CODEX) Guidelines on Nutrition Labeling CAC/GL 2-1985. 2017. Available from: [http://www.fao.org/fao-who-codexalimentarius/sh-proxy/es/?lnk=1&url=https%253A%252F%252Fworkspace.fao.org%252Fsites%252Fcodex%252FStandards%252FCAC%2BGL%2B2-1985%252FCXG\\_002e.pdf](http://www.fao.org/fao-who-codexalimentarius/sh-proxy/es/?lnk=1&url=https%253A%252F%252Fworkspace.fao.org%252Fsites%252Fcodex%252FStandards%252FCAC%2BGL%2B2-1985%252FCXG_002e.pdf)
  27. Federalregister.gov [Internet]. Food labeling: revision of the nutrition and supplement facts labels (21 CFR 101) [Internet]. c.May 2016 [cited 2018 Jan 10]. Washington (DC): Food and Drug Administration, Health and Human Services. Federal Register. Available from: <https://www.federalregister.gov/documents/2016/05/27/2016-11867/food-labelingrevision-of-the-nutrition-and-supplement-facts-labels>.
  28. Eswaran S, Muir J, Chey WD. Fiber and functional gastrointestinal disorders. *Am J Gastroenterol* 2013;108:718–27.
  29. Holscher HD. Dietary fiber and prebiotics and the gastrointestinal microbiota. *Gut Microbes* 2017;8:172–84.
  30. McRorie JW Jr, McKeown NM. Understanding the physics of functional fibers in the gastrointestinal tract: an evidence-based approach to resolving enduring misconceptions about insoluble and soluble fiber. *J Acad Nutr Diet* 2017;117:251–64.
  31. Wong JM, de Souza R, Kendall CW, Emam A, Jenkins DJ. Colonic health: fermentation and short chain fatty acids. *J Clin Gastroenterol* 2006;40:235–43.
  32. Gibson GR, Hutkins R, Sanders ME, Prescott SL, Reimer RA, Salminen SJ, Scott K, Stanton C, Swanson KS, Cani PD et al. Expert consensus document: the International Scientific Association for Probiotics and Prebiotics (ISAPP) consensus statement on the definition and scope of prebiotics. *Nat Rev Gastroenterol Hepatol* 2017;14:491–502.
  33. Gibson GR, Scott KP, Rastall RA, Tuohy KM, Hotchkiss A, Dubert-Ferrandon A, Gareau M, Murphy EF, Saulnier D, Loh G et al. Dietary prebiotics: current status and new definition. *Food Sci Technol Bull: Funct Foods* 2010;7:1–19.
  34. Roberfroid M, Gibson GR, Hoyle L, McCartney AL, Rastall R, Rowland I, Wolvers D, Watzl B, Szajewska H, Stahl B et al. Prebiotic effects: metabolic and health benefits. *Br J Nutr* 2010;104(Suppl 2):S1–63.
  35. David LA, Maurice CF, Carmody RN, Gootenberg DB, Button JE, Wolfe BE, Ling AV, Devlin AS, Varna Y, Fischbach MA et al. Diet rapidly and reproducibly alters the human gut microbiome. *Nature* 2014;505:559–63.
  36. Walker AW, Ince J, Duncan SH, Webster LM, Holtrop G, Ze X, Brown D, Stares MD, Scott P, Bergerat A et al. Dominant and diet-responsive groups of bacteria within the human colonic microbiota. *ISME J* 2011;5:220–30.
  37. Wu GD, Chen J, Hoffmann C, Bittinger K, Chen Y-Y, Keilbaugh SA, Bewtra M, Knights D, Walters WA, Knight R et al. Linking long-term dietary patterns with gut microbial enterotypes. *Science (New York, NY)* 2011;334:105–8.
  38. De Filippo C, Cavalieri D, Di Paola M, Ramazzotti M, Poullet JB, Massart S, Collini S, Pieraccini G, Lionetti P. Impact of diet in shaping gut microbiota revealed by a comparative study in children from Europe and rural Africa. *PNAS* 2010;107:14691–6.
  39. Schnorr SL, Candela M, Rampelli S, Centanni M, Consolandi C, Basaglia G, Turroni S, Biagi E, Peano C, Severgnini M et al. Gut microbiome of the Hadza hunter-gatherers. *Nat Commun* 2014;5:3654.
  40. Falony G, Joossens M, Vieira-Silva S, Wang J, Darzi Y, Faust K, Kurilshikov A, Bonder MJ, Valles-Colomer M, Vandeputte D et al. Population-level analysis of gut microbiome variation. *Science (New York, NY)* 2016;352:560–4.
  41. Moher D, Shamseer L, Clarke M, Ghersi D, Liberati A, Petticrew M, Shekelle P, Stewart LA. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Systematic Reviews* 2015;4:1–9.
  42. Higgins JPT, Green S, editors. *Cochrane handbook for systematic reviews of interventions*. Hoboken, NJ: Wiley, 2011.
  43. Elmagarmid A, Fedorowicz Z, Hammady H, Ilyas I, Khabsa M, Ouzzani M. Rayyan: a systematic reviews web app for exploring and filtering searches for eligible studies for Cochrane Reviews. In: *Evidence-informed public health: opportunities and challenges. Abstracts of the 22nd Cochrane Colloquium*. Hyderabad, India: John Wiley & Sons, 2014.
  44. Higgins JPT, Deeks JJ, Altman DG. Special topics in statistics. In: *Cochrane handbook for systematic reviews of interventions*. Hoboken, NJ: John Wiley & Sons, 2008. p. 481–529.
  45. Higgins JPT, Altman DG. Assessing risk of bias in included studies. In: *Cochrane handbook for systematic reviews of interventions*. Hoboken, NJ: John Wiley & Sons, 2008. p. 187–241.
  46. Higgins JPT, Deeks JJ. Selecting studies and collecting data. In: *Cochrane handbook for systematic reviews of interventions*. Hoboken, NJ: John Wiley & Sons, 2008. p. 151–85.
  47. Elbourne DR, Altman DG, Higgins JP, Curtin F, Worthington HV, Vail A. Meta-analyses involving cross-over trials: methodological issues. *Int J Epidemiol* 2002;31:140–9.
  48. Sterne JAC, Egger M, Moher D. Addressing reporting biases. In: *Cochrane handbook for systematic reviews of interventions*. Hoboken, NJ: John Wiley & Sons, 2008. p. 297–333.
  49. Hooda S, Vester Boler BM, Rossoni Serao MC, Brulc JM, Staeger MA, Boileau TW, Dowd SE, Fahey GC Jr, Swanson KS. 454 pyrosequencing reveals a shift in fecal microbiota of healthy adult men consuming polydextrose or soluble corn fiber. *J Nutr* 2012;142:1259–65.
  50. Ramirez-Farias C, Slezak K, Fuller Z, Duncan A, Holtrop G, Louis P. Effect of inulin on the human gut microbiota: stimulation of *Bifidobacterium adolescentis* and *Faecalibacterium prausnitzii*. *Br J Nutr* 2009;101:541–50.
  51. Vanegas SM, Meydani M, Barnett JB, Goldin B, Kane A, Rasmussen H, Brown C, Vangay P, Knights D, Jonnalagadda S et al. Substituting whole grains for refined grains in a 6-wk randomized trial has a modest effect on gut microbiota and immune and inflammatory markers of healthy adults. *Am J Clin Nutr* 2017;105:635–50.
  52. Alles MS, Hartemink R, Meyboom S, Harryvan JL, Van Laere KJM, Nagengast FM, Hautvast JGAJ. Effect of transgalactooligosaccharides on the composition of the human intestinal microflora and on putative risk markers for colon cancer. *Am J Clin Nutr* 1999;69:980–91.
  53. Beards E, Tuohy K, Gibson G. A human volunteer study to assess the impact of confectionery sweeteners on the gut microbiota composition. *Br J Nutr* 2010;104:701–8.



54. Bouhnik Y, Flourie B, Riottot M, Bisetti N, Gailing MF, Guibert A, Bornet F, Rambaud JC. Effects of fructo-oligosaccharides ingestion on fecal bifidobacteria and selected metabolic indexes of colon carcinogenesis in healthy humans. *Nutr Cancer* 1996;26:21–9.
55. Bouhnik Y, Raskine L, Champion K, Andrieux C, Penven S, Jacobs H, Simoneau G. Prolonged administration of low-dose inulin stimulates the growth of bifidobacteria in humans. *Nutr Res* 2007;27:187–93.
56. Bouhnik Y, Raskine L, Simoneau G, Paineau D, Bornet F. The capacity of short-chain fructo-oligosaccharides to stimulate faecal bifidobacteria: a dose-response relationship study in healthy humans. *Nutr J* 2006;5:8.
57. Bouhnik Y, Raskine L, Simoneau G, Vicaud E, Neut C, Flourie B, Brouns F, Bornet FR. The capacity of nondigestible carbohydrates to stimulate fecal bifidobacteria in healthy humans: a double-blind, randomized, placebo-controlled, parallel-group, dose-response relation study. *Am J Clin Nutr* 2004;80:1658–64.
58. Bouhnik Y, Vahedi K, Achour L, Attar A, Salfati J, Pochart P, Marteau P, Flourie B, Bornet F, Rambaud JC. Short-chain fructo-oligosaccharide administration dose-dependently increases fecal bifidobacteria in healthy humans. *J Nutr* 1999;129:113–16.
59. Brahe L, Chatelier E, Prifti E, Pons N, Kennedy S, Blædel T, Håkansson J, Dalsgaard T, Hansen T, Pedersen O et al. Dietary modulation of the gut microbiota – a randomised controlled trial in obese postmenopausal women. *Br J Nutr* 2015;114:406–17.
60. Calame W, Weseler AR, Viebke C, Flynn C, Siemensma AD. Gum arabic establishes prebiotic functionality in healthy human volunteers in a dose-dependent manner. *Br J Nutr* 2008;100:1269–75.
61. Dewulf EM, Cani PD, Claus SP, Fuentes S, Puylaert PGB, Neyrinck AM, Bindels LB, De Vos WM, Gibson GR, Thissen JP et al. Insight into the prebiotic concept: lessons from an exploratory, double blind intervention study with inulin-type fructans in obese women. *Gut* 2013;62:1112–21.
62. Elison E, Vignasa LK, Rindom Krogsgaard L, Rasmussen J, Sørensen N, McConnell B, Hennen T, Sommer MOA, Bytzer P. Oral supplementation of healthy adults with 2'-O-fucosyllactose and lacto-N-neotetraose is well tolerated and shifts the intestinal microbiota. *Br J Nutr* 2016;116:1356–68.
63. Fastinger ND, Karr-Lilienthal LK, Spears JK, Swanson KS, Zinn KE, Nava GM, Ohkuma K, Kanahori S, Gordon DT, Fahey GC Jr. A novel resistant maltodextrin alters gastrointestinal tolerance factors, fecal characteristics, and fecal microbiota in healthy adult humans. *J Am Coll Nutr* 2008;27:356–66.
64. Finegold S, Li Z, Summanen P, Downes J, Thames G, Corbett K, Dowd S, Krak M, Heber D. Xylooligosaccharide increases bifidobacteria but not lactobacilli in human gut microbiota. *Food Funct* 2014;5:436–45.
65. Gopal PK, Prasad J, Gill HS. Effects of the consumption of *Bifidobacterium lactis* HN019 (DR10™) and galacto-oligosaccharides on the microflora of the gastrointestinal tract in human subjects. *Nutr Res* 2003;23:1313–28.
66. Jie Z, Bang-Yao L, Ming-Jie X, Hai-Wei L, Zu-Kang Z, Ting-Song W, Craig S. Studies on the effects of polydextrose intake on physiologic functions in Chinese people. *Am J Clin Nutr* 2000;72:1503–9.
67. Kleessen B, Schwarz S, Boehm A, Fuhrmann H, Richter A, Henle T, Krueger M. Jerusalem artichoke and chicory inulin in bakery products affect faecal microbiota of healthy volunteers. *Br J Nutr* 2007;98:540–9.
68. Lecerf JM, Dépeint F, Clerc E, Dugenet Y, Niamba CN, Rhazi L, Cayzele A, Abdelnour G, Jaruga A, Younes H et al. Xylooligosaccharide (XOS) in combination with inulin modulates both the intestinal environment and immune status in healthy subjects, while XOS alone only shows prebiotic properties. *Br J Nutr* 2012;108:1847–58.
69. Lin SH, Chou LM, Chien YW, Chang JS, Lin CI. Prebiotic effects of xylooligosaccharides on the improvement of microbiota balance in human subjects. *Gastroenterol Res Pract* 2016;2016:5789232.
70. Lomax AR, Cheung L, Tuohy KM, Noakes PS, Miles EA, Calder PC.  $\beta$ -2-1 fructans have a bifidogenic effect in healthy middle-aged human subjects but do not alter immune responses examined in the absence of an in vivo immune challenge: results from a randomised controlled trial. *Br J Nutr* 2012;108:1818–28.
71. Nemoto H, Ikata K, Arimochi H, Iwasaki T, Ohnishi Y, Kuwahara T, Kataoka K. Effects of fermented brown rice on the intestinal environments in healthy adult. *J Med Invest* 2011;58:235–45.
72. Pallav K, Dowd SE, Villafuerte J, Yang X, Kabbani T, Hansen J, Dennis M, Leffler DA, Kelly CP. Effects of polysaccharopeptide from *Trametes versicolor* and amoxicillin on the gut microbiome of healthy volunteers: a randomized clinical trial. *Gut Microbes* 2014;5:458–67.
73. Pasman W, Wils D, Saniez MH, Kardinaal A. Long-term gastrointestinal tolerance of NUTRIOSE®FB in healthy men. *Eur J Clin Nutr* 2006;60:1024–34.
74. Ramnani P, Gaudier E, Bingham M, Van Bruggen P, Tuohy KM, Gibson GR. Prebiotic effect of fruit and vegetable shots containing Jerusalem artichoke inulin: a human intervention study. *Br J Nutr* 2010;104:233–40.
75. Tap J, Furet JP, Bensaada M, Philippe C, Roth H, Rabot S, Lakhdari O, Lombard V, Henrissat B, Corthier G et al. Gut microbiota richness promotes its stability upon increased dietary fibre intake in healthy adults. *Environ Microbiol* 2015;17:4954–64.
76. Wu WT, Cheng HC, Chen HL. Ameliorative effects of konjac glucomannan on human faecal beta-glucuronidase activity, secondary bile acid levels and faecal water toxicity towards Caco-2 cells. *Br J Nutr* 2011;105:593–600.
77. Alfa MJ, Strang D, Tappia PS, Graham M, Van Domselaar G, Forbes JD, Laminman V, Olson N, DeGagne P, Bray D et al. A randomized trial to determine the impact of a digestion resistant starch composition on the gut microbiome in older and mid-age adults. *Clin Nutr* 2018;37:797–807.
78. Cooper DN, Kable ME, Marco ML, De Leon A, Rust B, Baker JE, Horn W, Burnett D, Keim NL. The effects of moderate whole grain consumption on fasting glucose and lipids, gastrointestinal symptoms, and microbiota. *Nutrients* 2017;9:173.
79. Karl JP, Meydani M, Barnett JB, Vanegas SM, Goldin B, Kane A, Rasmussen H, Saltzman E, Vangay P, Knights D et al. Substituting whole grains for refined grains in a 6-wk randomized trial favorably affects energy-balance metrics in healthy men and postmenopausal women. *Am J Clin Nutr* 2017;105:589–99.
80. Salden BN, Troost FJ, Wilms E, Truchado P, Vilchez-Vargas R, Pieper DH, Jáuregui R, Marzorati M, van de Wiele T, Possemiers S et al. Reinforcement of intestinal epithelial barrier by arabinoxylans in overweight and obese subjects: a randomized controlled trial. *Arabinoxylans in gut barrier*. *Clin Nutr* 2018;37:471–80.
81. Abell GCJ, Cooke CM, Bennett CN, Conlon MA, McOrist AL. Phylotypes related to *Ruminococcus bromii* are abundant in the large bowel of humans and increase in response to a diet high in resistant starch. *FEMS Microbiol Ecol* 2008;66:505–15.
82. Ampatzoglou A, Atwal KK, Maidens CM, Williams CL, Ross AB, Thielecke F, Jonnalagadda SS, Kennedy OB, Yaqoob P. Increased whole grain consumption does not affect blood biochemistry, body composition, or gut microbiology in healthy, low-habitual whole grain consumers. *J Nutr* 2015;145:215–21.
83. Baer DJ, Stote KS, Henderson T, Paul DR, Okuma K, Tagami H, Kanahori S, Gordon DT, Rumpler WV, Ukhanova M et al. The metabolizable energy of dietary resistant maltodextrin is variable and alters fecal microbiota composition in adult men. *J Nutr* 2014;144:1023–9.
84. Boler B, Serao M, Bauer L, Staeger M, Boileau T, Swanson K, Fahey G. Digestive physiological outcomes related to polydextrose and soluble maize fibre consumption by healthy adult men. *Br J Nutr* 2011;106:1864–71.
85. Carvalho-Wells AL, Helmolz K, Nodet C, Molzer C, Leonard C, McKevith B, Thielecke F, Jackson KG, Tuohy KM. Determination of the in vivo prebiotic potential of a maize-based whole grain breakfast cereal: a human feeding study. *Br J Nutr* 2010;104:1353–6.
86. Clarke S, Green-Johnson J, Brooks S, Ramdath D, Bercik P, Avila C, Inglis G, Green J, Yanke L, Selinger L et al. beta-2-1 fructan supplementation alters host immune responses in a manner consistent with increased exposure to microbial components: results from a double-blinded, randomised, cross-over study in healthy adults. *Br J Nutr* 2016;115:1748–59.
87. Cloetens L, Broekaert WF, Delaet Y, Ollevier F, Courtin CM, Delcour JA, Rutgeerts P, Verbeke K. Tolerance of arabinoxylan-oligosaccharides and their prebiotic activity in healthy subjects: a randomised, placebo-controlled cross-over study. *Br J Nutr* 2010;103:703–13.
88. Costabile A, Fava F, Røytö H, Forssten SD, Olli K, Klievink J, Rowland IR, Ouwehand AC, Rastall RA, Gibson GR et al. Impact of polydextrose on the faecal microbiota: a double-blind, crossover,

- placebo-controlled feeding study in healthy human subjects. *Br J Nutr* 2012;108:471–81.
89. Costabile A, Klinder A, Fava F, Napolitano A, Fogliano V, Leonard C, Gibson GR, Tuohy KM. Whole-grain wheat breakfast cereal has a prebiotic effect on the human gut microbiota: a double-blind, placebo-controlled, crossover study. *Br J Nutr* 2008;99:110–20.
  90. Costabile A, Kolida S, Klinder A, Gietl E, Buerlein M, Frohberg C, Landschütze V, Gibson GR. A double-blind, placebo-controlled, cross-over study to establish the bifidogenic effect of a very-long-chain inulin extracted from globe artichoke (*Cynara scolymus*) in healthy human subjects. *Br J Nutr* 2010;104:1007–17.
  91. Damen B, Cloetens L, Broekaert WF, François I, Lescroart O, Trogh I, Arnaut F, Welling GW, Wijffels J, Delcour JA et al. Consumption of breads containing in situ-produced arabinoxylan oligosaccharides alters gastrointestinal effects in healthy volunteers. *J Nutr* 2012;142:470–7.
  92. Depeint F, Tzortzis G, Vulevic J, l'Anson K, Gibson GR. Prebiotic evaluation of a novel galactooligosaccharide mixture produced by the enzymatic activity of *Bifidobacterium bifidum* NCIMB 41171, in healthy humans: a randomized, double-blind, crossover, placebo-controlled intervention study. *Am J Clin Nutr* 2008;87:785–91.
  93. Fernando W, Hill J, Zello G, Tyler R, Dahl W, Kessel A. Diets supplemented with chickpea or its main oligosaccharide component raffinose modify faecal microbial composition in healthy adults. *Benef Microbes* 2010;1:197–207.
  94. François IEJA, Lescroart O, Veraverbeke WS, Marzorati M, Possemiers S, Evenepoel P, Hamer H, Houben E, Windey K, Welling GW et al. Effects of a wheat bran extract containing arabinoxylan oligosaccharides on gastrointestinal health parameters in healthy adult human volunteers: a double-blind, randomised, placebo-controlled, cross-over trial. *Br J Nutr* 2012;108:2229–42.
  95. Fuller Z, Louis P, Mihajlovski A, Rungapamestry V, Ratcliffe B, Duncan AJ. Influence of cabbage processing methods and prebiotic manipulation of colonic microflora on glucosinolate breakdown in man. *Br J Nutr* 2007;98:364–72.
  96. Gråsten SM, Juntunen KS, Mättö J, Mykkänen OT, El-Nezami H, Adlercreutz H, Poutanen KS, Mykkänen HM. High-fiber rye bread improves bowel function in postmenopausal women but does not cause other putatively positive changes in the metabolic activity of intestinal microbiota. *Nutr Res* 2007;27:454–61.
  97. Holscher HD, Bauer LL, Gourineni V, Pelkman CL, Fahey GC, Swanson KS. Agave inulin supplementation affects the fecal microbiota of healthy adults participating in a randomized, double-blind, placebo-controlled, crossover trial. *J Nutr* 2015;145:2025–32.
  98. Jenkins DJA, Vuksan V, Rao AV, Vidgen E, Kendall CWC, Tariq N, Würsch P, Koellreutter B, Shiwonari N, Jeffcoat R. Colonic bacterial activity and serum lipid risk factors for cardiovascular disease. *Metabolism* 1999;48:264–8.
  99. Maki KC, Gibson GR, Dickmann RS, Kendall CWC, Chen CYO, Costabile A, Comelli EM, McKay DL, Almeida NG, Jenkins D et al. Digestive and physiologic effects of a wheat bran extract, arabinoxylan-oligosaccharide, in breakfast cereal. *Nutrition* 2012;28:1115–21.
  100. Maneerat S, Lehtinen MJ, Childs CE, Forssten SD, Alhoniemi E, Tiphaine M, Yaqoob P, Ouwehand AC, Rastall RA. Consumption of *Bifidobacterium lactis* Bi-07 by healthy elderly adults enhances phagocytic activity of monocytes and granulocytes. *J Nutr Sci* 2013;2:e44.
  101. Martínez I, Kim J, Duffy PR, Schlegel VL, Walter J. Resistant starches types 2 and 4 have differential effects on the composition of the fecal microbiota in human subjects. *PLoS One* 2010;5:e15046.
  102. Petry N, Egli I, Chassard C, Lacroix C, Hurrell R. Inulin modifies the bifidobacteria population, fecal lactate concentration, and fecal pH but does not influence iron absorption in women with low iron status. *Am J Clin Nutr* 2012;96:325–31.
  103. Ramnani P, Costabile A, Bustillo AGR, Gibson GR. A randomised, double-blind, cross-over study investigating the prebiotic effect of agave fructans in healthy human subjects. *J Nutr Sci* 2015;4:e10.
  104. Ross AB, Bruce SJ, Blondel-Lubrano A, Oguey-Araymon S, Beaumont M, Bourgeois A, Nielsen-Moennoz C, Vigo M, Fay LB, Kochhar S et al. A whole-grain cereal-rich diet increases plasma betaine, and tends to decrease total and LDL-cholesterol compared with a refined-grain diet in healthy subjects. *Br J Nutr* 2011;105:1492–502.
  105. Slavin J, Feirtag J. Chicory inulin does not increase stool weight or speed up intestinal transit time in healthy male subjects. *Food Funct* 2011;2:72–7.
  106. Smith SC, Choy R, Johnson SK, Hall RS, Wildeboer-Veloo ACM, Welling GW. Lupin kernel fiber consumption modifies fecal microbiota in healthy men as determined by rRNA gene fluorescent in situ hybridization. *Eur J Nutr* 2006;45:335–41.
  107. Ten Bruggencate SJM, Bovee-Oudenhoven IMJ, Lettink-Wissink MLG, Katan MB, Van Der Meer R. Dietary fructooligosaccharides affect intestinal barrier function in healthy men. *J Nutr* 2006;136:70–4.
  108. Tuohy KM, Kolida S, Lustenberger AM, Gibson GR. The prebiotic effects of biscuits containing partially hydrolysed guar gum and fructooligosaccharides – a human volunteer study. *Br J Nutr* 2001;86:341–8.
  109. Vulevic J, Drakoularakou A, Yaqoob P, Tzortzis G, Gibson GR. Modulation of the fecal microflora profile and immune function by a novel trans-galactooligosaccharide mixture (B-GOS) in healthy elderly volunteers. *Am J Clin Nutr* 2008;88:1438–46.
  110. Vulevic J, Juric A, Walton GE, Claus SP, Tzortzis G, Toward RE, Gibson GR. Influence of galacto-oligosaccharide mixture (B-GOS) on gut microbiota, immune parameters and metabolomics in elderly persons. *Br J Nutr* 2015;114:586–95.
  111. Walton G, Heuvel E, Kusters M, Rastall R, Tuohy K, Gibson G. A randomised crossover study investigating the effects of galacto-oligosaccharides on the faecal microbiota in men and women over 50 years of age. *Br J Nutr* 2012;107:1466–75.
  112. Walton GE, Lu C, Trogh I, Arnaut F, Gibson GR. A randomised, double-blind, placebo controlled cross-over study to determine the gastrointestinal effects of consumption of arabinoxylan-oligosaccharides enriched bread in healthy volunteers. *Nutr J* 2012;11:36.
  113. Walton GE, Rastall RA, Rastall RA, Martini MC, Williams CE, Jeffries RL, Gibson GR. A double-blind, placebo controlled human study investigating the effects of coffee derived manno-oligosaccharides on the faecal microbiota of a healthy adult population. *Int J Probiotics Prebiotics* 2010;5:75–83.
  114. Zeng Y, Huang S, Mu G, Zeng X, Zhou X. Effects of whole grain-bean mixed staple food on intestinal microecology and metabolic parameters of obese people. *Chinese J Clin Nutr* 2015;23:27–34.
  115. Blädel T, Holm JB, Sundekilde UK, Schmedes MS, Hess AL, Lorenzen JK, Kristiansen K, Dalsgaard TK, Astrup A, Larsen LH. A randomised, controlled, crossover study of the effect of diet on angiotensin-like protein 4 (ANGPTL4) through modification of the gut microbiome. *J Nutr Sci* 2016;5:e45.
  116. Claesson MJ, Jeffery IB, Conde S, Power SE, O'Connor EM, Cusack S, Harris HMB, Coakley M, Lakshminarayanan B, O'Sullivan O et al. Gut microbiota composition correlates with diet and health in the elderly. *Nature* 2012;488:178–84.
  117. Coyte KZ, Schluter J, Foster KR. The ecology of the microbiome: networks, competition, and stability. *Science (New York, NY)* 2015;350:663–6.
  118. Johnson KVA, Burnet PWJ. Microbiome: should we diversify from diversity? *Gut Microbes* 2016;7:455–8.
  119. Van der Meulen R, Adriani T, Verbrugghe K, De Vuyst L. Kinetic analysis of bifidobacterial metabolism reveals a minor role for succinic acid in the regeneration of NAD<sup>+</sup> through its growth-associated production. *Appl Environ Microbiol* 2006;72:5204–10.
  120. Ganzle MG, Follador R. Metabolism of oligosaccharides and starch in lactobacilli: a review. *Front Microbiol* 2012;3:340.
  121. Gueimonde M, Debor L, Tolko S, Jokisalo E, Salminen S. Quantitative assessment of faecal bifidobacterial populations by real-time PCR using lanthanide probes. *J Appl Microbiol* 2007;102:1116–22.
  122. Matsuki T, Watanabe K, Fujimoto J, Miyamoto Y, Takada T, Matsumoto K, Oyaizu H, Tanaka R. Development of 16S rRNA-gene-targeted group-specific primers for the detection and identification of predominant bacteria in human feces. *Appl Environ Microbiol* 2002;68:5445–51.
  123. Davis LMG, Martínez I, Walter J, Goin C, Hutkins RW. Barcoded pyrosequencing reveals that consumption of galactooligosaccharides results in a highly specific bifidogenic response in humans. *PLoS One* 2011;6:e25200.



124. Whelan K, Judd PA, Preedy VR, Simmering R, Jann A, Taylor MA. Fructooligosaccharides and fiber partially prevent the alterations in fecal microbiota and short-chain fatty acid concentrations caused by standard enteral formula in healthy humans. *J Nutr* 2005;135:1896–902.
125. Moreno-Indias I, Sanchez-Alcoholado L, Perez-Martinez P, Andres-Lacueva C, Cardona F, Tinahones F, Queipo-Ortuno MI. Red wine polyphenols modulate fecal microbiota and reduce markers of the metabolic syndrome in obese patients. *Food Funct* 2016;7:1775–87.
126. Pokusaeva K, Fitzgerald GF, van Sinderen D. Carbohydrate metabolism in *Bifidobacteria*. *Genes Nutr* 2011;6:285–306.
127. Canani RB, Costanzo MD, Leone L, Pedata M, Meli R, Calignano A. Potential beneficial effects of butyrate in intestinal and extraintestinal diseases. *World J Gastroenterol* 2011;17:1519–28.
128. den Besten G, van Eunen K, Groen AK, Venema K, Reijngoud DJ, Bakker BM. The role of short-chain fatty acids in the interplay between diet, gut microbiota, and host energy metabolism. *J Lipid Res* 2013;54:2325–40.