



**Relationship Between Facility Type and Bulk Tank Milk  
Bacteriology, Udder Health, Udder Hygiene, and Milk  
Production on Vermont Organic Dairy Farms**

Journal:	<i>Journal of Dairy Science</i>
Manuscript ID	JDS.2023-24576.R1
Article Type:	Research
Date Submitted by the Author:	n/a
Complete List of Authors:	Jeffrey, Caitlin; University of Vermont, Department of Animal and Veterinary Sciences Andrews, Tucker; University of Vermont, Department of Plant and Soil Science Godden, Sandra; University of Minnesota, VPM Neher, Deborah; University of Vermont College of Agriculture and Life Sciences, Dept Plant and Soil Science Barlow, John; University of Vermont, Department of Animal Science
Key Words:	Mastitis, organic dairy cattle, bedded pack, milk quality

SCHOLARONE™  
Manuscripts

The STROBE-Vet statement checklist.

	Item	STROBE-Vet recommendation	Page #
Title and Abstract	1	(a) Indicate that the study was an observational study and, if applicable, use a common study design term	2
		(b) Indicate why the study was conducted, the design, the results, the limitations, and the relevance of the findings	2-3
Background / rationale	2	Explain the scientific background and rationale for the investigation being reported	4-5
Objectives	3	(a) State specific objectives, including any primary or secondary prespecified hypotheses or their absence	6
		(b) Ensure that the level of organization <sup>a</sup> is clear for each objective and hypothesis	6
Study design	4	Present key elements of study design early in the paper	2
Setting	5	(a) Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	7-8
		(b) If applicable, include information at each level of organization	NA
Participants <sup>b</sup>	6	(a) Describe the eligibility criteria for the owners/managers and for the animals, at each relevant level of organization	7-8
		(b) Describe the sources and methods of selection for the owners/managers and for the animals, at each relevant level of organization	7-8
		(c) Describe the method of follow-up	8
		(d) For matched studies, describe matching criteria and the number of matched individuals per subject (e.g., number of controls per case)	NA
Variables	7	(a) Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. If applicable, give diagnostic criteria	9-12
		(b) Describe the level of organization at which each variable was measured	9
		(c) For hypothesis-driven studies, the putative causal-structure among variables should be described (a diagram is strongly encouraged)	4-5

<b>Data sources / measurement</b>	8*	(a) For each variable of interest, give sources of data and details of methods of assessment (measurement). If applicable, describe comparability of assessment methods among groups and over time	9-12
		(b) If a questionnaire was used to collect data, describe its development, validation, and administration	9-10
		(c) Describe whether or not individuals involved in data collection were blinded, when applicable	NA
		(d) Describe any efforts to assess the accuracy of the data (including methods used for “data cleaning” in primary research, or methods used for validating secondary data)	13
<b>Bias</b>	9	Describe any efforts to address potential sources of bias due to confounding, selection, or information bias	32
<b>Study size</b>	10	(a) Describe how the study size was arrived at for each relevant level of organization	8
		(b) Describe how non-independence of measurements was incorporated into sample-size considerations, if applicable	NA
		(c) If a formal sample-size calculation was used, describe the parameters, assumptions, and methods that were used, including a justification for the effect size selected	NA
<b>Quantitative variables</b>	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen, and why	14
<b>Statistical methods</b>	12	(a) Describe all statistical methods for each objective, at a level of detail sufficient for a knowledgeable reader to replicate the methods. Include a description of the approaches to variable selection, control of confounding, and methods used to control for non-independence of observations	13-15
		(b) Describe the rationale for examining subgroups and interactions and the methods used	NA
		(c) Explain how missing data were addressed	17-18
		(d) If applicable, describe the analytical approach to loss to follow-up, matching, complex sampling, and multiplicity of analyses	NA
		(e) Describe any methods used to assess the robustness of the analyses (e.g., sensitivity analyses or quantitative bias assessment)	NA
<b>Participants</b>	13*	(a) Report the numbers of owners/managers and animals at each stage of study and at each relevant level of organization - e.g., numbers eligible, included in the study, completing follow-up, and analyzed	7-9

		(b) Give reasons for non-participation at each stage and at each relevant level of organization	7-9
		(c) Consider use of a flow diagram and/or a diagram of the organizational structure	NA
<b>Descriptive data on exposures and potential confounders</b>	14*	(a) Give characteristics of study participants (e.g., demographic, clinical, social) and information on exposures and potential confounders by group and level of organization, if applicable	16, 42-43, table S1-S5
		(b) Indicate number of participants with missing data for each variable of interest and at all relevant levels of organization	17-18
		(c) Summarize follow-up time (e.g., average and total amount), if appropriate to the study design	NA
<b>Outcome data</b>	15*	(a) Report outcomes as appropriate for the study design and summarize at all relevant levels of organization	17-21, 42-48
		(b) For proportions and rates, report the numerator and denominator	NA
		(c) For continuous outcomes, report the number of observations and a measure of variability	17, 43
<b>Main results</b>	16	(a) Give unadjusted estimates and, if applicable, adjusted estimates and their precision (e.g., 95% confidence interval). Make clear which confounders and interactions were adjusted. Report all relevant parameters that were part of the model	43-48, 17-21
		(b) Report category boundaries when continuous variables were categorized	42
		(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	NA
<b>Other analyses</b>	17	Report other analyses done, such as sensitivity/robustness analysis and analysis of subgroups	NA
<b>Key results</b>	18	Summarize key results with reference to study objectives	21-22
<b>Strengths and Limitations</b>	19	Discuss strengths and limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	21-22, 26, 32
<b>Interpretation</b>	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	21-34
<b>Generalizability</b>	21	Discuss the generalizability (external validity) of the study results	32
<b>Funding Transparency</b>	22	(a) Funding- Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	(a) 35
		(b) Conflicts of interest-Describe any conflicts of interest, or lack thereof, for each author	(b) 35

		(c) Describe the authors' roles- Provision of an authors' declaration of transparency is recommended (d) Ethical approval- Include information on ethical approval for use of animal and human subjects (e) Quality standards-Describe any quality standards used in the conduct of the research	(c) NA (d) 10 (IRB), 11 (IACUC) (e) 6
--	--	--	--

<sup>a</sup> Level of organization recognizes that observational studies in veterinary research often deal with repeated measures (within an animal or herd) or animals that are maintained in groups (such as pens and herds); thus, the observations are not statistically independent. This non-independence has profound implications for the design, analysis, and results of these studies.

<sup>b</sup> The word "participant" is used in the STROBE statement. However, for the veterinary version, it is understood that "participant" should be addressed for both the animal owner/manager and for the animals themselves.

\*Give such information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.

**Interpretive summary**

Previous studies reported bedded packs improve cow welfare and comfort and have advantages for manure management, soil health, and water quality. Consensus is lacking on whether bulk tank milk quality, udder health, udder hygiene and milk production are compromised on bedded packs. In an observational study of 21 organic dairies in Vermont during the non-grazing season (November-May), bedded packs were similar to tiestalls and freestalls in their bulk tank milk quality, udder health, udder hygiene and milk production. ~~We conclude that~~ ~~For producers considering a transition from tie stalls,~~ bedded packs ~~are may be~~ a viable option for dairy cattle housing ~~during the non-grazing season~~ in the Northeastern US.

**Running head:**

Milk quality and udder hygiene on VT organic dairies

**Relationship Between Facility Type and Bulk Tank Milk Bacteriology, Udder Health, Udder Hygiene, and Milk Production on Vermont Organic Dairy Farms**

Caitlin E. Jeffrey,<sup>1</sup> Tucker Andrews<sup>2</sup>, Sandra M. Godden<sup>3</sup>, Deborah A. Neher<sup>2</sup>, John W. Barlow<sup>1</sup>

<sup>1</sup> Department of Animal and Veterinary Sciences, University of Vermont, Burlington, VT 05405

<sup>2</sup> Department of Plant and Soil Science, University of Vermont, Burlington, VT 05405

<sup>3</sup> Department of Veterinary Population Medicine, College of Veterinary Medicine, University of Minnesota, St. Paul, MN 55108.

Corresponding author: John Barlow

Department of Animal and Veterinary Sciences,  
202 Terrill Building,  
University of Vermont,  
Burlington, VT 05405  
Phone: 802-656-1395  
Email: [john.barlow@uvm.edu](mailto:john.barlow@uvm.edu)

## Abstract

The primary objective of this cross-sectional observational study ~~on organic dairies~~ was to determine whether bulk tank milk quality, udder health, udder hygiene and milk production outcomes were associated with facility type ~~on organic dairies~~. A secondary objective was to identify other management-related risk factors associated with bulk tank milk quality, udder health, udder hygiene, and milk production on organic dairy herds in Vermont. We aimed to ~~collect bulk tank milk samples, udder hygiene scores, and complete a questionnaire on mastitis risk and bedding management practices on~~enroll 40 farms, in order to compare herds using the two most common housing systems (freestalls, tiestalls) with those using a bedded pack, for organic dairy cattle in the state during the non-grazing season (typically November-May). Variation in bedded pack management practices was observed, with two general styles, cultivated bedded packs and untilled deep bedded packs. Due to the limited number of herds using bedded packs to house lactating dairy cattle in Vermont, we combined untilled and cultivated bedded pack systems to describe udder hygiene, milk quality, and udder health on these loose-housing systems deeply-bedded with organic material. The study was completed on

21 farms (5 bedded packs, 6 freestalls, 10 tiestalls) before interruption due to the COVID-19 pandemic. Data captured from Dairy Herd Improvement Association records from the test closest to the date of the farm visit included average somatic cell score (SCS), standardized 150-day milk (pounds), % cows with current high SCS ( $\text{SCS} \geq 4.0$ ), % cows with newly elevated SCS (previous SCS  $< 4.0$  to current  $\geq 4.0$ ), and % cows with chronically elevated SCS ( $\text{SCS} \geq 4.0$  last two tests). Multivariable linear regression models were performed to describe outcomes by facility type, but suffered from limited statistical power due to small group sample sizes. Final results from unconditional comparisons showed that farms grouped by facility type did not differ in metrics captured from Dairy Herd Improvement Association test data (cow-level udder health measures, milk production), bulk tank milk somatic cell count (BTSCC) and aerobic culture data, or udder hygiene scores. Subsequently, a secondary analysis was conducted using univariate linear regression to identify associations between herd management factors and outcomes for all 21 farms combined. Although not all differences found were statistically significant in this secondary analysis which combined all farms, numeric differences that may be biologically important are reported showing farms with deeper bedding had a lower BTSCC, lower newly elevated SCS, lower elevated current SCS, lower average SCS, and better udder hygiene metrics. Farms with lower mean udder hygiene scores had numerically lower chronically elevated SCS, lower elevated current SCS, and lower average SCS. The current study provides insight on factors affecting bulk tank milk quality, udder health and hygiene measures on organic dairy farms in Vermont. We could not reject the null hypothesis that milk quality and udder health outcomes did not differ by facility type ~~bedded packs were comparable to more frequently used indoor housing systems (tiestalls and freestalls), we conclude that~~ Bbedded pack facilities are may be a viable option for confinement housing during the winter non-grazing season for



pasture-based herds interested in a loose-housing system in the Northeastern US. [More research including a larger number of herds is needed to test this hypothesis.](#)

**Keywords:** Mastitis, organic dairy cattle, housing, bedded pack, milk quality

## Introduction

Mastitis due to environmental pathogens, such as those commonly found in bedding material, has now become the “most common and costly form of mastitis in modern dairy herds” that have implemented standard mastitis control practices limiting the effect of contagious pathogens (Klaas and Zadoks, 2018). Teats of dairy cattle may be in direct contact with bedding materials for 40 to 60% of the day, making this an important potential source of exposure to opportunistic environmental mastitis pathogens (Tucker and Weary, 2004; Cook et al., 2005; Hogan and Smith, 2012). Work exploring how bedding materials relate to a cow’s risk of contracting mastitis has understandably focused on the most frequently used bedding materials and housing systems in the dairy industry. Currently, the most common type of dairy cattle housing for organic farms in Vermont is a tiestall barn, with freestall barns a distant second (Andrews et al., 2021). As consumer opinion about confinement housing of dairy cattle evolves and influences dairy policy, both the dairy industry and consumers are looking to move away from traditional housing systems ~~which~~ that restrict cow movement (Barkema et al., 2015). Many smaller-scale organic dairy farmers in Vermont with aging facilities, and especially tiestall barns, may be looking to adopt a bedded pack system on their farms as a form of loose-housing (Andrews et al., 2021).

The term “bedded pack” encompasses a variety of management styles (Bewley et al., 2017), including compost bedded-packs (CBP), which utilize aerobic decomposition to break down a bedding material of fine wood sawdust or shavings, as well as “conventional,” “traditional,” or “deep bedded packs” (Thurgood, 2009; Benson, 2012; Bewley et al., 2017; The Dairyland Initiative, 2024). CBP can vary in depth, frequency and depth of aeration (tilling), type of bedding material used, and in some regions the inclusion of forced air systems to dry the bedding (Leso et al., 2020). In the Northeastern U.S., some producers are using deep bedded pack systems where large volumes of straw or hay are added daily to an untilled surface in which strata of bedding and waste accumulate throughout the period of time when cows are housed on it (Benson, 2012). Oxygen is retained in the system by the selection of bedding material and the timing of its application (Neher et al., 2022; Thurgood et al., 2009). A number of authors suggest deep bedded -pack barns are synonymous with traditional straw yard housing systems (Bewley et al., 2017; Leso et al., 2020; Ferraz et al., 2020). However, we find the deep bedded packs being constructed on dairy farms in the Northeastern U.S. differ from traditional straw yards, in which material is completely removed at approximately monthly intervals and the housing is used year-round— (The Dairyland Initiative, 2024; Thurgood et al., 2009; Benson, 2012).

Bedded packs (BP) ~~These loose housing structures~~ are perceived to integrate well into Northeastern US pasture-based farm systems, and state and federal agencies in the U.S. are providing financial incentives for dairies to build these structures as part of manure management practices which improve water quality and contribute to soil conservation (USDA-NRCS; Andrews et al., 2021; Thurgood et al., 2009).

As interest in bedded packsBP grows, it is important to better understand milk quality, udder health and hygiene on farms using these housing alternatives. Understanding mastitis risk

for cattle housed on [bedded-packsBP](#) is especially important for organic dairy farmers, as they have limited effective options for treating intramammary infections (Ruegg, 2009). As mastitis-causing bacteria may thrive in the conditions found in compost bedded-packs (Black et al., 2014), previous work studying mastitis risk and bedding would suggest [bedded-packsBP](#) could pose a relatively higher risk for intramammary infections. Loose-housed cows continually add manure to the [bedded](#)-pack, contributing both pathogenic bacteria (non-*aureus* staphylococci, Wuytak et. al., 2020; *E. coli*, *Klebsiella* spp., and *Enterobacter* spp., Eberhart, 1984; streptococci, Zadoks et al., 2005) and nutrients to the organic bedding material. Organic bedding material is more likely to have a higher **bacteria** count than inorganic bedding, such as sand, (Hogan et al., 1989; Rowbotham and Ruegg, 2016b), as it supplies nutrients and moisture which encourages bacterial growth. This could lead to higher concentrations of bacteria on teat skin for cows on [bedded-packsBP](#), because: **1) organic bedding (in general) is inherently associated with a higher number of bacteria on teat ends (Fairchild et al., 1982; Rowbotham and Ruegg, 2016b), and 2) a higher concentration of bacteria in bedding is related to associated with a higher concentration of bacteria on teat ends (Hogan and Smith, 1997; Zdanowicz et al., 2004; Rowbotham and Ruegg, 2016b).** This higher concentration of bacteria on teat ends may put the mammary gland at an increased risk of infection, although limited evidence exists for this relationship (Neave et al., 1966; Pankey, 1989; Rowbotham and Ruegg, 2016a).

Previous work describing mastitis risk and cow hygiene on [bedded-packBP](#) systems includes descriptive studies of [actively-managed composting bedded-packsCBP](#) (Barberg et al., 2007b; Black et al., 2013; Fávero et al., 2015; Eckelkamp et al., 2016b; Albino et al., 2018; Heins et al., 2019). However, research comparing milk quality and cow hygiene between [bedded pack-systems-BP](#) and more traditional housing types has so far been limited to freestalls with

sand, which is an uncommon housing type for organic farms in Vermont (Andrews et al. 2021). These include a study comparing ~~actively-managed composting bedded packs~~ (CBP) and sand-bedded freestalls for farms with a history of low bulk tank somatic cell counts (Eckelkamp et al., 2016a), work describing hygiene and bulk tank milk somatic cell count (BTSCC) for sand-bedded freestalls and CBP (Adkins et al., 2022), and a comparison of CBP and two types of freestall barns (Lobeck et al., 2011). It is unclear whether the herds included in these prior studies were conventionally-managed or organic dairies. To the best of our knowledge, no studies describe and compare bulk tank milk quality, udder health and hygiene ~~on~~ for bedded packBP farms and tiestall barns of similar size and management style on small to midsize organic dairies in the same geographic area.

To better inform organic dairy producers in the Northeastern US, who may be interested in using a ~~bedded pack barn~~BP for housing their cattle during the non-grazing season (i.e., for “winter housing,” typically the months of November-May), we conducted a cross-sectional, observational study on organic dairies in Vermont. This study aimed to quantify bulk tank milk bacteriology, udder health and udder hygiene measures for the two most common indoor housing systems (freestalls, tiestalls) and farms using a ~~bedded pack~~BP for organic farms in Vermont. The objectives of this project were to identify whether bulk tank milk quality, udder health and hygiene outcomes differed by facility type, with a view to determining if ~~bedded pack systems~~BP are a viable option for indoor housing of lactating cows in VT during the non-grazing season. We hypothesized that udder health, hygiene, and bulk tank milk bacteriology of ~~bedded pack~~BP herds is inferior to that of more traditional housing types, as has been suggested by some previous research (~~Peeler et al., 2000; Fregonesi and Leaver, 2001~~; Barberg et al., 2007b; Lobeck et al., 2011). Therefore, our null hypothesis was that there no association between facility type

and udder health, hygiene, and bulk tank milk bacteriology on organic dairy farms using bedded  
packs BP and other systems for winter housing of lactating cow in Vermont. A secondary  
 objective was to identify other (non-facility) management-related risk factors associated with  
 bulk tank milk quality, udder health, udder hygiene, and milk production for organic VT dairy  
 herds.

## Materials and Methods

STROBE-VET (Strengthening the Reporting of Observational Studies in Epidemiology–  
 Veterinary Extension) statement guidelines were followed in the reporting of this study (O'Connor  
 et al., 2016).

### Herd enrollment and selection

The source population for this study was the 145 farms that responded to a survey sent to all  
 certified organic dairy farms producing cow milk in Vermont in Winter 2018-2019 (all farms, n  
 = 177). Certified organic dairy farms in the United States are required to allow their cows daily  
 access to pasture during the grazing season, and cows must obtain 30% of their dry matter intake  
 from grazing (Rinehart and Baier, 2011). In Vermont and other Northeastern US states, forage is  
unavailable directly from pasture during winter months and the climate necessitates use of indoor  
housing. During the non-grazing season (typically November–May in Vermont) When cows have  
no access to pasture in the winter non-grazing season are not on pasture), organic farms in  
Vermont house cows in a variety of indoor facility types. The Our previous Winter 2018-2019  
industry survey aimed to quantify the frequency and diversity of indoor housing and bedding  
 types used by organic dairy farmers in the state when cows were not on pasture, and for the

current study farms were recruited from respondents to this survey (Andrews et al., 2021). Dairy farms were eligible for enrollment in the current study if they: 1) responded to the initial survey in the Winter 2018-2019, 2) indicated they met the enrollment criteria of testing with the Dairy Herd Improvement Association (DHIA) at least monthly, 3) milked between 35 and 120 cows, and 4) indicated they would be interested in further participation. Eligible farms were contacted from this source population in Spring 2019 if they responded that they were using one of four categories of bedding/housing combinations for their indoor housing system: 1) freestall-[\(FS\)](#) barn bedded with sand, 2) [freestall-FS](#) barn bedded with shavings or sawdust, 3) tiestall [\(TS\)](#) barn bedded with shavings or sawdust, or 4) [BP, an enclosed loose housing facility deeply bedded with organic material](#) (hereafter, “bedded pack”). The first three housing and bedding combinations are the most frequently used by organic dairies in Vermont to house cows during the non-grazing season, and were compared to [bedded packsBP](#) as they were the housing type of interest for this project. [For the purposes of this study, the inclusive term “bedded pack” is used to encompass both CBP and deep bedded packs, and was defined as an enclosed loose housing facility deeply bedded with organic material, in which bedding and waste accumulate throughout the 6–8-month period of time when cows are housed on it and which is only removed once a year. Both CBP and deep bedded packs use carbon-rich substrates to create a clean, comfortable surface which allows animals to move freely. Urine and manure are not removed when bedding material is renewed, in contrast with other housing systems.](#)

A convenience sample of farms was enrolled in Spring 2019 from a list of eligible farms (grouped by housing/bedding combination) using the phone number or email address provided in the 2018-2019 survey response. Our aim was to enroll 40 farms for the current study, with 10 farms from each of the four housing/bedding categories described above.

Prior to obtaining the 2018-2019 survey results, based on preliminary data collected by the University of Vermont Center for Sustainable Agriculture Extension group, the study was designed anticipating that it would be possible to enroll 10 organic Vermont dairies using a [bedded-pack-systemBP](#) as their primary indoor housing system. However, out of the 17 farms from the 2018-2019 survey which indicated at least some use of a [bedded-pack-systemBP](#), one farm was not interested in any further participation, five did not use DHIA testing, and six only used a [bedded-pack-systemBP](#) as a secondary housing system in conjunction with a [tiestalHTS](#) barn, or cows were only on the pack a few hours a day. Because the number of farms using [bedded-packsBP](#) was fewer than anticipated, the eligibility requirements were relaxed to include one farm where cows spend the majority (two-thirds) of their time in a [bedded-packBP](#), with the remaining time in a [tiestalHTS](#) with wood shavings. Additionally, two [bedded-packBP](#) farms were included that had limited DHIA information: one farm did not utilize cow-level testing, and cow-level data for a second farm was limited due to their seasonal lactation schedule. ~~This study was intended to study cows while they were in their indoor housing system, so all herds visits were completed before any grazing had begun for the season.~~ As the number of BP being used in the state to house lactating dairy cattle was less than anticipated, those that were enrolled and grouped together utilized a variety of management strategies. Of the 5 enrolled farms using a BP, two would be classified as “compost bedded-packs,” utilizing aerobic decomposition to break down a bedding material of dry, fine wood sawdust or shavings (The Dairyland Initiative, 2024; Bewley et al., 2017; Endres, 2021). These two farms bedded solely with shavings/sawdust adding new bedding only as needed, and cultivated the pack twice a day. Two other farms used a “traditional” or “deep bedded pack” system, where large volumes of fresh, dry straw (or poor-quality hay) sufficient to keep cows clean and dry was added daily to a mass of bedding that



accumulates over the 6-8 months cows are housed indoors (The Dairyland Initiative, 2024; Thurgood, 2009; Benson, 2012; Bewley et al., 2017). The one remaining farm fell somewhere between these two types of classically defined BP; this farm bedded with straw and woodchips and cultivated every 48 hrs., adding chopped hay and woodchips every time the pack was cultivated. All farms in the study grouped as “bedded packs” shared the qualities of being an enclosed loose housing facility, deeply bedded with organic material (0.9-1.7 meters), which accumulated over the period of time animals were housed indoors and was only removed once a year.

Of the intended 40 herds to be recruited in the study, 21 herds (1 ~~freestall~~FS bedded with sand, 5 ~~freestalls~~FS bedded with wood shavings/sawdust, 10 ~~tiestalls~~TS bedded with wood shavings/sawdust, 5 ~~bedded packs~~BP) agreed to participate and farm visits were completed April-May 2019. This study was intended to study cows while they were in their winter (non-grazing months) indoor housing system, so all herds visits were completed before any grazing had begun for the season. Each herd was visited once during the study period. All herds sampled during this period were housing their cows as they would in the non-grazing season. Farm visits were suspended in mid-May 2019 as farms began turning their cows out to pasture ~~for the grazing season~~, with the intention of resuming in April 2020 to complete the remaining 19 herds. Due to COVID-19 pandemic activity restrictions, the decision was made to not resume the study, and the final analysis included the 21 herds sampled in 2019. As there was only one farm sampled using a ~~freestall~~FS facility bedded with sand, the initial plan to group farms by the four housing/bedding combinations specified was abandoned in favor of grouping farms by the three facility types used. The single sand ~~freestall~~FS was combined with ~~freestalls~~FS bedded with



wood shavings/sawdust (FS; n = 6), there were 10 ~~TS tiestalls~~ bedded with wood shavings/sawdust (TS), and 5 ~~bedded packs~~BP (BP).

### **Questionnaire administration, sampling, and udder hygiene scoring**

At each farm visit, a questionnaire was administered to collect information about housing and bedding management, as well as other practices on the farm that could impact mastitis risk (Supplemental Data). The study questionnaire was largely adapted from a previously published survey (Stiglbauer et al., 2013), with additional questions specific to the current study. The questionnaire was reviewed by a social scientist experienced in gathering qualitative data and tested before use with herd managers at the University of Vermont teaching dairy. Questions about mastitis risk explored producer concerns about bedding/mastitis risk; mastitis control, identification and record keeping; milking facilities, procedures, and hygiene practices; information about diet, vitamin and mineral supplementation, and water source; typical calving and periparturient practices; and fly control. Questions about housing and bedding management included describing type of housing system used for both lactating and dry cows; classification and description of any bedding material used; and bedding management practices for each housing type used. The questionnaire also collected some basic herd information (production numbers; number of lactating, dry, and youngstock; breed; record-keeping systems). Farms using ~~bedded pack systems~~BP were asked additional questions to gather detailed information about ~~bedded~~ pack construction, management, monitoring practices, and perceptions comparing ~~bedded packs~~BP to any previously used systems. Completion of the questionnaire required 45 minutes on average, ranging from about 30 minutes to 1.5 hours. The questionnaire and interview protocols were registered with the University of Vermont Institutional Review Board

273 (IRB certification 19-0057). The questionnaire was created and administered on a tablet using  
274 KoboCollect software (KoboCollect, 2019).

275 At each farm visit, a bulk tank milk sample ~~and bedding samples were collected. The bulk~~  
276 ~~tank milk sample was~~ collected directly from the top of the ~~bulk~~ tank using a 250-mL sterile  
277 single-use vial (Blue Dippas™, Dynalon Products, England) after at least 5 minutes of agitation.  
278 Samples were kept on ice in a cooler during transport until they were processed fresh for SCC  
279 measurement or were frozen and stored at -20°C in the laboratory, before being sent to a  
280 diagnostic lab for microbiological analysis. An on-farm observation sheet was completed, which  
281 collected information about the bulk tank, cow identification, a subjective assessment of air  
282 quality, and any outdoor exercise area (Supplemental Data). Additionally, measurements of the  
283 housing facilities were recorded for ~~freestalls~~FS and ~~tiestalls~~TS where appropriate (stall sizes,  
284 pen sizes, bedding depth, stocking density, trainer use), as well as observations about ~~bedded~~  
285 ~~packs~~BP when applicable (~~temperature~~, depth, pen size, and stocking density in m<sup>2</sup> per animal).  
286 ~~If multiple pens were present (e.g., freestall barn), used bedding samples were collected from the~~  
287 ~~pen containing the largest group of lactating cows, or from the highest producing group of~~  
288 ~~animals if there were multiple pens of equal size.~~ Bedding depth of ~~freestalls~~FS and ~~tiestalls~~TS  
289 was included as a producer reported value in the questionnaire. **Bedding depth of ~~bedded pack~~BP**  
290 **facilities was measured by forcing a meter stick down to the level of the cement pad or gravel**  
291 **under the pack, where the pack met a cement knee wall, and recording the height of the pack at**  
292 **that point.** Udder hygiene scoring was completed by the same researcher at all farms for a  
293 minimum of 30 ~~randomly selected~~ cows on each farm **(the first 30 able to be evaluated in a loose**  
294 **pen, or the first 30 encountered in a tiestall).** ~~Udder hygiene scores were taken from cows~~  
295 ~~housed in the same pens from which used bedding samples were collected.~~ A four-point udder

hygiene scoring system was used, where 1 = free of dirt, 2 = slightly dirty (2–10% of surface area), 3 = moderately covered with dirt (10–30% of surface area), and 4 = covered with caked on-dirt (>30% of surface area) (Schreiner and Ruegg, 2002). Animal use for this project was approved by the University of Vermont Institutional Animal Care and Use Committee (IACUC; protocol #PROTO202000089).

### **Herd-level udder health measurements**

Herd-level DHIA test results for the test day closest in time to the farm visit (either preceding or following day of farm visit, whichever was shorter) were captured from the record processing center working with each herd (Lancaster DHIA, Manheim, PA; Dairy One Co-Op. Inc., Ithaca, NY). Information captured included test date, number of lactating cows, standardized 150-day milk production (STD 150-day milk), and test-day average cow-level somatic cell score (SCS). The following udder health measures were also captured from DHIA records: proportion of cows with an  $\text{SCC} \geq 200,000$  cells/mL on most recent test day (“elevSCS”), where elevated SCS was defined as a somatic cell score of  $\geq 4.0$ ; the proportion of cows with a newly elevated SCS (“newSCS”), which was defined as a SCS changing from  $< 4.0$  to  $\geq 4.0$  over the last 2 tests; and the proportion of cows with a chronically elevated SCS (“chronSCS”), which was defined as having a SCS  $\geq 4.0$  on the last two tests (Schukken et al., 2003).

### **Bulk tank milk culture and bulk tank somatic cell count measures**

An aliquot of the bulk tank milk sample was stored at  $-4^{\circ}\text{C}$  until it could be transported to the laboratory of a dairy processing plant (St. Alban’s Cooperative/Dairy Farmers of America, St. Albans, VT) within 48 hours of collection for determination of the bulk tank somatic cell count (BTSCC).

Frozen bulk tank milk samples were shipped on ice to the Laboratory for Udder Health (University of Minnesota Veterinary Diagnostic Laboratory, St. Paul) for analysis. Methodology for bulk tank milk cultures at the Laboratory of Udder Health are described elsewhere (Patel et al., 2019). Briefly, thawed, room-temperature bulk tank milk and a 10-fold dilution of each bulk tank milk sample were plated onto MacConkey, Factor (gram-positive selective agar; University of Minnesota), and Focus (selective for SSLO bacteria; University of Minnesota) media plates and incubated for two days at 37°C. Any lactose-fermenting colonies on MacConkey medium were counted and reported as coliform bacteria. Any  $\beta$ -hemolytic colonies on Focus medium were counted and identified to the species level using a MALDI Biotyper (suspect *Streptococcus agalactiae*). All remaining colonies on Focus medium that were not identified as *Strep. agalactiae* were counted and recorded as streptococci or strep-like organisms (SSLO). Hemolytic colonies on Factor medium were counted and identified to the species level using a MALDI Biotyper (suspect *Staph. aureus*). Any hemolytic colonies with a confidence score  $\geq 2.0$  for *Staph. aureus* were counted and reported as such. Remaining colonies of staphylococci on Factor media (based on colony morphology, catalase reaction, or Gram stain) were counted and reported as *Staph. spp.* Bulk tank samples were also cultured for *Mycoplasma spp.* (0.1 mL milk was swabbed across a Mycoplasma agar plate, then placed in a 7% CO<sub>2</sub> incubator at 37°C for 7 days, after which they were examined for *Mycoplasma spp.* by a trained microbiology technician). For each bulk tank milk sample, total colony-forming units (cfu) per mL were calculated for coliform organisms, *Staph. spp.*, streptococci and strep-like organisms (SSLO), *Staph. aureus*, *Strep. agalactiae*, and *Mycoplasma spp.* The lower threshold of detection for bacteria in this bulk tank milk culture protocol was 5 cfu/mL, and the upper threshold was 62,500 cfu/mL.

## Data management and analysis

Bulk tank milk culture results, BTSCC, DHIA test results, farm-level udder hygiene outcomes, questionnaire data, and farm observations were entered into an Excel database (Microsoft Corp., Redmond, WA). Udder hygiene scores for individual cows were used to calculate two farm-level udder hygiene measures: 1) mean udder hygiene score, and 2) proportion of cows with dirty udders (udder hygiene score  $\geq 3$ ), which were incorporated into the database. This Excel database was then imported into the R Statistical Programming Environment (R Development Core Team, 2023) for data cleaning, checking, and statistical analysis. The distribution of outcome variables was assessed to check for normality using a Shapiro-Wilk test with significance set at  $P \leq 0.05$ , visual assessment of distribution and residuals, skewness, and comparison of the median and mean values. Raw bulk tank somatic cell count (BTSCC) data was  $\log_{10}$  transformed for analyses. Descriptive statistics were calculated to evaluate the distribution of data, data integrity, and to identify missing data. Descriptive statistics generated included description of general herd characteristics and farm traits, lactating cow housing/facilities, lactating cow bedding material/bedding management practices, milking hygiene procedures, and mastitis control practices for all 21 herds included in the study.

*Objective 1. Evaluation of relationships between housing system and measures of milk quality, udder health, udder hygiene and milk production.* As most measures of aerobic culture data were not normally distributed even after log transformation, a Kruskal-Wallis test was used to compare cfu counts of bacteria from bulk tank milk between the three facility types. Statistical significance for this test was declared at  $P \leq 0.05$ . Multiple attempts were made using multivariable analysis to compare the four aerobic culture outcomes for bulk tank milk, but all

modeling approaches suffered from over-parametrization even when data was log transformed and were not pursued further.

Independent farm-level predictors from the herd-management questionnaire offered to the multivariable models are described in Table 1. Continuous variables underwent correlation analysis to identify predictor variables that were highly correlated (correlation coefficient  $\geq 0.60$ ), and unconditional associations among categorical variables were evaluated using a Pearson's chi-squared or Fischer's Exact test as appropriate ( $P \leq 0.05$ ). An ANOVA was used to check for correlation between numeric continuous variables and categorical variables ( $P \leq 0.05$ ). When a categorical variable had multiple groups with a small number of observations in each, groups were combined when biologically reasonable to have all categories of predictor variables contain at least five observations. If any predictor had only one observation in a group and there was no way to combine groups in a logical way, it was excluded from further analysis (but listed in descriptive statistic tables, Supplemental Tables S1-S4).

Univariate linear regression was performed in R using the "lme4" package to investigate the unconditional relationship between the six udder health and production outcomes (BTSCC, avg. SCS, newSCS, elevSCS, chronSCS, STD 150-day milk) and two hygiene outcomes (mean hygiene score, proportion of dirty udders) for each farm and the previously-described herd-level independent variables. The two udder hygiene metrics (proportion dirty udders and average udder hygiene score) were used as both predictor variables (in models for other outcome variables) and outcome variables in models of their own. Any explanatory variable that was unconditionally associated with 1 or more of the outcomes of interest at  $P < 0.20$  was then offered into a multivariable model investigating the relationship between the udder health and production or hygiene outcome and the herd-level predictor variables. If any predictor variables

were found to be correlated with each other at the previously described cut-offs, the one with the more highly significant relationship from univariate analysis was offered to the multivariable model when appropriate. The two udder hygiene metrics were highly correlated (derived from the same data), so whichever one had a smaller  $P$ -value from the univariate analysis was chosen for inclusion in the model-building process. Facility type was forced into these multivariable models, as it was the primary explanatory predictor of interest. A backward stepwise variable selection process was then used, with the least significant variables being removed one by one until all remaining predictors had  $P \leq 0.10$ . Final models were selected based on lowest Akaike information criteria, and an  $F$ -test to compare the final model to the model with facility type as the only predictor (significance declared at  $P \leq 0.05$ ). Overall statistical significance for facility type (the main predictor of interest) was declared at  $P \leq 0.05$ . The multivariable modelling approach described above aimed to investigate the conditional relationship between facility type and the eight outcomes of interest while controlling for different farm management practices, housing characteristics, milking procedures and mastitis control practices.

*Objective 2. Identify other (non-facility) management-related risk factors associated with bulk tank milk quality, udder health, and milk production in organic dairy herds.* After grouping all 21 farms together, we used linear regression in the same manner as described above in Objective 1 to explore associations between the independent predictors described in Table 1 and the six udder health and production outcomes (BTSCC, avg. SCS, newSCS, elevSCS, chronSCS, STD 150-day milk) and two hygiene outcomes (mean hygiene score, proportion of dirty udders). Unconditional relationships between the eight outcome variables and independent predictors are reported for a significance level of  $P \leq 0.20$  for an  $F$ -test, and only for predictor variables with group sizes of at least  $n = 5$ .

## Power analysis

A priori sample size calculations were not performed, as group size was determined by the number of organic dairy herds housing lactating cows on [bedded pack systemsBP](#) in our region.

## Results

### Description of study herds

Of the 21 herds enrolled, 5 used a [bedded pack systemBP](#), 1 used a [freestallFS](#) bedded with sand, 5 used a [freestallFS](#) bedded with shavings/sawdust, and 10 used a [tiestallTS](#) bedded with shavings/sawdust (Supplemental Table S1). ~~Of the 5 BP farms, two bedded with shavings/sawdust and cultivated 2 times a day to promote aerobic composting, 1 bedded with straw and woodchips and cultivated 2 times/week, and 2 bedded mainly with straw, adding woodchips as needed, and did not cultivate the pack at all.~~ The predominant breeds on all farms were Holstein (n = 8 farms), Jersey (n = 10), and mixed Holstein-Jersey crosses/other (n = 3). The median (mean; range) number of lactating cows was 68 (64.9; 32-99). The median annual rolling herd average milk production for the farms was 6,367 (6,424; 4,082-9,618) kg. Nineteen of the 21 farms tested with DHIA monthly while their cows were in milk, 1 farm tested 5-8 times/year, and 1 tested every other month. On average, DHIA data was captured from a test day 4 days before the farm visit (range: -28 days to +33). [The average depth of bedding in the 15 freestalls and tiestalls where producers provided an estimate was 4.5 cm \(SD: 3.5 cm; range: 1.3-12.7 cm\). The average depth of bedded packs \(measured by researchers\) was 130 cm \(SD: 31; range: 90-170 cm\).](#) Detailed descriptions further characterizing study farm management practices and housing characteristics for lactating animals (e.g., laying surface, ventilation, stocking



density), and details about bedding material and bedding management practices for lactating animals (e.g., bedding depth, frequency of adding new bedding, manure removal) are provided in Supplemental Tables S1 and S2, respectively. Detailed descriptions of routine milking procedures and mastitis control practices are provided in Supplemental Tables S3 and S4, respectively.

### Description of bulk tank milk quality, udder health measures, milk production, and udder hygiene scores

There was no difference in cfu count between the three facility types for any of the four bacterial groups measured using a nonparametric unconditional comparison (Table 2). The aerobic culture results for the four bacterial groups measured for bulk tank milk did not differ among facility types (Table 2). None of the 21 bulk tank milk samples were positive for *Strep. agalactiae* or *Mycoplasma* spp. Sixteen of the 21 samples were negative for coliforms on aerobic culture, while 5 farms had a coliform count of 5 cfu/mL. *Staph. aureus* was found in the bulk tank milk from 13/21 herds, with a median (range) cfu/mL of 50 (15-320) when present.

BTSCC, % cows with newly elevated SCS, % cows with chronically elevated SCS, % cows with elevated SCS, avg. SCS, and STD 150-day milk production did not differ by facility type (Table 3).

The overall mean (95% CI) of herd-level udder hygiene scores for all 21 farms was 2.32 (2.16-2.49). The mean hygiene score was 2.2 (1.91-2.44) for ~~bedded pack~~<sup>BP</sup> farms (n = 5), 2.5 (2.24-2.76) for ~~tie stall~~<sup>TS</sup> farms (n = 10), and 2.15 (1.93-2.37) for ~~freestall~~<sup>FS</sup> farms (n = 6). Mean udder hygiene score did not differ by facility type. The overall mean proportion of cows with

dirty udders in a herd (udder hygiene score  $\geq 3$ ) was 40% (31-48). The mean proportion of cows with dirty udders (95% CI) was 32% (18-46) for [bedded-packBP](#) farms, 49% (35-62) for [tiestallTS](#) farms, and 32% (20-44) for [freestallFS](#) farms. The proportion of cows with dirty udders did not differ by facility type.

**Objective 1. Analysis of relationship between facility type and measures of bulk tank milk quality, udder health, milk production, and udder hygiene scores**

Final multivariable models are summarized in Table 4. All 21 farms were able to be included in the models for BTSCC, average hygiene score, and proportion of dirty udders. For the models exploring newSCS, chronSCS, and elevSCS, two [bedded-packBP](#) farms did not have available DHIA data (n = 19; group sizes: FS = 6, TS = 10, BP = 3). One [bedded-packBP](#) farm did not have average cow-level SCS data (n = 20; group sizes: FS = 6, TS = 10, BP = 4). For STD 150-day milk, one [bedded-packBP](#) farm and two [tiestallTS](#) farms were missing DHIA data (n = 18; group sizes: FS = 6, TS = 8, BP = 4). Farms with missing data for a particular outcome were excluded for the analyses of that outcome.

*Bulk tank milk quality outcomes*

~~There was no difference in cfu count between the three facility types for any of the four bacterial groups measured using a nonparametric unconditional comparison (Table 2). Multiple attempts were made using multivariable analysis to compare the four aerobic culture outcomes for bulk tank milk, but all modeling approaches suffered from over parametrization even when data was log transformed and were not pursued further.~~

Variables that were associated at  $P < 0.20$  with BTSCC in univariate analysis included predominant breed, if herds ever performed culture of mastitic milk, glove use, and herd size.

The final multivariable included facility type (forced) and herd size. Facility type was not associated with BTSCC in the final model (Table 4).

#### *Udder health outcomes*

Herd size category, use of bedding amendment, air quality as assessed by researcher, glove use at milking, and clinical mastitis record keeping practices were offered to a multivariable model for newSCS. The final multivariable model included facility type (forced), bedding amendment use, air quality, glove use, and mastitis record keeping practices. Facility type was not associated with newSCS in the final model (Table 4).

Variables that were associated at  $P < 0.20$  with chronSCS in univariate analysis included feeding additional supplemental selenium, use of a bedding amendment, clipping/flaming udder hair, and proportion of dirty udders. The final multivariable model included all four variables from univariate analysis, as well as facility type (forced). Facility type was not found to be a significant predictor of the outcome chronSCS (Table 4).

Bedding amendment use and mean hygiene were offered to a multivariable model for elevSCS. Facility type (forced), bedding amendment, and mean hygiene were retained in the final multivariable model. Facility type was not associated with elevSCS in the final model (Table 4).

Feeding additional supplemental selenium, use of bedding amendment, OMRI-listed intramammary product at dry-off, injectable selenium and vitamin E product, and mean hygiene were offered to a multivariable model for herd average SCS. The final multivariable model for avg. SCS included facility type (forced), use of bedding amendment, dry product, injectable selenium, and mean hygiene score. Facility type was not found to be a significant predictor of avg. SCS (Table 4).

*Milk production outcome*

Variables that were associated at  $P < 0.20$  with STD 150-day milk included use of injectable selenium and vitamin E product, whether producers cultured high SCC cows, and herd size group. All three variables and facility type (forced) remained in the final multivariable model (Table 4). Facility type was not associated with STD 150-day milk in the final model (Table 4).

*Udder hygiene outcomes*

Air quality assessed by researcher was offered to the multivariable model for proportion of dirty udders. The final multivariable model included only facility type (forced), which was not associated with proportion of dirty udders.

Variables that were associated at  $P < 0.20$  with average hygiene score included whether the producer ever cultured quarter milk samples and whether they checked for cases of clinical mastitis by both examining the udder and forestripping. The final multivariable model included facility type (forced), and how the producer checked for clinical mastitis. Facility type was not associated with the outcome of mean udder hygiene (Table 4).

**Objective 2. Analysis of farm management factors (non-facility) associated with bulk tank milk quality, udder health, milk production, and udder hygiene scores for all farms combined**

Selected results of univariate linear regression models identifying management factors beyond facility type which were unconditionally associated with bulk tank milk quality, udder health, milk production and hygiene outcomes for all farms combined ( $n = 21$ ) at  $P < 0.20$  are presented in Table 5. We report the results of these univariate regression models as they may be

biologically important, even though many failed to reach threshold for declaring statistical significance at  $P \leq 0.05$ , possibly due to small sample size.

The depth of bedding in stalls for [freestallFS](#) and [tiestallTS](#) herds was unconditionally associated with multiple udder health outcomes. As the depth of bedding in [freestallFS](#) and [tiestallTS](#) herds increased, multiple udder health measures improved, including lower avg. SCS, BTSCC, elevSCS, and newSCS. Similarly, comparing farms where cows were on deep bedding (i.e., grouping all herds reporting deeply-bedded stalls plus [bedded-packBP](#) herds) to herds that had stalls with a smaller amount of bedding on top of a mattress or concrete, farms with deep bedding had a numerically lower BTSCC.

Udder hygiene measures were associated with several udder health outcomes. Higher mean hygiene scores and proportion of udders scored  $\geq 3$  were associated with higher chronSCS, elevSCS, and average SCS. A few specific management practices were also found to be unconditionally associated with udder health outcomes: consistent glove use was associated with lower newSCS and BTSCC, clipping or flaming udders was associated with fewer chronSCS, and both parenteral supplementation of vit. E/selenium and use of an OMRI-listed intramammary product at dry-off were associated with lower average SCS and higher STD 150-day milk.

Both udder hygiene outcomes were unconditionally associated with the same predictors, most of which were related to the depth of bedding for cows. For [the five](#) herds using a [bedded packBP](#), deeper bedding was associated with lower average hygiene scores and lower proportion of dirty udders. Farms with cows housed on some type of deep bedding (i.e., grouping [the three all herds freestalls and tiestalls](#) reporting deeply-bedded stalls, plus [the five bedded-packBP](#) herds) had numerically lower average udder hygiene scores and proportion dirty udders

compared to cows on stalls with bedding over a mattress or concrete surface. For the fifteen ~~tiestalls and freestalls farms~~ reporting bedding depth in stalls, increased bedding depth was associated with lower mean udder hygiene score and a numerically lower proportion of dirty udders.

## Discussion

This work presents the results of our observational study exploring the relationship between facility type and udder health and hygiene metrics, BTM quality (SCC and microbiology), and milk production on organic dairy farms in Vermont. The current study is to the authors' knowledge the first direct comparison of milk quality, udder health and udder hygiene on ~~bedded-packBP~~ farms to both ~~tiestallTS~~ and ~~freestallFS~~ herds of similar size and management styles, for a population of entirely small to midsize organic dairy farms. The major objective was to identify if milk quality, udder health and hygiene outcomes were associated with facility type, thereby exploring if ~~bedded-packBP~~ systems are a viable option for housing in Vermont during the non-grazing season compared to the two most common indoor housing systems in the state (~~freestallsFS~~, ~~tiestallsTS~~). This study is also the first to describe udder health and hygiene on ~~bedded-packsBP~~ in the Northeastern US, which is significant as the performance of these systems can be greatly influenced by climatic factors. As BTM bacteriology, udder health and hygiene metrics, and milk yield did not differ for BP herds compared to TS and FS herds, there was insufficient evidence to reject our null hypothesis that these metrics would not vary by facility type. ~~We conclude that bedded-pack systems can be considered a viable loose-housing option for organic dairy cattle during the non-grazing season in the Northeastern US.~~

**Objective 1: Comparison of bulk tank milk quality, udder health, milk production, and udder hygiene measures by facility type**

~~Previous work describing bulk tank milk aerobic culture data for farms using a bedded pack system has primarily been limited to descriptive studies enrolling only composting bedded pack herds (Barberg et al., 2007b; Shane et al., 2010), with only one study directly comparing bacterial counts between composting bedded packs and freestall barns (Lobeck et al., 2012). Although there is a substantial body of work describing udder health and milk quality for cows housed in straw yards (Astiz et. al, 2014; Fregonesi and Leaver, 2001; Fregonesi and Leaver, 2002; Ward et. al 2002; Peeler et al. 2000), description of these outcomes in the literature is limited for static deep bedded packs. As such, the focus of the discussion will compare BP in the current study (both static and composting) to the more recent body of work on compost bedded-pack farms. Previous work describing bulk tank milk aerobic culture data for farms using a BP system has primarily been descriptive studies of compost bedded-pack herds (Barberg et al., 2007b; Shane et al., 2010), with one study directly comparing bacterial counts between CBP and FS barns (Lobeck et al., 2012). The current study is the first the authors are aware of directly comparing bacterial counts of bulk tank milk between bedded packs (both composting and static) and freestall barns, and the first one to describe a population of exclusively organic dairies. The six farms included in Lobeck et al. (2012) used mainly wood sawdust as bedding material (with one using wheat straw by-product) as did the 12 farms in Barberg et al. (2007). This is similar to the current study, where three of five bedded packs used a combination of woodchips/shavings and straw/hay, and two used exclusively sawdust/shavings. The six farms included in Shane et al. (2010) bedded with a variety of “alternative” organic materials, including straw by-products, soybean stubble, and oat hulls. Although farms in these previous studies used a similar array of~~

bedding materials to those in the current study (wood sawdust, wheat straw by-product, Lobeck et al. 2012; wood sawdust, Barberg et al. 2007; “alternative” organic materials, Shane et al. 2010), the sampling period for these previous works differed from the present in seasonality, compounding the difficulty of direct comparison for milk quality outcomes (Pantoja et al., 2009). Barberg et al., 2007b evaluated milk culture results across the summer months, while Lobeck et al., 2012 sampled year-round; the current study focused solely on sampling during the winter, when organic pasture-based herds animals are primarily housed inside in Vermont. In contrast to previous work, which evaluated milk culture results across the summer months (Barberg et al., 2007b) and year-round (Lobeck et al., 2012), the current study focused solely on sampling during the period when animals are primarily housed inside in Vermont. We were most interested in studying bulk tank milk bacteriology for these organic herds during the non-grazing season, as this is when these pastured-based farms need to house their animals inside. All herds included had excellent bulk tank milk quality; most (19/21) fell into the “low BTSCC” category as defined by Jayarao et al. 2004 with the remaining 2 in the “medium BTSCC” category.

The *Staph.* spp. count for the five bedded pack BP farms included in this study (median: 40 cfu/mL, range: 0-130) was comparable to previous work describing bulk tank milk quality for CBP in Minnesota during the winter months. Lobeck et al. 2012 found a mean of 26.1 cfu/mL (95% CI: 2-443) and Shane et al. (2010) found a range of 0-108 cfu/mL for *Staph.* spp. from BTM collected just over in the winter months from six composting bedded pack CBP farms. “*Staph.* spp.” is comprised of a diverse group of different species, with 23 (Condas et al., 2017) or 25 (De Visscher et al., 2017) different species isolated from intramammary infections in dairy cattle. Within this highly heterogenous group of bacteria, some species are considered primarily host-adapted (colonizing the skin or udder), while others are primarily found in the cow’s



environment (reviewed in De Buck et al., 2021). Certain species have been associated with stall surfaces, air, and unused sawdust bedding material (Piessens et al., 2011), some with different facility types (Condas et al., 2017), and others with environmental contamination and poor teat hygiene at milking time (De Visscher et al., 2016; De Visscher et al., 2017). Although the specific source and routes of transmission for many *Staph.* spp. are still being elucidated, the importance of post-milking teat dip to control this group of bacteria has been established (Hogan et al., 1987), while the efficacy of pre-dipping to control *Staph.* spp. other than *S. aureus* remains controversial (Pankey, 1989). In general, the use of pre- and post- milking teat dip decreases contamination of bulk tank milk both by commensal skin organisms and environmental contamination at milking time (Hogan et al., 1987; Pankey et al., 1985; Pankey et al., 1987; Quirk et al., 2012). All but one farm in the current study would fall into the “low” category for *Staph.* spp. counts in the BTM (Jayarao et al., 2004), which is consistent with all 21 herds using both pre- and post-dip consistently at milking time.

Streptococci and strep-like organisms (SSLO) counts in BTM for bedded packsBP in the current study were much lower than those from Minnesota composting bedded packsCBP in the winter (98-48,400 cfu/mL, -Shane et al. 2010; mean: 911 cfu/mL, 95% CI: 138-6,011, reported a range of SSLO counts of 98-48,400 cfu/mL for six farms, and Lobeck et al. 2012 reported a mean of 911 cfu/mL (95% CI: 138-6,011). The median SSLO counts for bedded pack farms included in the current study was 35 cfu/mL (range: 10-80). Work from Barberg et al. (2007) describing milk quality on composting bedded packsCBP in Minnesota noted that 6 of 12 farms sampled had “high” levels of SSLO. SSLO count did not differ between tiestallsTS, freestallsFS, and bedded packsBP in the current study. The overall SSLO count for all 21 farms included in the current study (median: 45 cfu/mL, range: 10-1250) was lower than that for the overall *Strep.*

count for all ~~three~~ facility types studied in Lobeck et al. 2012 (445 cfu/mL, 95% CI: 116-1704).  
~~As the overall SSLO counts for all farm types included in the Minnesota studies are higher than~~  
~~that found for all 21 farms in the current study, b~~ Metter milking and bedding hygiene practices  
amongst herds included in the current study may best explain this difference in BTM pathogen  
profiles (Jayarao and Wolfgang, 2003).

All farms had low levels of coliforms in bulk tank milk ~~(median: 0 cfu/mL, range: 0-5)~~,  
indicating excellent hygiene practices at milking time (Jayarao and Wolfgang, 2003). Coliform  
~~counts did not differ between the three facility types. Bedded pack farms in the current study had~~  
The low very BTM low coliform counts in BTM for BP in the current study are ~~(median: 0~~  
~~cfu/mL, range: 0-5)~~, similar to those found for three compost bedded pack CBP farms in a  
Brazilian study Brazil (2.8 cfu/mL; Fávero et al. 2015). ~~These low coliform counts are~~ This is in  
contrast with previous work describing BTM quality for this kind of facility in the United  
~~States U.S.~~ Coliform counts for bedded packs in Minnesota (in the winter ranged from 15-1,128  
~~cfu/mL, (Shane et al., 2010); mean: 63.7 cfu/mL, 95% CI: 6-735, and the six bedded packs  
~~included in~~ Lobeck et al. 2012. ~~had a mean of 63.7 cfu/mL (95% CI: 6-735), although d-~~  
~~However,~~ direct comparison of coliform counts between studies may be potentially problematic  
due to variation in duration of freezer storage (Schukken et al., 1989). ~~Although sampled during~~  
~~summer months,~~ Barberg et al. 2007 found that 5 of 12 bedded packs BP sampled during the  
summer months had “high” levels of coliforms in BTM, contributing to their conclusion that  
“special attention to cow preparation procedures at milking time are a must for achieving  
satisfactory milk quality when cows are housed in compost dairy barns.”~~

Prevalence of *Staph. aureus* was similar between the five VT bedded pack BP farms in  
the current study ~~(median: 0 cfu/mL, range: 0-30)~~ and the six ~~bedded packs~~ described in Lobeck

et al. 2012 (6.2 cfu/mL, 95% CI: 1.3-30.1). Farm-level prevalence of *Staph. aureus* was also fairly low for [bedded-packsBP](#) studied in Shane et al. 2010 (3 of 6 farms BTM negative) and Barberg et al. 2007 (only 1 of 12 farms with a “high” level of *Staph. aureus*). Overall, the population of all 21 farms in the current study had a higher amount of *Staph. aureus* in BTM than the 18 Minnesota farms described in Shane et al. 2010 (median: 30 cfu/mL, range: 0-320; vs. 17.3 cfu/mL, 95% CI: 3.3-91.2). Although it is not clear how many herds included in previous work on [bedded-packsBP](#) were certified organic, [thise](#) higher prevalence of *Staph. aureus* [amongst-on](#) organic farms in the current study is consistent with work comparing organic and conventional dairy systems (Pol and Ruegg, 2007).

Analysis of a single bulk tank milk sample from a farm is a simple, convenient, and relatively inexpensive way to capture a snapshot of current milk quality and animal health on a farm, and can be a highly specific (albeit poorly sensitive) screening test for major contagious mastitis pathogens (*Staph. aureus* and *Strep. agalactiae*; Godkin and Leslie 1993). Our bulk tank sampling strategy (collecting a single sample) differed from previous work describing the bacteriology of milk from [bedded-packBP](#) farms, where four or five consecutive bulk tank milk pickups were collected and then pooled for analysis (Barberg et al., 2007b; Shane et al., 2010; Lobeck et al., 2012). We acknowledge that analysis of a single BTM sample in the current study comes with limitations. Bacterial groups traditionally considered to be primarily environmental in origin (non-ag. *Strep.*, *Staph* spp., coliforms), may enter BTM from cows with an intramammary infection, but also may originate from non-specific contamination (teat and udder skin, bedding, manure, or other environmental sources; Elmoslemany et al., 2009). Furthermore, a single bulk tank sample does not give insight into long-term, consistent patterns of a particular farm’s milk quality as is possible from repeated BTM samplings (Jayarao and Wolfgang, 2003).

With the financial constraints of research on commercial dairy farms, the limitations inherent in performing analysis of a single bulk tank milk sample from each farm were a trade-off for the ability to get a picture of milk quality on a larger number of farms included in the study.

Udder health outcomes included in the current study (percent cows with elevSCS, percent cows with chronSCS, percent cows with newSCS, BTSCC, and average avg. SCS) ~~did not differ~~ were not statistically different-significantly between facility types. For BTSCC, BP were numerically lower than the other two facility types; the difference in BTSCC for BP vs. FS and BP vs. TS equated to an increase of 34,628 and 28,105 cells/mL, respectively, which could amount to an important difference at the bulk tank level quality premiums under some systems. NewSCS was also numerically lowest for BP, at 2.3% lower than FS and more modestly at 0.43% lower than TS. Ruegg and Pantoja- (2013) propose a benchmark of having <8% of cows developing a new subclinical mastitis infections per month, whereas Schukken et al. (2003) suggest <10%. Using either of these figures, a difference of 2.3% between facility types would be substantial. BP had numerically lower chronSCS in comparison to FS (1.5%), but were equivalent to TS herds in this respect. As an industry benchmark is to have <10% of cows with chronic subclinical mastitis infections carrying over month to month (U. Minnesota Extension Dairy Team), the numeric difference seen between BP and FS for this outcome may be biologically important. ElevSCS was numerically lowest for TS herds, while FS herds had a higher proportion of cows with an  $SCS \geq 4.0$  on current test compared to BP farms. The relative magnitude of the difference for these estimates when compared to BP may be biologically significant (1.8% for FS, -2.4% for TS), as a suggested goal for herds is to have a <15% prevalence for cows with subclinical mastitis (Ruegg and Pantoja, 2013). With regards to numeric difference in avg. SCS, BP farms performed slightly better than FS, and were equivalent

to TS. The increase in estimated avg. SCS for FS equates to an increase of roughly 16,250 cells/mL at the cow level, which represents a slight to modest increase in SCC. Although some numeric differences for outcomes were observed in the current study between facility types for newSCS, chronSCS, elevSCS, and avg. LS, given the proportionately large standard errors for all estimates, interpretation of the effect of facility type for these outcomes is challenging.

Although some previous work has found BTSCC to be elevated for CBP farms (425,000 cells/mL over all four seasons, Black et. al 2013; 325,000 cells/mL during summer, Barberg et. al 2007b), other groups have also found udder health and milk quality measures on ~~bedded packBP~~ farms are similar to farms using more traditional facility types. Specifically, subclinical mastitis prevalence levels did not differ between ~~compost bedded packsCBP~~ and two types of ~~freestallsFS~~ housing in Minnesota and South Dakota, where the percent of cows in a herd with an SCC on test day  $\geq 200,000$  cells/mL was 33.4, 26.8, and 26.8% for ~~compost bedded packsCBP~~, cross-ventilated ~~freestallsFS~~, and naturally-vented ~~freestallsFS~~ (Lobeck et al., 2011). Eckelkamp et. al 2016a found no significant difference in subclinical mastitis prevalence in CBP vs. sand-bedded ~~freestallsFS~~ in Kentucky with a history of low BTSCC (21.8 and 19.4%, respectively), as well as no difference in BTSCC between the two facility types (229,582 and 205,131 cells/mL, respectively). Subclinical mastitis prevalence was 27.7% for 12 CBP farms in Minnesota (Barberg et. al 2007b), which may be more representative of the general population of ~~bedded packBP~~ farms in that state as there were no inclusion criteria around maintaining a low SCC ~~prior~~ to the start of the study. The prevalence of subclinical mastitis for herds in the current study (~~26% for bedded packs~~) is similar to previous work in the U.S. In contrast, Fávero et. al (2015) found a much higher prevalence of subclinical mastitis (43.8%) and percent new

infections (20.9%) for three ~~bedded-pack~~BP farms in Brazil than our study (26 and 7% respectively, for ~~the three~~3 bedded-packsBP with available data).

STD 150-day milk production did not differ significantly between facility type in the current study. Cows on BP farms numerically made slightly more milk than those in TS, and were equivalent to those in FS. This increase of 1.7 pounds for BP over TS represents roughly 3% of the average STD 150-day milk production for herds in the study, which is a relatively modest increase in milk production. However, the comparatively large standard errors for both STD 150-day milk estimates make it difficult to interpret the effect of facility type for this metric. ~~This aligns with~~ previous research which has found no significant differences in various production metrics of cows housed on ~~bedded-packs~~BP vs. in ~~freestall~~FS barns (Lobeck et al., 2011; Eckelkamp et al., 2016a; Costa et al., 2018). Varying production metrics for cows housed on ~~bedded-packs~~BP have been reported previously (kg/cow/day, fat-corrected milk/cow/day, average L/cow/day, ME-305, rolling herd average, energy-corrected milk), preventing direct comparisons of milk production between the ~~bedded-packs~~BP in the current study and other work. Additionally, many variables play a role in determining milk production (nutrition, breed, seasonality, DIM), so teasing out the effect of facility type alone on production in an observational study is difficult. However, as Leso et. al (2020) point out, ~~the~~ “results in the literature indicate that high levels of milk production are possible in CBP.” As ~~bedded-packs~~BP potentially improve cow comfort, one may even expect greater milk production than in more traditional housing systems (Calamari et al., 2009; Ruud et al., 2010).

~~Our finding no difference in~~ The two udder hygiene measures were not statistically different between the three facility types. TS farms had numerically higher proportion of dirty

udders and avg. udder hygiene score, while FS and BP systems were equivalent. However, interpretation of these numerical differences is difficult, given that the standard errors for all four estimates are large relative to the coefficient estimates. ~~is in accordance with p~~Previous work, ~~which~~ found that cow hygiene on ~~bedded pack~~BP systems ~~is~~was comparable to traditional facility types in the Upper Midwestern U.S., Southeastern U.S., and Brazil (Barberg et al., 2007b; Shane et al., 2010; Black et al., 2013; Eckelkamp et al., 2016b; a; Costa et al., 2018; Adkins et al., 2022; Andrade et al., 2022). Black (2013) and Eckelkamp (2016a) reported that increased pack moisture allows wet bedding material and manure to adhere more easily to animals, meaning that cow hygiene is highly dependent on conditions of the ~~bedded pack~~BP. This sentiment was echoed by the ~~bedded pack~~BP producers in the current study, who shared that keeping their cows clean during periods of wet or humid weather could be a challenge. However, all ~~bedded packs~~BP in the current study had an average udder hygiene score of less than 2.5, and the farm with the lowest mean average udder hygiene score overall was a ~~bedded pack farm~~BP. Although Cook (2002) as pointed out the challenges of comparing dairy cattle hygiene between different facility types, we chose to focus on gathering observations of udder hygiene. The relationship between udder hygiene and health is well-studied, and was a tractable observation to make during non-grazing season farm visits where individual animals were often roaming freely in a pen, or confined in a ~~tiestall~~TS barn.

**Objective 2: Analysis of farm management factors (non-facility) associated with bulk tank milk quality, udder health, milk production, and udder hygiene scores for all farms combined**

~~As results from the multivariable models exploring the relationship between facility type and outcomes of interest suffered from limited statistical power due to small sample sizes, the~~



focus of the discussion will be on trends that emerged from the univariate analysis which combined all 21 farms.

One finding emerging from the univariate analysis combining all 21 farms from this work is that farms with deeper bedding had more favorable udder hygiene metrics (deeper bedding begets cleaner cows). When comparing farms that housed cows with a deep bedding system (deeply-bedded stalls or a bedded-packBP) to those that housed cows on stalls with a smaller amount of bedding (over a mattress or concrete surface), the deeply-bedded systems tended to have better hygiene scores. This agrees with previous observational field studies of freestallsFS barns, including: Cook et al. 2016 (prevalence of dirty udders was 13% lower for farms using deep bedding vs. stalls with mats), de Vries et al. 2015 (deep-bedding vs. mat/mattress reduced the likelihood of a cow having a dirty hindquarter by half), and Robles et al. 2020 (farms with mattress-based stalls had a higher prevalence of cows with dirty upper legs/flanks vs. those using a deep bedding system, often inorganic sand). In contrast, an experimental study looking at the effect of bedding depth in tiestallsTS over 28-day periods found no difference between leg, flank, and udder hygiene of cows using deeply-bedded stalls (14 cm) and the control treatment (2-3 cm; Wolfe et al., 2018).

Beyond comparing udder hygiene of cows housed on a deep-bedding system to cows that were not, there was a linear association between bedding depth (depth of bedded-packBP, depth of bedding in freestallsFS and tiestallsTS) and hygiene score. As the measured height of bedding got deeper (height of bedded-packBP, or amount of bedding material in stall), cows tended to have cleaner udders. To the best of our knowledge, work exploring this direct relationship between measured bedding depth and hygiene is limited to a single study by de Vries et al. 2015, who found no relationship between prevalence of dirty hindquarters and three different



freestallsFS bedding height groups (<0.56 cm, 0.56–1.75 cm, >1.75 cm). ~~In our study,~~ this relationship between bedding depth and udder hygiene was especially strong for ~~bedded packsBP~~ in particular, ~~despite the~~ ~~although limited~~ sample size ~~was limited at~~ ~~of~~ five herds. To the best of our knowledge, this specific association has not previously been explored for ~~bedded packBP~~ herds. There is ~~clearly~~ opportunity for future research looking at this relationship between increased amount of bedding used in deep-bedded systems (or more deeply-bedded stalls) and the benefit of improved udder hygiene and milk quality.

Multiple measures of udder health in this study were associated with udder hygiene, in accordance with the well-supported tenet that better cow hygiene is associated with better milk quality. ~~(cleaner cows beget better milk).~~ The association between hygiene and udder health has been well-documented, both at the cow level (for IMI presence: de Pinho et al. 2012; for SCS/SCC: Reneau et al. 2005; Dohmen et al. 2010; and Sant'anna et al. 2011; for both SCS and IMI: Schreiner and Ruegg, 2003) and at the herd-level (BTSCC: Barkema et al. 1998; new IMI rate: Cook et al. 2002; average herd SCC, incidence clinical mastitis, and % new high SCC: Dohmen et al. 2010). Of particular relevance to the current work, a study carried out on three ~~bedded packBP~~ farms in Brazil found the odds of a new case of subclinical mastitis (SCC  $\geq 200,000$  cells/mL) and of a cow having subclinical mastitis on test day increased 32% and 16% for each one-unit increase in leg cleanliness score, respectively (Fávero et al., 2015). Curiously, although leg cleanliness score was associated with both mastitis outcomes on Brazilian ~~bedded packsBP~~, udder hygiene score was not.

A third interesting finding to emerge from the univariate regression results is that farms using deeper bedding had better milk quality outcomes ~~(deeper bedding begets better milk).~~ Although there is an established recommendation of 15 cm for deep bedding of ~~freestallsFS~~

(Bickert, 2000; Cook, 2002) ~~and this depth appears to be based on optimizing cow comfort in deep-bedded freestalls with no reference to udder hygiene or health. There is very limited work study~~ exploring ideal bedding material depth for ~~tiestall~~<sup>TS</sup> barns (Tucker and Weary, 2004; Tucker et al., 2009), ~~and this is this work is again solely~~ focused on the important concern of cow comfort. As ~~stated is the experience of the authors, and is stated elsewhere~~ in a literature review by McPherson (2020), "very little research has investigated the effect of bedding depth on cow cleanliness" or considerations around udder health outcomes. It is likely that the effect seen in the current work of deeper bedding and better udder health outcomes is mediated through the presumed causal pathway of (1) deeper bedding leading to improved hygiene, and (2) improved hygiene resulting in better udder health. ~~Although recommending a particular depth may prove difficult as there are many contributing factors which are particular to a producer's barn and bedding source~~ Even still, the opportunity ~~still~~ exists for research exploring optimal stall bedding depths of different organic materials ~~in tiestall barns~~ with a focus on mastitis and udder health outcomes. ~~It may be that recommending a particular depth of bedding to use for different types of organic material would not prove feasible, as the ideal amount would vary with many factors particular to a producer's barn and bedding source (type of stall surface, presence/type of stall mat used, type of organic material, particle size, compressibility, percent dry matter, etc.).~~

~~to the goal was to~~ Recent previous work has exclusively focused on describing bedded packs that are actively managed for aerobic composting (Leso et al., 2020). Leso et al. contrasted composting bedded packs managed with daily cultivation with conventional static bedded packs, such as straw yards, noting the reduced cow cleanliness and increased risk of mastitis associated with the latter. While bedded pack systems are not common for housing lactating cows in Vermont, both composting and static systems are used (Andrews et al., 2021). This infrequent

use of bedded packs in our state created a challenge for enrolling ten herds using this kind of system in our observational study. Despite this limitation, by including bedded pack farms managed in a variety of ways, the current work sheds light on a broader spectrum of options used within this loose-housing system. Our current study shows that farms can achieve excellent milk quality using either a static or aerobically composting bedded pack system for indoor housing, e.g., three of the five bedded pack farms had a BTSCC  $\leq 99,000$  cells/mL, and the remaining two were  $\leq 160,000$  cells/mL. Furthermore, the lowest BTSCC in the study (54,000 cells/mL) was a static bedded pack farm using woodchips and straw. This low BTSCC was not just from selectively dumping milk from high SCC cows; this farm also had the lowest overall % cows with elevated SCS (8.6%; data not shown).

As for any observational study, there is the potential for bias to have influenced the observed results. Most importantly, participating herds were not a random sample of organic farms in the state, possibly resulting in selection bias. Participating herds were a convenience sample of a subset who responded to our initial survey in Winter 2018-2019 ([source population](#)). [The potential exists that producers who volunteered to participate in the current study are systematically different in some way with regards to their management practices compared to the general population of organic farms in Vermont.](#) In 2021, there were 147 organic dairy farms in Vermont selling milk, with an average herd size of 87 cows making 6,627 kg milk/cow/year (USDA, 2022). Herds in the current study were slightly smaller, averaging 65 cows per farm, but with higher-producing cows (7,828 kg milk/cow/year, estimated from captured DHIA records). [For comparison, the average dairy cow in the U.S. produced an average of 10,926 kg of milk in 2022 \(Progressive Dairy, 2017\). It may be interesting to acknowledge that organic cows on average produce less milk \(Stiglbauer et al., 2013\), and with decreased milk production comes](#)

decreased susceptibility to mastitis (Grohn, 2000). This relationship may in part explain the relatively low prevalence of mastitis occurring on these farms in comparison to the general population of dairy farms. The potential exists that producers who volunteered to participate in the current study are systematically more progressive or somehow different in their management practices than the general population of organic farms in Vermont. Additionally, cross-sectional studies are unable to demonstrate causality for associations presented between management practices and outcomes. However, these limitations are inherent to every observational study, and all attempts were made to control for potential confounding with the multivariable models presented.

One Perhaps the biggest limitation of the current study is the small number of farms in each facility type. As state agencies had been promoting the use of bedded pack BP systems for years in Vermont, we had anticipated it would be feasible to enroll 10 farms using this system to house their lactating animals. This turned out not to be the case; the Winter 2018-2019 survey showed that many dairy farms were instead using these systems for non-lactating animals (heifers, dry cows; Andrews et al. 2021). Furthermore, the COVID-19 pandemic precluded resumption of the study in Spring 2020, limiting the number of farms included to herds sampled in 2019, and not all farms had DHIA data for every outcome of interest. A related limitation is that well-established mastitis control practices ~~were (i.e., teat dipping, forestripping, using separate towels for individual cows)~~ were widely adapted by participating herds, so we were unable to analyze associations between certain practices and BTM quality, udder health, and hygiene. A large body of work exists showing consistent udder health benefits from using these and other practices, so lack of association between these fundamental mastitis control practices and desirable outcomes in the current study should not be taken as evidence that they provide no

benefit. As group sizes for each facility type were limited, we would caution against making inferences from the findings beyond the source population of this study. The potential still exists for future studies with a larger number of farms enrolled to further characterize milk quality and udder health on bedded-pack BP systems in the Northeastern US. Studies enrolling a larger number of bedded-pack farms by covering enrolling farms from a larger geographic area, future studies may be able to enroll a larger number of BP farms, increasing the statistical power needed to may have sufficient power to identify particular management factors which are beneficial on bedded-packs BP specifically.

While BP systems are not common for housing lactating cows in Vermont, farms using this system in the state are using both compost bedded-packs managed with daily cultivation and untilled deep bedded pack systems. As untilled and cultivated bedded pack systems differ in numerous regards (Leso et al., 2020), the initial goal was to enroll enough farms using each type and treat them as separate groups in the analysis. As the relatively small number of BP used in our state to house lactating dairy cattle created a challenge for enrolling ten herds using this kind of system in our observational study, it was necessary to combine both types of system in order to achieve our objective of describing udder hygiene, milk quality, and udder health on these loose-housing systems deeply-bedded with organic material. While we acknowledge that grouping them together is not ideal, this diversity is a reflection of how the target population (small-medium, pasture-based organic dairy farms) are actually using them in the Northeastern U.S. (Benson, 2012). Despite this limitation, including bedded pack farms managed in a variety of ways sheds light on a broader spectrum of options used within this loose-housing system. Our current study demonstrates that farms can achieve excellent milk quality using either an untilled, deep bedded pack system or an aerobically composting bedded pack system for indoor housing;

three of the five BP farms had a BTSCC  $\leq 99,000$  cells/mL, and the remaining two were  $\leq 160,000$  cells/mL. Furthermore, the lowest BTSCC in the study (54,000 cells/mL) was a static BP farm using woodchips and straw. This low BTSCC was not just from selectively dumping milk from high-SCC cows; this farm also had the lowest overall % cows with elevated SCS (8.6%; data not shown).

~~Bedded pack~~BP systems have a number of advantages for producers considering updating their facilities, including a smaller initial investment when compared to a new ~~freestall~~FS or ~~tiestall~~TS barn (Barberg et al., 2007a; Janni et al., 2007; Black et al., 2013), although the cost year-over-year for bedding is substantial (Shane et al., 2010). Bedded packs are designed for cow comfort (Barberg et al., 2007b; Bewley et al., 2012), and prevalence of lameness, foot, and leg injuries in these systems has been found to be less than ~~tiestall~~TS and ~~freestall~~FS barns (Barberg et al., 2007b; Lobeck et al., 2011; Burgstaller et al., 2016). Lastly, manure management and environmental stewardship is a top concern for both dairy producers and the general public (Holly et al., 2018). Anecdotally, the ~~five~~BP producers enrolled in the study were pleased with their systems of manure management, viewing their used bedding material and manure as a valuable soil amendment and an integral part of their nutrient management plan. Bedded pack systems decrease the amount of liquid manure waste when compared to conventional barns, and the used bedding with manure is more easily composted before use as a soil amendment. As ~~composted aged-bedded~~ pack material is drier before it is spread on fields, it poses less of a risk for run-off into waterways, increases soil infiltration of nutrients, and creates flexibility around timing of manure application to fields (Rushmann). ~~With no obvious disadvantages for udder health or hygiene when properly managed on farms with excellent milking hygiene practices already in place, bedded packs~~Bedded packs may be ~~a an~~

936 especially good housing option for small, pasture-based farms in the Northeastern U.S. -when  
937 properly managed on farms with excellent milking hygiene practices already in place. However,  
938 more research is needed to confirm that udder health, milk quality, udder hygiene and milk  
939 production compares favorably to more traditional housing systems.

## 941 Conclusion

942 For five of the six studied udder health and production metrics, and both udder hygiene  
943 measures, numerically BP either performed slightly better or were equivalent in comparison to  
944 the most commonly-used facility types for organic dairy cows in Vermont. However, the  
945 relatively large standard errors for most of these estimates make it difficult to rule out  
946 biologically important effects of facility type for these outcomes. This is likely a result of the  
947 small group size for each facility type. Bedded packs may therefore be a viable option for  
948 pasture-based herds looking for a loose-housing system, but future studies enrolling larger  
949 number of farms using each type of housing are needed to more definitively explore these  
950 relationships. Bedded pack systems did not differ significantly in their milk quality, udder health,  
951 udder hygiene measures, or milk production, as compared to the more commonly used indoor  
952 housing systems (freestall or tiestall) for organic cows in Vermont. Bedded packs can therefore  
953 be considered as a viable option for pasture-based herds looking for a loose-housing system.

954 Findings from the secondary analysis of results found evidence of the well-supported tenets that  
955 better cow hygiene is associated with better milk quality, and farms with deeper bedding had  
956 more favorable udder hygiene metrics. Additionally, farms using deeper bedding had better milk  
957 quality outcomes, which may likely be mediated through improved hygiene resulting in better  
958 udder health outcomes.



**Acknowledgements**

This project was funded by the USDA NIFA OREI project 2018-51300-28561. We would like to thank all the organic dairy producers who agreed to participate in this study, for giving us their time and allowing us to collect samples from their farms. We would also like to thank Jennifer Timmerman and the Laboratory for Udder Health at Veterinary Diagnostic Lab (University of Minnesota), as well as the laboratory staff at St. Alban’s Cooperative/Dairy Farmers of America, for their advice and analyses of bulk tank milk samples collected in the study. The authors have not stated any conflicts of interest.

**References**

Adkins, P. R. F., L. M. Placheta, M. R. Borchers, J. M. Bewley, and J. R. Middleton. 2022. Distribution of staphylococcal and mammaliicoccal species from compost-bedded pack or sand-bedded freestall dairy farms. *J Dairy Sci* 105(7):6261-6270.

Albino, R. L., J. L. Taraba, M. I. Marcondes, E. A. Eckelkamp, and J. M. Bewley. 2018. Comparison of bacterial populations in bedding material, on teat ends, and in milk of cows housed in compost bedded pack barns *J. Animal Production Science* 58(9):1686-1691.

Andrade, R. R., I. F. F. Tinôco, F. A. Damasceno, G. Ferraz, L. Freitas, C. F. S. Ferreira, M. Barbari, F. J. F. Baptista, and D. J. R. Coelho. 2022. Spatial distribution of bed variables, animal welfare indicators, and milk production in a closed compost-bedded pack barn with a negative tunnel ventilation system. *J Therm Biol* 105:103111.

Andrews, T., C. E. Jeffrey, R. E. Gilker, D. A. Neher, and J. W. Barlow. 2021. Design and implementation of a survey quantifying winter housing and bedding types used on Vermont organic dairy farms. *J. Dairy Sci.* 104(7):8326-8337.

Astiz, S., F. Sebastian, O. Fargas, M. Fernández, and E. Calvet. 2014. Enhanced udder health and milk yield of dairy cattle on compost bedding systems during the dry period: A comparative study. *Livestock Science* 159:161-164.



- 987 Barberg, A., M. Endres, and K. Janni. 2007a. Compost Dairy Barns in Minnesota: A Descriptive  
988 Study. *Applied Engineering in Agriculture* 23:231-238.
- 989 Barberg, A. E., M. I. Endres, J. A. Salfer, and J. K. Reneau. 2007b. Performance and welfare of  
990 dairy cows in an alternative housing system in Minnesota. *J Dairy Sci* 90(3):1575-1583.
- 991 Barkema, H. W., Y. H. Schukken, T. J. Lam, M. L. Beiboer, G. Benedictus, and A. Brand. 1998.  
992 Management practices associated with low, medium, and high somatic cell counts in bulk milk.  
993 *J. Dairy Sci* 81(7):1917-1927.
- 994 Barkema, H. W., M. A. von Keyserlingk, J. P. Kastelic, T. J. Lam, C. Luby, J. P. Roy, S. J.  
995 LeBlanc, G. P. Keefe, and D. F. Kelton. 2015. Invited review: Changes in the dairy industry  
996 affecting dairy cattle health and welfare. *J Dairy Sci* 98(11):7426-7445.
- 997 Benson, A. F. 2012. Consider deep pack barns for cow comfort and manure management.  
998 Accessed March 18, 2024. Cornell University, Ithaca, NY.  
999 [https://smallfarms.cornell.edu/2012/04/consider-deep-pack-barns-for-cow-comfort-and-manure-](https://smallfarms.cornell.edu/2012/04/consider-deep-pack-barns-for-cow-comfort-and-manure-management/)  
1000 [management/](https://smallfarms.cornell.edu/2012/04/consider-deep-pack-barns-for-cow-comfort-and-manure-management/).
- 1001 Bewley, J., J. Taraba, G. Day, R. Black, and F. Damasceno. 2012. Compost Bedded Pack Barn  
1002 Design: Features and Management Considerations. University of Kentucky Cooperative  
1003 Extension Service Publication ID.
- 1004 Bewley, J. M., L. M. Robertson, and E. A. Eckelkamp. 2017. A 100-Year Review: Lactating  
1005 dairy cattle housing management. *J. Dairy Sci.* 100(12):10418-10431.
- 1006 Bickert, W. G., B. Holmes, K. A. Janni, D. Kammel, R. Stowell, and J. M. Zulovich. 2000. Dairy  
1007 freestall housing and equipment. Pages 27–45 in *Designing Facilities for the Milking Herd*. 7<sup>th</sup>  
1008 ed., Mid-West Plan Service, Iowa State University, Ames.
- 1009 Black, R. A., J. L. Taraba, G. B. Day, F. A. Damasceno, and J. M. Bewley. 2013. Compost  
1010 bedded pack dairy barn management, performance, and producer satisfaction. *J Dairy Sci*  
1011 96(12):8060-8074.
- 1012 Black, R. A., J. L. Taraba, G. B. Day, F. A. Damasceno, M. C. Newman, K. A. Akers, C. L.  
1013 Wood, K. J. McQuerry, and J. M. Bewley. 2014. The relationship between compost bedded pack  
1014 performance, management, and bacterial counts. *J Dairy Sci* 97(5):2669-2679.
- 1015 Burgstaller, J., J. Raith, S. Kuchling, V. Mandl, A. Hund, and J. Kofler. 2016. Claw health and  
1016 prevalence of lameness in cows from compost bedded and cubicle freestall dairy barns in  
1017 Austria. *The Veterinary Journal* 216.
- 1018 Calamari, L., F. Calegari, and L. Stefanini. 2009. Effect of different free stall surfaces on  
1019 behavioural, productive and metabolic parameters in dairy cows. *Applied Animal Behaviour*  
1020 *Science* 120:9-17.
- 1021 Condas, L. A. Z., J. De Buck, D. B. Nobrega, D. A. Carson, S. Naushad, S. De Vlieghe, R. N.  
1022 Zadoks, J. R. Middleton, S. Dufour, J. P. Kastelic, and H. W. Barkema. 2017. Prevalence of non-

- 1023 aureus staphylococci species causing intramammary infections in Canadian dairy herds. *J Dairy*  
1024 *Sci* 100(7):5592-5612.
- 1025 Cook, N. B. 2002. Influence of Barn Design on Dairy Cow Hygiene, Lameness and Udder  
1026 Health. American Association of Bovine Practitioners Conference Proceedings: 97-103.
- 1027 Cook, N. B., T. B. Bennett, and K. V. Nordlund. 2005. Monitoring Indices of Cow Comfort in  
1028 Free-Stall-Housed Dairy Herds. *J. Dairy Sci.* 88(11):3876-3885.
- 1029 Cook, N. B., J. P. Hess, M. R. Foy, T. B. Bennett, and R. L. Brotzman. 2016. Management  
1030 characteristics, lameness, and body injuries of dairy cattle housed in high-performance dairy  
1031 herds in Wisconsin. *J Dairy Sci* 99(7):5879-5891.
- 1032 Costa, J. H. C., T. A. Burnett, M. A. G. von Keyserlingk, and M. J. Hötzel. 2018. Prevalence of  
1033 lameness and leg lesions of lactating dairy cows housed in southern Brazil: Effects of housing  
1034 systems. *J Dairy Sci* 101(3):2395-2405.
- 1035 The Dairyland Initiative: School of Veterinary Medicine, University of Wisconsin-Madison.  
1036 Housing Module: Adult Cow Housing, Bedded Packs. University of Wisconsin-Madison.  
1037 Accessed March 18, 2024. [https://thedairylandinitiative.vetmed.wisc.edu/home/housing-](https://thedairylandinitiative.vetmed.wisc.edu/home/housing-module/adult-cow-housing/bedded-pack/)  
1038 [module/adult-cow-housing/bedded-pack/](https://thedairylandinitiative.vetmed.wisc.edu/home/housing-module/adult-cow-housing/bedded-pack/).
- 1039 de Pinho Manzi, M., D. B. Nóbrega, P. Y. Faccioli, M. Z. Troncarelli, B. D. Menozzi, and H.  
1040 Langoni. 2012. Relationship between teat-end condition, udder cleanliness and bovine  
1041 subclinical mastitis. *Res Vet Sci* 93(1):430-434.
- 1042 De Visscher, A., S. Piepers, F. Haesebrouck, and S. De Vliegher. 2016. Intramammary infection  
1043 with coagulase-negative staphylococci at parturition: Species-specific prevalence, risk factors,  
1044 and effect on udder health. *J Dairy Sci* 99(8):6457-6469.
- 1045 De Visscher, A., S. Piepers, F. Haesebrouck, K. Supre, and S. De Vliegher. 2017. Coagulase-  
1046 negative *Staphylococcus* species in bulk milk: Prevalence, distribution, and associated subgroup-  
1047 and species-specific risk factors. *J Dairy Sci* 100(1):629-642.
- 1048 de Vries, M., E. A. Bokkers, C. G. van Reenen, B. Engel, G. van Schaik, T. Dijkstra, and I. J. de  
1049 Boer. 2015. Housing and management factors associated with indicators of dairy cattle welfare.  
1050 *Prev Vet Med* 118(1):80-92.
- 1051 Dohmen, W., F. Neijenhuis, and H. Hogeveen. 2010. Relationship between udder health and  
1052 hygiene on farms with an automatic milking system. *J Dairy Sci* 93(9):4019-4033.
- 1053 Eberhart, R. J. 1984. Coliform Mastitis. *Veterinary Clinics of North America: Large Animal*  
1054 *Practice* 6(2):287-300.
- 1055 Eckelkamp, E. A., J. L. Taraba, K. A. Akers, R. J. Harmon, and J. M. Bewley. 2016a. Sand  
1056 bedded freestall and compost bedded pack effects on cow hygiene, locomotion, and mastitis  
1057 indicators. *Livestock Science* 190:48-57.

- 1058 Eckelkamp, E. A., J. L. Taraba, K. A. Akers, R. J. Harmon, and J. M. Bewley. 2016b.  
1059 Understanding compost bedded pack barns: Interactions among environmental factors, bedding  
1060 characteristics, and udder health. *Livestock Science* 190:35-42.
- 1061 Elmoslemany, A. M., G. P. Keefe, I. R. Dohoo, and B. M. Jayarao. 2009. Risk factors for  
1062 bacteriological quality of bulk tank milk in Prince Edward Island dairy herds. Part 1: overall risk  
1063 factors. *J Dairy Sci* 92(6):2634-2643.
- 1064 Endres, M., K. Janni. 2021. Compost-bedded pack barns for dairy cows. University of Minnesota  
1065 Extension. Minneapolis, MN. Accessed March 18, 2024. [https://extension.umn.edu/dairy-](https://extension.umn.edu/dairy-milking-cows/compost-bedded-pack-barns-dairy-cows#a-wall-borders-the-pack-727910)  
1066 [milking-cows/compost-bedded-pack-barns-dairy-cows#a-wall-borders-the-pack-727910](https://extension.umn.edu/dairy-milking-cows/compost-bedded-pack-barns-dairy-cows#a-wall-borders-the-pack-727910).
- 1067 Fairchild, T. P., B. J. McArthur, J. H. Moore, and W. E. Hylton. 1982. Coliform Counts in  
1068 Various Bedding Materials. *J. Dairy Sci.* 65(6):1029-1035.
- 1069 Fávero, S., F. V. R. Portilho, A. C. R. Oliveira, H. Langoni, and J. C. F. Pantoja. 2015. Factors  
1070 associated with mastitis epidemiologic indexes, animal hygiene, and bulk milk bacterial  
1071 concentrations in dairy herds housed on compost bedding. *Livestock Science* 181:220-230.
- 1072 Ferraz, P. F. P., G. A. e. S. Ferraz, L. Leso, M. Klopčič, M. Barbari, and G. Rossi. 2020.  
1073 Properties of conventional and alternative bedding materials for dairy cattle. *J. Dairy Sci.*  
1074 103(9):8661-8674.
- 1075 Fregonesi, J. A. and J. D. Leaver. 2001. Behaviour, performance and health indicators of welfare  
1076 for dairy cows housed in strawyard or cubicle systems. *Livestock Production Science* 68(2):205-  
1077 216.
- 1078 Fregonesi, J. A. and J. D. Leaver. 2002. Influence of space allowance and milk yield level on  
1079 behaviour, performance and health of dairy cows housed in strawyard and cubicle systems.  
1080 *Livestock Production Science* 78(3):245-257.
- 1081 Godkin, M. A. and K. E. Leslie. 1993. Culture of bulk tank milk as a mastitis screening test: A  
1082 brief review. *Can Vet J* 34(10):601-605.
- 1083 Grohn, Y. 2000. Milk Yield and Disease: Towards Optimizing Dairy Herd Health and  
1084 Management Decisions. *Bovine Practice* 34:32-40.
- 1085 Heins, B. J., L. S. Sjostrom, M. I. Endres, M. R. Carillo, R. King, R. D. Moon, and U. S. Sorge.  
1086 2019. Effects of winter housing systems on production, economics, body weight, body condition  
1087 score, and bedding cultures for organic dairy cows. *J Dairy Sci* 102(1):706-714.
- 1088 Hogan, J. and K. L. Smith. 2012. Managing environmental mastitis. *Vet Clin North Am Food*  
1089 *Anim Pract* 28(2):217-224.
- 1090 Hogan, J. S. and K. L. Smith. 1997. Bacteria counts in sawdust bedding. *J Dairy Sci* 80(8):1600-  
1091 1605.

- 1092 Hogan, J. S., K. L. Smith, K. H. Hoblet, D. A. Todhunter, P. S. Schoenberger, W. D. Hueston, D.  
 1093 E. Pritchard, G. L. Bowman, L. E. Heider, B. L. Brockett, and H. R. Conrad. 1989. Bacterial  
 1094 Counts in Bedding Materials Used on Nine Commercial Dairies. *J. Dairy Sci.* 72(1):250-258.
- 1095 Hogan, J. S., D. G. White, and J. W. Pankey. 1987. Effects of teat dipping on intramammary  
 1096 infections by staphylococci other than *Staphylococcus aureus*. *J Dairy Sci* 70(4):873-879.
- 1097 Holly, M. A., P. J. Kleinman, R. B. Bryant, D. L. Bjorneberg, C. A. Rotz, J. M. Baker, M. V.  
 1098 Boggess, D. K. Brauer, R. Chintala, G. W. Feyereisen, J. D. Gamble, A. B. Leytem, K. F. Reed,  
 1099 P. A. Vadas, and H. M. Waldrip. 2018. Short communication: Identifying challenges and  
 1100 opportunities for improved nutrient management through the USDA's Dairy Agroecosystem  
 1101 Working Group. *J Dairy Sci* 101(7):6632-6641.
- 1102 Janni, K., M. Endres, J. Reneau, and W. Schoper. 2007. Compost Dairy Barn Layout and  
 1103 Management Recommendations. *Applied Engineering in Agriculture* 23(1):97-102.
- 1104 Jayarao, B. M., S. R. Pillai, A. A. Sawant, D. R. Wolfgang, and N. V. Hegde. 2004. Guidelines  
 1105 for monitoring bulk tank milk somatic cell and bacterial counts. *J Dairy Sci* 87(10):3561-3573.
- 1106 Jayarao, B. M. and D. R. Wolfgang. 2003. Bulk-tank milk analysis. A useful tool for improving  
 1107 milk quality and herd udder health. *Vet Clin North Am Food Anim Pract* 19(1):75-92, vi.
- 1108 Klaas, I. C. and R. N. Zadoks. 2018. An update on environmental mastitis: Challenging  
 1109 perceptions. *Transbound Emerg Dis* 65 Suppl 1:166-185.
- 1110 KoboCollect: Simple, Robust and Powerful Tools for Data Collection. 2019  
 1111 <http://www.kobotoolbox.org>.
- 1112 Leso, L., M. Barbari, M. A. Lopes, F. A. Damasceno, P. Galama, J. L. Taraba, and A. Kuipers.  
 1113 2020. Invited review: Compost-bedded pack barns for dairy cows. *J Dairy Sci* 103(2):1072-1099.
- 1114 Lobeck, K., M. Endres, K. Janni, S. Godden, and J. Fetrow. 2012. Environmental Characteristics  
 1115 and Bacterial Counts in Bedding and Milk Bulk Tank of Low Profile Cross-Ventilated, Naturally  
 1116 Ventilated, and Compost Bedded Pack Dairy Barns. *Applied Engineering in Agriculture* 28:117-  
 1117 128.
- 1118 Lobeck, K. M., M. I. Endres, E. M. Shane, S. M. Godden, and J. Fetrow. 2011. Animal welfare  
 1119 in cross-ventilated, compost-bedded pack, and naturally ventilated dairy barns in the upper  
 1120 Midwest. *J Dairy Sci* 94(11):5469-5479.
- 1121 McPherson, S. E. and E. Vasseur. 2020. Graduate Student Literature Review: The effects of  
 1122 bedding, stall length, and manger wall height on common outcome measures of dairy cow  
 1123 welfare in stall-based housing systems. *J Dairy Sci* 103(11):10940-10950.
- 1124 Neave, F. K., F. H. Dodd, and R. G. Kingwill. 1966. A method of controlling udder disease. *Vet*  
 1125 *Rec* 78(15):521-523.

- 1126 Neher, D. A., T. D. Andrews, T. R. Weicht, A. Hurd, and J. W. Barlow. 2022. Organic Farm  
1127 Bedded Pack System Microbiomes: A Case Study with Comparisons to Similar and Different  
1128 Bedded Packs. *Dairy*. doi:10.3390/dairy3030042.
- 1129 O'Connor, A. M., J. M. Sargeant, I. R. Dohoo, H. N. Erb, M. Cevallos, M. Egger, A. K. Ersbøll,  
1130 S. W. Martin, L. R. Nielsen, D. L. Pearl, D. U. Pfeiffer, J. Sanchez, M. E. Torrence, H. Vigre, C.  
1131 Waldner, and M. P. Ward. 2016. Explanation and Elaboration Document for the STROBE-Vet  
1132 Statement: Strengthening the Reporting of Observational Studies in Epidemiology-Veterinary  
1133 Extension. *J Vet Intern Med* 30(6):1896-1928.
- 1134 Pankey, J. W., R. L. Boddie, and S. C. Nickerson. 1985. Efficacy evaluation of two new teat dip  
1135 formulations under experimental challenge. *J Dairy Sci* 68(2):462-465.
- 1136 Pankey, J. W., E. E. Wildman, P. A. Drechsler, and J. S. Hogan. 1987. Field trial evaluation of  
1137 premilking teat disinfection. *J Dairy Sci* 70(4):867-872.
- 1138 Pankey, J. W. 1989. Premilking Udder Hygiene. *J. Dairy Sci.* 72(5):1308-1312.
- 1139 Pantoja, J. C. F., D. J. Reinemann, and P. L. Ruegg. 2009. Associations among milk quality  
1140 indicators in raw bulk milk. *J. Dairy Sci.* 92(10):4978-4987.
- 1141 Patel, K., S. M. Godden, E. Royster, B. A. Crooker, J. Timmerman, and L. Fox. 2019.  
1142 Relationships among bedding materials, bedding bacteria counts, udder hygiene, milk quality,  
1143 and udder health in US dairy herds. *J. Dairy Sci.* 102(11):10213-10234.
- 1144 Peeler, E. J., M. J. Green, J. L. Fitzpatrick, K. L. Morgan, and L. E. Green. 2000. Risk Factors  
1145 Associated with Clinical Mastitis in Low Somatic Cell Count British Dairy Herds. *J. Dairy Sci.*  
1146 83(11):2464-2472.
- 1147 Piessens, V., E. Van Coillie, B. Verbist, K. Supre, G. Braem, A. Van Nuffel, L. De Vuyst, M.  
1148 Heyndrickx, and S. De Vliegher. 2011. Distribution of coagulase-negative *Staphylococcus*  
1149 species from milk and environment of dairy cows differs between herds. *J Dairy Sci* 94(6):2933-  
1150 2944.
- 1151 Pol, M. and P. L. Ruegg. 2007. Relationship between antimicrobial drug usage and antimicrobial  
1152 susceptibility of gram-positive mastitis pathogens. *J Dairy Sci* 90(1):262-273.
- 1153 Progressive Dairy. 2022. U.S. Dairy Statistics. Accessed March 19, 2024.  
1154 <https://www.progressivepublish.com/downloads/2023/general/2022-pd-stats-highres.pdf>.
- 1155 Quirk, T., L. K. Fox, D. D. Hancock, J. Capper, J. Wenz, and J. Park. 2012. Intramammary  
1156 infections and teat canal colonization with coagulase-negative staphylococci after postmilking  
1157 teat disinfection: species-specific responses. *J Dairy Sci* 95(4):1906-1912.
- 1158 R Development Core Team. 2023. R: A Language and Environment for Statistical Computing. R  
1159 Foundation for Statistical Computing, Vienna, Austria.

- 1160 Reneau, J. K., A. J. Seykora, B. J. Heins, M. I. Endres, R. J. Farnsworth, and R. F. Bey. 2005.  
 1161 Association between hygiene scores and somatic cell scores in dairy cattle. *J Am Vet Med Assoc*  
 1162 227(8):1297-1301.
- 1163 Rinehart, L. and A. Baier. 2011. U.S. Department of Agriculture; National Center for  
 1164 Appropriate Technology (NCAT), National Organic Program. Pasture for Organic Ruminant  
 1165 Livestock: Understanding and Implementing the National Organic Program (NOP) Pasture Rule.  
 1166 Accessed Oct. 30, 2023. [https://www.ams.usda.gov/sites/default/files/media/NOP-](https://www.ams.usda.gov/sites/default/files/media/NOP-UnderstandingOrganicPastureRule.pdf)  
 1167 [UnderstandingOrganicPastureRule.pdf](https://www.ams.usda.gov/sites/default/files/media/NOP-UnderstandingOrganicPastureRule.pdf).
- 1168 Robles, I., D. F. Kelton, H. W. Barkema, G. P. Keefe, J. P. Roy, M. A. G. von Keyserlingk, and  
 1169 T. J. DeVries. 2020. Bacterial concentrations in bedding and their association with dairy cow  
 1170 hygiene and milk quality. *Animal* 14(5):1052-1066.
- 1171 Rowbotham, R. F. and P. L. Ruegg. 2016a. Associations of selected bedding types with  
 1172 incidence rates of subclinical and clinical mastitis in primiparous Holstein dairy cows. *J Dairy*  
 1173 *Sci* 99(6):4707-4717.
- 1174 Rowbotham, R. F. and P. L. Ruegg. 2016b. Bacterial counts on teat skin and in new sand,  
 1175 recycled sand, and recycled manure solids used as bedding in freestalls. *J Dairy Sci* 99(8):6594-  
 1176 6608.
- 1177 Ruegg, P. L. 2009. Management of mastitis on organic and conventional dairy farms. *J Anim Sci*  
 1178 87(13 Suppl):43-55.
- 1179 Ruegg, P. L. and J. C. F. Pantoja. 2013. Understanding and using somatic cell counts to improve  
 1180 milk quality. *Irish Journal of Agricultural and Food Research* 52(2):101-117.
- 1181 Rushmann, R. University of Wisconsin-Madison; Division of Extension: Agriculture Water  
 1182 Quality. Managing manure to reduce negative water quality impacts: Composting on Wisconsin  
 1183 farms. Accessed Aug. 1, 2023. [https://agwater.extension.wisc.edu/articles/managing-manure-to-](https://agwater.extension.wisc.edu/articles/managing-manure-to-reduce-negative-water-quality-impacts-composting-on-wisconsin-farms/)  
 1184 [reduce-negative-water-quality-impacts-composting-on-wisconsin-farms/](https://agwater.extension.wisc.edu/articles/managing-manure-to-reduce-negative-water-quality-impacts-composting-on-wisconsin-farms/).
- 1185 Ruud, L. E., K. E. Bøe, and O. Osterås. 2010. Associations of soft flooring materials in free stalls  
 1186 with milk yield, clinical mastitis, teat lesions, and removal of dairy cows. *J Dairy Sci*  
 1187 93(4):1578-1586.
- 1188 Sant'anna, A. C. and M. J. Paranhos da Costa. 2011. The relationship between dairy cow hygiene  
 1189 and somatic cell count in milk. *J Dairy Sci* 94(8):3835-3844.
- 1190 Schreiner, D. A. and P. L. Ruegg. 2002. Effects of tail docking on milk quality and cow  
 1191 cleanliness. *J Dairy Sci* 85(10):2503-2511.
- 1192 Schreiner, D. A. and P. L. Ruegg. 2003. Relationship between udder and leg hygiene scores and  
 1193 subclinical mastitis. *J Dairy Sci* 86(11):3460-3465.
- 1194 Schukken, Y. H., F. J. Grommers, J. A. Smit, D. Vandegheer, and A. Brand. 1989. Effect of  
 1195 freezing on bacteriologic culturing of mastitis milk samples. *J Dairy Sci* 72(7):1900-1906.



- 1196 Schukken, Y. H., D. J. Wilson, F. Welcome, L. Garrison-Tikofsky, and R. N. Gonzalez. 2003.  
1197 Monitoring udder health and milk quality using somatic cell counts. *Vet Res* 34(5):579-596.
- 1198 Shane, E., M. Endres, and K. Janni. 2010. Alternative Bedding Materials for Compost Bedded  
1199 Pack Barns in Minnesota: A Descriptive Study. *Applied Engineering in Agriculture* 26:465-473.
- 1200 Stiglbauer, K. E., K. M. Cicconi-Hogan, R. Richert, Y. H. Schukken, P. L. Ruegg, and M.  
1201 Gamroth. 2013. Assessment of herd management on organic and conventional dairy farms in the  
1202 United States. *J. Dairy Sci.* 96(2):1290-1300.
- 1203 Thurgood, J. M., C. M. Comer, D. J. Flaherty, and M. Kiraly. 2009. Bedded pack management  
1204 system case study. Pages 184–188 in *Proc. 5th National Small Farm Conference*,  
1205 Springfield, IL. Accessed March 18, 2024.  
1206 [https://conferences.illinois.edu/resources/20033/Proceedings\\_8-12-13.pdf](https://conferences.illinois.edu/resources/20033/Proceedings_8-12-13.pdf).
- 1207 Tucker, C. B., D. Weary, M. Keyserlingk, and K. Beauchemin. 2009. Cow comfort in tie-stalls:  
1208 Increased depth of shavings or straw bedding increases lying time. *J. Dairy Sci.* 92:2684-2690.
- 1209 Tucker, C. B. and D. M. Weary. 2004. Bedding on geotextile mattresses: how much is needed to  
1210 improve cow comfort? *J Dairy Sci* 87(9):2889-2895.
- 1211 University of Minnesota Extension Dairy Team. Using DHIA Records to Benchmark Herd SCC.  
1212 Accessed Apr. 1, 2024. [https://qualitycounts.umn.edu/sites/qualitycounts.umn.edu/files/2022-](https://qualitycounts.umn.edu/sites/qualitycounts.umn.edu/files/2022-01/w-mp-5.pdf)  
1213 [01/w-mp-5.pdf](https://qualitycounts.umn.edu/sites/qualitycounts.umn.edu/files/2022-01/w-mp-5.pdf)
- 1214 USDA-NRCS. NRCS Climate-Smart Mitigation Activities. Accessed Dec. 14, 2023.  
1215 [https://www.nrcs.usda.gov/conservation-basics/natural-resource-concerns/climate/climate-smart-](https://www.nrcs.usda.gov/conservation-basics/natural-resource-concerns/climate/climate-smart-mitigation-activities)  
1216 [mitigation-activities](https://www.nrcs.usda.gov/conservation-basics/natural-resource-concerns/climate/climate-smart-mitigation-activities).
- 1217 USDA. 2022. Certified Organic Survey, 2021 Summary. Accessed Nov. 10, 2023.  
1218 [https://downloads.usda.library.cornell.edu/usda-](https://downloads.usda.library.cornell.edu/usda-esmis/files/zg64tk92g/2z10z137s/bn99bh97r/cenorg22.pdf)  
1219 [esmis/files/zg64tk92g/2z10z137s/bn99bh97r/cenorg22.pdf](https://downloads.usda.library.cornell.edu/usda-esmis/files/zg64tk92g/2z10z137s/bn99bh97r/cenorg22.pdf).
- 1220 Ward, W. R., J. W. Hughes, W. B. Faull, P. J. Cripps, J. P. Sutherland, and J. E. Sutherst. 2002.  
1221 Observational study of temperature, moisture, pH and bacteria in straw bedding, and faecal  
1222 consistency, cleanliness and mastitis in cows in four dairy herds. *Vet Rec* 151(7):199-206.
- 1223 Wolfe, T., E. Vasseur, T. J. DeVries, and R. Bergeron. 2018. Effects of alternative deep bedding  
1224 options on dairy cow preference, lying behavior, cleanliness, and teat end contamination. *J Dairy*  
1225 *Sci* 101(1):530-536.
- 1226 Wuytack, A., A. De Visscher, S. Piepers, F. Haesebrouck, and S. De Vliegher. 2020. Fecal non-  
1227 aureus Staphylococci are a potential cause of bovine intramammary infection. *Vet Res* 51(1):32.
- 1228 Zadoks, R. N., L. L. Tikofsky, and K. J. Boor. 2005. Ribotyping of *Streptococcus uberis* from a  
1229 dairy's environment, bovine feces and milk. *Veterinary Microbiology* 109(3):257-265.

1230 Zdanowicz, M., J. A. Shelford, C. B. Tucker, D. M. Weary, and M. A. G. von Keyserlingk. 2004.  
1231 Bacterial Populations on Teat Ends of Dairy Cows Housed in Free Stalls and Bedded with Either  
1232 Sand or Sawdust. J. Dairy Sci. 87(6):1694-1701.

1233

1234 **Tables**

Table 1. Predictors offered to multivariable models for each of the eight different outcomes of interest along with facility type (forced)

Predictor	Level of parameter, if categorical:
Farm demographics/lactating cow housing	
Facility type	Bedded pack; Freestall; Tiestall
Predominant breed	Holstein; Jersey/Other
Herd size (lactating cows)	
Herd size group (lactating cows)	30-55; 56-69; 70-100
Subjective assessment of air quality (producer)	Excellent; Good; Fair/Poor
Subjective assessment of air quality (researcher)	Good; Fair
Age of facility (years)	
Feed supplemental vit. E and selenium	Yes; No
Lactating bedding management practices	
Lying surface for cows <sup>1</sup> (deeply-bedded vs. not)	Deeply-bedded stalls or bedded pack; Stalls with bedding on a mattress or concrete surface
<i>If use shavings/sawdust/woodchips for bedding material:</i>	
Moisture-content	Kiln-dried; Fresh/raw
Bedding amendment (e.g., hydrated lime) used on surface	Yes; No
<i>If facility is freestall or tiestall:</i>	
Freq. adding new bedding to stalls (times per week)	
Freq. scraping stalls (times per week)	
Depth bedding in stalls (cm)	
Mastitis control and milking hygiene practices	
Clip/flame udder hair	Yes; No
Keep record of clinical mastitis events	Always; Sometimes/Temp.; Never
Routinely culture mastitic milk	Always/Sometimes; Never
Routinely culture high somatic cell count cows	Always/Sometimes; Never
Ever perform culture of mastitic cows	Yes; Never culture
Use intramammary product at dry-off (OMRI-listed)	Yes; No
Parenteral supplementation with vit. E and selenium	All lactating cows regularly/ Occasionally as needed; No
Glove use at milking	All milkers consistently; Inconsistently/No



Check for clinical mastitis by noticing abnormal  
cow/abnormal udder and forestripping Yes; No  
Type of milking system used<sup>3</sup> Parlor; Tiestall  
Farm-level udder hygiene metrics  
Average udder hygiene score  
Prop. dirty udders (%; udder hygiene score  $\geq 3$ )  
<sup>1</sup> If freestall or tiestall, producer asked if used deeply-bedded stalls  
<sup>2</sup> OMRI: Organic Materials Review Institute  
<sup>3</sup> One freestall farm used an automated milking system

1235

1236

Table 2. Objective 1: Descriptive and univariable results for bulk tank milk aerobic culture outcomes by facility type [median (range)]. *P*-value is for Kruskal-Wallis test by facility type grouping

Bacteria group (cfu/mL)	Overall (n = 21)	Bedded packs (n = 5)	Tiestalls (n = 10)	Freestalls (n = 6)	<i>P</i> -value
<i>Staph. spp.</i>	65 (0-665)	40 (0-130)	85 (15-665)	67.5 (5-125)	0.62
<i>Strep.</i> and strep-like orgs.	45 (10-1250)	35 (10-80)	167.5 (20-1250)	32.5 (25-260)	0.10
<i>Staph. aureus</i>	30 (0-320)	0 (0-30)	47.5 (0-320)	42.5 (0-100)	0.19
Coliforms	0 (0-5)	0 (0-5)	0 (0-5)	0 (0-5)	0.82

1237

1238

1239

Table 3. Objective 1: Descriptive results for milk quality, udder health and production outcomes by facility type [mean (95%CI)]

Outcome	Overall	Bedded packs	Tiestalls	Freestalls
BTSCC (log <sub>10</sub> cells/mL)	n = 21	n = 5	n = 10	n = 6
	5.13 (5.06-5.20)	5.00 (4.84-5.17)	5.14 (5.05-5.23)	5.21 (5.09-5.33)
% newly elevated SCS <sup>1</sup>	n = 19	n = 3	n = 10	n = 6
	5.7 (4.2-7.3)	7.0 (2.8-11.2)	5.4 (3.0-7.8)	5.6 (3.0-8.3)
% chronically elevated SCS <sup>1</sup>	n = 19	n = 3	n = 10	n = 6
	13.6 (11.2-16.1)	14.5 (5.4-23.7)	14.3 (11.9-16.7)	12.0 (6.7-17.3)
% SCS $\geq 4.0$ current test <sup>1</sup>	n = 19	n = 3	n = 10	n = 6
	24.9 (21.6-28.3)	26.0 (12.6-39.3)	25.4 (22.1-28.6)	23.7 (16.9-30.5)
Avg. SCS <sup>2</sup>	n = 20	n = 4	n = 10	n = 6
	2.44 (2.26-2.62)	2.38 (1.84-2.91)	2.45 (2.31-2.59)	2.50 (2.00-2.93)

Standardized 150-day milk (pounds) <sup>3</sup>	n = 18	n = 4	n = 8	n = 6
	50.0 (45.7-54.3)	46.9 (39.8-53.9)	49.4 (43.1-55.7)	53.0 (43.5-62.5)

<sup>1</sup> DHIA data not available for 2 bedded pack farms  
<sup>2</sup> DHIA data not available for 1 bedded pack farm  
<sup>3</sup> DHIA data not available for 1 bedded pack farms and 2 tiestall farms

1240

For Peer Review

1241

Table 4. Objective 1: Final multivariable models describing the relationship between facility type (forced) and milk quality, udder health, production, and udder hygiene outcomes

Outcome	Explanatory variable	Group (sample size)	Coefficient estimate (SE)	P-value
BTSCC (log <sub>10</sub> cells/mL)	Intercept		4.8 (0.15)	
	Facility type (forced)	Freestall (n = 6)	0.19 (0.09)	0.05
		Tiestall (n = 10)	0.16 (0.08)	0.07
		Bedded pack (n = 5)	Ref.	Ref.
	Herd size	All herds (n = 21)	0.003 (0.002)	0.15
% newly elevated SCS	Intercept		-1.6 (2.5)	
	Facility type (forced)	Freestall (n = 6)	2.3 (2.2)	0.33
		Tiestall (n = 10)	0.43 (1.9)	0.82
		Bedded pack (n = 3)	Ref.	Ref.
	Use bedding amendment	Yes (n = 5)	3.9 (1.8)	0.05
		No (n = 14)	Ref.	Ref.
Subjective assessment air quality (researcher)		Good (n = 14)	3.6 (1.5)	0.04
		Fair (n = 5)	Ref.	Ref.
Glove use at milking <sup>1</sup>		Never/Inconsistently (n = 9)	2.0 (1.3)	0.17
		Always (n = 9)	Ref.	Ref.
Clinical mastitis events record keeping		Never kept records (n = 6)	4.4 (1.8)	0.03
		Sometimes/Temporarily kept records (n = 6)	1.1 (1.6)	0.52
		Always kept records (n = 7)	Ref.	Ref.
% chronically elevated SCS	Intercept		5.3 (3.6)	
	Facility type (forced)	Freestall (n = 6)	1.5 (4.6)	0.75
		Tiestall (n = 10)	-0.08 (3.5)	0.98
		Bedded pack (n = 3)	Ref.	Ref.
	Feed supplemental vit. E and selenium <sup>2</sup>	Yes (n = 11)	2.1 (2.8)	0.48
		No (n = 7)	Ref.	Ref.

	Use bedding amendment	Yes (n = 5)	5.7 (3.4)	0.12
		No (n = 14)	Ref.	Ref.
	Clip/flame udder hair	Yes (n = 5)	-6.3 (3.1)	0.07
		No (n = 14)	Ref.	Ref.
% udder hygiene scores $\geq 3$		Herds with available data (n = 19)	17.0 (6.1)	0.02
% SCS $\geq 4.0$ current test	Intercept		0.85 (10.6)	
	Facility type (forced)			
		Freestall (n = 6)	1.8 (5.7)	0.75
		Tiestall (n = 10)	-2.4 (5.3)	0.66
		Bedded pack (n = 3)	Ref.	Ref.
	Use bedding amendment	Yes (n = 5)	8.0 (4.2)	0.07
		No (n = 14)	Ref.	Ref.
	Mean hygiene	Herds with available data (n = 19)	9.8 (4.7)	0.06
Avg. SCS				20
	Intercept		0.93 (0.44)	
	Facility type (forced)			
		Freestall (n = 6)	0.38 (0.21)	0.09
		Tiestall (n = 10)	0.03 (0.19)	0.86
		Bedded pack (n = 4)	Ref.	Ref.
	Use intramammary product at dry-off (OMRI-listed)	Yes (n = 5)	-0.30 (0.16)	0.08
		No (n = 15)	Ref.	Ref.
	Use bedding amendment	Yes (n = 5)	0.52 (0.16)	0.007
		No (n = 15)	Ref.	Ref.
	Parenteral supplementation vit. E/selenium	Regularly or occasionally (n = 9)	-0.36 (0.14)	0.02
		No supplementation (n = 11)	Ref.	Ref.
	Mean hygiene	Herds with available data (n = 20)	0.64 (0.19)	0.005
Standardized 150-day milk (pounds)				18
	Intercept		41.2 (6.1)	
	Facility type (forced)			
		Freestall (n = 6)	-0.06 (7.0)	0.99
		Tiestall (n = 8)	-1.7 (6.6)	0.80
		Bedded pack (n = 4)	Ref.	Ref.
	Parenteral supplementation vit. E/selenium	Regularly or occasionally (n = 7)	7.0 (5.2)	0.20

	No supplementation (n = 11)	Ref.	Ref.
Culture high SCC cows	Always/Sometimes (n = 8)	9.3 (5.9)	0.14
	Never (n = 10)	Ref.	Ref.
Herd size grp. (lact. cows)	70-100 (n = 8)	-0.18 (7.3)	0.98
	56-69 (n = 5)	10.3 (6.2)	0.12
	30-55 (n = 5)	Ref.	Ref.
% udder hygiene scores $\geq 3$			
Intercept		0.32 (0.08)	
Facility type (forced)	Freestall (n = 6)	0.002 (0.11)	0.99
	Tiestall (n = 10)	0.17 (0.10)	0.12
	Bedded pack (n = 5)	Ref.	Ref.
Avg. udder hygiene score			
Intercept		2.3 (0.17)	
Facility type (forced)	Freestall (n = 6)	-0.04 (0.21)	0.84
	Tiestall (n = 10)	0.33 (0.19)	0.11
	Bedded pack (n = 5)	Ref.	Ref.
Check for clinical mastitis by noticing abnormal cow/abnormal udder and forestripping	Yes (n = 8)	-0.25 (0.16)	0.14
	No (n = 13)	Ref.	Ref.

<sup>1</sup> One farm used automatic milking system

<sup>2</sup> One farm unable to provide response

Table 5. Objective 2: Selected models of univariate analysis identifying (non-facility type) factors unconditionally associated with milk quality, udder health, production, and udder hygiene outcomes at  $P < 0.20$

Outcome	Explanatory Variable	Group (sample size)	Coefficient estimate (SE)	P-value	Intercept
BTSCC (log <sub>10</sub> cells/mL)					
Model 1	Lying surface	Mattress or concrete (n = 13)	0.12 (0.07)	0.12	5.1
		Deep bedding (n = 8)	Ref.	Ref.	
Model 2	Depth of bedding in stalls (cm) <sup>1</sup>	Tiestalls and freestalls (n = 15)	-0.02 (0.01)	0.11	5.2
Model 3	Glove use at milking <sup>2</sup>	Never/Inconsistently (n = 9)	0.10 (0.07)	0.19	5.1
		Always (n = 11)	Ref.	Ref.	
% newly elevated SCS <sup>3</sup>					
Model 4	Glove use at milking	Never/Inconsistently (n = 9)	2.83 (1.7)	0.11	4.3
		Always (n = 9)	Ref.	Ref.	
Model 5	Depth of bedding in stalls (cm) <sup>1</sup>	Tiestalls and freestalls (n = 15)	-0.62 (0.24)	0.02	8.3
% chronically elevated SCS <sup>3</sup>					
Model 6	Clip/flame udder hair	Yes (n = 5)	-4.31 (2.9)	0.16	14.8
		No (n = 14)	Ref.	Ref.	
Model 7	% udder hygiene scores ≥3	Herds with available data (n = 19)	12.7 (6.1)	0.05	8.6
Model 8	Avg. udder hygiene score	Herds with available data (n = 19)	6.39 (3.1)	0.05	-1.2
% SCS ≥ 4.0 current test <sup>3</sup>					
Model 9	Depth of bedding in stalls (cm) <sup>1</sup>	Tiestalls and freestalls (n = 15)	-1.2 (0.42)	0.01	30
Model 10	% udder hygiene scores ≥3	Herds with available data (n = 19)	13.6 (8.5)	0.13	19.6
Model 11	Avg. udder hygiene score	Herds with available data (n = 19)	7.7 (4.3)	0.09	7.1
Average SCS <sup>4</sup>					
Model 12	Parenteral supplementation vit. E and selenium	Regularly or occasionally (n = 9)	-0.27 (0.18)	0.15	2.6
		No supplementation (n = 11)	Ref.	Ref.	
Model 13	Use intramammary product at dry-off (OMRI-listed)	Yes (n = 5)	-0.29 (0.21)	0.18	2.5
		No (n = 15)	Ref.	Ref.	
Model 14	Depth of bedding in stalls (cm) <sup>1</sup>	Tiestalls and freestalls (n = 15)	-0.05 (0.03)	0.10	2.6

Model 15	% udder hygiene scores $\geq 3$	Herds with available data (n = 20)	0.75 (0.45)	0.12	2.1
Model 16	Avg. udder hygiene score	Herds with available data (n = 20)	0.39 (0.23)	0.11	1.5
Standardized 150-day milk (pounds) <sup>5</sup>					
Model 17	Parenteral supplementation vit. E and selenium	Regularly or occasionally (n = 7)	9.0 (4.5)	0.06	46.5
		No supplementation (n = 11)	Ref.	Ref.	
Model 18	Herd size	Herds with available data (n = 18)	0.26 (0.14)	0.07	33.1
% udder hygiene scores $\geq 3$					
Model 19	Depth of bedded pack (m)	Bedded pack herds (n = 5)	-0.5 (0.06)	0.004	0.97
Model 20	Lying surface	Mattress or concrete (n = 13)	0.17 (0.08)	0.06	0.30
		Deep bedding (n = 8)	Ref.	Ref.	
Model 21	Depth of bedding in stalls (cm) <sup>1</sup>	Tiestalls and freestalls (n = 15)	-0.02 (0.02)	0.13	0.54
Avg. udder hygiene score					
Model 22	Depth of bedded pack (m)	Bedded pack herds (n = 5)	-0.96 (0.15)	0.008	3.4
Model 23	Lying surface	Mattress or concrete (n = 13)	0.33 (0.16)	0.06	2.1
		Deep bedding (n = 8)	Ref.	Ref.	
Model 24	Depth of bedding in stalls (cm) <sup>1</sup>	Tiestalls and freestalls (n = 15)	-0.06 (0.03)	0.07	2.6

<sup>1</sup> Stall bedding depth for freestalls and tiestalls bedded with wood shavings or sawdust

<sup>2</sup> One farm used automatic milking system

<sup>3</sup> DHIA data available for n = 19 herds.

<sup>4</sup> DHIA data available for n = 20 herds.

<sup>5</sup> DHIA data available for n = 18 herds.

1243

1244

1245

**Interpretive summary**

Previous studies reported bedded packs improve cow welfare and comfort and have advantages for manure management, soil health, and water quality. Consensus is lacking on whether bulk tank milk quality, udder health, udder hygiene and milk production are compromised on bedded packs. In an observational study of 21 organic dairies in Vermont during the non-grazing season (November-May), bedded packs were similar to tiestalls and freestalls in their bulk tank milk quality, udder health, udder hygiene and milk production. For producers considering a transition from tiestalls, bedded packs may be a viable option for dairy cattle housing in the Northeastern US.

**Running head:**

Milk quality and udder hygiene on VT organic dairies

**Relationship Between Facility Type and Bulk Tank Milk Bacteriology, Udder Health, Udder Hygiene, and Milk Production on Vermont Organic Dairy Farms**

Caitlin E. Jeffrey,<sup>1</sup> Tucker Andrews<sup>2</sup>, Sandra M. Godden<sup>3</sup>, Deborah A. Neher<sup>2</sup>, John W. Barlow<sup>1</sup>

<sup>1</sup> Department of Animal and Veterinary Sciences, University of Vermont, Burlington, VT 05405

<sup>2</sup> Department of Plant and Soil Science, University of Vermont, Burlington, VT 05405

<sup>3</sup> Department of Veterinary Population Medicine, College of Veterinary Medicine, University of Minnesota, St. Paul, MN 55108.

Corresponding author: John Barlow



Department of Animal and Veterinary Sciences,  
202 Terrill Building,  
University of Vermont,  
Burlington, VT 05405  
Phone: 802-656-1395  
Email: [john.barlow@uvm.edu](mailto:john.barlow@uvm.edu)

## Abstract

The primary objective of this cross-sectional observational study on was to determine whether bulk tank milk quality, udder health, udder hygiene and milk production outcomes were associated with facility type on organic dairies. A secondary objective was to identify other management-related risk factors associated with bulk tank milk quality, udder health, udder hygiene, and milk production on organic dairy herds in Vermont. We aimed to enroll 40 farms, to compare herds using the two most common housing systems (freestalls, tiestalls) with those using a bedded pack, for organic dairy cattle in the state during the non-grazing season (typically November-May). Variation in bedded pack management practices was observed, with two general styles, cultivated bedded packs and untilled deep bedded packs. Due to the limited number of herds using bedded packs to house lactating dairy cattle in Vermont, we combined untilled and cultivated bedded pack systems to describe udder hygiene, milk quality, and udder health on these loose-housing systems deeply-bedded with organic material. The study was completed on 21 farms (5 bedded packs, 6 freestalls, 10 tiestalls) before interruption due to the COVID-19 pandemic. Data captured from Dairy Herd Improvement Association records from the test closest to the date of the farm visit

included average somatic cell score (SCS), standardized 150-day milk (pounds), % cows with current high SCS ( $\text{SCS} \geq 4.0$ ), % cows with newly elevated SCS (previous SCS  $< 4.0$  to current  $\geq 4.0$ ), and % cows with chronically elevated SCS ( $\text{SCS} \geq 4.0$  last two tests). Multivariable linear regression models were performed to describe outcomes by facility type, but suffered from limited statistical power due to small group sample sizes. Unconditional comparisons showed that farms grouped by facility type did not differ in metrics captured from Dairy Herd Improvement Association test data, bulk tank milk somatic cell count (BTSCC) and aerobic culture data, or udder hygiene scores. A secondary analysis was conducted using univariate linear regression to identify associations between herd management factors and outcomes for all 21 farms combined. Although not all differences found were statistically significant in this secondary analysis which combined all farms, numeric differences that may be biologically important are reported showing farms with deeper bedding had a lower BTSCC, lower newly elevated SCS, lower elevated current SCS, lower average SCS, and better udder hygiene metrics. Farms with lower mean udder hygiene scores had numerically lower chronically elevated SCS, lower elevated current SCS, and lower average SCS. The current study provides insight on factors affecting bulk tank milk quality, udder health and hygiene measures on organic dairy farms in Vermont. We could not reject the null hypothesis that milk quality and udder health outcomes did not differ by facility type. Bedded packs were comparable to more frequently used indoor housing systems (tiestalls and freestalls). Bedded pack facilities are may be a viable option for confinement housing during the winter non-grazing season for pasture-based herds interested in a loose-housing system in the Northeastern US. More research including a larger number of herds is needed to test this hypothesis.

**Keywords:** Mastitis, organic dairy cattle, housing, bedded pack, milk quality

## Introduction

Mastitis due to environmental pathogens, such as those commonly found in bedding material, has now become the “most common and costly form of mastitis in modern dairy herds” that have implemented standard mastitis control practices limiting the effect of contagious pathogens (Klaas and Zadoks, 2018). Teats of dairy cattle may be in direct contact with bedding materials for 40 to 60% of the day, making this an important potential source of exposure to opportunistic environmental mastitis pathogens (Tucker and Weary, 2004; Cook et al., 2005; Hogan and Smith, 2012). Work exploring how bedding materials relate to a cow’s risk of contracting mastitis has understandably focused on the most frequently used bedding materials and housing systems in the dairy industry. Currently, the most common type of dairy cattle housing for organic farms in Vermont is a tiestall barn, with freestall barns a distant second (Andrews et al., 2021). As consumer opinion about confinement housing of dairy cattle evolves and influences dairy policy, both the dairy industry and consumers are looking to move away from traditional housing systems that restrict cow movement (Barkema et al., 2015). Many smaller-scale organic dairy farmers in Vermont with aging facilities, and especially tiestall barns, may be looking to adopt a bedded pack system on their farms as a form of loose-housing (Andrews et al., 2021).

The term “bedded pack” encompasses a variety of management styles (Bewley et al., 2017), including compost bedded-packs (CBP), which utilize aerobic decomposition to break down a bedding material of fine wood sawdust or shavings, as well as “conventional,” “traditional,” or “deep bedded packs” (Thurgood, 2009; Benson, 2012; Bewley et al., 2017; The Dairyland Initiative, 2024). CBP can vary in depth, frequency and depth of aeration (tilling), type of bedding material used, and in some regions the inclusion of forced air systems to dry the

bedding (Leso et al., 2020). In the Northeastern U.S., some producers are using deep bedded pack systems where large volumes of straw or hay are added daily to an untilled surface in which strata of bedding and waste accumulate throughout the period of time when cows are housed on it (Benson, 2012). Oxygen is retained in the system by the selection of bedding material and the timing of its application (Neher et al., 2022; Thurgood et al., 2009). A number of authors suggest deep bedded pack barns are synonymous with traditional straw yard housing systems (Bewley et al., 2017; Leso et al., 2020; Ferraz et al., 2020). However, we find the deep bedded packs being constructed on dairy farms in the Northeastern U.S. differ from traditional straw yards, in which material is completely removed at approximately monthly intervals and the housing is used year-round (The Dairyland Initiative, 2024; Thurgood et al., 2009; Benson, 2012).

Bedded packs (BP) are perceived to integrate well into Northeastern US pasture-based farm systems, and state and federal agencies in the U.S. are providing financial incentives for dairies to build these structures as part of manure management practices which improve water quality and contribute to soil conservation (USDA-NRCS; Andrews et al., 2021; Thurgood et al., 2009). As interest in BP grows, it is important to better understand milk quality, udder health and hygiene on farms using these housing alternatives. Understanding mastitis risk for cattle housed on BP is especially important for organic dairy farmers, as they have limited effective options for treating intramammary infections (Ruegg, 2009). As mastitis-causing bacteria may thrive in the conditions found in compost bedded-packs (Black et al., 2014), previous work studying mastitis risk and bedding would suggest BP could pose a relatively higher risk for intramammary infections. Loose-housed cows continually add manure to the pack, contributing both pathogenic bacteria (non-*aureus* staphylococci, Wuytak et. al., 2020; *E. coli*, *Klebsiella* spp., and *Enterobacter* spp., Eberhart, 1984; streptococci, Zadoks et al., 2005) and nutrients to the organic

bedding material. Organic bedding material is more likely to have a higher bacterial count than inorganic bedding, such as sand, (Hogan et al., 1989; Rowbotham and Ruegg, 2016b), as it supplies nutrients and moisture which encourages bacterial growth. This could lead to higher concentrations of bacteria on teat skin for cows on BP, because: 1) organic bedding (in general) is inherently associated with a higher number of bacteria on teat ends (Fairchild et al., 1982; Rowbotham and Ruegg, 2016b), and 2) a higher concentration of bacteria in bedding is associated with a higher concentration of bacteria on teat ends (Hogan and Smith, 1997; Zdanowicz et al., 2004; Rowbotham and Ruegg, 2016b). This higher concentration of bacteria on teat ends may put the mammary gland at an increased risk of infection, although limited evidence exists for this relationship (Neave et al., 1966; Pankey, 1989; Rowbotham and Ruegg, 2016a).

Previous work describing mastitis risk and cow hygiene on BP systems includes descriptive studies of CBP (Barberg et al., 2007b; Black et al., 2013; Fávero et al., 2015; Eckelkamp et al., 2016b; Albino et al., 2018; Heins et al., 2019). However, research comparing milk quality and cow hygiene between BP and more traditional housing types has so far been limited to freestalls with sand, which is an uncommon housing type for organic farms in Vermont (Andrews et al. 2021). These include a study comparing CBP and sand-bedded freestalls for farms with a history of low bulk tank somatic cell counts (Eckelkamp et al., 2016a), work describing hygiene and bulk tank milk somatic cell count (BTSCC) for sand-bedded freestalls and CBP (Adkins et al., 2022), and a comparison of CBP and two types of freestall barns (Lobeck et al., 2011). It is unclear whether the herds included in these prior studies were conventionally-managed or organic dairies. To the best of our knowledge, no studies describe and compare bulk tank milk quality, udder health and hygiene for BP and freestall barns on small to midsize organic dairies in the same geographic area.

To better inform organic dairy producers in the Northeastern US, who may be interested in using a BP for housing their cattle during the non-grazing season (i.e., for “winter housing,” typically the months of November-May), we conducted a cross-sectional, observational study on organic dairies in Vermont. This study aimed to quantify bulk tank milk bacteriology, udder health and udder hygiene measures for the two most common indoor housing systems (freestalls, tiestalls) and farms using a BP for organic farms in Vermont. The objectives of this project were to identify whether bulk tank milk quality, udder health and hygiene outcomes differed by facility type, with a view to determining if BP are a viable option for indoor housing of lactating cows in VT during the non-grazing season. We hypothesized that udder health, hygiene, and bulk tank milk bacteriology of BP herds is inferior to that of more traditional housing types, as has been suggested by some previous research (Barberg et al., 2007b; Lobeck et al., 2011). Therefore, our null hypothesis was that there no association between facility type and udder health, hygiene, and bulk tank milk bacteriology on organic dairy farms using BP and other systems for winter housing of lactating cow in Vermont. A secondary objective was to identify other (non-facility) management-related risk factors associated with bulk tank milk quality, udder health, udder hygiene, and milk production for organic VT dairy herds.

## **Materials and Methods**

STROBE-VET (Strengthening the Reporting of Observational Studies in Epidemiology–Veterinary Extension) statement guidelines were followed in the reporting of this study (O'Connor et al., 2016).

### **Herd enrollment and selection**

The source population for this study was the 145 farms that responded to a survey sent to all certified organic dairy farms producing cow milk in Vermont in Winter 2018-2019 (all farms,  $n = 177$ ). Certified organic dairy farms in the United States are required to allow their cows daily access to pasture during the grazing season, and cows must obtain 30% of their dry matter intake from grazing (Rinehart and Baier, 2011). In Vermont and other Northeastern US states, forage is unavailable directly from pasture during winter months and the climate necessitates use of indoor housing. When cows have no access to pasture in the winter non-grazing season, organic farms in Vermont house cows in a variety of indoor facility types. Our previous Winter 2018-2019 industry survey quantified the frequency and diversity of indoor housing and bedding types used by organic dairy farmers in the state when cows were not on pasture, and for the current study farms were recruited from respondents to this survey (Andrews et al., 2021). Dairy farms were eligible for enrollment in the current study if they: 1) responded to the initial survey in the Winter 2018-2019, 2) indicated they met the enrollment criteria of testing with the Dairy Herd Improvement Association (DHIA) at least monthly, 3) milked between 35 and 120 cows, and 4) indicated they would be interested in further participation. Eligible farms were contacted from this source population in Spring 2019 if they responded that they were using one of four categories of bedding/housing combinations for their indoor housing system: 1) freestall (FS) barn bedded with sand, 2) FS barn bedded with shavings or sawdust, 3) tiestall (TS) barn bedded with shavings or sawdust, or 4) BP. The first three housing and bedding combinations are the most frequently used by organic dairies in Vermont to house cows during the non-grazing season, and were compared to BP as they were the housing type of interest for this project. For the purposes of this study, the inclusive term “bedded pack” is used to encompass both CBP and deep bedded packs, and was defined as an enclosed loose housing facility deeply bedded with

185 organic material, in which bedding and waste accumulate throughout the 6–8-month period of  
186 time when cows are housed on it and which is only removed once a year. Both CBP and deep  
187 bedded packs use carbon-rich substrates to create a clean, comfortable surface which allows  
188 animals to move freely. Urine and manure are not removed when bedding material is renewed, in  
189 contrast with other housing systems.

190 A convenience sample of farms was enrolled in Spring 2019 from a list of eligible farms  
191 (grouped by housing/bedding combination) using the phone number or email address provided in  
192 the 2018-2019 survey response. Our aim was to enroll 40 farms for the current study, with 10  
193 farms from each of the four housing/bedding categories described above.

194 Prior to obtaining the 2018-2019 survey results, based on preliminary data collected by  
195 the University of Vermont Center for Sustainable Agriculture Extension group, the study was  
196 designed anticipating that it would be possible to enroll 10 organic Vermont dairies using a BP  
197 as their primary indoor housing system. However, out of the 17 farms from the 2018-2019  
198 survey which indicated at least some use of a BP, one farm was not interested in any further  
199 participation, five did not use DHIA testing, and six only used a BP as a secondary housing  
200 system in conjunction with a TS barn, or cows were only on the pack a few hours a day. Because  
201 the number of farms using BP was fewer than anticipated, the eligibility requirements were  
202 relaxed to include one farm where cows spend the majority (two-thirds) of their time in a BP,  
203 with the remaining time in a TS with wood shavings. Additionally, two BP farms were included  
204 that had limited DHIA information: one farm did not utilize cow-level testing, and cow-level  
205 data for a second farm was limited due to their seasonal lactation schedule. As the number of BP  
206 being used in the state to house lactating dairy cattle was less than anticipated, those that were  
207 enrolled and grouped together utilized a variety of management strategies. Of the 5 enrolled



farms using a BP, two would be classified as “compost bedded-packs,” utilizing aerobic decomposition to break down a bedding material of dry, fine wood sawdust or shavings (The Dairyland Initiative, 2024; Bewley et al., 2017; Endres, 2021). These two farms bedded solely with shavings/sawdust, adding new bedding only as needed, and cultivated the pack twice a day. Two other farms used a “traditional” or “deep bedded pack” system, where large volumes of fresh, dry straw (or poor-quality hay) sufficient to keep cows clean and dry was added daily to a mass of bedding that accumulates over the 6-8 months cows are housed indoors (The Dairyland Initiative, 2024; Thurgood, 2009; Benson, 2012; Bewley et al., 2017). The one remaining farm fell somewhere between these two types of classically defined BP; this farm bedded with straw and woodchips and cultivated every 48 hrs., adding chopped hay and woodchips every time the pack was cultivated. All farms in the study grouped as “bedded packs” shared the qualities of being an enclosed loose housing facility, deeply bedded with organic material (0.9-1.7 meters), which accumulated over the period of time animals were housed indoors and was only removed once a year.

Of the intended 40 herds to be recruited in the study, 21 herds (1 FS bedded with sand, 5 FS bedded with wood shavings/sawdust, 10 TS bedded with wood shavings/sawdust, 5 BP) agreed to participate and farm visits were completed April-May 2019. This study was intended to study cows while they were in their winter (non-grazing months) indoor housing system, so all herds visits were completed before any grazing had begun for the season. Each herd was visited once during the study period. All herds sampled during this period were housing their cows as they would in the non-grazing season. Farm visits were suspended in mid-May 2019 as farms began turning their cows out to pasture, with the intention of resuming in April 2020 to complete the remaining 19 herds. Due to COVID-19 pandemic activity restrictions, the decision was made

to not resume the study, and the final analysis included the 21 herds sampled in 2019. As there was only one farm sampled using a FS facility bedded with sand, the initial plan to group farms by the four housing/bedding combinations specified was abandoned in favor of grouping farms by the three facility types used. The single sand FS was combined with FS bedded with wood shavings/sawdust (FS;  $n = 6$ ), there were 10 TS bedded with wood shavings/sawdust (TS), and 5 BP.

### **Questionnaire administration, sampling, and udder hygiene scoring**

At each farm visit, a questionnaire was administered to collect information about housing and bedding management, as well as other practices on the farm that could impact mastitis risk (Supplemental Data). The study questionnaire was largely adapted from a previously published survey (Stiglbauer et al., 2013), with additional questions specific to the current study. The questionnaire was reviewed by a social scientist experienced in gathering qualitative data and tested before use with herd managers at the University of Vermont teaching dairy. Questions about mastitis risk explored producer concerns about bedding/mastitis risk; mastitis control, identification and record keeping; milking facilities, procedures, and hygiene practices; information about diet, vitamin and mineral supplementation, and water source; typical calving and periparturient practices; and fly control. Questions about housing and bedding management included describing type of housing system used for both lactating and dry cows; classification and description of any bedding material used; and bedding management practices for each housing type used. The questionnaire also collected some basic herd information (production numbers; number of lactating, dry, and youngstock; breed; record-keeping systems). Farms using BP were asked additional questions to gather detailed information about pack construction, management, monitoring practices, and perceptions comparing BP to any previously used

systems. Completion of the questionnaire required 45 minutes on average, ranging from about 30 minutes to 1.5 hours. The questionnaire and interview protocols were registered with the University of Vermont Institutional Review Board (IRB certification 19-0057). The questionnaire was created and administered on a tablet using KoboCollect software (KoboCollect, 2019).

At each farm visit, a bulk tank milk sample collected directly from the top of the tank using a 250-mL sterile single-use vial (Blue Dippas™, Dynalox Products, England) after at least 5 minutes of agitation. Samples were kept on ice in a cooler during transport until they were processed fresh for SCC measurement or were frozen and stored at  $-20^{\circ}\text{C}$  in the laboratory, before being sent to a diagnostic lab for microbiological analysis. An on-farm observation sheet was completed, which collected information about the bulk tank, cow identification, a subjective assessment of air quality, and any outdoor exercise area (Supplemental Data). Additionally, measurements of the housing facilities were recorded for FS and TS where appropriate (stall sizes, pen sizes, bedding depth, stocking density, trainer use), as well as observations about BP when applicable (depth, pen size, and stocking density in  $\text{m}^2$  per animal). Bedding depth of FS and TS was included as a producer reported value in the questionnaire. Bedding depth of BP facilities was measured by forcing a meter stick down to the level of the cement pad or gravel under the pack, where the pack met a cement knee wall, and recording the height of the pack at that point. Udder hygiene scoring was completed by the same researcher at all farms for a minimum of 30 cows on each farm (the first 30 able to be evaluated in a loose pen, or the first 30 encountered in a tiestall). A four-point udder hygiene scoring system was used, where 1 = free of dirt, 2 = slightly dirty (2–10% of surface area), 3 = moderately covered with dirt (10–30% of surface area), and 4 = covered with caked on-dirt (>30% of surface area) (Schreiner and Ruegg,

2002). Animal use for this project was approved by the University of Vermont Institutional Animal Care and Use Committee (IACUC; protocol #PROTO202000089).

### **Herd-level udder health measurements**

Herd-level DHIA test results for the test day closest in time to the farm visit (either preceding or following day of farm visit, whichever was shorter) were captured from the record processing center working with each herd (Lancaster DHIA, Manheim, PA; Dairy One Co-Op. Inc., Ithaca, NY). Information captured included test date, number of lactating cows, standardized 150-day milk production (STD 150-day milk), and test-day average cow-level somatic cell score (SCS). The following udder health measures were also captured from DHIA records: proportion of cows with an  $SCC \geq 200,000$  cells/mL on most recent test day (“elevSCS”), where elevated SCS was defined as a somatic cell score of  $\geq 4.0$ ; the proportion of cows with a newly elevated SCS (“newSCS”), which was defined as a SCS changing from  $< 4.0$  to  $\geq 4.0$  over the last 2 tests; and the proportion of cows with a chronically elevated SCS (“chronSCS”), which was defined as having a SCS  $\geq 4.0$  on the last two tests (Schukken et al., 2003).

### **Bulk tank milk culture and bulk tank somatic cell count measures**

An aliquot of the bulk tank milk sample was stored at  $-4^{\circ}\text{C}$  until it could be transported to the laboratory of a dairy processing plant (St. Alban’s Cooperative/Dairy Farmers of America, St. Albans, VT) within 48 hours of collection for determination of the bulk tank somatic cell count (BTSCC).

Frozen bulk tank milk samples were shipped on ice to the Laboratory for Udder Health (University of Minnesota Veterinary Diagnostic Laboratory, St. Paul) for analysis. Methodology for bulk tank milk cultures at the Laboratory of Udder Health are described elsewhere (Patel et

al., 2019). Briefly, thawed, room-temperature bulk tank milk and a 10-fold dilution of each bulk tank milk sample were plated onto MacConkey, Factor (gram-positive selective agar; University of Minnesota), and Focus (selective for SSLO bacteria; University of Minnesota) media plates and incubated for two days at 37°C. Any lactose-fermenting colonies on MacConkey medium were counted and reported as coliform bacteria. Any  $\beta$ -hemolytic colonies on Focus medium were counted and identified to the species level using a MALDI Biotyper (suspect *Streptococcus agalactiae*). All remaining colonies on Focus medium that were not identified as *Strep. agalactiae* were counted and recorded as streptococci or strep-like organisms (SSLO). Hemolytic colonies on Factor medium were counted and identified to the species level using a MALDI Biotyper (suspect *Staph. aureus*). Any hemolytic colonies with a confidence score  $\geq 2.0$  for *Staph. aureus* were counted and reported as such. Remaining colonies of staphylococci on Factor media (based on colony morphology, catalase reaction, or Gram stain) were counted and reported as *Staph. spp.* Bulk tank samples were also cultured for *Mycoplasma spp.* (0.1 mL milk was swabbed across a Mycoplasma agar plate, then placed in a 7% CO<sub>2</sub> incubator at 37°C for 7 days, after which they were examined for *Mycoplasma spp.* by a trained microbiology technician). For each bulk tank milk sample, total colony-forming units (cfu) per mL were calculated for coliform organisms, *Staph. spp.*, streptococci and strep-like organisms (SSLO), *Staph. aureus*, *Strep. agalactiae*, and *Mycoplasma spp.* The lower threshold of detection for bacteria in this bulk tank milk culture protocol was 5 cfu/mL, and the upper threshold was 62,500 cfu/mL.

### **Data management and analysis**

Bulk tank milk culture results, BTSCC, DHIA test results, farm-level udder hygiene outcomes, questionnaire data, and farm observations were entered into an Excel database

(Microsoft Corp., Redmond, WA). Udder hygiene scores for individual cows were used to calculate two farm-level udder hygiene measures: 1) mean udder hygiene score, and 2) proportion of cows with dirty udders (udder hygiene score  $\geq 3$ ), which were incorporated into the database. This Excel database was then imported into the R Statistical Programming Environment (R Development Core Team, 2023) for data cleaning, checking, and statistical analysis. The distribution of outcome variables was assessed to check for normality using a Shapiro-Wilk test with significance set at  $P \leq 0.05$ , visual assessment of distribution and residuals, skewness, and comparison of the median and mean values. Raw bulk tank somatic cell count (BTSCC) data was  $\log_{10}$  transformed for analyses. Descriptive statistics were calculated to evaluate the distribution of data, data integrity, and to identify missing data. Descriptive statistics generated included description of general herd characteristics and farm traits, lactating cow housing/facilities, lactating cow bedding material/bedding management practices, milking hygiene procedures, and mastitis control practices for all 21 herds included in the study.

*Objective 1. Evaluation of relationships between housing system and measures of milk quality, udder health, udder hygiene and milk production.* As most measures of aerobic culture data were not normally distributed even after log transformation, a Kruskal-Wallis test was used to compare cfu counts of bacteria from bulk tank milk between the three facility types. Statistical significance for this test was declared at  $P \leq 0.05$ . Multiple attempts were made using multivariable analysis to compare the four aerobic culture outcomes for bulk tank milk, but all modeling approaches suffered from over-parametrization even when data was log transformed and were not pursued further.

Independent farm-level predictors from the herd-management questionnaire offered to the multivariable models are described in Table 1. Continuous variables underwent correlation

analysis to identify predictor variables that were highly correlated (correlation coefficient  $\geq 0.60$ ), and unconditional associations among categorical variables were evaluated using a Pearson's chi-squared or Fischer's Exact test as appropriate ( $P \leq 0.05$ ). An ANOVA was used to check for correlation between numeric continuous variables and categorical variables ( $P \leq 0.05$ ). When a categorical variable had multiple groups with a small number of observations in each, groups were combined when biologically reasonable to have all categories of predictor variables contain at least five observations. If any predictor had only one observation in a group and there was no way to combine groups in a logical way, it was excluded from further analysis (but listed in descriptive statistic tables, Supplemental Tables S1-S4).

Univariate linear regression was performed in R using the "lme4" package to investigate the unconditional relationship between the six udder health and production outcomes (BTSCC, avg. SCS, newSCS, elevSCS, chronSCS, STD 150-day milk) and two hygiene outcomes (mean hygiene score, proportion of dirty udders) for each farm and the previously-described herd-level independent variables. The two udder hygiene metrics (proportion dirty udders and average udder hygiene score) were used as both predictor variables (in models for other outcome variables) and outcome variables in models of their own. Any explanatory variable that was unconditionally associated with 1 or more of the outcomes of interest at  $P < 0.20$  was then offered into a multivariable model investigating the relationship between the udder health and production or hygiene outcome and the herd-level predictor variables. If any predictor variables were found to be correlated with each other at the previously described cut-offs, the one with the more highly significant relationship from univariate analysis was offered to the multivariable model when appropriate. The two udder hygiene metrics were highly correlated (derived from the same data), so whichever one had a smaller  $P$ -value from the univariate analysis was chosen

for inclusion in the model-building process. Facility type was forced into these multivariable models, as it was the primary explanatory predictor of interest. A backward stepwise variable selection process was then used, with the least significant variables being removed one by one until all remaining predictors had  $P \leq 0.10$ . Final models were selected based on lowest Akaike information criteria, and an  $F$ -test to compare the final model to the model with facility type as the only predictor (significance declared at  $P \leq 0.05$ ). Overall statistical significance for facility type (the main predictor of interest) was declared at  $P \leq 0.05$ . The multivariable modelling approach described above aimed to investigate the conditional relationship between facility type and the eight outcomes of interest while controlling for different farm management practices, housing characteristics, milking procedures and mastitis control practices.

*Objective 2. Identify other (non-facility) management-related risk factors associated with bulk tank milk quality, udder health, and milk production in organic dairy herds.* After grouping all 21 farms together, we used linear regression in the same manner as described above in Objective 1 to explore associations between the independent predictors described in Table 1 and the six udder health and production outcomes (BTSCC, avg. SCS, newSCS, elevSCS, chronSCS, STD 150-day milk) and two hygiene outcomes (mean hygiene score, proportion of dirty udders). Unconditional relationships between the eight outcome variables and independent predictors are reported for a significance level of  $P \leq 0.20$  for an  $F$ -test, and only for predictor variables with group sizes of at least  $n = 5$ .

### **Power analysis**

A priori sample size calculations were not performed, as group size was determined by the number of organic dairy herds housing lactating cows on BP in our region.



## Results

### Description of study herds

Of the 21 herds enrolled, 5 used a BP, 1 used a FS bedded with sand, 5 used a FS bedded with shavings/sawdust, and 10 used a TS bedded with shavings/sawdust (Supplemental Table S1). The predominant breeds on all farms were Holstein ( $n = 8$  farms), Jersey ( $n = 10$ ), and mixed Holstein-Jersey crosses/other ( $n = 3$ ). The median (mean; range) number of lactating cows was 68 (64.9; 32-99). The median annual rolling herd average milk production for the farms was 6,367 (6,424; 4,082-9,618) kg. Nineteen of the 21 farms tested with DHIA monthly while their cows were in milk, 1 farm tested 5-8 times/year, and 1 tested every other month. On average, DHIA data was captured from a test day 4 days before the farm visit (range: -28 days to +33). The average depth of bedding in the 15 freestalls and tiestalls where producers provided an estimate was 4.5 cm (SD: 3.5 cm; range: 1.3-12.7 cm). The average depth of bedded packs (measured by researchers) was 130 cm (SD: 31; range: 90-170 cm). Detailed descriptions further characterizing study farm management practices and housing characteristics for lactating animals (e.g., laying surface, ventilation, stocking density), and details about bedding material and bedding management practices for lactating animals (e.g., bedding depth, frequency of adding new bedding, manure removal) are provided in Supplemental Tables S1 and S2, respectively. Detailed descriptions of routine milking procedures and mastitis control practices are provided in Supplemental Tables S3 and S4, respectively.

### Description of bulk tank milk quality, udder health measures, milk production, and udder hygiene scores

There was no difference in cfu count between the three facility types for any of the four bacterial groups measured using a nonparametric unconditional comparison (Table 2). None of the 21 bulk tank milk samples were positive for *Strep. agalactiae* or *Mycoplasma* spp. Sixteen of the 21 samples were negative for coliforms on aerobic culture, while 5 farms had a coliform count of 5 cfu/mL. *Staph. aureus* was found in the bulk tank milk from 13/21 herds, with a median (range) cfu/mL of 50 (15-320) when present.

BTSCC, % cows with newly elevated SCS, % cows with chronically elevated SCS, % cows with elevated SCS, avg. SCS, and STD 150-day milk production did not differ by facility type (Table 3).

The overall mean (95% CI) of herd-level udder hygiene scores for all 21 farms was 2.32 (2.16-2.49). The mean hygiene score was 2.2 (1.91-2.44) for BP farms (n = 5), 2.5 (2.24-2.76) for TS farms (n = 10), and 2.15 (1.93-2.37) for FS farms (n = 6). Mean udder hygiene score did not differ by facility type. The overall mean proportion of cows with dirty udders in a herd (udder hygiene score  $\geq 3$ ) was 40% (31-48). The mean proportion of cows with dirty udders (95% CI) was 32% (18-46) for BP farms, 49% (35-62) for TS farms, and 32% (20-44) for FS farms. The proportion of cows with dirty udders did not differ by facility type.

**Objective 1. Analysis of relationship between facility type and measures of bulk tank milk quality, udder health, milk production, and udder hygiene scores**

Final multivariable models are summarized in Table 4. All 21 farms were able to be included in the models for BTSCC, average hygiene score, and proportion of dirty udders. For the models exploring newSCS, chronSCS, and elevSCS, two BP farms did not have available DHIA data (n = 19; group sizes: FS = 6, TS = 10, BP = 3). One BP farm did not have average cow-level SCS data (n = 20; group sizes: FS = 6, TS = 10, BP = 4). For STD 150-day milk, one

BP farm and two TS farms were missing DHIA data ( $n = 18$ ; group sizes: FS = 6, TS = 8, BP = 4). Farms with missing data for a particular outcome were excluded for the analyses of that outcome.

#### *Bulk tank milk quality outcomes*

Variables that were associated at  $P < 0.20$  with BTSCC in univariate analysis included predominant breed, if herds ever performed culture of mastitic milk, glove use, and herd size. The final multivariable included facility type (forced) and herd size. Facility type was not associated with BTSCC in the final model (Table 4).

#### *Udder health outcomes*

Herd size category, use of bedding amendment, air quality as assessed by researcher, glove use at milking, and clinical mastitis record keeping practices were offered to a multivariable model for newSCS. The final multivariable model included facility type (forced), bedding amendment use, air quality, glove use, and mastitis record keeping practices. Facility type was not associated with newSCS in the final model (Table 4).

Variables that were associated at  $P < 0.20$  with chronSCS in univariate analysis included feeding additional supplemental selenium, use of a bedding amendment, clipping/flaming udder hair, and proportion of dirty udders. The final multivariable model included all four variables from univariate analysis, as well as facility type (forced). Facility type was not found to be a significant predictor of the outcome chronSCS (Table 4).

Bedding amendment use and mean hygiene were offered to a multivariable model for elevSCS. Facility type (forced), bedding amendment, and mean hygiene were retained in the final multivariable model. Facility type was not associated with elevSCS in the final model (Table 4).

Feeding additional supplemental selenium, use of bedding amendment, OMRI-listed intramammary product at dry-off, injectable selenium and vitamin E product, and mean hygiene were offered to a multivariable model for herd average SCS. The final multivariable model for avg. SCS included facility type (forced), use of bedding amendment, dry product, injectable selenium, and mean hygiene score. Facility type was not found to be a significant predictor of avg. SCS (Table 4).

#### *Milk production outcome*

Variables that were associated at  $P < 0.20$  with STD 150-day milk included use of injectable selenium and vitamin E product, whether producers cultured high SCC cows, and herd size group. All three variables and facility type (forced) remained in the final multivariable model (Table 4). Facility type was not associated with STD 150-day milk in the final model (Table 4).

#### *Udder hygiene outcomes*

Air quality assessed by researcher was offered to the multivariable model for proportion of dirty udders. The final multivariable model included only facility type (forced), which was not associated with proportion of dirty udders.

Variables that were associated at  $P < 0.20$  with average hygiene score included whether the producer ever cultured quarter milk samples and whether they checked for cases of clinical mastitis by both examining the udder and forestripping. The final multivariable model included facility type (forced), and how the producer checked for clinical mastitis. Facility type was not associated with the outcome of mean udder hygiene (Table 4).

**Objective 2. Analysis of farm management factors (non-facility) associated with bulk tank milk quality, udder health, milk production, and udder hygiene scores for all farms combined**

Selected results of univariate linear regression models identifying management factors beyond facility type which were unconditionally associated with bulk tank milk quality, udder health, milk production and hygiene outcomes for all farms combined ( $n = 21$ ) at  $P < 0.20$  are presented in Table 5. We report the results of these univariate regression models as they may be biologically important, even though many failed to reach threshold for declaring statistical significance at  $P \leq 0.05$ , possibly due to small sample size.

The depth of bedding in stalls for FS and TS herds was unconditionally associated with multiple udder health outcomes. As the depth of bedding in FS and TS herds increased, multiple udder health measures improved, including lower avg. SCS, BTSCC, elevSCS, and newSCS. Similarly, comparing farms where cows were on deep bedding (i.e., grouping all herds reporting deeply-bedded stalls plus BP herds) to herds that had stalls with a smaller amount of bedding on top of a mattress or concrete, farms with deep bedding had a numerically lower BTSCC.

Udder hygiene measures were associated with several udder health outcomes. Higher mean hygiene scores and proportion of udders scored  $\geq 3$  were associated with higher chronSCS, elevSCS, and average SCS. A few specific management practices were also found to be unconditionally associated with udder health outcomes: consistent glove use was associated with lower newSCS and BTSCC, clipping or flaming udders was associated with fewer chronSCS, and both parenteral supplementation of vit. E/selenium and use of an OMRI-listed intramammary product at dry-off were associated with lower average SCS and higher STD 150-day milk.

Both udder hygiene outcomes were unconditionally associated with the same predictors, most of which were related to the depth of bedding for cows. For the five herds using a BP, deeper bedding was associated with lower average hygiene scores and lower proportion of dirty udders. Farms with cows housed on some type of deep bedding (i.e., grouping the three freestalls and tiestalls reporting deeply-bedded stalls, plus the five BP herds) had numerically lower average udder hygiene scores and proportion dirty udders compared to cows on stalls with bedding over a mattress or concrete surface. For the fifteen tiestalls and freestalls reporting bedding depth in stalls increased bedding depth was associated with lower mean udder hygiene score and a numerically lower proportion of dirty udders.

## Discussion

This work presents the results of our observational study exploring the relationship between facility type and udder health and hygiene metrics, BTM quality (SCC and microbiology), and milk production on organic dairy farms in Vermont. The current study is to the authors' knowledge the first direct comparison of milk quality, udder health and udder hygiene on BP farms to both TS and FS herds of similar size and management styles, for a population of entirely small to midsize organic dairy farms. The major objective was to identify if milk quality, udder health and hygiene outcomes were associated with facility type, thereby exploring if BP systems are a viable option for housing in Vermont during the non-grazing season compared to the two most common indoor housing systems in the state (FS, TS). This study is also the first to describe udder health and hygiene on BP in the Northeastern US, which is significant as the performance of these systems can be greatly influenced by climatic factors. As BTM bacteriology, udder health and hygiene metrics, and milk yield did not differ for BP

herds compared to TS and FS herds, there was insufficient evidence to reject our null hypothesis that these metrics would not vary by facility type.

**Objective 1: Comparison of bulk tank milk quality, udder health, milk production, and udder hygiene measures by facility type**

Although there is a substantial body of work describing udder health and milk quality for cows housed in straw yards (Astiz et. al, 2014; Fregonesi and Leaver, 2001; Fregonesi and Leaver, 2002; Ward et. al 2002; Peeler et al. 2000), description of these outcomes in the literature is limited for static deep bedded packs. As such, the focus of the discussion will compare BP in the current study (both static and composting) to the more recent body of work on compost bedded-pack farms. Previous work describing bulk tank milk aerobic culture data for farms using a BP system has primarily been descriptive studies of compost bedded-pack herds (Barberg et al., 2007b; Shane et al., 2010), with one study directly comparing bacterial counts between CBP and FS barns (Lobeck et al., 2012). Although farms in these previous studies used a similar array of bedding materials to those in the current study (wood sawdust, wheat straw by-product, Lobeck et al. 2012; wood sawdust, Barberg et al. 2007; “alternative” organic materials, Shane et al. 2010), the sampling period for these previous works differed from the present in seasonality, compounding the difficulty of direct comparison for milk quality outcomes (Pantoja et al., 2009). Barberg et al., 2007b evaluated milk culture results across the summer months, while Lobeck et al., 2012 sampled year-round; the current study focused solely on sampling during the winter, when organic pasture-based herds are primarily housed inside in Vermont.

The *Staph.* spp. count for the five BP farms included in this study was comparable to previous work describing bulk tank milk quality for CBP in Minnesota during the winter months. Lobeck et al. 2012 found a mean of 26.1 cfu/mL (95% CI: 2-443) and Shane et al. (2010) found

a range of 0-108 cfu/mL for *Staph.* spp. from BTM in the winter months from six CBP farms. Within this highly heterogenous group of bacteria, some species are considered primarily host-adapted (colonizing the skin or udder), while others have been associated with stall surfaces, air, and unused sawdust (Piessens et al., 2011), different facility types (Condas et al., 2017), and environmental contamination and poor teat hygiene at milking time (De Visscher et al., 2016; De Visscher et al., 2017). In general, the use of pre- and post- milking teat dip decreases contamination of bulk tank milk both by commensal skin organisms and environmental contamination at milking time (Hogan et al., 1987, Pankey et al., 1985; Pankey et al., 1987; Quirk et al., 2012). All but one farm in the current study would fall into the “low” category for *Staph.* spp. counts in the BTM (Jayarao et al., 2004), which is consistent with all 21 herds using both pre- and post-dip consistently at milking time.

Streptococci and strep-like organisms (SSLO) counts in BTM for BP in the current study were much lower than those from Minnesota CBP in the winter (98-48,400 cfu/mL, Shane et al. 2010; mean: 911 cfu/mL, 95% CI: 138-6,01, Lobeck et al. 2012). Work from Barberg et al. (2007) describing milk quality on CBP in Minnesota noted that 6 of 12 farms sampled had “high” levels of SSLO. SSLO count did not differ between TS, FS, and BP in the current study. The overall SSLO count for all 21 farms included in the current study was lower than that for the overall *Strep.* count for all facility types studied in Lobeck et al. 2012 (445 cfu/mL, 95% CI: 116-1704). Milking and bedding hygiene practices amongst herds included in the current study may best explain this difference in BTM pathogen profiles (Jayarao and Wolfgang, 2003).

All farms had low levels of coliforms in bulk tank milk, indicating excellent hygiene practices at milking time (Jayarao and Wolfgang, 2003). The low BTM coliform counts for BP in the current study are similar to those found for three CBP farms in Brazil (2.8 cfu/mL; Fávero



et al. 2015). This is in contrast with previous work describing BTM quality for this kind of facility in the U.S. (15-1,128 cfu/mL, Shane et al., 2010; mean: 63.7 cfu/mL, 95% CI: 6-735, Lobeck et al. 2012), although direct comparison of coliform counts between studies may be potentially problematic due to variation in duration of freezer storage (Schukken et al., 1989). Barberg et al. 2007 found that 5 of 12 BP sampled during the summer months had “high” levels of coliforms in BTM, contributing to their conclusion that “special attention to cow preparation procedures at milking time are a must for achieving satisfactory milk quality when cows are housed in compost dairy barns.”

Prevalence of *Staph. aureus* was similar between the five VT BP farms in the current study and the six described in Lobeck et al. 2012 (6.2 cfu/mL, 95% CI: 1.3-30.1). Farm-level prevalence of *Staph. aureus* was also fairly low for BP studied in Shane et al. 2010 (3 of 6 farms BTM negative) and Barberg et al. 2007 (only 1 of 12 farms with a “high” level of *Staph. aureus*). Overall, the population of all 21 farms in the current study had a higher amount of *Staph. aureus* in BTM than the 18 Minnesota farms described in Shane et al. 2010 (median: 30 cfu/mL, range: 0-320; vs. 17.3 cfu/mL, 95% CI: 3.3-91.2). Although it is not clear how many herds included in previous work on BP were certified organic, this higher prevalence of *Staph. aureus* on organic farms in the current study is consistent with work comparing organic and conventional dairy systems (Pol and Ruegg, 2007).

Analysis of a single bulk tank milk sample from a farm is a simple, convenient, and relatively inexpensive way to capture a snapshot of current milk quality and animal health on a farm, and can be a highly specific (albeit poorly sensitive) screening test for major contagious mastitis pathogens (*Staph. aureus* and *Strep. agalactiae*; Godkin and Leslie 1993). Our bulk tank sampling strategy (collecting a single sample) differed from previous work describing the

bacteriology of milk from BP farms, where four or five consecutive bulk tank milk pickups were collected and then pooled for analysis (Barberg et al., 2007b; Shane et al., 2010; Lobeck et al., 2012). We acknowledge that analysis of a single BTM sample in the current study comes with limitations. Bacterial groups traditionally considered to be primarily environmental in origin (non-ag. *Strep.*, *Staph* spp., coliforms), may enter BTM from cows with an intramammary infection, but also may originate from non-specific contamination (teat and udder skin, bedding, manure, or other environmental sources; Elmoslemany et al., 2009). Furthermore, a single bulk tank sample does not give insight into long-term, consistent patterns of a particular farm's milk quality as is possible from repeated BTM samplings (Jayarao and Wolfgang, 2003). With the financial constraints of research on commercial dairy farms, the limitations inherent in performing analysis of a single bulk tank milk sample from each farm were a trade-off for the ability to get a picture of milk quality on a larger number of farms included in the study.

Udder health outcomes included in the current study (percent cows with elevSCS, percent cows with chronSCS, percent cows with newSCS, BTSCC, and avg. SCS) were not statistically different between facility types. For BTSCC, BP were numerically lower than the other two facility types; the difference in BTSCC for BP vs. FS and BP vs. TS equated to an increase of 34,628 and 28,105 cells/mL, respectively, which could amount to an important difference at the bulk tank level quality premiums under some systems. NewSCS was also numerically lowest for BP, at 2.3% lower than FS and more modestly at 0.43% lower than TS. Ruegg and Pantoja (2013) propose a benchmark of having <8% of cows developing a new subclinical mastitis infections per month, whereas Schukken et al. (2003) suggest <10%. Using either of these figures, a difference of 2.3% between facility types would be substantial. BP had numerically lower chronSCS in comparison to FS (1.5%), but were equivalent to TS herds in this respect. As

an industry benchmark is to have <10% of cows with chronic subclinical mastitis infections carrying over month to month (U. Minnesota Extension Dairy Team), the numeric difference seen between BP and FS for this outcome may be biologically important. ElevSCS was numerically lowest for TS herds, while FS herds had a higher proportion of cows with an SCS  $\geq$  4.0 on current test compared to BP farms. The relative magnitude of the difference for these estimates when compared to BP may be biologically significant (1.8% for FS, -2.4% for TS), as a suggested goal for herds is to have a <15% prevalence for cows with subclinical mastitis (Ruegg and Pantoja, 2013). With regards to numeric difference in avg. SCS, BP farms performed slightly better than FS, and were equivalent to TS. The increase in estimated avg. SCS for FS equates to an increase of roughly 16,250 cells/mL at the cow level, which represents a slight to modest increase in SCC. Although some numeric differences for outcomes were observed in the current study between facility types for newSCS, chronSCS, elevSCS, and avg. LS, given the proportionately large standard errors for all estimates, interpretation of the effect of facility type for these outcomes is challenging.

Although some previous work has found BTSCC to be elevated for CBP farms (425,000 cells/mL over all four seasons, Black et. al 2013; 325,000 cells/mL during summer, Barberg et. al 2007b), other groups have found udder health and milk quality measures on BP farms are similar to farms using more traditional facility types. Specifically, subclinical mastitis prevalence levels did not differ between CBP and two types of FS housing in Minnesota and South Dakota, where the percent of cows in a herd with an SCC on test day  $\geq$  200,000 cells/mL was 33.4, 26.8, and 26.8% for CBP, cross-ventilated FS, and naturally-vented FS (Lobeck et al., 2011). Eckelkamp et. al 2016a found no significant difference in subclinical mastitis prevalence in CBP vs. sand-bedded FS in Kentucky with a history of low BTSCC (21.8 and 19.4%, respectively), as

well as no difference in BTSCC between the two facility types (229,582 and 205,131 cells/mL, respectively). Subclinical mastitis prevalence was 27.7% for 12 CBP farms in Minnesota (Barberg et. al 2007b), which may be more representative of the general population of BP farms in that state as there were no inclusion criteria around maintaining a low SCC prior to the start of the study. The prevalence of subclinical mastitis for herds in the current study is similar to previous work in the U.S. In contrast, Fávero et. al (2015) found a much higher prevalence of subclinical mastitis (43.8%) and percent new infections (20.9%) for three BP farms in Brazil than our study (26 and 7% respectively, for 3 BP with available data).

STD 150-day milk production did not differ significantly between facility type in the current study. Cows on BP farms numerically made slightly more milk than those in TS, and were equivalent to those in FS. This increase of 1.7 pounds for BP over TS represents roughly 3% of the average STD 150-day milk production for herds in the study, which is a relatively modest increase in milk production. However, the comparatively large standard errors for both STD 150-day milk estimates make it difficult to interpret the effect of facility type for this metric. Previous research has found no significant differences in various production metrics of cows housed on BP vs. in FS barns (Lobeck et al., 2011; Eckelkamp et al., 2016a; Costa et al., 2018). Varying production metrics for cows housed on BP have been reported previously (kg/cow/day, fat-corrected milk/cow/day, average L/cow/day, ME-305, rolling herd average, energy-corrected milk), preventing direct comparisons of milk production between the BP in the current study and other work. Additionally, many variables play a role in determining milk production (nutrition, breed, seasonality, DIM), so teasing out the effect of facility type alone on production in an observational study is difficult. However, as Leso et. al (2020) point out, “results in the literature indicate that high levels of milk production are possible in CBP.” As BP

potentially improve cow comfort, one may even expect greater milk production than in more traditional housing systems (Calamari et al., 2009; Ruud et al., 2010).

The two udder hygiene measures were not statistically different between the three facility types. TS farms had numerically higher proportion of dirty udders and avg. udder hygiene score, while FS and BP systems were equivalent. However, interpretation of these numerical differences is difficult, given that the standard errors for all four estimates are large relative to the coefficient estimates. Previous work found that cow hygiene on BP systems was comparable to traditional facility types in the Upper Midwestern U.S., Southeastern U.S., and Brazil (Barberg et al., 2007b; Shane et al., 2010; Black et al., 2013; Eckelkamp et al., 2016b; a; Costa et al., 2018; Adkins et al., 2022; Andrade et al., 2022). Black (2013) and Eckelkamp (2016a) reported that increased pack moisture allows wet bedding material and manure to adhere more easily to animals, meaning that cow hygiene is highly dependent on conditions of the BP. This sentiment was echoed by the BP producers in the current study, who shared that keeping their cows clean during periods of wet or humid weather could be a challenge. However, all BP in the current study had an average udder hygiene score of less than 2.5, and the farm with the lowest mean average udder hygiene score overall was a BP. Although Cook (2002) as pointed out the challenges of comparing dairy cattle hygiene between different facility types, we chose to focus on gathering observations of udder hygiene. The relationship between udder hygiene and health is well-studied, and was a tractable observation to make during non-grazing season farm visits where individual animals were often roaming freely in a pen, or confined in a TS barn.

**Objective 2: Analysis of farm management factors (non-facility) associated with bulk tank milk quality, udder health, milk production, and udder hygiene scores for all farms combined**

One finding from the univariate analysis combining all 21 farms is that farms with deeper bedding had more favorable udder hygiene metrics. When comparing farms that housed cows with a deep bedding system (deeply-bedded stalls or a BP) to those that housed cows on stalls with a smaller amount of bedding (over a mattress or concrete surface), the deeply-bedded systems tended to have better hygiene scores. This agrees with previous observational field studies of FS barns, including: Cook et al. 2016 (prevalence of dirty udders 13% lower for farms using deep bedding vs. stalls with mats), de Vries et al. 2015 (deep-bedding vs. mat/mattress reduced the likelihood of a cow having a dirty hindquarter by half), and Robles et al. 2020 (farms with mattress-based stalls had a higher prevalence of cows with dirty upper legs/flanks vs. those using a deep bedding system, often inorganic sand). In contrast, an experimental study looking at the effect of bedding depth in TS over 28-day periods found no difference between leg, flank, and udder hygiene of cows using deeply-bedded stalls (14 cm) and the control treatment (2-3 cm; Wolfe et al., 2018).

Beyond comparing udder hygiene of cows housed on a deep-bedding system to cows that were not, there was a linear association between bedding depth (depth of BP, depth of bedding in FS and TS) and hygiene score. As the measured height of bedding got deeper (height of BP, or amount of bedding material in stall), cows tended to have cleaner udders. To the best of our knowledge, work exploring this direct relationship between measured bedding depth and hygiene is limited to a single study by de Vries et al. 2015, who found no relationship between prevalence of dirty hindquarters and three different FS bedding height groups (<0.56 cm, 0.56–1.75 cm, >1.75 cm). This relationship between bedding depth and udder hygiene was especially strong for BP in particular, although sample size was limited at five herds. To the best of our knowledge, this specific association has not previously been explored for BP herds. There is

712 opportunity for future research looking at this relationship between increased amount of bedding  
713 used in deep-bedded systems (or more deeply-bedded stalls) and the benefit of improved udder  
714 hygiene and milk quality.

715 Multiple measures of udder health in this study were associated with udder hygiene, in  
716 accordance with the well-supported tenet that better cow hygiene is associated with better milk  
717 quality. The association between hygiene and udder health has been well-documented, both at  
718 the cow level (for IMI presence: de Pinho et al. 2012; for SCS/SCC: Reneau et al. 2005, Dohmen  
719 et al. 2010, and Sant'anna et al. 2011; for both SCS and IMI: Schreiner and Ruegg, 2003) and at  
720 the herd-level (BTSCC: Barkema et al. 1998; new IMI rate: Cook et al. 2002; average herd SCC,  
721 incidence clinical mastitis, and % new high SCC: Dohmen et al. 2010). Of particular relevance to  
722 the current work, a study carried out on three BP farms in Brazil found the odds of a new case of  
723 subclinical mastitis ( $\text{SCC} \geq 200,000$  cells/mL) and of a cow having subclinical mastitis on test  
724 day increased 32% and 16% for each one-unit increase in leg cleanliness score, respectively  
725 (Fávero et al., 2015). Curiously, although leg cleanliness score was associated with both mastitis  
726 outcomes on Brazilian BP, udder hygiene score was not.

727 A third interesting finding to emerge from the univariate regression results is that farms  
728 using deeper bedding had better milk quality outcomes. Although there is an established  
729 recommendation of 15 cm for deep bedding of FS (Bickert, 2000; Cook, 2002) and limited study  
730 exploring ideal bedding material depth for TS barns (Tucker and Weary, 2004; Tucker et al.,  
731 2009), this work is focused on the important concern of cow comfort. As stated in a literature  
732 review by McPherson (2020), "very little research has investigated the effect of bedding depth on  
733 cow cleanliness" or considerations around udder health outcomes. It is likely that the effect seen  
734 in the current work of deeper bedding and better udder health outcomes is mediated through the

presumed causal pathway of (1) deeper bedding leading to improved hygiene, and (2) improved hygiene resulting in better udder health. Although recommending a particular depth may prove difficult as there are many contributing factors which are particular to a producer's barn and bedding source, the opportunity still exists for research exploring optimal stall bedding depths of different organic materials with a focus on mastitis and udder health outcomes.

As for any observational study, there is the potential for bias to have influenced the observed results. Most importantly, participating herds were not a random sample of organic farms in the state, possibly resulting in selection bias. Participating herds were a convenience sample of a subset who responded to our initial survey in Winter 2018-2019 (source population). The potential exists that producers who volunteered to participate in the current study are systematically different in some way with regards to their management practices compared to the general population of organic farms in Vermont. In 2021, there were 147 organic dairy farms in Vermont selling milk, with an average herd size of 87 cows making 6,627 kg milk/cow/year (USDA, 2022). Herds in the current study were slightly smaller, averaging 65 cows per farm, but with higher-producing cows (7,828 kg milk/cow/year, estimated from captured DHIA records). For comparison, the average dairy cow in the U.S. produced an average of 10,926 kg of milk in 2022 (Progressive Dairy, 2017). It may be interesting to acknowledge that organic cows on average produce less milk (Stiglbauer et al., 2013), and with decreased milk production comes decreased susceptibility to mastitis (Grohn, 2000). This relationship may in part explain the relatively low prevalence of mastitis occurring on these farms in comparison to the general population of dairy farms. Lastly, cross-sectional studies are unable to demonstrate causality for associations presented between management practices and outcomes. However, these limitations



are inherent to every observational study, and all attempts were made to control for potential confounding with the multivariable models presented.

Perhaps the biggest limitation of the current study is the small number of farms in each facility type. As state agencies had been promoting the use of BP systems for years in Vermont, we had anticipated it would be feasible to enroll 10 farms using this system to house their lactating animals. This turned out not to be the case; the Winter 2018-2019 survey showed that many dairy farms were instead using these systems for non-lactating animals (heifers, dry cows; Andrews et al. 2021). Furthermore, the COVID-19 pandemic precluded resumption of the study in Spring 2020, limiting the number of farms included to herds sampled in 2019, and not all farms had DHIA data for every outcome of interest. A related limitation is that well-established mastitis control practices were widely adapted by participating herds, so we were unable to analyze associations between certain practices and BTM quality, udder health, and hygiene. A large body of work exists showing consistent udder health benefits from using these and other practices, so lack of association between these fundamental mastitis control practices and desirable outcomes in the current study should not be taken as evidence that they provide no benefit. As group sizes for each facility type were limited, we would caution against making inferences from the findings beyond the source population of this study. The potential still exists for future studies with a larger number of farms enrolled to further characterize milk quality and udder health on BP systems in the Northeastern US. By enrolling farms from a larger geographic area, future studies may be able to enroll a larger number of BP farms, increasing the statistical power needed to identify particular management factors which are beneficial on BP specifically.

While BP systems are not common for housing lactating cows in Vermont, farms using this system in the state are using both compost bedded-packs managed with daily cultivation and

untilled deep bedded pack systems. As untilled and cultivated bedded pack systems differ in numerous regards (Leso et al., 2020), the initial goal was to enroll enough farms using each type and treat them as separate groups in the analysis. As the relatively small number of BP used in our state to house lactating dairy cattle created a challenge for enrolling ten herds using this kind of system in our observational study, it was necessary to combine both types of system in order to achieve our objective of describing udder hygiene, milk quality, and udder health on these loose-housing systems deeply-bedded with organic material. While we acknowledge that grouping them together is not ideal, this diversity is a reflection of how the target population (small-medium, pasture-based organic dairy farms) are actually using them in the Northeastern U.S. (Benson, 2012). Despite this limitation, including bedded pack farms managed in a variety of ways sheds light on a broader spectrum of options used within this loose-housing system. Our current study demonstrates that farms can achieve excellent milk quality using either an untilled, deep bedded pack system or an aerobically composting bedded pack system for indoor housing; three of the five BP farms had a BTSCC  $\leq 99,000$  cells/mL, and the remaining two were  $\leq 160,000$  cells/mL. Furthermore, the lowest BTSCC in the study (54,000 cells/mL) was a static BP farm using woodchips and straw. This low BTSCC was not just from selectively dumping milk from high-SCC cows; this farm also had the lowest overall % cows with elevated SCS (8.6%; data not shown).

BP systems have a number of advantages for producers considering updating their facilities, including a smaller initial investment when compared to a new FS or TS barn (Barberg et al., 2007a; Janni et al., 2007; Black et al., 2013), although the cost year-over-year for bedding is substantial (Shane et al., 2010). Bedded packs are designed for cow comfort (Barberg et al., 2007b; Bewley et al., 2012), and prevalence of lameness, foot, and leg injuries in these systems

has been found to be less than TS and FS barns (Barberg et al., 2007b; Lobeck et al., 2011; Burgstaller et al., 2016). Lastly, manure management and environmental stewardship is a top concern for both dairy producers and the general public (Holly et al., 2018). Anecdotally, the BP producers enrolled in the study were pleased with their systems of manure management, viewing their used bedding material and manure as a valuable soil amendment and an integral part of their nutrient management plan. Bedded pack systems decrease the amount of liquid manure waste when compared to conventional barns, and the used bedding with manure is more easily composted before use as a soil amendment. As aged pack material is drier before it is spread on fields, it poses less of a risk for run-off into waterways, increases soil infiltration of nutrients, and creates flexibility around timing of manure application to fields (Rushmann). Bedded packs may be a good housing option for small, pasture-based farms in the Northeastern U.S. when properly managed on farms with excellent milking hygiene practices already in place. However, more research is needed to confirm that udder health, milk quality, udder hygiene and milk production compares favorably to more traditional housing systems.

## Conclusion

For five of the six studied udder health and production metrics, and both udder hygiene measures, numerically BP either performed slightly better or were equivalent in comparison to the most commonly-used facility types for organic dairy cows in Vermont. However, the relatively large standard errors for most of these estimates make it difficult to rule out biologically important effects of facility type for these outcomes. This is likely a result of the small group size for each facility type. Bedded packs may therefore be a viable option for pasture-based herds looking for a loose-housing system, but future studies enrolling larger

number of farms using each type of housing are needed to more definitively explore these relationships. Findings from the secondary analysis of results found evidence of the well-supported tenets that better cow hygiene is associated with better milk quality, and farms with deeper bedding had more favorable udder hygiene metrics. Additionally, farms using deeper bedding had better milk quality outcomes, which may likely be mediated through improved hygiene resulting in better udder health outcomes.

**Acknowledgements**

This project was funded by the USDA NIFA OREI project 2018-51300-28561. We would like to thank all the organic dairy producers who agreed to participate in this study, for giving us their time and allowing us to collect samples from their farms. We would also like to thank Jennifer Timmerman and the Laboratory for Udder Health at Veterinary Diagnostic Lab (University of Minnesota), as well as the laboratory staff at St. Alban's Cooperative/Dairy Farmers of America, for their advice and analyses of bulk tank milk samples collected in the study. The authors have not stated any conflicts of interest.

**References**

- Adkins, P. R. F., L. M. Placheta, M. R. Borchers, J. M. Bewley, and J. R. Middleton. 2022. Distribution of staphylococcal and mammaliicoccal species from compost-bedded pack or sand-bedded freestall dairy farms. *J Dairy Sci* 105(7):6261-6270.
- Albino, R. L., J. L. Taraba, M. I. Marcondes, E. A. Eckelkamp, and J. M. Bewley. 2018. Comparison of bacterial populations in bedding material, on teat ends, and in milk of cows housed in compost bedded pack barns *J. Animal Production Science* 58(9):1686-1691.

- Andrade, R. R., I. F. F. Tinôco, F. A. Damasceno, G. Ferraz, L. Freitas, C. F. S. Ferreira, M. Barbari, F. J. F. Baptista, and D. J. R. Coelho. 2022. Spatial distribution of bed variables, animal welfare indicators, and milk production in a closed compost-bedded pack barn with a negative tunnel ventilation system. *J Therm Biol* 105:103111.
- Andrews, T., C. E. Jeffrey, R. E. Gilker, D. A. Neher, and J. W. Barlow. 2021. Design and implementation of a survey quantifying winter housing and bedding types used on Vermont organic dairy farms. *J. Dairy Sci.* 104(7):8326-8337.
- Astiz, S., F. Sebastian, O. Fargas, M. Fernández, and E. Calvet. 2014. Enhanced udder health and milk yield of dairy cattle on compost bedding systems during the dry period: A comparative study. *Livestock Science* 159:161-164.
- Barberg, A., M. Endres, and K. Janni. 2007a. Compost Dairy Barns in Minnesota: A Descriptive Study. *Applied Engineering in Agriculture* 23:231-238.
- Barberg, A. E., M. I. Endres, J. A. Salfer, and J. K. Reneau. 2007b. Performance and welfare of dairy cows in an alternative housing system in Minnesota. *J Dairy Sci* 90(3):1575-1583.
- Barkema, H. W., Y. H. Schukken, T. J. Lam, M. L. Beiboer, G. Benedictus, and A. Brand. 1998. Management practices associated with low, medium, and high somatic cell counts in bulk milk. *J. Dairy Sci* 81(7):1917-1927.
- Barkema, H. W., M. A. von Keyserlingk, J. P. Kastelic, T. J. Lam, C. Luby, J. P. Roy, S. J. LeBlanc, G. P. Keefe, and D. F. Kelton. 2015. Invited review: Changes in the dairy industry affecting dairy cattle health and welfare. *J Dairy Sci* 98(11):7426-7445.
- Benson, A. F. 2012. Consider deep pack barns for cow comfort and manure management. Accessed March 18, 2024. Cornell University, Ithaca, NY. <https://smallfarms.cornell.edu/2012/04/consider-deep-pack-barns-for-cow-comfort-and-manure-management/>.
- Bewley, J., J. Taraba, G. Day, R. Black, and F. Damasceno. 2012. Compost Bedded Pack Barn Design: Features and Management Considerations. University of Kentucky Cooperative Extension Service Publication ID.
- Bewley, J. M., L. M. Robertson, and E. A. Eckelkamp. 2017. A 100-Year Review: Lactating dairy cattle housing management. *J. Dairy Sci.* 100(12):10418-10431.
- Bickert, W. G., B. Holmes, K. A. Janni, D. Kammel, R. Stowell, and J. M. Zulovich. 2000. Dairy freestall housing and equipment. Pages 27–45 in *Designing Facilities for the Milking Herd*. 7<sup>th</sup> ed., Mid-West Plan Service, Iowa State University, Ames.
- Black, R. A., J. L. Taraba, G. B. Day, F. A. Damasceno, and J. M. Bewley. 2013. Compost bedded pack dairy barn management, performance, and producer satisfaction. *J Dairy Sci* 96(12):8060-8074.

- 885 Black, R. A., J. L. Taraba, G. B. Day, F. A. Damasceno, M. C. Newman, K. A. Akers, C. L.  
 886 Wood, K. J. McQuerry, and J. M. Bewley. 2014. The relationship between compost bedded pack  
 887 performance, management, and bacterial counts. *J Dairy Sci* 97(5):2669-2679.
- 888 Burgstaller, J., J. Raith, S. Kuchling, V. Mandl, A. Hund, and J. Kofler. 2016. Claw health and  
 889 prevalence of lameness in cows from compost bedded and cubicle freestall dairy barns in  
 890 Austria. *The Veterinary Journal* 216.
- 891 Calamari, L., F. Calegari, and L. Stefanini. 2009. Effect of different free stall surfaces on  
 892 behavioural, productive and metabolic parameters in dairy cows. *Applied Animal Behaviour*  
 893 *Science* 120:9-17.
- 894 Condas, L. A. Z., J. De Buck, D. B. Nobrega, D. A. Carson, S. Naushad, S. De Vliegher, R. N.  
 895 Zadoks, J. R. Middleton, S. Dufour, J. P. Kastelic, and H. W. Barkema. 2017. Prevalence of non-  
 896 aureus staphylococci species causing intramammary infections in Canadian dairy herds. *J Dairy*  
 897 *Sci* 100(7):5592-5612.
- 898 Cook, N. B. 2002. Influence of Barn Design on Dairy Cow Hygiene, Lameness and Udder  
 899 Health. American Association of Bovine Practitioners Conference Proceedings: 97-103.
- 900 Cook, N. B., T. B. Bennett, and K. V. Nordlund. 2005. Monitoring Indices of Cow Comfort in  
 901 Free-Stall-Housed Dairy Herds. *J. Dairy Sci.* 88(11):3876-3885.
- 902 Cook, N. B., J. P. Hess, M. R. Foy, T. B. Bennett, and R. L. Brotzman. 2016. Management  
 903 characteristics, lameness, and body injuries of dairy cattle housed in high-performance dairy  
 904 herds in Wisconsin. *J Dairy Sci* 99(7):5879-5891.
- 905 Costa, J. H. C., T. A. Burnett, M. A. G. von Keyserlingk, and M. J. Hötzel. 2018. Prevalence of  
 906 lameness and leg lesions of lactating dairy cows housed in southern Brazil: Effects of housing  
 907 systems. *J Dairy Sci* 101(3):2395-2405.
- 908 The Dairyland Initiative: School of Veterinary Medicine, University of Wisconsin-Madison.  
 909 Housing Module: Adult Cow Housing, Bedded Packs. University of Wisconsin-Madison.  
 910 Accessed March 18, 2024. [https://thedairylandinitiative.vetmed.wisc.edu/home/housing-](https://thedairylandinitiative.vetmed.wisc.edu/home/housing-module/adult-cow-housing/bedded-pack/)  
 911 [module/adult-cow-housing/bedded-pack/](https://thedairylandinitiative.vetmed.wisc.edu/home/housing-module/adult-cow-housing/bedded-pack/).
- 912 de Pinho Manzi, M., D. B. Nóbrega, P. Y. Faccioli, M. Z. Troncarelli, B. D. Menozzi, and H.  
 913 Langoni. 2012. Relationship between teat-end condition, udder cleanliness and bovine  
 914 subclinical mastitis. *Res Vet Sci* 93(1):430-434.
- 915 De Visscher, A., S. Piepers, F. Haesebrouck, and S. De Vliegher. 2016. Intramammary infection  
 916 with coagulase-negative staphylococci at parturition: Species-specific prevalence, risk factors,  
 917 and effect on udder health. *J Dairy Sci* 99(8):6457-6469.
- 918 De Visscher, A., S. Piepers, F. Haesebrouck, K. Supre, and S. De Vliegher. 2017. Coagulase-  
 919 negative *Staphylococcus* species in bulk milk: Prevalence, distribution, and associated subgroup-  
 920 and species-specific risk factors. *J Dairy Sci* 100(1):629-642.

- 921 de Vries, M., E. A. Bokkers, C. G. van Reenen, B. Engel, G. van Schaik, T. Dijkstra, and I. J. de  
 922 Boer. 2015. Housing and management factors associated with indicators of dairy cattle welfare.  
 923 *Prev Vet Med* 118(1):80-92.
- 924 Dohmen, W., F. Neijenhuis, and H. Hogeveen. 2010. Relationship between udder health and  
 925 hygiene on farms with an automatic milking system. *J Dairy Sci* 93(9):4019-4033.
- 926 Eberhart, R. J. 1984. Coliform Mastitis. *Veterinary Clinics of North America: Large Animal*  
 927 *Practice* 6(2):287-300.
- 928 Eckelkamp, E. A., J. L. Taraba, K. A. Akers, R. J. Harmon, and J. M. Bewley. 2016a. Sand  
 929 bedded freestall and compost bedded pack effects on cow hygiene, locomotion, and mastitis  
 930 indicators. *Livestock Science* 190:48-57.
- 931 Eckelkamp, E. A., J. L. Taraba, K. A. Akers, R. J. Harmon, and J. M. Bewley. 2016b.  
 932 Understanding compost bedded pack barns: Interactions among environmental factors, bedding  
 933 characteristics, and udder health. *Livestock Science* 190:35-42.
- 934 Elmoslemany, A. M., G. P. Keefe, I. R. Dohoo, and B. M. Jayarao. 2009. Risk factors for  
 935 bacteriological quality of bulk tank milk in Prince Edward Island dairy herds. Part 1: overall risk  
 936 factors. *J Dairy Sci* 92(6):2634-2643.
- 937 Endres, M., K. Janni. 2021. Compost-bedded pack barns for dairy cows. University of Minnesota  
 938 Extension. Minneapolis, MN. Accessed March 18, 2024. [https://extension.umn.edu/dairy-](https://extension.umn.edu/dairy-milking-cows/compost-bedded-pack-barns-dairy-cows#a-wall-borders-the-pack-727910)  
 939 [milking-cows/compost-bedded-pack-barns-dairy-cows#a-wall-borders-the-pack-727910](https://extension.umn.edu/dairy-milking-cows/compost-bedded-pack-barns-dairy-cows#a-wall-borders-the-pack-727910).
- 940 Fairchild, T. P., B. J. McArthur, J. H. Moore, and W. E. Hylton. 1982. Coliform Counts in  
 941 Various Bedding Materials. *J. Dairy Sci.* 65(6):1029-1035.
- 942 Fávero, S., F. V. R. Portilho, A. C. R. Oliveira, H. Langoni, and J. C. F. Pantoja. 2015. Factors  
 943 associated with mastitis epidemiologic indexes, animal hygiene, and bulk milk bacterial  
 944 concentrations in dairy herds housed on compost bedding. *Livestock Science* 181:220-230.
- 945 Ferraz, P. F. P., G. A. e. S. Ferraz, L. Leso, M. Klopčič, M. Barbari, and G. Rossi. 2020.  
 946 Properties of conventional and alternative bedding materials for dairy cattle. *J. Dairy Sci.*  
 947 103(9):8661-8674.
- 948 Fregonesi, J. A. and J. D. Leaver. 2001. Behaviour, performance and health indicators of welfare  
 949 for dairy cows housed in strawyard or cubicle systems. *Livestock Production Science* 68(2):205-  
 950 216.
- 951 Fregonesi, J. A. and J. D. Leaver. 2002. Influence of space allowance and milk yield level on  
 952 behaviour, performance and health of dairy cows housed in strawyard and cubicle systems.  
 953 *Livestock Production Science* 78(3):245-257.
- 954 Godkin, M. A. and K. E. Leslie. 1993. Culture of bulk tank milk as a mastitis screening test: A  
 955 brief review. *Can Vet J* 34(10):601-605.



- 956 Grohn, Y. 2000. Milk Yield and Disease: Towards Optimizing Dairy Herd Health and  
957 Management Decisions. *Bovine Practice* 34:32-40.
- 958 Heins, B. J., L. S. Sjostrom, M. I. Endres, M. R. Carillo, R. King, R. D. Moon, and U. S. Sorge.  
959 2019. Effects of winter housing systems on production, economics, body weight, body condition  
960 score, and bedding cultures for organic dairy cows. *J Dairy Sci* 102(1):706-714.
- 961 Hogan, J. and K. L. Smith. 2012. Managing environmental mastitis. *Vet Clin North Am Food*  
962 *Anim Pract* 28(2):217-224.
- 963 Hogan, J. S. and K. L. Smith. 1997. Bacteria counts in sawdust bedding. *J Dairy Sci* 80(8):1600-  
964 1605.
- 965 Hogan, J. S., K. L. Smith, K. H. Hoblet, D. A. Todhunter, P. S. Schoenberger, W. D. Hueston, D.  
966 E. Pritchard, G. L. Bowman, L. E. Heider, B. L. Brockett, and H. R. Conrad. 1989. Bacterial  
967 Counts in Bedding Materials Used on Nine Commercial Dairies. *J. Dairy Sci.* 72(1):250-258.
- 968 Hogan, J. S., D. G. White, and J. W. Pankey. 1987. Effects of teat dipping on intramammary  
969 infections by staphylococci other than *Staphylococcus aureus*. *J Dairy Sci* 70(4):873-879.
- 970 Holly, M. A., P. J. Kleinman, R. B. Bryant, D. L. Bjorneberg, C. A. Rotz, J. M. Baker, M. V.  
971 Boggess, D. K. Brauer, R. Chintala, G. W. Feyereisen, J. D. Gamble, A. B. Leytem, K. F. Reed,  
972 P. A. Vadas, and H. M. Waldrip. 2018. Short communication: Identifying challenges and  
973 opportunities for improved nutrient management through the USDA's Dairy Agroecosystem  
974 Working Group. *J Dairy Sci* 101(7):6632-6641.
- 975 Janni, K., M. Endres, J. Reneau, and W. Schoper. 2007. Compost Dairy Barn Layout and  
976 Management Recommendations. *Applied Engineering in Agriculture* 23(1):97-102.
- 977 Jayarao, B. M., S. R. Pillai, A. A. Sawant, D. R. Wolfgang, and N. V. Hegde. 2004. Guidelines  
978 for monitoring bulk tank milk somatic cell and bacterial counts. *J Dairy Sci* 87(10):3561-3573.
- 979 Jayarao, B. M. and D. R. Wolfgang. 2003. Bulk-tank milk analysis. A useful tool for improving  
980 milk quality and herd udder health. *Vet Clin North Am Food Anim Pract* 19(1):75-92, vi.
- 981 Klaas, I. C. and R. N. Zadoks. 2018. An update on environmental mastitis: Challenging  
982 perceptions. *Transbound Emerg Dis* 65 Suppl 1:166-185.
- 983 KoboCollect: Simple, Robust and Powerful Tools for Data Collection. 2019  
984 <http://www.kobotoolbox.org>.
- 985 Leso, L., M. Barbari, M. A. Lopes, F. A. Damasceno, P. Galama, J. L. Taraba, and A. Kuipers.  
986 2020. Invited review: Compost-bedded pack barns for dairy cows. *J Dairy Sci* 103(2):1072-1099.
- 987 Lobeck, K., M. Endres, K. Janni, S. Godden, and J. Fetrow. 2012. Environmental Characteristics  
988 and Bacterial Counts in Bedding and Milk Bulk Tank of Low Profile Cross-Ventilated, Naturally  
989 Ventilated, and Compost Bedded Pack Dairy Barns. *Applied Engineering in Agriculture* 28:117-  
990 128.



- 991 Lobeck, K. M., M. I. Endres, E. M. Shane, S. M. Godden, and J. Fetrow. 2011. Animal welfare  
992 in cross-ventilated, compost-bedded pack, and naturally ventilated dairy barns in the upper  
993 Midwest. *J Dairy Sci* 94(11):5469-5479.
- 994 McPherson, S. E. and E. Vasseur. 2020. Graduate Student Literature Review: The effects of  
995 bedding, stall length, and manger wall height on common outcome measures of dairy cow  
996 welfare in stall-based housing systems. *J Dairy Sci* 103(11):10940-10950.
- 997 Neave, F. K., F. H. Dodd, and R. G. Kingwill. 1966. A method of controlling udder disease. *Vet*  
998 *Rec* 78(15):521-523.
- 999 Neher, D. A., T. D. Andrews, T. R. Weicht, A. Hurd, and J. W. Barlow. 2022. Organic Farm  
1000 Bedded Pack System Microbiomes: A Case Study with Comparisons to Similar and Different  
1001 Bedded Packs. *Dairy*. doi:10.3390/dairy3030042.
- 1002 O'Connor, A. M., J. M. Sargeant, I. R. Dohoo, H. N. Erb, M. Cevallos, M. Egger, A. K. Ersbøll,  
1003 S. W. Martin, L. R. Nielsen, D. L. Pearl, D. U. Pfeiffer, J. Sanchez, M. E. Torrence, H. Vigre, C.  
1004 Waldner, and M. P. Ward. 2016. Explanation and Elaboration Document for the STROBE-Vet  
1005 Statement: Strengthening the Reporting of Observational Studies in Epidemiology-Veterinary  
1006 Extension. *J Vet Intern Med* 30(6):1896-1928.
- 1007 Pankey, J. W., R. L. Boddie, and S. C. Nickerson. 1985. Efficacy evaluation of two new teat dip  
1008 formulations under experimental challenge. *J Dairy Sci* 68(2):462-465.
- 1009 Pankey, J. W., E. E. Wildman, P. A. Drechsler, and J. S. Hogan. 1987. Field trial evaluation of  
1010 premilking teat disinfection. *J Dairy Sci* 70(4):867-872.
- 1011 Pankey, J. W. 1989. Premilking Udder Hygiene. *J. Dairy Sci.* 72(5):1308-1312.
- 1012 Pantoja, J. C. F., D. J. Reinemann, and P. L. Ruegg. 2009. Associations among milk quality  
1013 indicators in raw bulk milk. *J. Dairy Sci.* 92(10):4978-4987.
- 1014 Patel, K., S. M. Godden, E. Royster, B. A. Crooker, J. Timmerman, and L. Fox. 2019.  
1015 Relationships among bedding materials, bedding bacteria counts, udder hygiene, milk quality,  
1016 and udder health in US dairy herds. *J. Dairy Sci.* 102(11):10213-10234.
- 1017 Peeler, E. J., M. J. Green, J. L. Fitzpatrick, K. L. Morgan, and L. E. Green. 2000. Risk Factors  
1018 Associated with Clinical Mastitis in Low Somatic Cell Count British Dairy Herds. *J. Dairy Sci.*  
1019 83(11):2464-2472.
- 1020 Piessens, V., E. Van Coillie, B. Verbist, K. Supre, G. Braem, A. Van Nuffel, L. De Vuyst, M.  
1021 Heyndrickx, and S. De Vliegher. 2011. Distribution of coagulase-negative *Staphylococcus*  
1022 species from milk and environment of dairy cows differs between herds. *J Dairy Sci* 94(6):2933-  
1023 2944.
- 1024 Pol, M. and P. L. Ruegg. 2007. Relationship between antimicrobial drug usage and antimicrobial  
1025 susceptibility of gram-positive mastitis pathogens. *J Dairy Sci* 90(1):262-273.

- 1026 Progressive Dairy. 2022. U.S. Dairy Statistics. Accessed March 19, 2024.  
1027 <https://www.progressivepublish.com/downloads/2023/general/2022-pd-stats-highres.pdf>.
- 1028 Quirk, T., L. K. Fox, D. D. Hancock, J. Capper, J. Wenz, and J. Park. 2012. Intramammary  
1029 infections and teat canal colonization with coagulase-negative staphylococci after postmilking  
1030 teat disinfection: species-specific responses. *J Dairy Sci* 95(4):1906-1912.
- 1031 R Development Core Team. 2023. R: A Language and Environment for Statistical Computing. R  
1032 Foundation for Statistical Computing, Vienna, Austria.
- 1033 Reneau, J. K., A. J. Seykora, B. J. Heins, M. I. Endres, R. J. Farnsworth, and R. F. Bey. 2005.  
1034 Association between hygiene scores and somatic cell scores in dairy cattle. *J Am Vet Med Assoc*  
1035 227(8):1297-1301.
- 1036 Rinehart, L. and A. Baier. 2011. U.S. Department of Agriculture; National Center for  
1037 Appropriate Technology (NCAT), National Organic Program. Pasture for Organic Ruminant  
1038 Livestock: Understanding and Implementing the National Organic Program (NOP) Pasture Rule.  
1039 Accessed Oct. 30, 2023. [https://www.ams.usda.gov/sites/default/files/media/NOP-](https://www.ams.usda.gov/sites/default/files/media/NOP-UnderstandingOrganicPastureRule.pdf)  
1040 [UnderstandingOrganicPastureRule.pdf](https://www.ams.usda.gov/sites/default/files/media/NOP-UnderstandingOrganicPastureRule.pdf).
- 1041 Robles, I., D. F. Kelton, H. W. Barkema, G. P. Keefe, J. P. Roy, M. A. G. von Keyserlingk, and  
1042 T. J. DeVries. 2020. Bacterial concentrations in bedding and their association with dairy cow  
1043 hygiene and milk quality. *Animal* 14(5):1052-1066.
- 1044 Rowbotham, R. F. and P. L. Ruegg. 2016a. Associations of selected bedding types with  
1045 incidence rates of subclinical and clinical mastitis in primiparous Holstein dairy cows. *J Dairy*  
1046 *Sci* 99(6):4707-4717.
- 1047 Rowbotham, R. F. and P. L. Ruegg. 2016b. Bacterial counts on teat skin and in new sand,  
1048 recycled sand, and recycled manure solids used as bedding in freestalls. *J Dairy Sci* 99(8):6594-  
1049 6608.
- 1050 Ruegg, P. L. 2009. Management of mastitis on organic and conventional dairy farms. *J Anim Sci*  
1051 87(13 Suppl):43-55.
- 1052 Ruegg, P. L. and J. C. F. Pantoja. 2013. Understanding and using somatic cell counts to improve  
1053 milk quality. *Irish Journal of Agricultural and Food Research* 52(2):101-117.
- 1054 Rushmann, R. University of Wisconsin-Madison; Division of Extension: Agriculture Water  
1055 Quality. Managing manure to reduce negative water quality impacts: Composting on Wisconsin  
1056 farms. Accessed Aug. 1, 2023. [https://agwater.extension.wisc.edu/articles/managing-manure-to-](https://agwater.extension.wisc.edu/articles/managing-manure-to-reduce-negative-water-quality-impacts-composting-on-wisconsin-farms/)  
1057 [reduce-negative-water-quality-impacts-composting-on-wisconsin-farms/](https://agwater.extension.wisc.edu/articles/managing-manure-to-reduce-negative-water-quality-impacts-composting-on-wisconsin-farms/).
- 1058 Ruud, L. E., K. E. Bøe, and O. Osterås. 2010. Associations of soft flooring materials in free stalls  
1059 with milk yield, clinical mastitis, teat lesions, and removal of dairy cows. *J Dairy Sci*  
1060 93(4):1578-1586.

- 1061 Sant'anna, A. C. and M. J. Paranhos da Costa. 2011. The relationship between dairy cow hygiene  
1062 and somatic cell count in milk. *J Dairy Sci* 94(8):3835-3844.
- 1063 Schreiner, D. A. and P. L. Ruegg. 2002. Effects of tail docking on milk quality and cow  
1064 cleanliness. *J Dairy Sci* 85(10):2503-2511.
- 1065 Schreiner, D. A. and P. L. Ruegg. 2003. Relationship between udder and leg hygiene scores and  
1066 subclinical mastitis. *J Dairy Sci* 86(11):3460-3465.
- 1067 Schukken, Y. H., F. J. Grommers, J. A. Smit, D. Vandegeer, and A. Brand. 1989. Effect of  
1068 freezing on bacteriologic culturing of mastitis milk samples. *J Dairy Sci* 72(7):1900-1906.
- 1069 Schukken, Y. H., D. J. Wilson, F. Welcome, L. Garrison-Tikofsky, and R. N. Gonzalez. 2003.  
1070 Monitoring udder health and milk quality using somatic cell counts. *Vet Res* 34(5):579-596.
- 1071 Shane, E., M. Endres, and K. Janni. 2010. Alternative Bedding Materials for Compost Bedded  
1072 Pack Barns in Minnesota: A Descriptive Study. *Applied Engineering in Agriculture* 26:465-473.
- 1073 Stiglbauer, K. E., K. M. Cicconi-Hogan, R. Richert, Y. H. Schukken, P. L. Ruegg, and M.  
1074 Gamroth. 2013. Assessment of herd management on organic and conventional dairy farms in the  
1075 United States. *J. Dairy Sci.* 96(2):1290-1300.
- 1076 Thurgood, J. M., C. M. Comer, D. J. Flaherty, and M. Kiraly. 2009. Bedded pack management  
1077 system case study. Pages 184–188 in *Proc. 5th National Small Farm Conference*,  
1078 Springfield, IL. Accessed March 18, 2024.  
1079 [https://conferences.illinois.edu/resources/20033/Proceedings\\_8-12-13.pdf](https://conferences.illinois.edu/resources/20033/Proceedings_8-12-13.pdf).
- 1080 Tucker, C. B., D. Weary, M. Keyserlingk, and K. Beauchemin. 2009. Cow comfort in tie-stalls:  
1081 Increased depth of shavings or straw bedding increases lying time. *J. Dairy Sci.* 92:2684-2690.
- 1082 Tucker, C. B. and D. M. Weary. 2004. Bedding on geotextile mattresses: how much is needed to  
1083 improve cow comfort? *J Dairy Sci* 87(9):2889-2895.
- 1084 University of Minnesota Extension Dairy Team. Using DHIA Records to Benchmark Herd SCC.  
1085 Accessed Apr. 1, 2024. [https://qualitycounts.umn.edu/sites/qualitycounts.umn.edu/files/2022-](https://qualitycounts.umn.edu/sites/qualitycounts.umn.edu/files/2022-01/w-mp-5.pdf)  
1086 [01/w-mp-5.pdf](https://qualitycounts.umn.edu/sites/qualitycounts.umn.edu/files/2022-01/w-mp-5.pdf)
- 1087 USDA-NRCS. NRCS Climate-Smart Mitigation Activities. Accessed Dec. 14, 2023.  
1088 [https://www.nrcs.usda.gov/conservation-basics/natural-resource-concerns/climate/climate-smart-](https://www.nrcs.usda.gov/conservation-basics/natural-resource-concerns/climate/climate-smart-mitigation-activities)  
1089 [mitigation-activities](https://www.nrcs.usda.gov/conservation-basics/natural-resource-concerns/climate/climate-smart-mitigation-activities).
- 1090 USDA. 2022. Certified Organic Survey, 2021 Summary. Accessed Nov. 10, 2023.  
1091 [https://downloads.usda.library.cornell.edu/usda-](https://downloads.usda.library.cornell.edu/usda-esmis/files/zg64tk92g/2z10z137s/bn99bh97r/cenorg22.pdf)  
1092 [esmis/files/zg64tk92g/2z10z137s/bn99bh97r/cenorg22.pdf](https://downloads.usda.library.cornell.edu/usda-esmis/files/zg64tk92g/2z10z137s/bn99bh97r/cenorg22.pdf).
- 1093 Ward, W. R., J. W. Hughes, W. B. Faull, P. J. Cripps, J. P. Sutherland, and J. E. Sutherst. 2002.  
1094 Observational study of temperature, moisture, pH and bacteria in straw bedding, and faecal  
1095 consistency, cleanliness and mastitis in cows in four dairy herds. *Vet Rec* 151(7):199-206.

1096 Wolfe, T., E. Vasseur, T. J. DeVries, and R. Bergeron. 2018. Effects of alternative deep bedding  
1097 options on dairy cow preference, lying behavior, cleanliness, and teat end contamination. J Dairy  
1098 Sci 101(1):530-536.

1099 Wuytack, A., A. De Visscher, S. Piepers, F. Haesebrouck, and S. De Vliegheer. 2020. Fecal non-  
1100 aureus Staphylococci are a potential cause of bovine intramammary infection. Vet Res 51(1):32.

1101 Zadoks, R. N., L. L. Tikofsky, and K. J. Boor. 2005. Ribotyping of Streptococcus uberis from a  
1102 dairy's environment, bovine feces and milk. Veterinary Microbiology 109(3):257-265.

1103 Zdanowicz, M., J. A. Shelford, C. B. Tucker, D. M. Weary, and M. A. G. von Keyserlingk. 2004.  
1104 Bacterial Populations on Teat Ends of Dairy Cows Housed in Free Stalls and Bedded with Either  
1105 Sand or Sawdust. J. Dairy Sci. 87(6):1694-1701.

1106

1107 **Tables**

Table 1. Predictors offered to multivariable models for each of the eight different outcomes of interest along with facility type (forced)

Predictor	Level of parameter, if categorical:
Farm demographics/lactating cow housing	
Facility type	Bedded pack; Freestall; Tiestall
Predominant breed	Holstein; Jersey/Other
Herd size (lactating cows)	
Herd size group (lactating cows)	30-55; 56-69; 70-100
Subjective assessment of air quality (producer)	Excellent; Good; Fair/Poor
Subjective assessment of air quality (researcher)	Good; Fair
Age of facility (years)	
Feed supplemental vit. E and selenium	Yes; No
Lactating bedding management practices	
Lying surface for cows <sup>1</sup> (deeply-bedded vs. not)	Deeply-bedded stalls or bedded pack; Stalls with bedding on a mattress or concrete surface
<i>If use shavings/sawdust/woodchips for bedding material:</i>	
Moisture-content	Kiln-dried; Fresh/raw
Bedding amendment (e.g., hydrated lime) used on surface	Yes; No
<i>If facility is freestall or tiestall:</i>	
Freq. adding new bedding to stalls (times per week)	
Freq. scraping stalls (times per week)	
Depth bedding in stalls (cm)	
Mastitis control and milking hygiene practices	
Clip/flame udder hair	Yes; No

Keep record of clinical mastitis events	Always; Sometimes/Temp.; Never
Routinely culture mastitic milk	Always/Sometimes; Never
Routinely culture high somatic cell count cows	Always/Sometimes; Never
Ever perform culture of mastitic cows	Yes; Never culture
Use intramammary product at dry-off (OMRI-listed)	Yes; No
Parenteral supplementation with vit. E and selenium	All lactating cows regularly/ Occasionally as needed; No
Glove use at milking	All milkers consistently; Inconsistently/No
Check for clinical mastitis by noticing abnormal cow/abnormal udder and forestripping	Yes; No
Type of milking system used <sup>3</sup>	Parlor; Tiestall
Farm-level udder hygiene metrics	
Average udder hygiene score	
Prop. dirty udders (%; udder hygiene score $\geq 3$ )	
<sup>1</sup> If freestall or tiestall, producer asked if used deeply-bedded stalls	
<sup>2</sup> OMRI: Organic Materials Review Institute	
<sup>3</sup> One freestall farm used an automated milking system	

Table 2. Objective 1: Descriptive and univariable results for bulk tank milk aerobic culture outcomes by facility type [median (range)]. *P*-value is for Kruskal-Wallis test by facility type grouping

Bacteria group (cfu/mL)	Overall (n = 21)	Bedded packs (n = 5)	Tiestalls (n = 10)	Freestalls (n = 6)	<i>P</i> -value
<i>Staph. spp.</i>	65 (0-665)	40 (0-130)	85 (15-665)	67.5 (5-125)	0.62
<i>Strep.</i> and strep-like orgs.	45 (10-1250)	35 (10-80)	167.5 (20-1250)	32.5 (25-260)	0.10
<i>Staph. aureus</i>	30 (0-320)	0 (0-30)	47.5 (0-320)	42.5 (0-100)	0.19
Coliforms	0 (0-5)	0 (0-5)	0 (0-5)	0 (0-5)	0.82

Table 3. Objective 1: Descriptive results for milk quality, udder health and production outcomes by facility type [mean (95%CI)]

Outcome	Overall n = 21	Bedded packs n = 5	Tiestalls n = 10	Freestalls n = 6
BTSCC (log <sub>10</sub> cells/mL)	5.13 (5.06-5.20)	5.00 (4.84-5.17)	5.14 (5.05-5.23)	5.21 (5.09-5.33)
% newly elevated SCS <sup>1</sup>	n = 19 5.7 (4.2-7.3)	n = 3 7.0 (2.8-11.2)	n = 10 5.4 (3.0-7.8)	n = 6 5.6 (3.0-8.3)

% chronically elevated SCS <sup>1</sup>	n = 19	n = 3	n = 10	n = 6
	13.6 (11.2-16.1)	14.5 (5.4-23.7)	14.3 (11.9-16.7)	12.0 (6.7-17.3)
% SCS ≥ 4.0 current test <sup>1</sup>	n = 19	n = 3	n = 10	n = 6
	24.9 (21.6-28.3)	26.0 (12.6-39.3)	25.4 (22.1-28.6)	23.7 (16.9-30.5)
Avg. SCS <sup>2</sup>	n = 20	n = 4	n = 10	n = 6
	2.44 (2.26-2.62)	2.38 (1.84-2.91)	2.45 (2.31-2.59)	2.50 (2.00-2.93)
Standardized 150-day milk (pounds) <sup>3</sup>	n = 18	n = 4	n = 8	n = 6
	50.0 (45.7-54.3)	46.9 (39.8-53.9)	49.4 (43.1-55.7)	53.0 (43.5-62.5)

<sup>1</sup> DHIA data not available for 2 bedded pack farms

<sup>2</sup> DHIA data not available for 1 bedded pack farm

<sup>3</sup> DHIA data not available for 1 bedded pack farms and 2 tiestall farms

1114

Table 4. Objective 1: Final multivariable models describing the relationship between facility type (forced) and milk quality, udder health, production, and udder hygiene outcomes

Outcome	Explanatory variable	Group (sample size)	Coefficient estimate (SE)	P-value
BTSCC (log <sub>10</sub> cells/mL)	Intercept		4.8 (0.15)	
	Facility type (forced)	Freestall (n = 6)	0.19 (0.09)	0.05
		Tiestall (n = 10)	0.16 (0.08)	0.07
		Bedded pack (n = 5)	Ref.	Ref.
	Herd size	All herds (n = 21)	0.003 (0.002)	0.15
% newly elevated SCS	Intercept		-1.6 (2.5)	
	Facility type (forced)	Freestall (n = 6)	2.3 (2.2)	0.33
		Tiestall (n = 10)	0.43 (1.9)	0.82
		Bedded pack (n = 3)	Ref.	Ref.
	Use bedding amendment	Yes (n = 5)	3.9 (1.8)	0.05
No (n = 14)		Ref.	Ref.	
Subjective assessment air quality (researcher)	Good (n = 14)	3.6 (1.5)	0.04	
	Fair (n = 5)	Ref.	Ref.	
Glove use at milking <sup>1</sup>	Never/Inconsistently (n = 9)	2.0 (1.3)	0.17	
	Always (n = 9)	Ref.	Ref.	
Clinical mastitis events record keeping	Never kept records (n = 6)	4.4 (1.8)	0.03	
	Sometimes/Temporarily kept records (n = 6)	1.1 (1.6)	0.52	
	Always kept records (n = 7)	Ref.	Ref.	
% chronically elevated SCS	Intercept		5.3 (3.6)	
	Facility type (forced)	Freestall (n = 6)	1.5 (4.6)	0.75
		Tiestall (n = 10)	-0.08 (3.5)	0.98
		Bedded pack (n = 3)	Ref.	Ref.
	Feed supplemental vit. E and selenium <sup>2</sup>	Yes (n = 11)	2.1 (2.8)	0.48
		No (n = 7)	Ref.	Ref.

	Use bedding amendment	Yes (n = 5)	5.7 (3.4)	0.12
		No (n = 14)	Ref.	Ref.
	Clip/flame udder hair	Yes (n = 5)	-6.3 (3.1)	0.07
		No (n = 14)	Ref.	Ref.
% udder hygiene scores $\geq 3$		Herds with available data (n = 19)	17.0 (6.1)	0.02
% SCS $\geq 4.0$ current test	Intercept		0.85 (10.6)	
	Facility type (forced)			
		Freestall (n = 6)	1.8 (5.7)	0.75
		Tiestall (n = 10)	-2.4 (5.3)	0.66
		Bedded pack (n = 3)	Ref.	Ref.
	Use bedding amendment	Yes (n = 5)	8.0 (4.2)	0.07
		No (n = 14)	Ref.	Ref.
	Mean hygiene	Herds with available data (n = 19)	9.8 (4.7)	0.06
Avg. SCS				20
	Intercept		0.93 (0.44)	
	Facility type (forced)			
		Freestall (n = 6)	0.38 (0.21)	0.09
		Tiestall (n = 10)	0.03 (0.19)	0.86
		Bedded pack (n = 4)	Ref.	Ref.
	Use intramammary product at dry-off (OMRI-listed)	Yes (n = 5)	-0.30 (0.16)	0.08
		No (n = 15)	Ref.	Ref.
	Use bedding amendment	Yes (n = 5)	0.52 (0.16)	0.007
		No (n = 15)	Ref.	Ref.
	Parenteral supplementation vit. E/selenium	Regularly or occasionally (n = 9)	-0.36 (0.14)	0.02
		No supplementation (n = 11)	Ref.	Ref.
	Mean hygiene	Herds with available data (n = 20)	0.64 (0.19)	0.005
Standardized 150-day milk (pounds)				18
	Intercept		41.2 (6.1)	
	Facility type (forced)			
		Freestall (n = 6)	-0.06 (7.0)	0.99
		Tiestall (n = 8)	-1.7 (6.6)	0.80
		Bedded pack (n = 4)	Ref.	Ref.
	Parenteral supplementation vit. E/selenium	Regularly or occasionally (n = 7)	7.0 (5.2)	0.20



	No supplementation (n = 11)	Ref.	Ref.
Culture high SCC cows	Always/Sometimes (n = 8)	9.3 (5.9)	0.14
	Never (n = 10)	Ref.	Ref.
Herd size grp. (lact. cows)	70-100 (n = 8)	-0.18 (7.3)	0.98
	56-69 (n = 5)	10.3 (6.2)	0.12
	30-55 (n = 5)	Ref.	Ref.
% udder hygiene scores $\geq 3$			
Intercept		0.32 (0.08)	
Facility type (forced)	Freestall (n = 6)	0.002 (0.11)	0.99
	Tiestall (n = 10)	0.17 (0.10)	0.12
	Bedded pack (n = 5)	Ref.	Ref.
Avg. udder hygiene score			
Intercept		2.3 (0.17)	
Facility type (forced)	Freestall (n = 6)	-0.04 (0.21)	0.84
	Tiestall (n = 10)	0.33 (0.19)	0.11
	Bedded pack (n = 5)	Ref.	Ref.
Check for clinical mastitis by noticing abnormal cow/abnormal udder and forestripping			
	Yes (n = 8)	-0.25 (0.16)	0.14
	No (n = 13)	Ref.	Ref.

<sup>1</sup> One farm used automatic milking system

<sup>2</sup> One farm unable to provide response

Table 5. Objective 2: Selected models of univariate analysis identifying (non-facility type) factors unconditionally associated with milk quality, udder health, production, and udder hygiene outcomes at  $P < 0.20$

Outcome	Explanatory Variable	Group (sample size)	Coefficient estimate (SE)	P-value	Intercept
BTSCC (log <sub>10</sub> cells/mL)					
Model 1	Lying surface	Mattress or concrete (n = 13)	0.12 (0.07)	0.12	5.1
		Deep bedding (n = 8)	Ref.	Ref.	
Model 2	Depth of bedding in stalls (cm) <sup>1</sup>	Tiestalls and freestalls (n = 15)	-0.02 (0.01)	0.11	5.2
Model 3	Glove use at milking <sup>2</sup>	Never/Inconsistently (n = 9)	0.10 (0.07)	0.19	5.1
		Always (n = 11)	Ref.	Ref.	
% newly elevated SCS <sup>3</sup>					
Model 4	Glove use at milking	Never/Inconsistently (n = 9)	2.83 (1.7)	0.11	4.3
		Always (n = 9)	Ref.	Ref.	
Model 5	Depth of bedding in stalls (cm) <sup>1</sup>	Tiestalls and freestalls (n = 15)	-0.62 (0.24)	0.02	8.3
% chronically elevated SCS <sup>3</sup>					
Model 6	Clip/flame udder hair	Yes (n = 5)	-4.31 (2.9)	0.16	14.8
		No (n = 14)	Ref.	Ref.	
Model 7	% udder hygiene scores ≥3	Herds with available data (n = 19)	12.7 (6.1)	0.05	8.6
Model 8	Avg. udder hygiene score	Herds with available data (n = 19)	6.39 (3.1)	0.05	-1.2
% SCS ≥ 4.0 current test <sup>3</sup>					
Model 9	Depth of bedding in stalls (cm) <sup>1</sup>	Tiestalls and freestalls (n = 15)	-1.2 (0.42)	0.01	30
Model 10	% udder hygiene scores ≥3	Herds with available data (n = 19)	13.6 (8.5)	0.13	19.6
Model 11	Avg. udder hygiene score	Herds with available data (n = 19)	7.7 (4.3)	0.09	7.1
Average SCS <sup>4</sup>					
Model 12	Parenteral supplementation vit. E and selenium	Regularly or occasionally (n = 9)	-0.27 (0.18)	0.15	2.6
		No supplementation (n = 11)	Ref.	Ref.	
Model 13	Use intramammary product at dry-off (OMRI-listed)	Yes (n = 5)	-0.29 (0.21)	0.18	2.5
		No (n = 15)	Ref.	Ref.	
Model 14	Depth of bedding in stalls (cm) <sup>1</sup>	Tiestalls and freestalls (n = 15)	-0.05 (0.03)	0.10	2.6

Model 15	% udder hygiene scores $\geq 3$	Herds with available data (n = 20)	0.75 (0.45)	0.12	2.1
Model 16	Avg. udder hygiene score	Herds with available data (n = 20)	0.39 (0.23)	0.11	1.5
Standardized 150-day milk (pounds) <sup>5</sup>					
Model 17	Parenteral supplementation vit. E and selenium	Regularly or occasionally (n = 7)	9.0 (4.5)	0.06	46.5
		No supplementation (n = 11)	Ref.	Ref.	
Model 18	Herd size	Herds with available data (n = 18)	0.26 (0.14)	0.07	33.1
% udder hygiene scores $\geq 3$					
Model 19	Depth of bedded pack (m)	Bedded pack herds (n = 5)	-0.5 (0.06)	0.004	0.97
Model 20	Lying surface	Mattress or concrete (n = 13)	0.17 (0.08)	0.06	0.30
		Deep bedding (n = 8)	Ref.	Ref.	
Model 21	Depth of bedding in stalls (cm) <sup>1</sup>	Tiestalls and freestalls (n = 15)	-0.02 (0.02)	0.13	0.54
Avg. udder hygiene score					
Model 22	Depth of bedded pack (m)	Bedded pack herds (n = 5)	-0.96 (0.15)	0.008	3.4
Model 23	Lying surface	Mattress or concrete (n = 13)	0.33 (0.16)	0.06	2.1
		Deep bedding (n = 8)	Ref.	Ref.	
Model 24	Depth of bedding in stalls (cm) <sup>1</sup>	Tiestalls and freestalls (n = 15)	-0.06 (0.03)	0.07	2.6

<sup>1</sup> Stall bedding depth for freestalls and tiestalls bedded with wood shavings or sawdust

<sup>2</sup> One farm used automatic milking system

<sup>3</sup> DHIA data available for n = 19 herds.

<sup>4</sup> DHIA data available for n = 20 herds.

<sup>5</sup> DHIA data available for n = 18 herds.

1116

1117

1118