

# Analysis 1A — Supremum Example

Christian Jones: University of Bath

October 2022

## Contents

<b>Question</b>	<b>1</b>
Method 1 — Contradiction . . . . .	1
Method 2 — Alternative Characterisation of Suprema . . . . .	2
Method 3 — Limits . . . . .	2

## Question

Here is an example of finding the supremum of a set taken from an old problem sheet, together with three possible methods to find it.

### Example 0.1.

Let

$$B = \left\{ \frac{2n-1}{n+1} \mid n \in \mathbb{N} \right\}.$$

Show that  $B$  is bounded above and find  $\sup(B)$ .

### Method 1 — Contradiction

#### Solution.

First, note for  $n \in \mathbb{N}$ :

$$\frac{2n-1}{n+1} = \frac{2n+2-3}{n+1} = 2 - \frac{3}{n+1} < 2.$$

Hence  $B$  is bounded above by 2. Therefore, by the completeness axiom, as  $B \neq \emptyset$ ,  $\sup(B)$  exists and  $\sup(B) \leq 2$ .

Next, suppose for contradiction that  $\sup(B) < 2$ . Now, for any  $x < 2$ ,

$$2 - \frac{3}{n+1} > x \Leftrightarrow n+1 > \frac{3}{2-x} \Leftrightarrow n > \frac{3}{2-x} - 1.$$

Taking  $x = \sup(B)$  and applying Archimedes' Postulate,  $\exists N \in \mathbb{N}$  such that

$$N > \frac{3}{2 - \sup(B)} - 1,$$

$$\Leftrightarrow 2 - \frac{3}{N+1} > \sup(B),$$

which is a contradiction as  $2 - \frac{3}{N+1} \in B$ . Hence  $\sup(B) \geq 2$ , and by combining our found inequalities,  $\sup(B) = 2$ .

## Method 2 — Alternative Characterisation of Suprema

### Solution.

First, note for  $n \in \mathbb{N}$ :

$$\frac{2n-1}{n+1} = \frac{2n+2-3}{n+1} = 2 - \frac{3}{n+1} < 2.$$

Hence  $B$  is bounded above by 2. Therefore as  $B \neq \emptyset$ , the completeness axiom says that  $\sup(B)$  exists.

We claim that  $\sup(B) = 2$ . Fix  $\epsilon > 0$ . Then, for  $n \in \mathbb{N}$ :

$$2 - \frac{3}{n+1} > 2 - \epsilon,$$

$$\Leftrightarrow \epsilon > \frac{3}{n+1},$$

$$\Leftrightarrow n\epsilon > 3 - \epsilon,$$

$$\Leftrightarrow n > \frac{3-\epsilon}{\epsilon}.$$

Now, by Archimedes' Postulate,  $\exists N \in \mathbb{N}$  such that  $N > \frac{3-\epsilon}{\epsilon}$ , from which

$$2 - \frac{3}{N+1} > 2 - \epsilon.$$

At this stage, take  $b = 2 - \frac{3}{N+1} \in B$ . Since  $\epsilon > 0$  was arbitrary, we have that  $\forall \epsilon > 0, \exists b \in B$  such that  $b > 2 - \epsilon$ . So, by the alternative characterisation of suprema (Theorem 2.1),  $\sup(B) = 2$ .

## Method 3 — Limits

Note that this doesn't work in general, but it might be quicker when you can use it. It relies on the following theorem (which we'll eventually cover):

**Theorem 0.1.**

A bounded, increasing sequence  $(b_n)_{n \in \mathbb{N}}$  is convergent, and its limit is given by

$$\lim_{n \rightarrow \infty} b_n = \sup\{b_n \mid n \in \mathbb{N}\}.$$

**Solution.**

Define  $b_n = \frac{2n-1}{n+1}$  for  $n \in \mathbb{N}$ .

**Step 1 — Show  $(b_n)_n$  is bounded above:**

First, note for  $n \in \mathbb{N}$ :

$$b_n = \frac{2n+2-3}{n+1} = 2 - \frac{3}{n+1} < 2.$$

Hence  $B$  is bounded above by 2. Therefore, as  $B \neq \emptyset$ , the completeness axiom says that  $\sup(B)$  exists.

**Step 2 — Show  $(b_n)_n$  is increasing (i.e. show  $b_{n+1} \geq b_n \forall n \in \mathbb{N}$ ):**

We have for  $n \in \mathbb{N}$ ,

$$\begin{aligned} b_{n+1} - b_n &= 2 - \frac{3}{n+2} - \left(2 - \frac{3}{n+1}\right), \\ &= \frac{3(n+2) - 3(n+1)}{(n+1)(n+2)}, \\ &= \frac{3}{(n+1)(n+2)}, \\ &\geq 0. \end{aligned}$$

So  $(b_n)$  is increasing. Hence, by the above theorem,  $(b_n)$  converges, and by the **Algebra of Limits**,

$$\sup(B) = \lim_{n \rightarrow \infty} b_n = \lim_{n \rightarrow \infty} \left(2 - \frac{\frac{3}{n}}{1 + \frac{1}{n}}\right) = 2,$$

as expected!