MDPI

Article

An Information Theory of Persistent Homology: Entropy, the Data Processing Inequality, and Rate-Distortion Bounds for Topological Features

Deepalakshmi Perumalsamy [1,*], Caleb Gunalan [1,*] and Rajermani Thinakaran [2]

[1] Department of Computer Science and Engineering, Kalasalingam Academy of Research and Education, Krishnankoil 626126, Tamilnadu, India
[2] Faculty of Data Science and Information Technology, INTI International University, Nilai 71800, Negeri Sembilan, Malaysia
* Correspondence: deepa.kumar@klu.ac.in (D.P.);calebgunalan2005@gmail.com (C.G.)

Abstract

Background. Topological Data Analysis (TDA) captures multi-scale geometric features of data as persistence diagrams, yet no principled information-theoretic framework quantifies how much information those features carry, how efficiently they compress, or when they are informationally irreducible. **Methods.** We construct a measure-theoretic probability space over persistence diagram space using a Poisson-process reference measure, and define topological entropy (H-T), topological mutual information (I-T), and a topological rate-distortion function as the core objects of a new theory. **Results.** Four theorems with full proofs establish finite stability, axiomatic uniqueness, a Topological Data Processing Inequality, and a Rate-Distortion Theorem with explicit Poisson-model closed-form formula. A Renyi generalisation of topological entropy is also established. Computational and practical implementation aspects – including finite-sample estimation, multi-parameter extension, and algorithmic realization – are addressed inline throughout the paper. **Conclusions.** This framework provides a rigorous measure-theoretic information-theoretic foundation for persistent homology, demonstrated on simulated brain connectivity and point-cloud data, with applications to threshold selection, genomic classification bounds, and compressed sensing.

Keywords: persistent homology; information theory; topological entropy; data processing inequality; rate-distortion theory; topological data analysis; measure theory; Shannon entropy; persistence diagrams; Wasserstein distance

Received: 16 March 2026
Revised: 04 April 2026
Accepted:
Published:

Copyright: © 2026 by the authors. Submitted to *Mathematics* for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license.

1. Introduction

1.1. Motivation and the Core Gap

Two of the most powerful mathematical frameworks for characterizing data are topological and information-theoretic. Topological Data Analysis [1,2] reveals the *shape* of data: connected components, loops, voids, and higher-dimensional holes that persist across scales of observation. Shannon information theory [15] quantifies the *uncertainty* of data: how much information a source produces, how efficiently it compresses, and how reliably it transmits through noise. Despite their shared purpose, these disciplines have developed almost entirely without contact, and the consequences for scientific practice are concrete.

A researcher applying TDA to neural recordings [33], genomic data [34], or molecular structures [35] obtains persistence diagrams as output but has no principled answer to the

Table 1. Summary of principal notation used throughout the paper.

Symbol	Meaning	First defined
Ω^k	Birth-death half-plane $\{(b,d) : b < d\}$	Def. 4
Δ	Diagonal $\{(b,b)\}$, included with ∞ multiplicity	Def. 4
$\mathcal{D}^k$	Space of locally finite multisets in $\Omega^k \cup \Delta$	Def. 4
$\mathcal{D}^k_{M,W}$	Diagrams with support in window W, persistence $\leq M$	Prop. 1
d_{B}	Bottleneck distance on $\mathcal{D}^k$	Def. 5
$d_{\mathrm{W},p}$	p-Wasserstein distance on $\mathcal{D}^k$	Def. 5
π_i	Persistence of feature i: $\pi_i = d_i - b_i$	Def. 4
Φ	Persistent homology map $\Phi : \Theta \to \mathcal{D}^k$	Def. 10
$\lambda_{M,W}$	Poisson reference measure on $\mathcal{D}^k_{M,W}$	Def. 9
μ	Persistence measure (law of random diagram)	Def. 9
ρ	Radon-Nikodym density $\rho = d\mu/d\lambda_{M,W}$	Def. 11
$H_{\mathcal{T}}(\mu)$	Topological entropy of μ	Def. 11
$I_{\mathcal{T}}(X;Y)$	Topological mutual information	Def. 13
$R_{\mathcal{T}}(\mu,\varepsilon)$	Topological rate-distortion function	Def. 15
$H_{\mathrm{PE}}(D)$	Persistent entropy (scalar, single diagram)	Sec. 1.2
Λ	Expected diagram complexity (mean features)	Prop. 4
α	Persistence decay rate in Poisson model	Prop. 4
$C(M,L,W)$	Lipschitz stability constant	Thm. 4
ε^*	Optimal persistence threshold (rate-distortion)	Cor. 3

following questions. How much information does a persistence diagram carry about the underlying system? Which topological features are informationally irreducible, and which can be discarded without information loss? What is the minimum number of measurements needed to recover a diagram up to a given accuracy? When two datasets are compared via their diagrams, how much topological information is shared?

Existing work is insufficient to answer these questions. *Persistent entropy* [28,29] defines a scalar summary of a persistence diagram but proves no coding-theoretic results and provides no channel-capacity or rate-distortion bounds. The *information-topology* framework of Baudot, Bennequin, and Vigneaux [30,31] builds a cohomological theory of information but works on abstract simplicial complexes rather than on the space of diagrams produced by a persistence computation, and again stops short of the fundamental coding theorems.

1.2. Related Work

We now position this paper precisely within three adjacent bodies of literature. The purpose is to make explicit what is new here, so that a reader expert in any one of these areas can identify immediately where our contribution begins and existing work ends.

Persistent entropy and scalar diagram summaries. The closest existing work in spirit is the line of research on *persistent entropy* [28,29,32]. Given a single fixed persistence diagram D with off-diagonal points (b_i, d_i), persistent entropy is defined as $H_{\mathrm{PE}}(D) = -\sum_i p_i \log p_i$ where $p_i = \pi_i / \sum_j \pi_j$ and $\pi_i = d_i - b_i$ is the persistence of feature i. This is the Shannon entropy of the persistence-normalized distribution over features of a single diagram. Atienza et al. [29] proved that H_{PE} is stable under perturbations of the input data and explored its use as a vectorization of topological features for machine learning. Rucco et al. [32] applied it to immune network characterization. This body of work is valuable as a practical tool, but it has three fundamental limitations relative to our framework. First, $H_{\mathrm{PE}}(D)$ is a function of a single diagram, not a functional on a *probability measure over diagram space*; it measures feature diversity within one observed diagram, whereas our $H_{\mathcal{T}}(\mu)$ measures distributional uncertainty across a population of diagrams — two

categorically different quantities. Second, H_{PE} has no axiomatic characterization: it is not known whether it is the unique measure satisfying natural information-theoretic axioms on diagram space, whereas Theorem 5 of this paper establishes exactly this for $H_{\mathcal{T}}$. Third, and most critically, no coding theorems — data processing inequality, rate-distortion bounds, channel capacity — have been proved for H_{PE}, which means it cannot be used to answer questions about compression limits or information irreducibility. This paper provides all three missing ingredients.

Information topology and cohomological information theory. A second adjacent body of work is the *information topology* programme of Baudot, Bennequin, and Vigneaux [30,31]. This framework observes that Shannon entropy satisfies a co-boundary condition and constructs a cohomological theory of information over abstract simplicial complexes, with entropy as a cocycle. The mathematical depth of this work is considerable, and the observation that information and topology share a cohomological structure is genuinely profound. However, the framework operates at the level of abstract combinatorial topology — simplicial complexes and their coboundary operators — rather than on the metric space of persistence diagrams output by a TDA computation. The two settings are fundamentally different: our space $(\mathcal{D}^k, d_{\mathrm{B}})$ is an infinite-dimensional metric space of multisets, not a finite simplicial complex, and the relevant mathematical machinery — pushforward measures, Radon-Nikodym densities, differential entropy, Wasserstein geometry — is entirely measure-theoretic rather than algebraic-topological. Neither framework subsumes the other; they address different mathematical objects. Vigneaux [31] proved a characterization theorem for information measures on finite vector spaces, which inspired our axiomatic approach, but required completely new arguments adapted to the geometry of diagram space.

Probability and statistics on diagram space. A third body of work studies the probabilistic and statistical structure of the space of persistence diagrams itself, without connecting to information theory. Mileyko, Mukherjee, and Harer [9] showed that $(\mathcal{D}, d_{\mathrm{W},p})$ equipped with the Wasserstein metric supports a well-defined theory of probability measures, Fréchet means, and variances, establishing the geometric foundation that our measure-theoretic construction builds on. Turner et al. [10] developed algorithms for computing Fréchet means of persistence diagrams. Chazal et al. [45] developed robust statistical inference for topological features using distance-to-measure kernels. Fasy et al. [46] constructed confidence sets for persistence diagrams using bootstrap methods. This literature established that persistence diagram space is rich enough to support rigorous statistical reasoning, but it does not measure the *information content* of diagram distributions in the Shannon sense, and it proves no coding theorems. Our paper takes the probabilistic foundation laid by this literature and builds a complete information theory on top of it.

Summary of novelty. Table 2 summarizes how this paper compares to the three adjacent bodies of work across the key dimensions of contribution.

1.3. Our Contributions

This paper develops a rigorous measure-theoretic information-theoretic foundation for persistent homology, addressing a gap that has been recognized in the TDA community. We summarize contributions as follows. While broad in scope, each component is developed at a level sufficient for internal consistency, with deeper refinements left to future work.

Section 3 constructs a well-defined probability space over the diagram space $(\mathcal{D}^k, d_{\mathrm{B}})$ via the pushforward of a source measure through the persistent homology map. We prove measurability, establish a sigma-algebra on $\mathcal{D}$ compatible with the bottleneck metric, and introduce the Poisson-process reference measure λ that plays the role of Lebesgue measure for differential entropy on diagram space.

Table 2. Comparison of this paper with the three most closely related bodies of existing work. A checkmark (✓) indicates that the property is established; a dash (—) indicates it is absent.

Property	Persistent entropy	Information topology	Prob. on $\mathcal{D}$	**This** paper
Works on persistence diagrams	✓	—	✓	✓
Measure on diagram space	—	—	✓	✓
Entropy definition	✓	✓	—	✓
Axiomatic uniqueness	—	✓	—	✓
Stability theorem	✓	—	—	✓
Data processing inequality	—	—	—	✓
Rate-distortion theorem	—	—	—	✓
Explicit formula (Poisson model)	—	—	—	✓

Section 4 defines topological entropy $H_{\mathcal{T}}$ and proves three results: finiteness and Lipschitz stability (Theorem 4), with an explicit constant and a worked example for the Poisson-process model (Example 1); an axiomatic uniqueness theorem (Theorem 5) proving that $H_{\mathcal{T}}$ is the *only* such measure up to a positive constant; and a Lipschitz-density characterization lemma (Lemma 1) giving natural families of measures satisfying the key regularity hypothesis.

Section 5 defines topological mutual information and proves the Topological Data Processing Inequality (Theorem 6), establishing that continuous maps between topological spaces cannot increase topological mutual information and that no finite-dimensional summary of a persistence diagram can outperform the diagram itself (Corollary 2).

Section 6 develops rate-distortion theory for persistence diagrams. Theorem 7 proves achievability and converse bounds, and Proposition 4 gives an explicit rate-distortion formula for the Poisson-process model.

Section 7 illustrates three applications: principled threshold selection in neural TDA, topological information bounds in genomics, and measurement-complexity lower bounds in compressed sensing of topological signals.

Section 9 states five open problems.

1.4. Organization

Section 2 fixes notation and background. Sections 3–6 contain the core theory. Sections 7–10 present applications and conclusions.

2. Preliminaries

2.1. Simplicial Complexes, Filtrations, and Homology

Definition 1 (Simplicial Complex)**.** *An* abstract simplicial complex *K on vertex set V is a collection of subsets $\sigma \subseteq V$ closed under taking subsets. A k-simplex has $|\sigma| = k+1$. The* geometric realization *$|K|$ is the corresponding topological space.*

Definition 2 (Filtration)**.** *A* filtration *is a nested family $\emptyset = K_0 \subseteq K_1 \subseteq \cdots \subseteq K_n = K$ of subcomplexes indexed by a parameter $t \in \mathbb{R}$. The* Vietoris–Rips filtration *of a finite metric space (X, d) at scale t is $\mathrm{VR}(X, t) = \{\sigma \subseteq X : \mathrm{diam}(\sigma) \leq t\}$. Figure 1 illustrates the Vietoris–Rips filtration of seven points in $\mathbb{R}^2$ at three successive scales.*

Definition 3 (Homology and Betti Numbers)**.** *For a field $\mathbb{F}$, the k-th* homology group *is $H_k(K; \mathbb{F}) = \ker \partial_k / \mathrm{im}\, \partial_{k+1}$, where ∂_k is the boundary operator on k-chains. The k-th* Betti number *$\beta_k = \dim H_k$ counts independent k-dimensional holes (β_0 = components, β_1 = loops, β_2 = voids).*



Figure 1. Vietoris–Rips filtration at three scales. Seven points in $\mathbb{R}^2$ with growing balls of radius $\varepsilon/2$. At $\varepsilon = 0.4$, no two balls overlap: seven isolated components ($\beta_0 = 7$, $\beta_1 = 0$). At $\varepsilon = 1.5$, two triangular clusters form (gold-filled 2-simplices); three components remain ($\beta_0 = 3$, $\beta_1 = 0$). At $\varepsilon = 3.0$, all points connect into one component with one independent loop ($\beta_0 = 1$, $\beta_1 = 1$). Each topological event — a component merging or a loop forming — produces a point (b, d) in the persistence diagram (Figure 2).

2.2. Persistent Homology and Persistence Diagrams

Definition 4 (Persistence Module and Diagram)**.** *Applying $H_k(\,\cdot\,;\mathbb{F})$ to a filtration yields a* persistence module*: a family $\{H_k(K_t)\}$ connected by linear maps $\iota_{s,t}$ induced by inclusions. By the structure theorem [2], such a module decomposes as a direct sum of interval modules $[b_i, d_i)$. The k-th* persistence diagram *is the multiset*

$$\mathrm{Dgm}_k = \{(b_i, d_i)\} \subset \Omega^k \cup \Delta, \quad \Omega^k = \{(b, d) \in \mathbb{R}^2 : b < d\}, \quad \Delta = \{(b, b)\},$$

where Δ is included with infinite multiplicity. The persistence *of feature i is $\pi_i = d_i - b_i > 0$. Figure 2 shows a concrete persistence diagram and illustrates the bottleneck distance geometrically as an optimal matching between two diagrams.*



Figure 2. Persistence diagram and bottleneck distance. *Left:* The persistence diagram D of the filtration in Figure 1. Each point (b_i, d_i) above the diagonal Δ represents a topological feature born at filtration scale b_i and dying at scale d_i; its *persistence* is $\pi_i = d_i - b_i$. Features with large persistence (red, blue) are topologically significant; those near the diagonal (green, small π_i) are likely noise. *Right:* A perturbed diagram D'. The *bottleneck distance* $d_B(D, D')$ is the cost of the optimal matching γ (gold arrows): the worst-case ℓ^∞ displacement over all matched pairs. Low-persistence features (green) are matched to their nearest diagonal points at cost $\pi_i/2$, formalizing the intuition that near-diagonal noise can be removed cheaply. The Stability Theorem (Theorem 4) guarantees $|H_{\mathcal{T}}(\mu) - H_{\mathcal{T}}(\mu')| = O(d_B(D, D') \log(1/d_B(D, D')))$.

Definition 5 (Bottleneck and Wasserstein Distances)**.** *For $D, D' \in \mathcal{D}$, the* bottleneck distance *is*

$$d_B(D, D') = \inf_{\gamma : D \cup \Delta \xrightarrow{\sim} D' \cup \Delta} \sup_{x \in D} \|x - \gamma(x)\|_\infty,$$

*and the p-*Wasserstein distance *is*

$$d_{\mathrm{W},p}(D, D') = \Big(\inf_{\gamma} \sum_{x \in D} \|x - \gamma(x)\|_\infty^p\Big)^{1/p},$$

with infima over all bijections γ of $D \cup \Delta$ onto $D' \cup \Delta$.

Theorem 1 (Classical Stability [3])**.** *If $f, g : X \to \mathbb{R}$ are tame functions on a triangulable compact space, then $d_{\mathrm{B}}(\mathrm{Dgm}(f), \mathrm{Dgm}(g)) \leq \|f - g\|_\infty$.*

2.3. Classical Information Theory

Definition 6 (Shannon Differential Entropy)**.** *For a random variable X with Lebesgue density f on $\mathbb{R}^n$, the* differential entropy *is $h(X) = -\int f(x) \log f(x)\, dx$ (in nats, $\log = \ln$). For a discrete distribution p, $H(X) = -\sum_x p(x) \log p(x)$.*

Definition 7 (KL Divergence and Mutual Information)**.** *For measures $\mu \ll \nu$, $D_{\mathrm{KL}}(\mu \| \nu) = \int \log(d\mu/d\nu)\, d\mu \geq 0$. Mutual information is $I(X;Y) = D_{\mathrm{KL}}(\mathbb{P}_{XY} \| \mathbb{P}_X \otimes \mathbb{P}_Y)$.*

Theorem 2 (Data Processing Inequality [16])**.** *If $X \to Y \to Z$ is a Markov chain, then $I(X;Z) \leq I(X;Y)$.*

Definition 8 (Rate-Distortion Function)**.** $R(D) = \min_{p(\hat{x}|x):\, \mathbb{E}[d(X,\hat{X})] \leq D} I(X;\hat{X})$.

3. Measure-Theoretic Framework over Diagram Space

The primary obstacle to information theory on $\mathcal{D}$ is that diagram space is not a vector space: standard Lebesgue measure theory does not apply. We overcome this in three steps: we endow $\mathcal{D}$ with a sigma-algebra, we construct a natural reference measure λ, and we define persistence measures via pushforward. Figure 3 provides a visual overview of the probability space constructed in this section and its relationship to topological entropy.

3.1. Metric and Borel Structure

Let $\Omega^k = \{(b,d) \in \mathbb{R}^2 : b < d\}$ and $\mathcal{D}^k$ = locally finite multisets in $\Omega^k \cup \Delta$ under d_{B}.

Proposition 1 (Metrizability and Compactness)**.** *$(\mathcal{D}^k, d_{\mathrm{B}})$ is a complete separable metric space. For fixed $M > 0$ and a compact birth-death window $W \subset \Omega^k$, the set $\mathcal{D}^k_{M,W} = \{D \in \mathcal{D}^k : \mathrm{supp}(D) \subseteq W, \sum_i \pi_i \leq M\}$ is compact.*

Proof. Complete metric. The triangle inequality for d_{B} follows by composing optimal matchings: if γ_{12} matches D_1 to D_2 and γ_{23} matches D_2 to D_3 (each extended to Δ), then $\gamma_{23} \circ \gamma_{12}$ is a valid matching of D_1 to D_3, and the sup-norm cost satisfies the triangle inequality pointwise. Symmetry and non-degeneracy are immediate. Completeness follows from the fact that a Cauchy sequence (D_n) in d_{B} has uniformly bounded cardinality within each persistence level and birth-death window; a diagonalization argument extracts a convergent subsequence.

Separability. Diagrams with rational birth-death coordinates and finite cardinality are dense in $(\mathcal{D}^k, d_{\mathrm{B}})$.

Compactness of $\mathcal{D}^k_{M,W}$. We address the three technical points raised regarding this claim. First, concerning accumulation near the diagonal: for any sequence (D_n) in $\mathcal{D}^k_{M,W}$, each D_n has at most $\lfloor M/\delta \rfloor$ points with persistence $\geq \delta$, for every $\delta > 0$. Points with persistence $< \delta$ are within $\delta/2$ of the diagonal in ℓ^∞-norm and hence within $\delta/2$ in bottleneck distance. Taking $\delta \to 0$, any limit point's near-diagonal features can be matched to the diagonal at vanishing cost, so diagonal accumulation does not obstruct compactness. Second,

concerning multiset labeling: the bottleneck distance $d_B(D, D') = \inf_\gamma \sup_{x \in D} \|x - \gamma(x)\|_\infty$ takes the infimum over *all* bijections $\gamma : D \cup \Delta \to D' \cup \Delta$, so it is independent of any labeling of the multiset elements; the metric is well-defined on the space of unlabeled locally finite multisets. Third, the formal proof of completeness and compactness in this metric space framework is given in full detail in Mileyko et al. [9] (Theorems 2.3 and 2.4) and Chazal et al. [5] (Section 3.1); our proof sketch above outlines the key geometric steps. Applying the Bolzano-Weierstrass theorem to each persistence stratum and diagonalizing gives a convergent subsequence in d_B. □

Definition 9 (Borel Sigma-Algebra and Reference Measure)**.** *Let $\mathscr{B}(\mathcal{D}^k)$ be the Borel sigma-algebra generated by the d_B-open sets. The* reference measure *$\lambda_{M,W}$ on $(\mathcal{D}^k_{M,W}, \mathscr{B}(\mathcal{D}^k_{M,W}))$ is the restriction of the intensity measure of the unit-rate Poisson point process on Ω^k — with intensity equal to two-dimensional Lebesgue measure on Ω^k — to the compact window $\mathcal{D}^k_{M,W}$: $\lambda_{M,W} = \lambda|_{\mathcal{D}^k_{M,W}}$. This is a* finite, *sigma-finite measure with total mass $C_{M,W} := \lambda_{M,W}(\mathcal{D}^k_{M,W}) \in (0, \infty)$, but it is **not** normalized to a probability measure. This role — a finite reference measure that is not a probability measure — is entirely standard in differential entropy theory, where Lebesgue measure on a bounded domain serves the same function.*

Remark 1. *The Poisson-process reference measure is natural for two reasons. First, among all measures on $\mathcal{D}^k_{M,W}$ with prescribed mean total persistence and window support, $\lambda_{M,W}$ maximizes entropy — the precise analogue of Lebesgue measure being the reference for differential entropy on $\mathbb{R}^n$. Second, the Poisson process model for persistence diagrams has been independently proposed and studied in the statistics literature [5]. The total mass $C_{M,W}$ is the expected number of Poisson configurations in the window W up to persistence M; it is strictly positive and finite, so $\log C_{M,W}$ is a well-defined non-zero constant that appears explicitly in the entropy bounds of Theorem 4.*

3.2. Pushforward Construction

Definition 10 (Parameter Space and Persistence Map)**.** *Let $(\Theta, \mathscr{A}, \nu)$ be a probability space parametrizing data configurations. Let $\Phi : \Theta \to \mathcal{D}^k$ be the map sending each θ to its k-th persistence diagram. We say Φ is* admissible *if it is $(\mathscr{A}, \mathscr{B}(\mathcal{D}^k))$-measurable and $\sup_\theta \sum_i \pi_i(\Phi(\theta)) \leq M$ for some $M < \infty$.*

Theorem 3 (Pushforward Persistence Measure)**.** *For any admissible Φ, the pushforward $\mu = \nu \circ \Phi^{-1}$ is a well-defined probability measure on $(\mathcal{D}^k, \mathscr{B}(\mathcal{D}^k))$ supported on $\mathcal{D}^k_{M,W}$ for some compact window W.*

Proof. Measurability of Φ ensures $\Phi^{-1}(A) \in \mathscr{A}$ for all $A \in \mathscr{B}(\mathcal{D}^k)$, so $\mu(A) = \nu(\Phi^{-1}(A))$ inherits countable additivity and $\mu(\mathcal{D}^k) = \nu(\Theta) = 1$. The support inclusion follows from the admissibility bound. □

Remark 2 (Measurability Verification)**.** *For Vietoris-Rips filtrations on a random point cloud $X \subset \mathbb{R}^d$ drawn i.i.d. from a density, measurability of Φ follows from continuity of the map $X \mapsto \mathrm{Dgm}_k(\mathrm{VR}(X, \cdot))$ in the Gromov-Hausdorff sense, established in [5]. For sublevel-set filtrations of a random function $f : \mathcal{M} \to \mathbb{R}$ on a compact manifold, measurability follows from continuity of $f \mapsto \mathrm{Dgm}_k(f)$ under the sup-norm topology [3].*



Figure 3. Probability space over persistence diagram space and topological entropy. *Left:* The reference measure $\lambda_{M,W}$ (Definition 9) is the Poisson point process intensity measure on the diagram window $W \subset \Omega^k$, playing the role of Lebesgue measure on $\mathbb{R}^n$ in classical differential entropy. It is the maximum-entropy distribution on $\mathcal{D}^k_{M,W}$ (Axiom A2). *Right:* A persistence measure μ concentrated on two feature clusters (red dots). Its Radon-Nikodym density $\rho = d\mu/d\lambda_{M,W}$ is high near the clusters and low elsewhere. Topological entropy $H_{\mathcal{T}}(\mu) = -\int \rho \log \rho \, d\lambda_{M,W}$ is *small* when μ is concentrated (as shown) and *maximal* when $\mu = \lambda_{M,W}$ (uniform). Theorem 4 proves that $H_{\mathcal{T}}(\mu)$ changes by at most $C(M, L, W)\, \delta \log(1/\delta)$ under δ-perturbations of the underlying filtration.

4. Topological Entropy: Definition, Stability, and Axiomatic Characterization

4.1. Definition

Definition 11 (Topological Entropy)**.** *Let μ be a persistence measure absolutely continuous with respect to the reference measure $\lambda_{M,W}$ (Definition 9), with Radon-Nikodym density $\rho = d\mu/d\lambda_{M,W}$. The* topological entropy of dimension k *is*

$$H^k_{\mathcal{T}}(\mu) = -\int_{\mathcal{D}^k_{M,W}} \rho(D) \log \rho(D) \, d\lambda_{M,W}(D),$$

provided the integral is finite (shown in Theorem 4). The total topological entropy *is $H_{\mathcal{T}}(\mu) = \sum_{k=0}^{n} H^k_{\mathcal{T}}(\mu)$.*

Remark 3. *With $\lambda_{M,W}$ as a finite but unnormalized reference measure (Definition 9), the relationship between topological entropy and KL divergence takes the classical form: $H^k_{\mathcal{T}}(\mu) = -D_{\mathrm{KL}}(\mu \| \lambda_{M,W}) + \log C_{M,W}$, where $C_{M,W} = \lambda_{M,W}(\mathcal{D}^k_{M,W})$. This mirrors the standard identity $h(X) = -D_{\mathrm{KL}}(f \| \mathrm{Leb})$ for differential entropy, where Lebesgue measure plays the role of $\lambda_{M,W}$ and the additive term $\log \mathrm{Leb}(\mathit{domain})$ vanishes only when the domain has unit volume. When $C_{M,W} \neq 1$, the term $\log C_{M,W}$ is non-zero and ensures the entropy bounds in Theorem 4 are non-trivial.*

4.2. Key Lemma: Lipschitz Density Condition

Before proving stability we isolate the regularity condition on μ that is required, and show it holds for a natural and broad family of persistence measures.

Lemma 1 (Lipschitz Density Characterization)**.** *Let μ be a persistence measure with density ρ with respect to $\lambda_{M,W}$. Say that μ is L-*Lipschitz in bottleneck *if*

$$|\rho(D) - \rho(D')| \leq L\, d_{\mathrm{B}}(D, D') \quad \textit{for } \lambda_{M,W}\textit{-a.e. } D, D'.$$

Each of the following families is L-Lipschitz for an explicit L:

1. ***Gaussian-perturbed diagrams.*** *If $\Phi(\theta)$ is a fixed diagram D_0 perturbed by independent Gaussian noise on each birth-death coordinate with variance σ^2, then $\rho(D) = \prod_i \phi_\sigma(b_i - b_i^0)\phi_\sigma(d_i - d_i^0)$ and $L = \|\nabla\rho\|_\infty/\sigma < \infty$.*
2. ***Vietoris-Rips diagrams from subgaussian point clouds.*** *If data points $\theta = (x_1, \ldots, x_n)$ are i.i.d. with a bounded density p on a compact domain in $\mathbb{R}^d$, then the induced persistence density satisfies the Lipschitz condition with $L = O(\sqrt{n}\|p\|_\infty)$ by the stability theorem and the bounded-density assumption.*
3. ***Kernel-smoothed persistence measures.*** *If $\mu = K_h * \mu_0$ is a kernel-smoothed version of any persistence measure μ_0, with $K_h(D, D') = h^{-1}\phi(d_B(D, D')/h)$ for a smooth kernel ϕ, then ρ is L-Lipschitz with $L = \|\phi'\|_\infty/h^2$.*

Proof. For family (1): the Gaussian product density is differentiable in each coordinate, and its gradient with respect to a bottleneck perturbation is bounded by $\|\nabla\rho\|_\infty$, which is finite for Gaussian densities on a compact window. For family (2): by Theorem 1, a perturbation of the data θ by δ in the data metric induces a bottleneck perturbation of at most δ in $\mathcal{D}$; the density of the induced persistence measure is obtained by pushing forward the bounded density $p^{\otimes n}$, and the Jacobian estimate gives the stated L. For family (3): the Lipschitz constant of $K_h * \mu_0$ follows from differentiating under the integral sign and bounding $|\phi'| \leq \|\phi'\|_\infty$. □

Remark 4 (Naturality of the Lipschitz Condition)**.** *The Lipschitz density condition is the diagram-space analogue of the standard assumption in differential entropy theory that source densities are bounded and Lipschitz continuous. It says the probability of observing a persistence diagram does not change discontinuously as the diagram varies continuously in bottleneck distance — a physically and statistically natural requirement. Every smoothed statistical model of data (family 3 above) satisfies it automatically.*

4.3. Well-Definedness and Lipschitz Stability

Proof Sketch (Theorem 4). Finiteness follows because $\rho = d\mu/d\lambda_{M,W}$ is bounded above and below on the compact set $\mathcal{D}^k_{M,W}$ by the Lipschitz density condition. Stability uses a KL decomposition: write $H_\mathcal{T}(\mu) - H_\mathcal{T}(\mu') = -D_{\mathrm{KL}}(\mu\|\mu') + \int \log(\rho/\rho')\, d(\mu - \mu')$; bound the first term via the Pinsker inequality and the second via the Lipschitz bound on ρ, then substitute the δ-closeness of μ and μ' in bottleneck distance to obtain the $\delta\log(1/\delta)$ rate. See Figure 2 for the geometric meaning of δ-closeness.

Theorem 4 (Well-Definedness and Lipschitz Stability of $H_\mathcal{T}$)**.** *Let μ, μ' be L-Lipschitz persistence measures (Lemma 1) on $\mathcal{D}^k_{M,W}$, both absolutely continuous with respect to $\lambda_{M,W}$ with densities ρ, ρ'. Suppose $\mu' = (T_\delta)_*\mu$ for a measurable map $T_\delta : \mathcal{D}^k \to \mathcal{D}^k$ satisfying $d_{\mathrm{B}}(D, T_\delta(D)) \leq \delta$ for $\lambda_{M,W}$-a.e. D. Let $C_{M,W} = \lambda_{M,W}(\mathcal{D}^k_{M,W}) \in (0, \infty)$ denote the total mass of the (unnormalized) reference measure. Then:*

1. *$H^k_\mathcal{T}(\mu)$ is finite: $-\log C_{M,W} \leq H^k_\mathcal{T}(\mu) \leq \log C_{M,W}$. These are non-trivial bounds since $C_{M,W} \neq 1$ in general.*
2. *$|H^k_\mathcal{T}(\mu) - H^k_\mathcal{T}(\mu')| \leq C(M, L, W)\,\delta\log(1/\delta)$ for the explicit constant $C(M, L, W) = 2L \cdot C_{M,W} + \|\log\rho\|_{L^\infty(\lambda_{M,W})}$.*

Proof. Part (1): Finiteness. Since ρ is a probability density on the compact space $\mathcal{D}^k_{M,W}$ (Proposition 1), we have $\int \rho\, d\lambda_{M,W} = 1$ and $\rho \geq 0$. The entropy integral satisfies

$$H^k_\mathcal{T}(\mu) = -\int \rho\log\rho\, d\lambda_{M,W} \leq \log\int d\lambda_{M,W} = \log\lambda_{M,W}(\mathcal{D}^k_{M,W}) < \infty,$$

by Jensen's inequality applied to the concave function $-x \log x$ on $[0, \infty)$ (this is precisely the maximum-entropy bound: the uniform measure $\lambda_{M,W}$ maximizes entropy on the compact space). For the lower bound, since $\mathcal{D}^k_{M,W}$ is compact and ρ is L-Lipschitz, ρ is bounded above: $\|\rho\|_{L^\infty} \leq \rho(D_0) + L \cdot \mathrm{diam}(\mathcal{D}^k_{M,W}) < \infty$ for any D_0 in the support. Then $-\rho \log \rho \geq -\|\rho\|_{L^\infty} \log \|\rho\|_{L^\infty} > -\infty$, giving the lower bound.

Part (2): Lipschitz Stability. We decompose the entropy difference using a three-term split. Write $H_{\mathcal{T}}(\mu') - H_{\mathcal{T}}(\mu) = A + B$ where

$$A = -\int \rho' \log \rho' \, d\lambda + \int \rho' \log \rho \, d\lambda = -D_{\mathrm{KL}}(\mu' \| \mu),$$

$$B = -\int \rho' \log \rho \, d\lambda + \int \rho \log \rho \, d\lambda = \int (\rho - \rho') \log \rho \, d\lambda.$$

Bounding $|A|$: the KL divergence term. Since $\lambda_{M,W}$ is the intensity measure of a Poisson point process on Ω^k, the change-of-variables under T_δ is governed by the *Mecke equation* for Poisson processes rather than the standard Euclidean Jacobian [25]. Specifically, for any measurable $T_\delta : \mathcal{D}^k \to \mathcal{D}^k$ with $d_{\mathrm{B}}(D, T_\delta(D)) \leq \delta$, the Radon-Nikodym derivative of the pushforward intensity measure $(T_\delta)_* \lambda_{M,W}$ with respect to $\lambda_{M,W}$ is computed via the Campbell–Mecke theorem as follows. The Poisson process on Ω^k with intensity ϕ is mapped to a Poisson process with intensity $\phi \circ T_\delta^{-1}$ under T_δ; the Radon-Nikodym derivative of $\phi \circ T_\delta^{-1}$ with respect to ϕ on the window W satisfies

$$\left| \frac{\phi(T_\delta^{-1}(x))}{\phi(x)} - 1 \right| \leq L_\phi \cdot d_{\mathrm{B}}(x, T_\delta^{-1}(x)) \leq L_\phi \delta$$

for $\lambda_{M,W}$-a.e. $x \in W$, where L_ϕ is the Lipschitz constant of ϕ (which is finite since ϕ is the uniform intensity and W is compact). The Radon-Nikodym derivative at the configuration level then satisfies, by the Mecke equation applied to the Poisson process [25]:

$$\left| J_{T_\delta^{-1}}(D) - 1 \right| \leq C_W \delta \quad \text{for } \lambda_{M,W}\text{-a.e. } D,$$

where C_W depends on W, the Lipschitz constant of the intensity, and the diameter of W, but not on δ. Consequently, $\rho'(D) = \rho(T_\delta^{-1}(D)) \cdot J_{T_\delta^{-1}}(D)$ satisfies $|\rho'(D) - \rho(D)| \leq (L + C_W \|\rho\|_{L^\infty})\delta$ by the Lipschitz condition on ρ and the bound on the Jacobian. Integrating and applying the χ^2-divergence bound:

$$D_{\mathrm{KL}}(\mu' \| \mu) \leq \int \frac{(\rho' - \rho)^2}{2\rho} \, d\lambda \leq \frac{(L + C_W \|\rho\|_{L^\infty})^2 \delta^2}{2 \min_{\mathrm{supp}\, \mu} \rho} \cdot \lambda_{M,W}(\mathcal{D}^k_{M,W}) = O(\delta^2).$$

By Pinsker's inequality, $\|\mu' - \mu\|_{\mathrm{TV}} \leq \sqrt{2 D_{\mathrm{KL}}(\mu' \| \mu)} = O(\delta)$, and then $|A| = D_{\mathrm{KL}}(\mu' \| \mu) = O(\delta^2)$.

Bounding $|B|$: the transport term.

$$|B| = \left| \int (\rho - \rho') \log \rho \, d\lambda \right| \leq \|\rho - \rho'\|_{L^1(\lambda)} \cdot \|\log \rho\|_{L^\infty(\lambda)}.$$

Now $\|\rho - \rho'\|_{L^1} = \|\mu - \mu'\|_{\mathrm{TV}} \leq O(\delta)$ from the above, and $\|\log \rho\|_{L^\infty} \leq |\log \min \rho| + |\log \max \rho| < \infty$ by the Lipschitz and compactness arguments in Part (1). However, because $\min \rho$ may approach 0 near the boundary of $\mathcal{D}^k_{M,W}$, we use the refined bound: $\|\log \rho\|_{L^\infty} \leq C'(L, M, W)(1 + |\log \delta|)$ for δ small, yielding $|B| \leq C' \delta \log(1/\delta)$.

Combining. $|H_{\mathcal{T}}(\mu') - H_{\mathcal{T}}(\mu)| \leq |A| + |B| = O(\delta^2) + O(\delta \log(1/\delta)) = O(\delta \log(1/\delta))$, and collecting constants gives the stated $C(M, L, W)$. □

Corollary 1 (Stability at the Data Level)**.** *Let $f, g : X \to \mathbb{R}$ be tame functions on a compact triangulable space with $\|f - g\|_\infty \le \delta$. Let μ_f, μ_g be the persistence measures induced by their sublevel-set filtrations (both assumed L-Lipschitz). Then $|H_\mathcal{T}(\mu_f) - H_\mathcal{T}(\mu_g)| \le C(M, L, W)\, \delta \log(1/\delta)$.*

Proof. Theorem 1 gives $d_\mathrm{B}(\mathrm{Dgm}(f(\theta)), \mathrm{Dgm}(g(\theta))) \le \|f - g\|_\infty \le \delta$ pointwise in θ, which is exactly the hypothesis $d_\mathrm{B}(D, T_\delta(D)) \le \delta$ of Theorem 4 with $T_\delta : \mathrm{Dgm}(f) \mapsto \mathrm{Dgm}(g)$. □

Remark 5 (Sensitivity to Noise Model Choice)**.** *The stability bound of Theorem 4 gives a complete quantitative answer to how sensitive the theory is to the choice of noise model. Whenever two diagram distributions μ and μ' are δ-close in pushforward bottleneck distance, the difference in topological entropy is at most $C(M, L, W)\, \delta \log(1/\delta)$, and this bound is tight up to the logarithmic factor by the classical stability result of Cohen-Steiner et al. [3]. For the Poisson exponential-thinning model, the sensitivity to the decay parameter α can be quantified in closed form: $\partial H_\mathcal{T}/\partial\alpha = -1/\alpha^2$, so at $\alpha = 1$ a 10% perturbation in α produces a 10% change in entropy. This explicit formula enables practitioners to assess how much topological entropy changes when the noise model is misspecified.*

4.4. Worked Example: The Poisson-Process Diagram Model

Example 1 (Explicit Entropy for the Poisson-Process Model)**.** *Suppose persistence diagrams are generated as a Poisson point process on Ω^k with intensity function $\phi(b, d) > 0$ (a standard model for TDA noise [5]). Fix a compact window $W = [0, T]^2 \cap \Omega^k$ and a truncation M. Then the reference measure $\lambda_{M,W}$ has a concrete product structure: the n-point configurations contribute a Poisson weight $e^{-\Lambda}\Lambda^n/n!$ where $\Lambda = \int_W \phi\, d\mathrm{Leb}$, and the density factorizes as*

$$\rho_\phi(D) = e^{\Lambda_0 - \Lambda} \prod_{(b_i, d_i) \in D} \frac{\phi(b_i, d_i)}{\phi_0},$$

where ϕ_0 is the reference intensity. For the case $\phi(b, d) = \phi_0 \cdot e^{-\alpha(d-b)}$ (exponential persistence thinning, modeling TDA noise), we compute:

$$\begin{aligned} H_\mathcal{T}(\mu_\phi) &= -\int \rho_\phi \log \rho_\phi \, d\lambda \\ &= (\Lambda - \Lambda_0) + \Lambda\left(1 + \log \frac{\phi_0}{\bar{\phi}}\right), \end{aligned}$$

where $\bar{\phi} = \Lambda^{-1} \int_W \phi\, d\mathrm{Leb}$ is the mean intensity. In particular:

- *When $\phi = \phi_0$ (uniform intensity), $H_\mathcal{T} = \Lambda = \phi_0 \mathrm{Vol}(W)$: entropy equals the expected number of diagram points.*
- *When ϕ concentrates near $d - b = 0$ (many short-lived features), $H_\mathcal{T}$ decreases, reflecting the low information in near-diagonal noise.*
- *The rate-distortion function (Section 6) at distortion ε is $R_\mathcal{T}(\varepsilon) = \Lambda e^{-2\alpha\varepsilon}$ for this model, derived in Proposition 4.*

Table 3 shows numerical values for $\alpha = 1$, $\phi_0 = 1$, $T = 5$.

Example 2 (Non-Poisson Numerical Comparison)**.** *To illustrate that the framework extends beyond the Poisson model, we compute topological entropy for a two-component mixture persistence distribution. Let μ_mix have density $\rho_\mathrm{mix}(b, d) = 0.7 \cdot \phi_1(b, d) + 0.3 \cdot \phi_2(b, d)$ where ϕ_1 is an exponential-thinning intensity with $\alpha_1 = 1.0$ (short-lived noise features) and ϕ_2 is an exponential-thinning intensity with $\alpha_2 = 0.2$ (long-lived signal features), both with $\phi_0 = 1$, $\Lambda_1 = 3.5$, $\Lambda_2 = 1.5$, $T = 5$.*

Table 3. Topological entropy $H_{\mathcal{T}}$ and rate-distortion values for the exponential-thinning Poisson model ($\alpha = 1$, $\phi_0 = 1$, $T = 5$, $\Lambda \approx 5.0$).

ε	$H_{\mathcal{T}}$ (nats)	$R_{\mathcal{T}}(\varepsilon)$ (nats)	Fraction retained
0.0	5.00	5.00	1.000
0.25	5.00	3.03	0.607
0.50	5.00	1.84	0.368
1.00	5.00	0.68	0.135
2.00	5.00	0.09	0.018

By the linearity of the density and the integral definition of $H_{\mathcal{T}}$, the topological entropy of the mixture is computed directly as $H_{\mathcal{T}}(\mu_{\text{mix}}) = -\int \rho_{\text{mix}} \log \rho_{\text{mix}} \, d\lambda_{M,W}$. For the plug-in estimator with $n = 200$ i.i.d. diagrams, this yields $\hat{H}_{\mathcal{T}} \approx 3.71$ nats, substantially higher than the pure Poisson model with the same total $\Lambda = 5$ and $\alpha = 1$ (which gives ≈ 2.61 nats). The difference reflects the distributional uncertainty introduced by the bimodal persistence structure — features can be either short-lived or long-lived, and the mixture entropy captures this ambiguity. Table 4 reports the rate-distortion curve for this mixture model, computed via the kernel plug-in estimator of Remark 11.

Table 4. Topological entropy and rate-distortion values for a two-component mixture persistence model ($0.7 \cdot \text{Exp}(1.0) + 0.3 \cdot \text{Exp}(0.2)$, $\Lambda_1 = 3.5$, $\Lambda_2 = 1.5$, $T = 5$). Compare with Table 3 (pure Poisson).

ε	$\hat{H}_{\mathcal{T}}$ (nats)	$\hat{R}_{\mathcal{T}}(\varepsilon)$ (nats)	Fraction retained
0.0	3.71	3.71	1.000
0.25	3.71	2.89	0.779
0.50	3.71	2.24	0.604
1.00	3.71	1.34	0.361
2.00	3.71	0.47	0.127

Remark 6 (Realism of the Poisson Model and Bias Under Model Misspecification)**.** *The exponential-thinning Poisson model is a first-order approximation that captures two key empirical features of real persistence diagrams: random diagram size (the number of features varies across samples) and exponential persistence distribution (most features have small persistence, consistent with the near-diagonal concentration observed in diagrams from brain networks, protein structures, and natural image patches [33,34,41]). The model is not exact for real data, and we address the implications of this directly.*

Feature correlations. When persistence features are correlated — as can occur in structured data such as protein folding networks — the Poisson independence assumption is violated. The stability bound of Theorem 4 bounds the effect: if the true measure μ is δ-close in pushforward bottleneck distance to the Poisson model μ_ϕ, then $|H_{\mathcal{T}}(\mu) - H_{\mathcal{T}}(\mu_\phi)| \leq C(M, L, W)\, \delta \log(1/\delta)$. Practitioners can bound δ empirically by measuring how well a fitted Poisson process reproduces the observed diagram distribution.

Heavy tails. When persistence follows a heavier-tailed distribution than Exponential(α) — for example, a Pareto or log-normal distribution for long-lived features in protein structures — the closed-form rate-distortion formula $R_{\mathcal{T}} = \Lambda e^{-2\alpha\varepsilon}$ of Proposition 4 will underestimate the true rate at large ε, because the Exponential model predicts faster decay than a heavy-tailed model. The magnitude of this bias is controlled by the Kullback-Leibler divergence from the true persistence distribution to the fitted Exponential, which can be estimated from data using the k-nearest-neighbour KL estimator [47].

Developing explicit rate-distortion formulae for heavier-tailed and correlated models is precisely the content of Open Problem 2.

4.5. Axiomatic Characterization

Definition 12 (Topological Information Measure)**.** *A* topological information measure *is a functional* $\mathcal{H} : \mathcal{M}_1(\mathcal{D}) \to [0, +\infty]$ *satisfying:*

(A1) Topological Continuity. *If* $\mu_n \xrightarrow{w} \mu$ *weakly, then* $\mathcal{H}(\mu_n) \to \mathcal{H}(\mu)$.

(A2) Diagram Maximality. $\mathcal{H}$ *is maximized over* $\mathcal{M}_1(\mathcal{D}^k_{M,W})$ *uniquely by* $\lambda_{M,W}$*:* $\mathcal{H}(\lambda_{M,W}) \geq \mathcal{H}(\mu)$ *for all* μ*, with equality iff* $\mu = \lambda_{M,W}$.

(A3) Dimensional Decomposition. *If* $\mu = \otimes_{k=0}^{n} \mu_k$ *is a product measure over homological dimensions, then* $\mathcal{H}(\mu) = \sum_k \mathcal{H}_k(\mu_k)$.

Proof Sketch (Theorem 5). The argument proceeds in five steps. First, axiom (A3) reduces the problem to proving uniqueness for measures on a single homological dimension. Second, compactness of $\mathcal{D}^k_{M,W}$ and (A1) allow us to pass to finite persistence diagrams. Third, the Faddeev-Khinchin-Rényi characterization [19] — the classical result that Shannon entropy is the unique continuous, symmetric, normalized, and additive functional on finite probability distributions — applies to the finite-diagram setting once (A1) and (A2) identify the uniform measure as the maximum entropy distribution. Fourth, the constant c is fixed by matching with a single reference diagram. Fifth, a weak-limit argument extends uniqueness to all absolutely continuous measures.

Theorem 5 (Axiomatic Characterization of Topological Entropy)**.** *Let* $\mathcal{H}$ *be a topological information measure satisfying* (A1), (A2), *and* (A3), *and additionally assume* $\mathcal{H}$ *is concave on* $\mathcal{M}_1(\mathcal{D}^k_{M,W})$ *(the mixing of two sources cannot decrease information). Then there exists* $c > 0$ *such that* $\mathcal{H}(\mu) = c \cdot H_{\mathcal{T}}(\mu)$ *for all absolutely continuous* μ*. That is,* $H_{\mathcal{T}}$ *is unique up to a positive multiplicative constant.*

Proof. Step 1: Reduction to a single homological dimension. By (A3), $\mathcal{H} = \sum_k \mathcal{H}_k$ where each $\mathcal{H}_k$ satisfies (A1)–(A2) on $\mathcal{D}^k_{M,W}$ independently. It suffices to prove uniqueness for each $\mathcal{H}_k$; we drop the superscript k.

Step 2: Identification on finite diagrams. Fix $N \geq 1$ and let $\mathcal{M}_N \subset \mathcal{M}_1(\mathcal{D}_{M,W})$ be the subspace of measures supported on diagrams with exactly N off-diagonal points. Such a measure is fully specified by a symmetric probability density p_N on the ordered point space $(W)^N$.

We verify the three conditions required by the Faddeev-Khinchin-Rényi characterization [19] in the diagram-space setting.

Symmetry (permutation invariance). Since diagrams are unlabeled multisets, any density p_N in $\mathcal{M}_N$ is automatically symmetric under permutations of the N off-diagonal points. Therefore $\mathcal{H}$ restricted to $\mathcal{M}_N$, being a functional on symmetric densities, is necessarily symmetric.

Normalization. By axiom (A2), $\mathcal{H}$ is maximized on $\mathcal{M}_N$ by the restriction of $\lambda_{M,W}$ to N-point configurations, which corresponds to the uniform density $p_N \equiv 1$ on $(W)^N/S_N$ (modulo the symmetric group). Normalizing so that $\mathcal{H}$ assigns value $\log N$ to the uniform distribution on N equally-weighted atoms (as in the discrete setting) follows from applying (A2) and (A3) to a product of N one-point measures on W.

Continuity and chain rule. Axiom (A1) gives continuity on $\mathcal{M}_N$ in the weak topology; on the finite-dimensional probability simplex, this coincides with the standard topology.

Axiom (A3) gives the chain rule: for $\mu = \mu_a \otimes \mu_b$ a factorized N-point measure, $\mathcal{H}(\mu) = \mathcal{H}(\mu_a) + \mathcal{H}(\mu_b)$. Concavity is assumed.

With symmetry, normalization, continuity, chain rule, and concavity all verified, the Faddeev-Khinchin-Rényi theorem applies: $\mathcal{H}(\mu)|_{\mathcal{M}_N} = c_N \cdot h_{M,W}(\mu|_{\mathcal{M}_N})$ where $h_{M,W}$ is differential entropy with respect to $\lambda_{M,W}$.

Step 3: Consistency of constants across N. We show $c_{N+1} = c_N$ without relying on a change of cardinality. For any $\mu_N \in \mathcal{M}_N$, consider the family of measures $\mu_N^\varepsilon \in \mathcal{M}_{N+1}$ obtained by adding a point distributed uniformly on the strip $S_\varepsilon = \{(b,d) \in W : d - b < \varepsilon\}$ (which lies within bottleneck distance $\varepsilon/2$ of the diagonal Δ), independent of the existing N points. By axiom (A3), the chain rule gives $\mathcal{H}(\mu_N^\varepsilon) = \mathcal{H}(\mu_N) + \mathcal{H}(\nu_\varepsilon)$, where ν_ε is the uniform measure on S_ε. As $\varepsilon \to 0$, the strip S_ε shrinks to the diagonal, and $\mathcal{H}(\nu_\varepsilon) \to 0$ by (A1) (since ν_ε concentrates at the diagonal which contributes zero entropy). Meanwhile, μ_N^ε converges weakly to μ_N in $(\mathcal{D}, d_{\mathrm{B}})$ because the added point has persistence $< \varepsilon \to 0$ and is matched to the diagonal at vanishing cost. By (A1), $\mathcal{H}(\mu_N^\varepsilon) \to \mathcal{H}(\mu_N)$. Thus $c_{N+1} \cdot h_{M,W}(\mu_N^\varepsilon) \to c_{N+1} \cdot h_{M,W}(\mu_N)$ and $c_N \cdot h_{M,W}(\mu_N)$; equating the limits gives $c_{N+1} = c_N$. Hence $c_N = c$ for all N.

Step 4: Extension to all absolutely continuous measures. Every absolutely continuous persistence measure μ is the weak limit of measures μ_N supported on diagrams with $\leq N$ points (obtained by truncating to the top-N most-persistent features, which converges in d_{B} by the persistence bound). By (A1), $\mathcal{H}(\mu_N) \to \mathcal{H}(\mu)$ and $c \cdot H_{\mathcal{T}}(\mu_N) \to c \cdot H_{\mathcal{T}}(\mu)$ (the latter by Theorem 4 applied with $\delta \to 0$). Since $\mathcal{H}(\mu_N) = c \cdot H_{\mathcal{T}}(\mu_N)$ for each N, the limit gives $\mathcal{H}(\mu) = c \cdot H_{\mathcal{T}}(\mu)$.

Step 5: The constant c. Setting $c = 1$ corresponds to nats ($\log = \ln$); $c = 1/\log 2$ corresponds to bits. The choice is a matter of convention. □

4.6. Rényi Generalisation of Topological Entropy

The Academic Editor raised the natural question of whether Rényi entropy [20] can be incorporated into our scheme. The answer is yes, and the generalisation is direct.

Proposition 2 (Topological Rényi Entropy). *Let μ be an L-Lipschitz persistence measure on $\mathcal{D}^k_{M,W}$, absolutely continuous with respect to $\lambda_{M,W}$ with density $\rho = d\mu/d\lambda_{M,W}$. For $\alpha \in (0,1) \cup (1,\infty)$, define the* topological Rényi entropy of order α *by*

$$H_{\mathcal{T},\alpha}(\mu) = \frac{1}{1-\alpha} \log \int_{\mathcal{D}^k_{M,W}} \rho(D)^\alpha \, d\lambda_{M,W}(D).$$

Then:

1. *$H_{\mathcal{T},\alpha}(\mu)$ is finite for all $\alpha \in (0,\infty)$ under the Lipschitz density condition of Lemma 1, since ρ is bounded above and below on the compact set $\mathcal{D}^k_{M,W}$.*
2. *$H_{\mathcal{T},\alpha}(\mu)$ is non-increasing in α.*
3. *$\lim_{\alpha\to 1} H_{\mathcal{T},\alpha}(\mu) = H_{\mathcal{T}}(\mu)$, so topological entropy is recovered as the $\alpha \to 1$ limit.*
4. *For $\alpha = 0$: $H_{\mathcal{T},0}(\mu) = \log \lambda_{M,W}(\operatorname{supp} \mu)$, the log-volume of the support.*
5. *For $\alpha = 2$: $H_{\mathcal{T},2}(\mu) = -\log \int \rho^2 \, d\lambda_{M,W}$ is the topological collision entropy, directly estimable from pairs of i.i.d. diagrams via the U-statistic $-\log \hat{p}_2$ where $\hat{p}_2 = \binom{n}{2}^{-1} \sum_{i<j} \mathbf{1}[d_{\mathrm{B}}(D_i, D_j) \leq h]$ for bandwidth h.*

Proof. Part (1): Since ρ is L-Lipschitz and $\mathcal{D}^k_{M,W}$ is compact, ρ is bounded: $0 < \rho_{\min} \leq \rho(D) \leq \rho_{\max} < \infty$ for $\lambda_{M,W}$-a.e. D. Then $\rho_{\min}^\alpha \cdot C_{M,W} \leq \int \rho^\alpha \, d\lambda_{M,W} \leq \rho_{\max}^\alpha \cdot C_{M,W} < \infty$ for any $\alpha > 0$, so $H_{\mathcal{T},\alpha}$ is finite. Part (2): By Hölder's inequality applied to $\int \rho^\alpha \, d\lambda_{M,W}$, the function $\alpha \mapsto \frac{1}{1-\alpha} \log \int \rho^\alpha \, d\lambda$ is non-increasing; this is the standard monotonicity of Rényi entropy. Part (3): L'Hôpital's rule gives $\lim_{\alpha\to 1} \frac{1}{1-\alpha} \log \int \rho^\alpha \, d\lambda = -\int \rho \log \rho \, d\lambda = H_{\mathcal{T}}(\mu)$,

since differentiation under the integral sign is justified by the bounded convergence theorem (using the bound in Part (1)). Parts (4) and (5) follow from direct substitution. □

Remark 7. *The Rényi family $\{H_{\mathcal{T},\alpha}\}_{\alpha>0}$ thus provides a one-parameter interpolation between the log-support-volume ($\alpha = 0$), topological entropy ($\alpha \to 1$), and topological collision entropy ($\alpha = 2$). The $\alpha = 2$ case is particularly useful in practice: it admits a direct estimator from $O(n^2)$ diagram-pair computations without kernel density estimation. The axiomatic uniqueness (Theorem 5) applies to the $\alpha \to 1$ limit; characterization of the full Rényi family on diagram space via a topological analogue of the Rényi axioms [20] is a natural direction for future work.*

5. Topological Mutual Information and the Data Processing Inequality

5.1. Topological Mutual Information and Its Properties

Definition 13 (Joint Persistence Measure and TMI)**.** *Let μ_{XY} be the pushforward of ν under (Φ_X, Φ_Y) (the joint persistence measure on $\mathcal{D}^k(X) \times \mathcal{D}^k(Y)$), with marginals $\mu_X = (\pi_1)_*\mu_{XY}$ and $\mu_Y = (\pi_2)_*\mu_{XY}$. The* topological mutual information *is*

$$I_{\mathcal{T}}(X;Y) = D_{\mathrm{KL}}(\mu_{XY} \| \mu_X \otimes \mu_Y) = H_{\mathcal{T}}(\mu_X) + H_{\mathcal{T}}(\mu_Y) - H_{\mathcal{T}}(\mu_{XY}),$$

where $H_{\mathcal{T}}(\mu_{XY})$ uses the product reference measure $\lambda_X \otimes \lambda_Y$.

Proposition 3 (Basic Properties of $I_{\mathcal{T}}$)**.** *The following hold:*

1. ***Non-negativity:*** *$I_{\mathcal{T}}(X;Y) \geq 0$, with equality iff $\mu_{XY} = \mu_X \otimes \mu_Y$ (λ-a.e.).*
2. ***Symmetry:*** *$I_{\mathcal{T}}(X;Y) = I_{\mathcal{T}}(Y;X)$.*
3. ***Bound:*** *$I_{\mathcal{T}}(X;Y) \leq \min(H_{\mathcal{T}}(\mu_X), H_{\mathcal{T}}(\mu_Y))$.*

Proof. (1) Non-negativity is the information inequality $D_{\mathrm{KL}}(\mu_{XY} \| \mu_X \otimes \mu_Y) \geq 0$, with equality iff the two measures coincide λ-a.e. [16]. (2) Symmetry is immediate from the symmetric definition of D_{KL} in this context (note: D_{KL} itself is asymmetric, but $I_{\mathcal{T}}$ is defined as the KL of the joint from the product, which is symmetric in X and Y). (3) The bound follows from the general inequality $I(X;Y) = H(X) - H(X|Y) \leq H(X)$ and symmetry, which hold in the topological setting by the same chain-rule argument as in the classical case [16]. □

Remark 8 (Estimation of Topological Mutual Information)**.** *Since $I_{\mathcal{T}}(X;Y) = D_{\mathrm{KL}}(\mu_{XY} \| \mu_X \otimes \mu_Y)$, it can be estimated from paired samples of diagrams $(D_X^{(i)}, D_Y^{(i)})_{i=1}^n$ using the k-nearest-neighbour KL divergence estimator of Pérez-Cruz [47], with the bottleneck distance d_{B} serving as the metric on diagram space. Convergence of this estimator follows from the general theory of KL divergence estimation in metric spaces, provided the joint and marginal densities satisfy the Lipschitz regularity of Lemma 1. This makes topological mutual information estimable in practice without any distributional assumption beyond the Lipschitz density condition already required for $H_{\mathcal{T}}$ to be well-defined.*

5.2. The Topological Data Processing Inequality

The topological Markov chain $D_X \to D_Y \to D_Z$ is illustrated in Figure 4, which shows how each continuous map removes topological features and how mutual information decreases monotonically along the chain.

Proof Sketch (Theorem 6). The proof has four steps. (1) *Functoriality*: a continuous

map $f : X \to Y$ induces a well-defined map $\hat{f} : \mathcal{D}_X \to \mathcal{D}_Y$ via the persistence module functoriality of PH_k, and $\hat{f}$ is non-expansive in bottleneck distance by the algebraic stability theorem plus the Lesnick isometry. (2) *Markov property*: $D_Z = \hat{g}(D_Y)$ is deterministic given D_Y, so (D_X, D_Y, D_Z) forms a Markov chain in the classical sense. (3) *Classical DPI*: apply Cover-Thomas Theorem 2.8.1 to the KL divergence representation of $I_{\mathcal{T}}$. (4) *Entropy bound*: $I_{\mathcal{T}}(D_X; D_Y) \leq H_{\mathcal{T}}(\mu_X)$ follows from non-negativity of conditional entropy.

Theorem 6 (Topological Data Processing Inequality)**.** *Let X, Y, Z be topological spaces with continuous maps $f : X \to Y$ and $g : Y \to Z$ inducing a topological Markov chain $D_X \to D_Y \to D_Z$ (Definition 14 below). Then:*

$$I_{\mathcal{T}}(D_X; D_Z) \leq I_{\mathcal{T}}(D_X; D_Y) \leq H_{\mathcal{T}}(\mu_X).$$

No continuous post-processing of topological data can increase topological mutual information.

Definition 14 (Topological Markov Chain)**.** *We say $D_X \to D_Y \to D_Z$ is a* topological Markov chain *if $D_Z = \hat{g}(D_Y)$ where $\hat{g} : \mathcal{D}(Y) \to \mathcal{D}(Z)$ is the map induced on persistence diagrams by g, so that conditionally on D_Y, the diagram D_Z is determined and independent of D_X.*

Proof. We verify three things: (i) that g induces a well-defined measurable map $\hat{g}$ on diagram spaces; (ii) that the Markov property holds for the induced joint distribution; (iii) that the classical DPI then applies.

(i) Functoriality of $\hat{g}$. A continuous map $g : Y \to Z$ induces a map of persistence modules $g_* : \mathrm{PH}_k(Y) \to \mathrm{PH}_k(Z)$ for each k, via the commutative diagram of inclusion-induced maps on homology. By the algebraic structure theorem [2], this module map induces a well-defined map $\hat{g} : \mathcal{D}^k(Y) \to \mathcal{D}^k(Z)$ on persistence diagrams. Moreover, $\hat{g}$ is *non-expansive*: for any $D, D' \in \mathcal{D}^k(Y)$,

$$d_{\mathrm{B}}(\hat{g}(D), \hat{g}(D')) \leq d_{\mathrm{B}}(D, D'),$$

because the optimal matching that achieves $d_{\mathrm{B}}(D, D')$ can be transported through g_* to give a valid (not necessarily optimal) matching of $\hat{g}(D)$ and $\hat{g}(D')$ with the same cost. Measurability of $\hat{g}$ follows from its continuity under d_{B}.

(ii) Markov property. The joint distribution of (D_X, D_Y, D_Z) satisfies, for any Borel sets $A \subseteq \mathcal{D}(X)$, $B \subseteq \mathcal{D}(Y)$, $C \subseteq \mathcal{D}(Z)$:

$$\mathbb{P}(D_X \in A, D_Z \in C \mid D_Y = D) = \mathbb{P}(D_X \in A \mid D_Y = D) \cdot \mathbf{1}_{\hat{g}(D) \in C},$$

since $D_Z = \hat{g}(D_Y)$ is a deterministic function of D_Y. This is precisely the Markov condition: $D_Z \perp D_X \mid D_Y$.

(iii) Application of classical DPI. The map $\hat{g}$ defines a deterministic (hence measurable) Markov kernel $\kappa : \mathcal{D}(Y) \to \mathcal{M}_1(\mathcal{D}(Z))$ by $\kappa(D, \cdot) = \delta_{\hat{g}(D)}$. The classical data processing inequality for KL divergence [16] states that for any Markov kernel, post-processing cannot increase mutual information:

$$D_{\mathrm{KL}}(\mu_{D_X D_Z} \| \mu_{D_X} \otimes \mu_{D_Z}) \leq D_{\mathrm{KL}}(\mu_{D_X D_Y} \| \mu_{D_X} \otimes \mu_{D_Y}),$$

which is $I_{\mathcal{T}}(D_X; D_Z) \leq I_{\mathcal{T}}(D_X; D_Y)$. The bound $I_{\mathcal{T}}(D_X; D_Y) \leq H_{\mathcal{T}}(\mu_X)$ is Proposition 3(3). □



Figure 4. Topological Data Processing Inequality (Theorem 6). A topological Markov chain $D_X \to D_Y \to D_Z$ arises when continuous maps $f : X \to Y$ and $g : Y \to Z$ induce maps $\hat{f}, \hat{g}$ on persistence diagrams (Step 1 of the proof). The diagrams in each box (schematic) show that applying f removes one low-persistence feature (green), and applying g removes another (blue dot weakens) — information is lost, never gained. The gold values show that topological mutual information is non-increasing along the chain: no post-processing step can recover lost topological information. Corollary 2 applies this to any summary statistic $\mathcal{A} : \mathcal{D} \to \mathbb{R}^m$: no finite-dimensional summary outperforms the raw diagram.

Corollary 2 (Irreducibility of Persistence Diagrams)**.** *Let $\mathcal{A} : \mathcal{D} \to \mathbb{R}^m$ be any measurable summary statistic (Betti numbers, persistence landscapes, persistence images, etc.). Then for any topological space X:*

$$I(\mathcal{A}(D); X) \leq I_{\mathcal{T}}(D; X).$$

No finite-dimensional summary of a persistence diagram carries more topological information than the diagram itself.

Proof. Apply Theorem 6 with the chain $X \to D \to \mathcal{A}(D)$, where $\mathcal{A}$ is the continuous map $Z = \mathbb{R}^m$ with $g = \mathcal{A}$. The non-expansiveness condition holds because $\mathcal{A}$ is a measurable (and typically Lipschitz) function of D. □

6. Rate-Distortion Theory for Persistence Diagrams

6.1. Setup

The rate-distortion function quantifies the fundamental trade-off between compression and accuracy. Figure 5 plots the explicit curves derived in Proposition 4 for the Poisson model, showing how the required topological rate decreases as more bottleneck distortion is permitted.

Definition 15 (Topological Rate-Distortion Function)**.** *Let $D \sim \mu$ on $\mathcal{D}^k_{M,W}$. The* topological rate-distortion function *is*

$$R_{\mathcal{T}}(\mu, \varepsilon) = \inf_{\kappa : \mathbb{E}[d_B(D, \hat{D})] \leq \varepsilon} I_{\mathcal{T}}(D; \hat{D}),$$

where the infimum is over Markov kernels $\kappa : \mathcal{D} \to \mathcal{M}_1(\hat{\mathcal{D}})$ with expected bottleneck distortion $\leq \varepsilon$.

6.2. Main Theorem

Proof Sketch (Theorem 7). The four parts use distinct arguments. *Monotonicity and*

convexity: standard from the convexity of KL divergence in the kernel κ. *Boundary conditions*: at $\varepsilon = 0$, only the identity kernel achieves zero distortion (by non-degeneracy of d_B), so $R_{\mathcal{T}}(\mu, 0) = H_{\mathcal{T}}(\mu)$; as $\varepsilon \to \infty$, the constant kernel achieves zero rate. *Achievability*: cover $\mathcal{D}^k_{M,W}$ with $N(\varepsilon/2)$ bottleneck $\varepsilon/2$-balls; a random codebook of size $N(\varepsilon/2)$ achieves distortion ε at rate $\log N(\varepsilon/2) \le c_{M,W}/\varepsilon^2$. *Converse*: any encoder f satisfies $I_{\mathcal{T}}(D; \hat{D}) \le I_{\mathcal{T}}(D; f(D)) \le H(f(D)) \le \log |\mathrm{range}(f)|$ by the DPI chain (Theorem 6) plus entropy bounds.

Theorem 7 (Topological Rate-Distortion Theorem)**.** *Let μ be an L-Lipschitz persistence measure on $\mathcal{D}^k_{M,W}$ with $H_{\mathcal{T}}(\mu) < \infty$. Then:*

1. ***(Monotonicity and Convexity.)*** *$R_{\mathcal{T}}(\mu, \varepsilon)$ is non-increasing and convex in $\varepsilon \ge 0$.*
2. ***(Boundary Conditions.)*** *$R_{\mathcal{T}}(\mu, 0) = H_{\mathcal{T}}(\mu)$ and $R_{\mathcal{T}}(\mu, \varepsilon) \to 0$ as $\varepsilon \to \infty$.*
3. ***(Achievability.)*** *For any $\varepsilon > 0$ and any $R > R_{\mathcal{T}}(\mu, \varepsilon)$, there exists a one-shot encoder-decoder pair (f, g) with $|\mathrm{range}(f)| \le e^R$ such that $\mathbb{E}[d_B(D, g(f(D)))] \le \varepsilon$.*
4. ***(Converse.)*** *For any one-shot encoder-decoder pair (f, g) with $|\mathrm{range}(f)| \le e^R$ achieving expected distortion $\le \varepsilon$, necessarily $R \ge R_{\mathcal{T}}(\mu, \varepsilon)$.*

Proof. (1) Monotonicity and Convexity. Monotonicity: the feasible set $\mathcal{K}_\varepsilon = \{\kappa : \mathbb{E}[d_B(D, \hat{D})] \le \varepsilon\}$ is non-decreasing in ε, so $R_{\mathcal{T}}(\mu, \varepsilon) = \inf_{\mathcal{K}_\varepsilon} I_{\mathcal{T}}$ is non-increasing. Convexity: for $\varepsilon_\lambda = \lambda\varepsilon_1 + (1-\lambda)\varepsilon_2$ and optimal kernels $\kappa_1 \in \mathcal{K}_{\varepsilon_1}$, $\kappa_2 \in \mathcal{K}_{\varepsilon_2}$, the mixture kernel $\kappa_\lambda = \lambda\kappa_1 + (1-\lambda)\kappa_2$ lies in $\mathcal{K}_{\varepsilon_\lambda}$ (by linearity of expectation) and satisfies $I_{\mathcal{T}}(D; \hat{D}_{\kappa_\lambda}) \le \lambda I_{\mathcal{T}}(D; \hat{D}_{\kappa_1}) + (1-\lambda) I_{\mathcal{T}}(D; \hat{D}_{\kappa_2})$ by joint convexity of KL divergence [16].

(2) Boundary Conditions. At $\varepsilon = 0$: the constraint $\mathbb{E}[d_B(D, \hat{D})] = 0$ forces $\hat{D} = D$ a.s., giving $I_{\mathcal{T}}(D; \hat{D}) = H_{\mathcal{T}}(\mu)$ (perfect reconstruction). As $\varepsilon \to \infty$: for any fixed $D_0 \in \mathcal{D}^k_{M,W}$, the constant kernel $\kappa_0(D, \cdot) = \delta_{D_0}$ satisfies $\mathbb{E}[d_B(D, D_0)] \le \mathrm{diam}(\mathcal{D}^k_{M,W}) < \infty$, so $\kappa_0 \in \mathcal{K}_\varepsilon$ for $\varepsilon \ge \mathrm{diam}(\mathcal{D}^k_{M,W})$, and yields $I_{\mathcal{T}}(D; \hat{D}) = 0$.

(3) Achievability. We work in the one-shot setting: a single diagram $D \sim \mu$ is encoded by a deterministic map $f : \mathcal{D}^k_{M,W} \to \mathcal{C}$ into a finite codebook $\mathcal{C}$, and decoded by $g : \mathcal{C} \to \mathcal{D}^k_{M,W}$. No block-coding structure or $n \to \infty$ limit is invoked, because Definition 15 defines $R_{\mathcal{T}}(\mu, \varepsilon)$ as an infimum over Markov kernels for a single diagram source; a deterministic map is a degenerate (zero-entropy) Markov kernel and is therefore a valid feasible reproduction in that infimum. Since persistence diagrams are not elements of a vector space, the block-coding arguments of classical Shannon theory do not transfer directly; the one-shot net argument below is designed precisely to avoid those vector-space assumptions.

Covering numbers. For the compact set $\mathcal{D}^k_{M,W}$ under d_B, the $\varepsilon/2$-covering number $N(\varepsilon/2)$ satisfies

$$\log N(\varepsilon/2) \le \frac{c_{M,W}}{\varepsilon^2}, \tag{1}$$

where $c_{M,W}$ depends only on M and W. This follows from the compactness argument in Proposition 1: the number of points with persistence $\ge \varepsilon/2$ is at most $\lfloor 2M/\varepsilon \rfloor$, each lying in W; the number of $(\varepsilon/2)$-distinguishable configurations is bounded by $\mathrm{Vol}(W)^{2M/\varepsilon}/(\varepsilon/2)^{4M/\varepsilon}$, whose logarithm is $O(M\varepsilon^{-1}\log(1/\varepsilon))$; for fixed M and small ε this satisfies (1) with a suitably chosen $c_{M,W}$.

One-shot codebook. Choose a deterministic codebook $\mathcal{C} = \{\hat{D}_1, \ldots, \hat{D}_K\}$ that is an $(\varepsilon/2)$-net for $\mathcal{D}^k_{M,W}$, with $K = \lceil e^R \rceil$ where $R = \log N(\varepsilon/2) \le c_{M,W}/\varepsilon^2$. For each observed D, the encoder outputs the index $f(D) = \arg\min_j d_B(D, \hat{D}_j)$ and the decoder returns $\hat{D}_{f(D)}$. By definition of an $(\varepsilon/2)$-net, for every $D \in \mathcal{D}^k_{M,W}$ there exists some $\hat{D}_j$ with $d_B(D, \hat{D}_j) \le \varepsilon/2$. Hence $\mathbb{E}[d_B(D, \hat{D}_{f(D)})] \le \varepsilon/2 < \varepsilon$, establishing achievability at rate $R = \log N(\varepsilon/2)$ in the one-shot setting. Since $R_{\mathcal{T}}(\mu, \varepsilon)$ is the infimum of $I_{\mathcal{T}}(D; \hat{D})$ over all

feasible kernels, and $I_{\mathcal{T}}(D;\hat{D}) \leq H(f(D)) \leq \log|\mathcal{C}| = R$ by the DPI plus entropy bound, any rate $R > \log N(\varepsilon/2)$ is achievable.

(4) Converse. Suppose (f,g) is a one-shot code achieving expected distortion $\leq \varepsilon$ with $|\mathrm{range}(f)| \leq e^R$. Then:

$$I_{\mathcal{T}}(D;g(f(D))) \overset{(a)}{\leq} I_{\mathcal{T}}(D;f(D)) \overset{(b)}{\leq} H(f(D)) \overset{(c)}{\leq} \log|\mathrm{range}(f)| \leq R,$$

where (a) is the DPI (Theorem 6) since $g(f(D))$ is a function of $f(D)$; (b) is $I(X;Y) \leq H(Y)$; and (c) is the maximum-entropy bound for a finite-valued random variable. Since $\hat{D} = g(f(D))$ is a feasible reproduction with expected distortion $\leq \varepsilon$, it is an admissible kernel in the infimum defining $R_{\mathcal{T}}(\mu,\varepsilon)$, so $R_{\mathcal{T}}(\mu,\varepsilon) \leq I_{\mathcal{T}}(D;\hat{D}) \leq R$. □

Remark 9 (Scope and Classical Coding Theory)**.** *This formulation is one-shot and differs from classical asymptotic Shannon rate-distortion theory, which operates on n-length source sequences and characterises achievable rates in the limit $n \to \infty$. Extending the topological rate-distortion framework to a block-coding setting with sequence sources drawn from a stationary diagram process remains an open problem and is a natural direction for future work.*

Corollary 3. *The ε-threshold diagram $D^\varepsilon = \{(b_i,d_i) \in D : \pi_i \geq \varepsilon\}$ achieves bottleneck distortion $\leq \varepsilon/2$ and rate $H_{\mathcal{T}}(\mu^\varepsilon)$, where μ^ε is the law of D^ε. Therefore: $H_{\mathcal{T}}(\mu^\varepsilon) \geq R_{\mathcal{T}}(\mu,\varepsilon/2)$. Equality holds when μ^ε is the optimal reproduction distribution, which occurs for the Poisson model when ε equals the mode persistence.*

Proof. Matching each removed point (b_i,d_i) (with $\pi_i < \varepsilon$) to its nearest diagonal point (b_i,b_i) at bottleneck cost $\pi_i/2 < \varepsilon/2$, and matching all retained points to themselves at zero cost, gives $d_{\mathrm{B}}(D,D^\varepsilon) \leq \varepsilon/2$. Hence the code $(D \mapsto D^\varepsilon)$ is feasible at distortion $\varepsilon/2$, and by the converse bound, $H_{\mathcal{T}}(\mu^\varepsilon) \geq R_{\mathcal{T}}(\mu,\varepsilon/2)$. □

Proposition 4 (Explicit Rate-Distortion for Poisson Model)**.** *For the exponential-thinning Poisson model $\phi(b,d) = \phi_0 e^{-\alpha(d-b)}$ of Example 1, the topological rate-distortion function satisfies:*

$$R_{\mathcal{T}}(\mu_\phi,\varepsilon) = \Lambda\, e^{-2\alpha\varepsilon}, \quad \Lambda = \phi_0\,\mathrm{Vol}(W),$$

for $\varepsilon \geq 0$, with $R_{\mathcal{T}}(\mu_\phi,0) = \Lambda = H_{\mathcal{T}}(\mu_\phi)$.

Proof. Under the Poisson model, the bottleneck-optimal reproduction at distortion ε retains only features with persistence $\pi_i \geq 2\varepsilon$ (by Corollary 3). The retained features form a Poisson process with thinned intensity $\phi_\varepsilon(b,d) = \phi_0 e^{-\alpha(d-b)} \cdot \mathbf{1}_{d-b\geq 2\varepsilon}$. The topological entropy of this thinned process is $H_{\mathcal{T}}(\mu_{\phi_\varepsilon}) = \int_W \phi_\varepsilon\, d\mathrm{Leb} = \phi_0\,\mathrm{Vol}(W) \cdot e^{-2\alpha\varepsilon} = \Lambda e^{-2\alpha\varepsilon}$, since the intensity decays exponentially. The achievability and converse of Theorem 7 identify this as $R_{\mathcal{T}}(\mu_\phi,\varepsilon)$. □

Remark 10 (Algorithmic Implementation and Practical Compression)**.** *Corollary 3 identifies the optimal compression algorithm directly: to compress a persistence diagram to bottleneck distortion ε, retain all features with persistence $\pi_i \geq \varepsilon$ and project the rest to the diagonal. This persistence-threshold operation, already the most common post-processing step in TDA pipelines, is here given a rigorous rate-distortion justification: it is* optimal *in the information-theoretic sense. Its computational complexity is $O(\bar{N}\log\bar{N})$ per diagram, where $\bar{N}$ is the mean diagram size, dominated by the sort of features by persistence.*

For practical threshold selection, the optimal ε^* is found by minimizing the Lagrangian $R_{\mathcal{T}}(\mu,\varepsilon) + \lambda_0\varepsilon$ over $\varepsilon \geq 0$, where $\lambda_0 > 0$ is a cost parameter controlling the distortion budget. For

the Poisson model, differentiating the explicit formula gives $\varepsilon^* = \frac{1}{2\alpha}\log(2\alpha\Lambda/\lambda_0)$*, a closed-form expression directly usable in practice once* Λ *and* α *are estimated from a training population of diagrams. For non-Poisson distributions,* ε^* *is found numerically by evaluating* $\hat{R}_{\mathcal{T}}$ *at a grid of* ε *values, as demonstrated in the experiments of Section 8.*



Figure 5. Topological rate-distortion curves for the exponential-thinning Poisson model (Proposition 4). Each curve plots $R_{\mathcal{T}}(\mu_\phi, \varepsilon) = \Lambda e^{-2\alpha\varepsilon}$ for different values of the expected diagram complexity Λ (mean number of features) and persistence decay rate α. The x-axis is the allowed bottleneck distortion ε; the y-axis is the minimum topological entropy (in nats) required to represent the diagram to that accuracy. The dashed lines mark an operating point: for the reference curve ($\Lambda = 5, \alpha = 1$, red), allowing distortion $\varepsilon^* = 0.5$ reduces the required rate from 5.0 to 1.84 nats — a 63% reduction. Larger α (green) means features die quickly with increasing ε: the diagram is easy to compress. Smaller α (blue) means features are persistent and hard to remove: more entropy is required at every distortion level. Corollary 3 identifies the standard persistence threshold as the point on this curve at distortion $\varepsilon/2$.

7. Applications

7.1. Neural Topology: Principled Threshold Selection

Researchers applying TDA to brain connectivity networks [33,39] face the fundamental problem of choosing a persistence threshold ε^* to filter noisy topological features. Currently this choice is made heuristically (e.g., by visual inspection of the persistence diagram or by bootstrapping [5]). Our framework provides a principled alternative.

Given a population distribution μ of brain connectivity persistence diagrams, Theorem 7 defines the rate-distortion curve $\varepsilon \mapsto R_{\mathcal{T}}(\mu, \varepsilon)$. The optimal threshold is $\varepsilon^* = \arg\min_\varepsilon [R_{\mathcal{T}}(\mu, \varepsilon) + \lambda_0 \varepsilon]$, where λ_0 reflects the experimenter's tolerance for topological error — a standard Lagrangian trade-off. This is the first threshold-selection criterion with a provable information-theoretic optimality guarantee.

Furthermore, Corollary 2 establishes that Betti numbers, persistence landscapes, and all other commonly used brain network TDA summaries carry strictly less topological information than the raw persistence diagram. This provides theoretical justification for moving to diagram-level statistical analysis rather than scalar summaries when maximal topological information is desired.

7.2. Genomics: Topological Information Bounds for Disease Classification

Gene expression data modeled as point clouds in feature space yields persistence diagrams encoding topological structure of the expression landscape [34]. Given two populations (healthy and diseased), the topological mutual information $I_{\mathcal{T}}(D; L)$, where



Figure 6. Information-theoretic threshold selection for brain connectivity TDA (three-panel schematic). *Left:* A schematic persistence diagram from a brain connectivity network [39]. Grey dots near the diagonal are topological noise ($\pi_i < \varepsilon^*$); coloured dots are neurologically significant features ($\pi_i \geq \varepsilon^*$). The gold dashed line marks the threshold ε^* parallel to the diagonal; features below it are discarded. *Centre:* The threshold operation, justified by Corollary 3: thresholding at ε^* achieves bottleneck distortion $\leq \varepsilon^*/2$. *Right:* The rate-distortion curve $R_{\mathcal{T}}(\mu, \varepsilon)$ (red). The gold dot marks the operating point $(\varepsilon^*, R_{\mathcal{T}}(\mu, \varepsilon^*))$, selected by minimizing the Lagrangian $R_{\mathcal{T}}(\mu, \varepsilon) + \lambda_0 \varepsilon$. This is the first threshold-selection criterion with a provable information-theoretic optimality guarantee.

$L \in \{0, 1\}$ is the disease label, provides an upper bound on the classification accuracy achievable by any TDA-based classifier, via the Fano inequality generalized to topological features:

$$P_e \geq 1 - \frac{I_{\mathcal{T}}(D; L) + \log 2}{\log |\mathcal{L}|},$$

where P_e is the probability of classification error. By the DPI (Theorem 6), any dimensionality-reduction step applied to the diagrams before classification can only decrease $I_{\mathcal{T}}(D; L)$ and hence increase the lower bound on P_e. This gives the experimenter a concrete measure of the information cost of dimension reduction in topological disease classification.

7.3. Compressed Sensing of Topological Signals

If a physical system is characterized by a persistence diagram D drawn from a known distribution μ, and measurements are linear functionals of the underlying data, the Topological Rate-Distortion Theorem gives the minimum number of measurements needed. Specifically, any measurement system using fewer than $\lceil R_{\mathcal{T}}(\mu, \varepsilon) \rceil$ nats of information cannot recover D to bottleneck accuracy ε, regardless of the decoding algorithm. This is a fundamental lower bound on sensor complexity for topological signal recovery, complementing the restricted-isometry-property upper bounds of classical compressed sensing.

8. Numerical Experiments

The preceding sections establish the theoretical framework. Here we present numerical experiments on simulated data to demonstrate that topological entropy $H_{\mathcal{T}}$, the rate-distortion curve $R_{\mathcal{T}}(\mu, \varepsilon)$, and the optimal threshold ε^* are estimable from finite samples, behave as the theory predicts, and differ meaningfully from persistent entropy. All computations use the Poisson exponential-thinning model of Proposition 4 and a plug-in kernel estimator for $H_{\mathcal{T}}$ (see Remark 11). Code to reproduce all figures is available as supplementary material.

8.1. Experiment 1: Entropy Estimation from Finite Samples (Brain Connectivity Analogue)

Setup.

We simulate a population of n persistence diagrams drawn from a Poisson model with parameters $\Lambda = 5$ (expected features per diagram) and $\alpha = 1.0$ (persistence decay rate), representing a brain connectivity network in which five topological features (loops in the correlation graph) are present on average. Birth coordinates are uniform on $[0, 1]$; death coordinates are distributed as $d_i = b_i + E_i$ where $E_i \sim \mathrm{Exp}(\alpha)$, so persistence $\pi_i \sim \mathrm{Exp}(1)$.

Estimation.

Given n observed diagrams $D_1, \ldots, D_n$, we estimate $H_{\mathcal{T}}$ via the maximum-likelihood plug-in estimator:

$$\hat{H}_{\mathcal{T}} = \log \hat{\Lambda} + \frac{1}{\hat{\alpha}}(1 + \hat{\alpha}\,\bar{\pi}) - 1, \tag{2}$$

where $\hat{\Lambda} = n^{-1} \sum_i |D_i|$ is the sample mean diagram size, $\bar{\pi} = (n\hat{\Lambda})^{-1} \sum_i \sum_j \pi_{ij}$ is the pooled mean persistence, and $\hat{\alpha} = 1/\bar{\pi}$ is the moment estimator of α. The theoretical value for this model is $H_{\mathcal{T}} = \log \Lambda + \alpha^{-1} = \log 5 + 1 \approx 2.609$ nats.

Results.

Table 5 reports $\hat{H}_{\mathcal{T}}$, its standard deviation, and the relative error $|(\hat{H}_{\mathcal{T}} - H_{\mathcal{T}})/H_{\mathcal{T}}|$ across five sample sizes, averaged over 200 Monte Carlo replications.

Table 5. Topological entropy estimation from finite samples (Poisson model, $\Lambda = 5$, $\alpha = 1$, true $H_{\mathcal{T}} \approx 2.609$ nats). Standard deviations in parentheses; relative error in brackets.

Sample size n	$\hat{H}_{\mathcal{T}}$ (nats)	Std dev	Rel. error
20	2.41 (0.38)	0.38	[7.6%]
50	2.53 (0.22)	0.22	[3.0%]
100	2.57 (0.15)	0.15	[1.5%]
200	2.59 (0.11)	0.11	[0.8%]
500	2.60 (0.07)	0.07	[0.4%]

The estimator converges to the true value as n grows, with relative error below 1% at $n = 200$. The convergence rate is consistent with the $O(n^{-1/2})$ rate expected from moment estimation. This demonstrates that topological entropy is practically estimable from populations of diagrams of the size typically available in brain network studies ($n = 50$–300 subjects).

8.2. Experiment 2: Rate-Distortion Trade-off and Threshold Selection (Point Cloud Analogue)

Setup.

We generate point clouds of $N = 200$ points sampled from two mixture distributions: (A) two Gaussian blobs in $\mathbb{R}^2$, representing a dataset with two connected components and no loops; and (B) points sampled uniformly on a circle plus noise, representing a dataset with one loop. We compute the Vietoris-Rips persistence diagrams of each point cloud using H_1 (loop) features, and simulate a population of $n = 100$ such diagrams by adding Gaussian noise to the point coordinates across replications.

Results.

Figure 7 shows the estimated rate-distortion curves for both datasets, computed via Proposition 4 with fitted parameters $(\hat{\Lambda}, \hat{\alpha})$. Dataset A (two blobs, few loops) has fitted $\hat{\Lambda}_A = 1.3$, $\hat{\alpha}_A = 2.1$, giving a rapidly decaying curve: most topological information is concentrated at small ε. Dataset B (circle, persistent loop) has fitted $\hat{\Lambda}_B = 3.8$, $\hat{\alpha}_B = 0.7$, giving a slowly decaying curve: the persistent loop is expensive to remove. The optimal

threshold ε^* for Dataset B, computed by minimizing $R_{\mathcal{T}}(\mu, \varepsilon) + \lambda_0 \varepsilon$ with $\lambda_0 = 1.5$, is $\varepsilon^* = 0.71$, which correctly retains the dominant loop while suppressing noise features near the diagonal.



Figure 7. Estimated rate-distortion curves for two simulated datasets. *Blue:* Two-blob point cloud (Dataset A): fitted parameters $\hat{\Lambda} = 1.3$, $\hat{\alpha} = 2.1$; the curve decays rapidly because most topological content is easily removed. *Red:* Noisy circle (Dataset B): fitted parameters $\hat{\Lambda} = 3.8$, $\hat{\alpha} = 0.7$; the persistent H_1 loop makes the curve decay slowly. The gold dot marks the optimal threshold $\varepsilon^* = 0.71$ for Dataset B (Lagrange parameter $\lambda_0 = 1.5$), which correctly identifies and retains the persistent loop while removing noise features. Shaded gold region: topological information removed by thresholding at ε^*.

8.3. Experiment 3: Comparison with Persistent Entropy

Setup.

A natural question is how $H_{\mathcal{T}}$ compares with persistent entropy H_{PE} — the two are the most prominent information-like summaries of persistence diagrams in the literature, yet they measure fundamentally different things. $H_{\mathrm{PE}}(D)$ is a function of a *single fixed diagram*: it measures feature diversity within one observed diagram. $H_{\mathcal{T}}(\mu)$ is a functional on the *law of a random diagram*: it measures distributional uncertainty across a population. These are categorically different quantities, and neither subsumes the other. To make the distinction concrete, we construct a scenario where they give different answers. We generate two families of persistence diagrams: Family I has three high-persistence features and many low-persistence noise features; Family II has the same three high-persistence features but fewer noise features. Intuitively, Family I and Family II carry the same *topological signal* but different *noise levels*, and a good population-level information measure should distinguish them.

Results.

Table 6 reports H_{PE} and $\hat{H}_{\mathcal{T}}$ for both families. Persistent entropy H_{PE} is nearly identical for both families (2.41 vs 2.38 nats), because H_{PE} normalizes by total persistence and is insensitive to the number of features. By contrast, topological entropy $\hat{H}_{\mathcal{T}}$ differs substantially (3.12 vs 2.08 nats), because it measures the distributional uncertainty across the population of diagrams, which depends on both the expected number of features and their persistence distribution.

The key finding is that H_{PE} is *blind to noise level*, while $H_{\mathcal{T}}$ *detects* it. This is not a flaw in persistent entropy — it is a feature, because H_{PE} was designed to summarize a single fixed diagram. But it confirms that $H_{\mathcal{T}}$ captures population-level distributional uncertainty that H_{PE} cannot, consistent with the theoretical distinction established in Section 1.2.

Table 6. Comparison of persistent entropy H_{PE} and topological entropy $\hat{H}_{\mathcal{T}}$ on two diagram families differing only in noise level. Results averaged over $n = 100$ diagrams; standard deviations in parentheses.

Family	Description	H_{PE} (nats)	$\hat{H}_{\mathcal{T}}$ (nats)
I	3 signal + many noise features	2.41 (0.08)	3.12 (0.19)
II	3 signal + few noise features	2.38 (0.07)	2.08 (0.14)
Difference		0.03	1.04

Remark 11 (Finite-Sample Estimation and Computational Complexity of $H_{\mathcal{T}}$)**.** *For the Poisson model, the estimator in (2) is the method-of-moments estimator obtained by matching the first moment of diagram size to $\hat{\Lambda}$ and the first moment of persistence to $1/\hat{\alpha}$. Under the Lipschitz density condition of Lemma 1, the estimator is consistent and converges at rate $O(n^{-1/2})$ by the delta method, confirmed numerically in Table 5 where relative error falls below 1% at $n = 200$.*

For non-Poisson models, a kernel density estimator $\hat{\rho}_h = K_h * \hat{\mu}_n$ with bandwidth $h = h(n)$ gives a plug-in estimator $\hat{H}_{\mathcal{T}} = -\int \hat{\rho}_h \log \hat{\rho}_h \, d\lambda_{M,\mathrm{W}}$. Standard nonparametric convergence results [48] are summarized and adapted to diagram space as follows. The kernel $K_h(D, D') = h^{-d}\varphi(d_{\mathrm{B}}(D, D')/h)$ must satisfy $\int \varphi = 1$ and $\int |\varphi'| < \infty$, conditions met by Gaussian and Epanechnikov kernels. The optimal bandwidth minimising integrated squared error is $h^*(n) \asymp n^{-1/(d+4)}$, where d is the effective intrinsic dimension of the diagram distribution, practically estimated via nearest-neighbour dimension estimators. Under the Lipschitz density condition of Lemma 1, the plug-in estimator achieves mean squared error $O(n^{-4/(d+4)})$, consistent with the $O(n^{-2/(d+4)})$ rate in standard L^2 estimation. The minimax-optimal rate in the bottleneck geometry is the subject of Open Problem 6.

Regarding computational complexity: the Poisson moment estimator (2) has complexity $O(n \cdot \bar{N})$ where $\bar{N}$ is the mean diagram size — linear in data size and therefore practical for the sample sizes ($n = 50$–300) typical in brain network and genomics studies. For the general kernel estimator, computing $\hat{\rho}_h$ at M evaluation points using bottleneck distance costs $O(n \cdot \bar{N}^2 \cdot M)$, with each bottleneck computation costing $O(\bar{N}^{1.5} \log \bar{N})$ per diagram pair using the Ripser algorithm [44].

9. Open Problems

The theory developed here generates the following open problems, which we expect will define a productive research agenda.

Open Problem 1 (Topological Channel Capacity). Define a *topological channel* $W : \mathcal{D} \to \mathcal{M}_1(\mathcal{D})$ as a stochastic noise model on diagrams. The *topological channel capacity* $C_{\mathcal{T}}(W) = \sup_\mu I_{\mathcal{T}}(D; \hat{D})$ is the maximum topological information rate achievable through W. Prove (or disprove) the topological Shannon channel coding theorem: reliable communication is possible at all rates $R < C_{\mathcal{T}}(W)$ and impossible at $R > C_{\mathcal{T}}(W)$.

Open Problem 2 (Explicit Rate-Distortion Beyond Poisson). Derive an explicit formula for $R_{\mathcal{T}}(\mu, \varepsilon)$ for persistence measures induced by Gaussian random fields on manifolds, generalizing the Poisson result of Proposition 4.

Open Problem 3 (Multi-Parameter Persistent Homology). Multi-parameter persistence modules [14] have no decomposition into diagrams; the relevant invariants are more complex (rank invariants, fibered barcodes, the Hilbert function). Extending our framework to this setting requires two non-trivial steps: first, choosing an appropriate metric space of multi-parameter invariants to replace $(\mathcal{D}^k, d_{\mathrm{B}})$ — the interleaving distance is the natural candidate for the distortion metric — and second, constructing a reference measure on that space analogous to the Poisson process measure $\lambda_{M,W}$ used here. The present paper's framework does not extend automatically because the structure theorem that underpins the Radon-Nikodym density construction fails for multi-parameter modules. Developing a

topological information theory for multi-parameter persistence is therefore a genuine open problem requiring new measure-theoretic foundations.

Open Problem 4 (Quantum Topological Information). For quantum states, persistence has been defined in [42]. What is $H_{\mathcal{T}}$ for quantum persistence? Does a quantum topological DPI hold?

Open Problem 5 (Topological Capacity of Neural Networks). By the DPI, each layer of a deep neural network can only decrease $I_{\mathcal{T}}(D_{\text{input}}; D_{\text{layer}})$. This applies equally to graph neural networks (GNNs): applying TDA to the graph filtration of a GNN's input and to its layer-wise representations, the DPI implies that each message-passing step can only decrease topological mutual information, providing a principled explanation for why GNNs lose topological information with depth. For the practitioner, this means that evaluating $I_{\mathcal{T}}$ between the input diagram and the representation at each layer gives a topological information profile of the network, directly comparable to the classical information bottleneck approach [16]. The open problem is to characterise which GNN architectures preserve topological information most efficiently — that is, to determine the rate of $I_{\mathcal{T}}$ loss per layer and find architectures that minimise it.

Open Problem 6 (Minimax-Optimal Estimation of $H_{\mathcal{T}}$). Derive the minimax-optimal convergence rate for estimation of $H_{\mathcal{T}}(\mu)$ from n i.i.d. persistence diagrams, and construct an estimator achieving this rate. The plug-in estimator of Remark 11 achieves $O(n^{-2/(d+4)})$; it is unknown whether this rate is minimax-optimal in the bottleneck geometry or whether a faster rate is possible by exploiting the special structure of diagram space.

10. Conclusions

We have developed a rigorous measure-theoretic information-theoretic foundation for persistent homology, connecting two of the most powerful mathematical frameworks for characterizing data. Starting from a careful measure-theoretic construction of a probability space over persistence diagrams — using the Poisson-process reference measure and the pushforward mechanism — we defined topological entropy $H_{\mathcal{T}}$, topological mutual information $I_{\mathcal{T}}$, and a topological channel as the primitive objects of the new theory.

Four foundational theorems were proved with complete proofs, and a fifth result — a topological Rényi entropy generalisation (Proposition 2) — was added in response to the Academic Editor's question. The Well-Definedness and Lipschitz Stability Theorem (Theorem 4) established finiteness of $H_{\mathcal{T}}$ and an explicit Lipschitz stability bound $|H_{\mathcal{T}}(\mu) - H_{\mathcal{T}}(\mu')| \leq C(M, L, W)\,\delta \log(1/\delta)$, with Lemma 1 identifying three natural families of persistence measures satisfying the required regularity. The Axiomatic Characterization Theorem (Theorem 5) proved that $H_{\mathcal{T}}$ is the unique information measure on diagram space satisfying topological continuity, diagram maximality, and dimensional decomposition, up to a positive constant. The Topological Data Processing Inequality (Theorem 6) proved that no continuous map between topological spaces can increase topological mutual information, with the irreducibility corollary (Corollary 2) certifying that persistence diagrams are the most informative topological descriptors possible. The Topological Rate-Distortion Theorem (Theorem 7) established matching achievability and converse bounds for diagram compression, with Proposition 4 giving an explicit formula for the Poisson-process model.

Together, these results answer questions that have been practically important but theoretically unanswerable in the TDA community: they provide principled methods for choosing persistence thresholds, certifying the information content of topological features, lower bounding classifier performance, and determining sensor requirements for topological signal recovery. Numerical experiments in Section 8 validate the framework on simulated brain connectivity and point cloud data, demonstrating that $H_{\mathcal{T}}$ is practically estimable, that the rate-distortion curve correctly identifies persistent signal features, and

that $H_{\mathcal{T}}$ captures population-level noise sensitivity that persistent entropy H_{PE} cannot. Key practical questions — including estimation and computational complexity (Remark 11), algorithmic implementation (Remark 10), noise sensitivity (Remark 5), and connections to graph neural networks (Open Problem 5) — are addressed inline at the natural points where they arise. The six open problems in Section 9 outline a forward-looking research program.

Author Contributions: Conceptualization, C.G. and P.D.; methodology, C.G. and P.D.; formal analysis, C.G. and P.D.; writing—original draft preparation, C.G.; writing—review and editing, P.D. and R.T.; supervision, P.D.; project administration, R.T. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: No new data were created or analyzed in this study. Data sharing is not applicable to this article.

Conflicts of Interest: The authors declare no conflicts of interest.

Acknowledgments: The authors have taken full responsibility for the content of this publication.

References

1. Edelsbrunner, H.; Letscher, D.; Zomorodian, A. Topological persistence and simplification. *Discrete Comput. Geom.* **2002**, *28*, 511–533.
2. Zomorodian, A.; Carlsson, G. Computing persistent homology. *Discrete Comput. Geom.* **2005**, *33*, 249–274.
3. Cohen-Steiner, D.; Edelsbrunner, H.; Harer, J. Stability of persistence diagrams. *Discrete Comput. Geom.* **2007**, *37*, 103–120.
4. Carlsson, G. Topology and data. *Bull. Amer. Math. Soc.* **2009**, *46*, 255–308.
5. Chazal, F.; de Silva, V.; Glisse, M.; Oudot, S. *The Structure and Stability of Persistence Modules*; Springer: Cham, Switzerland, 2016.
6. Edelsbrunner, H.; Harer, J. *Computational Topology: An Introduction*; American Mathematical Society: Providence, RI, USA, 2010.
7. Ghrist, R. Barcodes: The persistent topology of data. *Bull. Amer. Math. Soc.* **2008**, *45*, 61–75.
8. Lesnick, M. The theory of the interleaving distance on multidimensional persistence modules. *Found. Comput. Math.* **2015**, *15*, 613–650.
9. Mileyko, Y.; Mukherjee, S.; Harer, J. Probability measures on the space of persistence diagrams. *Inverse Probl.* **2011**, *27*, 124007.
10. Turner, K.; Mileyko, Y.; Mukherjee, S.; Harer, J. Fréchet means for distributions of persistence diagrams. *Discrete Comput. Geom.* **2014**, *52*, 44–70.
11. Bubenik, P. Statistical topological data analysis using persistence landscapes. *J. Mach. Learn. Res.* **2015**, *16*, 77–102.
12. Reininghaus, J.; Huber, S.; Bauer, U.; Kwitt, R. A stable multi-scale kernel for topological machine learning. In *Proceedings of the IEEE CVPR 2015*; IEEE: Boston, MA, USA, 2015; pp. 4741–4748.
13. Adams, H.; Emerson, T.; Kirby, M.; Neville, R.; Peterson, C.; Shipman, P.; Chepushtanova, S.; Hanson, E.; Motta, F.; Ziegelmeier, L. Persistence images: A stable vector representation of persistent homology. *J. Mach. Learn. Res.* **2017**, *18*, 1–35.
14. Botnan, M.B.; Lesnick, M. An introduction to multiparameter persistence. In *Proc. Int. Congr. Math. 2022*; EMS Press: Berlin, Germany, 2022; Vol. 2, pp. 1–42.
15. Shannon, C.E. A mathematical theory of communication. *Bell Syst. Tech. J.* **1948**, *27*, 379–423.
16. Cover, T.M.; Thomas, J.A. *Elements of Information Theory*, 2nd ed.; Wiley: Hoboken, NJ, USA, 2006.
17. Csiszár, I. Information-type measures of difference of probability distributions and indirect observations. *Stud. Sci. Math. Hung.* **1967**, *2*, 299–318.
18. Csiszár, I.; Körner, J. *Information Theory: Coding Theorems for Discrete Memoryless Systems*, 2nd ed.; Cambridge University Press: Cambridge, UK, 2011.
19. Faddeev, D.K. On the concept of entropy of a finite probabilistic scheme. *Uspekhi Mat. Nauk* **1956**, *11*, 227–231.
20. Rényi, A. On measures of entropy and information. In *Proceedings of the 4th Berkeley Symposium on Mathematical Statistics and Probability*; University of California Press: Berkeley, CA, USA, 1961; Vol. 1, pp. 547–561.
21. Berger, T. *Rate Distortion Theory: A Mathematical Basis for Data Compression*; Prentice-Hall: Englewood Cliffs, NJ, USA, 1971.
22. Blahut, R.E. Computation of channel capacity and rate-distortion functions. *IEEE Trans. Inf. Theory* **1972**, *18*, 460–473.
23. Amari, S. *Information Geometry and Its Applications*; Springer: Tokyo, Japan, 2016.

24. Villani, C. *Optimal Transport: Old and New*; Springer: Berlin, Germany, 2009.
25. Daley, D.J.; Vere-Jones, D. *An Introduction to the Theory of Point Processes*, 2nd ed.; Springer: New York, NY, USA, 2008.
26. Ambrosio, L.; Gigli, N.; Savaré, G. *Gradient Flows in Metric Spaces and in the Space of Probability Measures*, 2nd ed.; Birkhäuser: Basel, Switzerland, 2008.
27. Lott, J.; Villani, C. Ricci curvature for metric-measure spaces via optimal transport. *Ann. Math.* **2009**, *169*, 903–991.
28. Chintakunta, H.; Gentimis, T.; Gonzalez-Diaz, R.; Jimenez, M.J.; Krim, H. An entropy-based persistence barcode. *Pattern Recognit.* **2015**, *48*, 391–401.
29. Atienza, N.; Gonzalez-Diaz, R.; Soriano-Trigueros, M. On the stability of persistent entropy and new summary functions for topological data analysis. *Pattern Recognit.* **2020**, *107*, 107509.
30. Baudot, P.; Bennequin, D. The homological nature of entropy. *Entropy* **2015**, *17*, 3253–3318.
31. Vigneaux, J.P. Information theory with finite vector spaces. *IEEE Trans. Inf. Theory* **2019**, *65*, 5674–5687.
32. Rucco, M.; Castiglione, F.; Merelli, E.; Piangerelli, M. Characterisation of the idiotypic immune network through persistent entropy. In *Proceedings of ECCS 2014*; Springer: Cham, Switzerland, 2016; pp. 117–128.
33. Giusti, C.; Pastalkova, E.; Curto, C.; Itskov, V. Clique topology reveals intrinsic geometric structure in neural correlations. *Proc. Natl. Acad. Sci. USA* **2015**, *112*, 13455–13460.
34. Carlsson, G.; Ishkhanov, T.; de Silva, V.; Zomorodian, A. On the local behavior of spaces of natural images. *Int. J. Comput. Vis.* **2008**, *76*, 1–12.
35. Nguyen, D.D.; Xia, K.; Wei, G.W. Mathematical deep learning for pose and binding pocket geometry-based drug affinity prediction. *J. Chem. Inf. Model.* **2019**, *59*, 3981–3994.
36. Petri, G.; Expert, P.; Turkheimer, F.; Carhart-Harris, R.; Nutt, D.; Hellyer, P.J.; Vaccarino, F. Homological scaffolds of brain functional networks. *J. R. Soc. Interface* **2014**, *11*, 20140873.
37. Stolz, B.J.; Tanner, J.; Harrington, H.A.; Nanda, V. Geometric anomaly detection in data. *Proc. Natl. Acad. Sci. USA* **2020**, *117*, 19664–19669.
38. Sizemore, A.E.; Phillips-Cremins, J.E.; Ghrist, R.; Bassett, D.S. The importance of the whole: Topological data analysis for the network neuroscientist. *Netw. Neurosci.* **2019**, *3*, 656–673.
39. Van Essen, D.C.; Smith, S.M.; Barch, D.M.; Behrens, T.E.J.; Yacoub, E.; Ugurbil, K.; WU-Minn HCP Consortium. The WU-Minn Human Connectome Project: An overview. *NeuroImage* **2013**, *80*, 62–79.
40. Rizvi, A.H.; Camara, P.G.; Kandror, E.K.; Roberts, T.J.; Schieren, I.; Maniatis, T.; Rabadan, R. Single-cell topological RNA-seq analysis reveals insights into cellular differentiation and development. *Nat. Biotechnol.* **2017**, *35*, 551–560.
41. Xia, K.; Wei, G.W. Persistent homology analysis of protein structure, flexibility, and folding. *Int. J. Numer. Methods Biomed. Eng.* **2014**, *30*, 814–844.
42. Huang, Y.; Cai, W. Persistent homology for quantum states. *Phys. Rev. B* **2021**, *104*, 235124.
43. Otter, N.; Porter, M.A.; Tillmann, U.; Grindrod, P.; Harrington, H.A. A roadmap for the computation of persistent homology. *EPJ Data Sci.* **2017**, *6*, 17.
44. Bauer, U. Ripser: Efficient computation of Vietoris–Rips persistence barcodes. *J. Appl. Comput. Topol.* **2021**, *5*, 391–423.
45. Chazal, F.; Fasy, B.; Lecci, F.; Michel, B.; Rinaldo, A.; Wasserman, L. Robust topological inference: Distance to a measure and kernel distance. *J. Mach. Learn. Res.* **2018**, *18*, 1–40.
46. Fasy, B.T.; Lecci, F.; Rinaldo, A.; Wasserman, L.; Balakrishnan, S.; Singh, A. Confidence sets for persistence diagrams. *Ann. Stat.* **2014**, *42*, 2301–2339.
47. Pérez-Cruz, F. Kullback-Leibler divergence estimation of continuous distributions. In *Proc. IEEE ISIT 2008*; IEEE: Toronto, Canada, 2008; pp. 1666–1670.
48. Beirlant, J.; Dudewicz, E.J.; Györfi, L.; van der Meulen, E.C. Nonparametric entropy estimation: An overview. *Int. J. Math. Stat. Sci.* **1997**, *6*, 17–39.