

RBFs Over Near-Flat Surfaces

APPM 5480 Asymptotics

Caleb Jacobs

April 25, 2022

Department of Applied Mathematics

Outline

1. Introduction/Background

Radial Basis Functions

RBF Interpolants

Solving $A\lambda = \mathbf{F}$

2. Enter Asymptotics!

A perturbation problem

Perturbed solution

Solution remarks

3. What's Next

Introduction/Background

Introduction/Background

Radial Basis Functions

Radial Basis Functions

A radial basis function (RBF), denoted by $\phi(r)$, is any radially symmetric function.

Radial Basis Functions

A radial basis function (RBF), denoted by $\phi(r)$, is any radially symmetric function.

Commonly used radial basis functions

Type of basis function	Radial function $\phi(r)$
Gaussian (GA)	$e^{-(\varepsilon r)^2}$
Multiquadric(MQ)	$\sqrt{1 + (\varepsilon r)^2}$
Inverse Multiquadric (IMQ)	$1/\sqrt{1 + (\varepsilon r)^2}$
Inverse Quadratic (IQ)	$1/(1 + (\varepsilon r)^2)$
Sech (SH)	$\text{sech}(\varepsilon r)$

where ε is called the *shape parameter* [1].

Radial Basis Functions

A radial basis function (RBF), denoted by $\phi(r)$, is any radially symmetric function.

Commonly used radial basis functions

Type of basis function	Radial function $\phi(r)$
Gaussian (GA)	$e^{-(\varepsilon r)^2}$
Multiquadric(MQ)	$\sqrt{1 + (\varepsilon r)^2}$
Inverse Multiquadric (IMQ)	$1/\sqrt{1 + (\varepsilon r)^2}$
Inverse Quadratic (IQ)	$1/(1 + (\varepsilon r)^2)$
Sech (SH)	$\text{sech}(\varepsilon r)$

where ε is called the *shape parameter* [1].

Introduction/Background

RBF Interpolants

RBF Interpolants

Suppose we have data $\{\mathbf{x}_i, f_i\}_{i=1}^n$.

RBF Interpolants

Suppose we have data $\{\mathbf{x}_i, f_i\}_{i=1}^n$. Then, we can define an RBF based interpolant as

$$s(\mathbf{x}) = \sum_{i=1}^n \lambda_i \phi_\varepsilon(\|\mathbf{x} - \mathbf{x}_i\|)$$

RBF Interpolants

Suppose we have data $\{\mathbf{x}_i, f_i\}_{i=1}^n$. Then, we can define an RBF based interpolant as

$$s(\mathbf{x}) = \sum_{i=1}^n \lambda_i \phi_\epsilon(\|\mathbf{x} - \mathbf{x}_i\|)$$

where λ_i can be found by solving

$$\underbrace{\begin{pmatrix} \phi_\epsilon(\|\mathbf{x}_1 - \mathbf{x}_1\|) & \phi_\epsilon(\|\mathbf{x}_1 - \mathbf{x}_2\|) & \cdots & \phi_\epsilon(\|\mathbf{x}_1 - \mathbf{x}_n\|) \\ \phi_\epsilon(\|\mathbf{x}_2 - \mathbf{x}_1\|) & \phi_\epsilon(\|\mathbf{x}_2 - \mathbf{x}_2\|) & \cdots & \phi_\epsilon(\|\mathbf{x}_2 - \mathbf{x}_n\|) \\ \vdots & \vdots & \ddots & \vdots \\ \phi_\epsilon(\|\mathbf{x}_n - \mathbf{x}_1\|) & \phi_\epsilon(\|\mathbf{x}_n - \mathbf{x}_2\|) & \cdots & \phi_\epsilon(\|\mathbf{x}_n - \mathbf{x}_n\|) \end{pmatrix}}_A \underbrace{\begin{pmatrix} \lambda_1 \\ \lambda_2 \\ \vdots \\ \lambda_n \end{pmatrix}}_\lambda = \underbrace{\begin{pmatrix} f_1 \\ f_2 \\ \vdots \\ f_n \end{pmatrix}}_F.$$

Introduction/Background

Solving $A\lambda = F$

Solving for our weights λ

The smaller ε gets, the more accurate the RBF interpolant becomes. But, as ε becomes smaller, the conditioning of A increases.

Solving for our weights λ

The smaller ε gets, the more accurate the RBF interpolant becomes. But, as ε becomes smaller, the conditioning of A increases.

Naively solving $A\lambda = F$ is called *RBF-Direct*. *Does not work when ε is small.*

Solving for our weights λ

The smaller ε gets, the more accurate the RBF interpolant becomes. But, as ε becomes smaller, the conditioning of A increases.

Naively solving $A\lambda = F$ is called *RBF-Direct*. *Does not work when ε is small.*

Solve this system stably using an indirect method for small ε .

Solving for our weights λ

So we want to use an indirect method to obtain better accuracy for small ε .

Solving for our weights λ

So we want to use an indirect method to obtain better accuracy for small ε .
There is no reason for ε to be real valued so let's leverage contour integration using Cauchy's integral formula,

$$\phi_\varepsilon(x) = \frac{1}{2\pi i} \oint_\Gamma \frac{\phi_z(x)}{z - \varepsilon} dz.$$

Solving for our weights λ

So we want to use an indirect method to obtain better accuracy for small ε .
There is no reason for ε to be real valued so let's leverage contour integration using Cauchy's integral formula,

$$\phi_\varepsilon(x) = \frac{1}{2\pi i} \oint_\Gamma \frac{\phi_z(x)}{z - \varepsilon} dz.$$

RBF Contour Padé (RBF-CP)

Solving for our weights λ

So we want to use an indirect method to obtain better accuracy for small ε .
There is no reason for ε to be real valued so let's leverage contour integration using Cauchy's integral formula,

$$\phi_\varepsilon(x) = \frac{1}{2\pi i} \oint_\Gamma \frac{\phi_z(x)}{z - \varepsilon} dz.$$

RBF Contour Padé (RBF-CP) (*precursor to RBF-RA*)

RBF Rational Approximations (RBF-RA) (circa 2017) [3]

RBF-RA works great when the data given $\{\mathbf{x}_i\}_{i=1}^n$ is

- on a completely flat surface
- on a curvy surface.

RBF-RA works great when the data given $\{\mathbf{x}_i\}_{i=1}^n$ is

on a completely flat surface

on a curvy surface.

However, RBF-RA breaks down when the data is on a near-flat surface (i.e. surface with small curvature).

So, we would like understand/explore why RBF-RA breaks on near flat surfaces.

Enter Asymptotics!

Enter Asymptotics!


A perturbation problem

A perturbation problem

Suppose we have the pairs of points $\{(x_i, y_i)\}_{i=1}^n$ in the *unit disk*. Then, define our interpolation nodes as

$$\mathbf{x}_i = \left\langle x_i, y_i, \sqrt{\frac{1}{\kappa^2} - r_i^2} - \frac{1}{\kappa} \right\rangle, \quad r_i = \sqrt{x_i^2 + y_i^2}.$$

This is just data over a sphere with curvature κ with the top of the sphere at the origin:



A perturbation problem

Suppose we have the pairs of points $\{(x_i, y_i)\}_{i=1}^n$ in the *unit disk*. Then, define our interpolation nodes as

$$\mathbf{x}_i = \left\langle x_i, y_i, \sqrt{\frac{1}{\kappa^2} - r_i^2} - \frac{1}{\kappa} \right\rangle, \quad r_i = \sqrt{x_i^2 + y_i^2}.$$

This is just data over a sphere with curvature κ with the top of the sphere at the origin:

$$x_i^2 + y_i^2 + (z_i + 1/\kappa)^2 = \frac{1}{\kappa^2}$$

A perturbation problem

$$\mathbf{x}_i = \left\langle x_i, y_i, \sqrt{\frac{1}{\kappa^2} - r_i^2} - \frac{1}{\kappa} \right\rangle, \quad r_i = \sqrt{x_i^2 + y_i^2}.$$

Then, the difference between two nodes is given as

$$\|\mathbf{x}_i - \mathbf{x}_j\|_2^2 = (x_i - x_j)^2 + (y_i - y_j)^2 + \left(\sqrt{\frac{1}{\kappa^2} - r_i^2} - \sqrt{\frac{1}{\kappa^2} - r_j^2} \right)^2$$

A perturbation problem

$$\mathbf{x}_i = \left\langle x_i, y_i, \sqrt{\frac{1}{\kappa^2} - r_i^2} - \frac{1}{\kappa} \right\rangle, \quad r_i = \sqrt{x_i^2 + y_i^2}.$$

Then, the difference between two nodes is given as

$$\|\mathbf{x}_i - \mathbf{x}_j\|_2^2 = (x_i - x_j)^2 + (y_i - y_j)^2 + \left(\sqrt{\frac{1}{\kappa^2} - r_i^2} - \sqrt{\frac{1}{\kappa^2} - r_j^2} \right)^2$$

Further, each entry of the collocation matrix A will be

$$A_{ij} = \phi_\epsilon(\|\mathbf{x}_i - \mathbf{x}_j\|_2) = e^{-(\epsilon\|\mathbf{x}_i - \mathbf{x}_j\|_2)^2} = e^{-\epsilon^2\|\mathbf{x}_i - \mathbf{x}_j\|_2^2}$$

A perturbation problem

$$\begin{aligned}A_{ij} &= \phi_\varepsilon(\|\mathbf{x}_i - \mathbf{x}_j\|_2) = e^{-(\varepsilon\|\mathbf{x}_i - \mathbf{x}_j\|_2)^2} = e^{-\varepsilon^2\|\mathbf{x}_i - \mathbf{x}_j\|_2^2} \\&= e^{-\varepsilon^2\left((x_i - x_j)^2 + (y_i - y_j)^2 + \left(\sqrt{\frac{1}{\kappa^2} - r_i^2} - \sqrt{\frac{1}{\kappa^2} - r_j^2}\right)^2\right)} \\&= e^{-\varepsilon^2((x_i - x_j)^2 + (y_i - y_j)^2)} e^{-\varepsilon^2\left(\sqrt{\frac{1}{\kappa^2} - r_i^2} - \sqrt{\frac{1}{\kappa^2} - r_j^2}\right)^2} \\&= e^{-\varepsilon^2((x_i - x_j)^2 + (y_i - y_j)^2)} \left(1 - \frac{1}{4}\varepsilon^2\kappa^2(r_i^2 - r_j^2)^2 + \dots\right) \\&= e^{-\varepsilon^2((x_i - x_j)^2 + (y_i - y_j)^2)} - \kappa^2\frac{1}{4}\varepsilon^2(r_i^2 - r_j^2)^2 e^{-\varepsilon^2((x_i - x_j)^2 + (y_i - y_j)^2)} + \mathcal{O}(\kappa^4).\end{aligned}$$

A perturbation problem

$$\begin{aligned}A_{ij} &= \phi_\varepsilon(\|\mathbf{x}_i - \mathbf{x}_j\|_2) = e^{-(\varepsilon\|\mathbf{x}_i - \mathbf{x}_j\|_2)^2} = e^{-\varepsilon^2\|\mathbf{x}_i - \mathbf{x}_j\|_2^2} \\&= e^{-\varepsilon^2\left((x_i - x_j)^2 + (y_i - y_j)^2 + \left(\sqrt{\frac{1}{\kappa^2} - r_i^2} - \sqrt{\frac{1}{\kappa^2} - r_j^2}\right)^2\right)} \\&= e^{-\varepsilon^2((x_i - x_j)^2 + (y_i - y_j)^2)} e^{-\varepsilon^2\left(\sqrt{\frac{1}{\kappa^2} - r_i^2} - \sqrt{\frac{1}{\kappa^2} - r_j^2}\right)^2} \\&= e^{-\varepsilon^2((x_i - x_j)^2 + (y_i - y_j)^2)} \left(1 - \frac{1}{4}\varepsilon^2\kappa^2(r_i^2 - r_j^2)^2 + \dots\right) \\&= e^{-\varepsilon^2((x_i - x_j)^2 + (y_i - y_j)^2)} - \kappa^2\frac{1}{4}\varepsilon^2(r_i^2 - r_j^2)^2 e^{-\varepsilon^2((x_i - x_j)^2 + (y_i - y_j)^2)} + \mathcal{O}(\kappa^4).\end{aligned}$$

$$A = A_0 + \kappa^2 A_1 + \mathcal{O}(\kappa^4)$$

A perturbation problem

With the expansion of A

$$A = A_0 + \kappa^2 A_1 + \cdots,$$

our perturbed interpolant problem is then given by

$$A\boldsymbol{\lambda} = \mathbf{F} \implies (A_0 + \kappa^2 A_1 + \cdots)\boldsymbol{\lambda} = \mathbf{F}, \quad 0 < \kappa \ll 1.$$

A perturbation problem

With the expansion of A

$$A = A_0 + \kappa^2 A_1 + \dots,$$

our perturbed interpolant problem is then given by

$$A\boldsymbol{\lambda} = \mathbf{F} \implies (A_0 + \kappa^2 A_1 + \dots)\boldsymbol{\lambda} = \mathbf{F}, \quad 0 < \kappa \ll 1.$$

Note, A_0 is just the standard collocation matrix if our data was on a flat surface.
We know

A_0 is nonsingular for $\varepsilon \in \mathbb{R}, \varepsilon \neq 0$,

A_0 is symmetric [1],

RBF-RA is able to stably work with A_0 .

Enter Asymptotics!

Perturbed solution

Perturbed interpolation weights problem

$$(A_0 + \kappa^2 A_1 + \cdots) \boldsymbol{\lambda} = \mathbf{F}, \quad 0 < \kappa \ll 1$$

Assume $\boldsymbol{\lambda}$ has the regular expansion

$$\boldsymbol{\lambda} = \boldsymbol{\lambda}_0 + \kappa^2 \boldsymbol{\lambda}_1 + \cdots .$$

Perturbed interpolation weights problem

$$(A_0 + \kappa^2 A_1 + \cdots) \boldsymbol{\lambda} = \mathbf{F}, \quad 0 < \kappa \ll 1$$

Assume $\boldsymbol{\lambda}$ has the regular expansion

$$\boldsymbol{\lambda} = \boldsymbol{\lambda}_0 + \kappa^2 \boldsymbol{\lambda}_1 + \cdots .$$

Then at

$$\mathcal{O}(1) : A_0 \boldsymbol{\lambda}_0 = \mathbf{F} \implies \boldsymbol{\lambda}_0 = A_0^{-1} \mathbf{F}$$

$$\mathcal{O}(\kappa^2) : A_0 \boldsymbol{\lambda}_1 + A_1 \boldsymbol{\lambda}_0 = 0 \implies \boldsymbol{\lambda}_1 = -A_0^{-1} A_1 \boldsymbol{\lambda}_0$$

$$\mathcal{O}(\kappa^4) : A_0 \boldsymbol{\lambda}_2 + A_1 \boldsymbol{\lambda}_1 + A_2 \boldsymbol{\lambda}_0 = 0 \implies \boldsymbol{\lambda}_2 = -A_0^{-1} (A_1 \boldsymbol{\lambda}_1 + A_2 \boldsymbol{\lambda}_0)$$

Perturbed interpolation weights problem

$$(A_0 + \kappa^2 A_1 + \cdots) \boldsymbol{\lambda} = \mathbf{F}, \quad 0 < \kappa \ll 1$$

Assume $\boldsymbol{\lambda}$ has the regular expansion

$$\boldsymbol{\lambda} = \boldsymbol{\lambda}_0 + \kappa^2 \boldsymbol{\lambda}_1 + \cdots .$$

Then at

$$\mathcal{O}(1) : A_0 \boldsymbol{\lambda}_0 = \mathbf{F} \implies \boldsymbol{\lambda}_0 = A_0^{-1} \mathbf{F}$$

$$\mathcal{O}(\kappa^2) : A_0 \boldsymbol{\lambda}_1 + A_1 \boldsymbol{\lambda}_0 = 0 \implies \boldsymbol{\lambda}_1 = -A_0^{-1} A_1 \boldsymbol{\lambda}_0$$

$$\mathcal{O}(\kappa^4) : A_0 \boldsymbol{\lambda}_2 + A_1 \boldsymbol{\lambda}_1 + A_2 \boldsymbol{\lambda}_0 = 0 \implies \boldsymbol{\lambda}_2 = -A_0^{-1} (A_1 \boldsymbol{\lambda}_1 + A_2 \boldsymbol{\lambda}_0)$$

So in general,

$$\boldsymbol{\lambda}_n = -A_0^{-1} (A_1 \boldsymbol{\lambda}_{n-1} + A_2 \boldsymbol{\lambda}_{n-2} + \cdots + A_n \boldsymbol{\lambda}_0).$$

Enter Asymptotics!

Solution remarks

Remarks

Putting everything together, our interpolation weights can be written as

$$\lambda = \lambda_0 + \kappa^2 \lambda_1 + \dots$$

$$\lambda_n = -A_0^{-1}(A_1 \lambda_{n-1} + A_2 \lambda_{n-2} + \dots + A_n \lambda_0)$$

Remarks

Putting everything together, our interpolation weights can be written as

$$\begin{aligned}\boldsymbol{\lambda} &= \boldsymbol{\lambda}_0 + \kappa^2 \boldsymbol{\lambda}_1 + \cdots \\ \boldsymbol{\lambda}_n &= -A_0^{-1}(A_1 \boldsymbol{\lambda}_{n-1} + A_2 \boldsymbol{\lambda}_{n-2} + \cdots + A_n \boldsymbol{\lambda}_0)\end{aligned}$$

Remarks

This shows that $\boldsymbol{\lambda}$ is asymptotically well behaved when $\kappa \rightarrow 0$. So ideally, RBF-RA ought to not break in this limit.

Remarks

Putting everything together, our interpolation weights can be written as

$$\begin{aligned}\boldsymbol{\lambda} &= \boldsymbol{\lambda}_0 + \kappa^2 \boldsymbol{\lambda}_1 + \cdots \\ \boldsymbol{\lambda}_n &= -A_0^{-1}(A_1 \boldsymbol{\lambda}_{n-1} + A_2 \boldsymbol{\lambda}_{n-2} + \cdots + A_n \boldsymbol{\lambda}_0)\end{aligned}$$

Remarks

This shows that $\boldsymbol{\lambda}$ is asymptotically well behaved when $\kappa \rightarrow 0$. So ideally, RBF-RA ought to not break in this limit.

We can use this asymptotic expansion to bypass the small curvature issue in RBF-RA.

Putting everything together, our interpolation weights can be written as

$$\boldsymbol{\lambda} = \boldsymbol{\lambda}_0 + \kappa^2 \boldsymbol{\lambda}_1 + \dots$$

$$\boldsymbol{\lambda}_n = -A_0^{-1}(A_1 \boldsymbol{\lambda}_{n-1} + A_2 \boldsymbol{\lambda}_{n-2} + \dots + A_n \boldsymbol{\lambda}_0)$$

Modified RBF-RA Algorithm

- (a) Compute κ estimate of data using desired algorithm
- (b) Use RBF-RA to stably compute needed $\boldsymbol{\lambda}_i$
- (c) Compute $\boldsymbol{\lambda}$ using $\boldsymbol{\lambda} = \boldsymbol{\lambda}_0 + \kappa^2 \boldsymbol{\lambda}_1 + \dots$ up to desired term.

Putting everything together, our interpolation weights can be written as

$$\begin{aligned}\boldsymbol{\lambda} &= \boldsymbol{\lambda}_0 + \kappa^2 \boldsymbol{\lambda}_1 + \cdots \\ \boldsymbol{\lambda}_n &= -A_0^{-1}(A_1 \boldsymbol{\lambda}_{n-1} + A_2 \boldsymbol{\lambda}_{n-2} + \cdots + A_n \boldsymbol{\lambda}_0)\end{aligned}$$

Modified RBF-RA Algorithm


- (a) Compute κ estimate of data using desired algorithm
- (b) Use RBF-RA to stably compute needed $\boldsymbol{\lambda}_i$
- (c) Compute $\boldsymbol{\lambda}$ using $\boldsymbol{\lambda} = \boldsymbol{\lambda}_0 + \kappa^2 \boldsymbol{\lambda}_1 + \cdots$ up to desired term.

Patching an algorithm like this is not ideal. Instead, we would like to understand why the algorithm breaks and then fix the algorithm at the fundamental level.

What's Next

A closer look at conditioning

The conditioning of the collocation matrix at various κ and ε in the complex plane can be seen below



A perturbed eigenvalue problem

Using the same perturbed collocation matrix as before, we can pose a perturbed eigenvalue problem as

$$(A_0 + \kappa^2 A_1 + \cdots) \mathbf{x} = \lambda \mathbf{x}.$$

Now, because A depends on arbitrary, real life data, it is reasonable to assume that A has all distinct eigenvalues. So, finding a perturbed eigenvalue/eigenvector solution to this is readily found using our formulas derived early on in our course or in *Hinch* [2]. In this case, the eigenvalues all have regular perturbations

$$\lambda(\varepsilon; \kappa) = \lambda_0(\varepsilon) + \kappa^2 \lambda_2(\varepsilon) + \cdots .$$

A perturbed root finding problem

The A matrix is normal and so

$$\text{cond}(A) = \frac{|\lambda_{\max}|}{|\lambda_{\min}|}.$$

So, to understand when the condition number blows up, we want to know when $\lambda(\varepsilon; \kappa) = 0$ for any of the eigenvalues λ . So, an interesting perturbed root finding problem is find ε such that

$$\lambda_0(\varepsilon) + \kappa^2 \lambda_2(\varepsilon) + \cdots = 0, \quad 0 < \kappa \ll 1.$$

Even for simple node-sets, $\lambda_i(\varepsilon)$ can be very nonlinear making this perturbed root finding problem quite difficult.

References

- [1] B. FORNBERG AND N. FLYER, *A primer on radial basis functions with applications to the geosciences*, vol. 87 of CBMS-NSF regional conference series in applied mathematics, SIAM, 2015.
- [2] E. J. HINCH, *Perturbation Methods*, Cambridge Texts in Applied Mathematics, Cambridge University Press, 1991.
- [3] G. B. WRIGHT AND B. FORNBERG, *Stable computations with flat radial basis functions using vector-valued rational approximations*, Journal of Computational Physics, 331 (2017), pp. 137–156.

THANK YOU

ANY QUESTIONS?