

Caleb Logemann
MATH 562 Numerical Analysis II
Final Exam

1. Let $A \in \mathbb{R}^{m \times m}$ be written in the form $A = L + D + U$, where L is strictly lower triangular, D is the diagonal of A , and U is the strictly upper triangular part of A . Assuming D is invertible, $A\mathbf{x} = \mathbf{b}$ is equivalent to $\mathbf{x} = -D^{-1}(L + U)\mathbf{x} + D^{-1}\mathbf{b}$. The Jacobi iteration method for solving $A\mathbf{x} = \mathbf{b}$ is defined by

$$\mathbf{x}^{(n+1)} = -D^{-1}(L + U)\mathbf{x}^{(n)} + D^{-1}\mathbf{b}$$

Show that if A is nonsingular and strictly row diagonally dominant:

$$0 < \sum_{j \neq i} (|a_{ij}|) < |a_{ii}|$$

then the Jacobi iteration converges to $\mathbf{x}_* = A^{-1}\mathbf{b}$ for each fixed $\mathbf{b} \in \mathbb{R}^m$.

Proof. Let \mathbf{e}_n be the error of the n th iteration of the Jacobi iteration from the actual solution, that is let

$$\mathbf{e}_n = \mathbf{x}^{(n)} - \mathbf{x}_*.$$

The Jacobi iteration converges to the real solution if

$$\lim_{n \rightarrow \infty} (\|\mathbf{e}_n\|_\infty) = 0$$

The error vector can be expressed recursively by noting that $\mathbf{x}^{(n)}$ is the Jacobi iteration evaluated on $\mathbf{x}^{(n-1)}$ and that \mathbf{x}_* is a fixed point of the Jacobi iteration as it is the true solution to the linear system. This means that

$$\begin{aligned}\mathbf{x}^{(n)} &= -D^{-1}(L + U)\mathbf{x}^{(n-1)} + D^{-1}\mathbf{b} \\ \mathbf{x}_* &= -D^{-1}(L + U)\mathbf{x}_* + D^{-1}\mathbf{b}.\end{aligned}$$

Therefore we can express the error recursively as

$$\begin{aligned}\mathbf{e}_n &= \mathbf{x}^{(n)} - \mathbf{x}_* \\ \mathbf{e}_n &= \left(-D^{-1}(L + U)\mathbf{x}^{(n-1)} + D^{-1}\mathbf{b}\right) - \left(-D^{-1}(L + U)\mathbf{x}_* + D^{-1}\mathbf{b}\right) \\ \mathbf{e}_n &= -D^{-1}(L + U)\mathbf{x}^{(n-1)} + D^{-1}(L + U)\mathbf{x}_* \\ \mathbf{e}_n &= -D^{-1}(L + U)\left(\mathbf{x}^{(n-1)} - \mathbf{x}_*\right) \\ \mathbf{e}_n &= -D^{-1}(L + U)\mathbf{e}_{n-1}.\end{aligned}$$

Extrapolating this backwards we see that \mathbf{e}_n can be expressed in terms of \mathbf{e}_0

$$\mathbf{e}_n = \left(-D^{-1}(L+U)\right)^n \mathbf{e}_0.$$

Now we can consider the limit of $\|\mathbf{e}_n\|_\infty$ as n goes to infinity.

$$\begin{aligned} \lim_{n \rightarrow \infty} (\|\mathbf{e}_n\|_\infty) &= \lim_{n \rightarrow \infty} \left(\left\| \left(-D^{-1}(L+U)\right)^n \mathbf{e}_0 \right\|_\infty \right) \\ \lim_{n \rightarrow \infty} (\|\mathbf{e}_n\|_\infty) &\leq \|\mathbf{e}_0\|_\infty \lim_{n \rightarrow \infty} \left(\left\| D^{-1}(L+U) \right\|_\infty^n \right) \end{aligned}$$

Now consider $\|D^{-1}(L+U)\|_\infty$. The infinity norm is the max row sum of the matrix, that is

$$\left\| D^{-1}(L+U) \right\|_\infty = \max_{1 \leq i \leq m} \sum_{j=1}^m \left| \left(D^{-1}(L+U) \right)_{ij} \right|$$

Because $L+U = A-D$, $(L+U)_{ij} = a_{ij}$ if $i \neq j$ and $(L+U)_{ii} = 0$. Also D^{-1} is diagonal with $(D^{-1})_{ii} = \frac{1}{D_{ii}} = \frac{1}{a_{ii}}$. Therefore the matrix product $D^{-1}(L+U)$ has entries $(D^{-1}(L+U))_{ij} = \frac{a_{ij}}{a_{ii}}$ if $i \neq j$ or if $i = j$, then $(D^{-1}(L+U))_{ii} = 0$. We can now say that

$$\begin{aligned} \left\| D^{-1}(L+U) \right\|_\infty &= \max_{1 \leq i \leq m} \sum_{j \neq k} \left(\left| \frac{a_{ij}}{a_{ii}} \right| \right) \\ \left\| D^{-1}(L+U) \right\|_\infty &= \max_{1 \leq i \leq m} \frac{1}{|a_{ii}|} \sum_{j \neq k} (|a_{ij}|) \end{aligned}$$

However since A is strictly row diagonally dominant $|a_{ii}| > \sum_{j \neq k} (|a_{ij}|)$, we can conclude that $\frac{1}{|a_{ii}|} \sum_{j \neq k} (|a_{ij}|) < 1$. Therefore

$$\left\| D^{-1}(L+U) \right\|_\infty < 1$$

Since $\left\| D^{-1}(L+U) \right\|_\infty < 1$, it is true that $\lim_{n \rightarrow \infty} \left(\left\| D^{-1}(L+U) \right\|_\infty^n \right) = 0$. Thus

$$\begin{aligned} \lim_{n \rightarrow \infty} (\|\mathbf{e}_n\|_\infty) &\leq \|\mathbf{e}_0\|_\infty \lim_{n \rightarrow \infty} \left(\left\| D^{-1}(L+U) \right\|_\infty^n \right) \\ \lim_{n \rightarrow \infty} (\|\mathbf{e}_n\|_\infty) &\leq 0 \end{aligned}$$

This shows that the error converges to zero, and this proves that the Jacobi iteration does converge to the true solution if A is strictly row diagonally dominant. \square

- Let $A \in \mathbb{R}^{m \times m}$ be symmetric positive definite (SPD), $\mathbf{b} \in \mathbb{R}^m$ and define $\phi : \mathbb{R}^m \rightarrow \mathbb{R}$ by

$$\phi(\mathbf{x}) = \frac{1}{2} \mathbf{x}^T A \mathbf{x} - \mathbf{x}^T \mathbf{b}$$

Suppose K is a subspace of \mathbb{R}^m . Show that $\hat{\mathbf{x}} \in K$ minimizes $\phi(\mathbf{x})$ over K if and only if $\nabla \phi(\hat{\mathbf{x}}) \perp K$.

Proof. First let me describe $\nabla\phi(\mathbf{x})$.

$$\begin{aligned}
\nabla\phi(\mathbf{x}) &= \begin{bmatrix} \frac{\partial\phi}{\partial x_1} \\ \vdots \\ \frac{\partial\phi}{\partial x_m} \end{bmatrix} \\
\frac{\partial\phi}{\partial x_i} &= \frac{\partial}{\partial x_i} \left(\frac{1}{2} \mathbf{x}^T A \mathbf{x} - \mathbf{x}^T \mathbf{b} \right) \\
&= \frac{\partial}{\partial x_i} \left(\frac{1}{2} \sum_{j=1}^m \left(x_j \sum_{k=1}^m (a_{jk} x_k) \right) - \sum_{j=1}^m (x_j b_j) \right) \\
&= \frac{1}{2} \frac{\partial}{\partial x_i} \sum_{j=1}^m \left(x_j \sum_{k=1}^m (a_{jk} x_k) \right) - b_i \\
&= \frac{1}{2} \frac{\partial}{\partial x_i} \left(x_i \sum_{k=1}^m (a_{ik} x_k) + \sum_{j \neq i} \left(x_j \sum_{k=1}^m (a_{jk} x_k) \right) \right) - b_i \\
&= \frac{1}{2} \left(\sum_{k \neq i} (a_{ik} x_k) + 2a_{ii} x_i + \sum_{j \neq i} (a_{ji} x_j) \right) - b_i \\
&= \frac{1}{2} \left(\sum_{k=1}^m (a_{ik} x_k) + \sum_{j=1}^m (a_{ji} x_j) \right) - b_i \\
&= \frac{1}{2} \left((A\mathbf{x})_i + (A^T \mathbf{x})_i \right) - b_i
\end{aligned}$$

This is one entry of the vector $\nabla\phi(\mathbf{x})$ therefore we can write the entire vector as

$$\nabla\phi(\mathbf{x}) = \frac{1}{2} (A\mathbf{x} + A^T \mathbf{x}) - \mathbf{b}$$

Since A is symmetric, $A = A^T$, this simplifies to

$$\nabla\phi(\mathbf{x}) = A\mathbf{x} - \mathbf{b}$$

Now assume that $\nabla\phi(\hat{\mathbf{x}}) \perp K$, therefore $\mathbf{x} \cdot (A\hat{\mathbf{x}} - \mathbf{b}) = 0$ for any $\mathbf{x} \in K$. Let $\mathbf{x} \in K$, then $\mathbf{x} = \hat{\mathbf{x}} + \mathbf{y}$ for some $\mathbf{y} \in K$.

$$\begin{aligned}
\phi(\mathbf{x}) &= \phi(\hat{\mathbf{x}} + \mathbf{y}) \\
&= \frac{1}{2} (\hat{\mathbf{x}} + \mathbf{y})^T A (\hat{\mathbf{x}} + \mathbf{y}) - (\hat{\mathbf{x}} + \mathbf{y})^T \mathbf{b} \\
&= \frac{1}{2} (\hat{\mathbf{x}}^T + \mathbf{y}^T) A (\hat{\mathbf{x}} + \mathbf{y}) - \hat{\mathbf{x}}^T \mathbf{b} - \mathbf{y}^T \mathbf{b} \\
&= \frac{1}{2} (\hat{\mathbf{x}}^T A + \mathbf{y}^T A) (\hat{\mathbf{x}} + \mathbf{y}) - \hat{\mathbf{x}}^T \mathbf{b} - \mathbf{y}^T \mathbf{b} \\
&= \frac{1}{2} (\hat{\mathbf{x}}^T A \hat{\mathbf{x}} + \hat{\mathbf{x}}^T A \mathbf{y} + \mathbf{y}^T A \hat{\mathbf{x}} + \mathbf{y}^T A \mathbf{y}) - \hat{\mathbf{x}}^T \mathbf{b} - \mathbf{y}^T \mathbf{b}
\end{aligned}$$

Note that $\hat{\mathbf{x}}^T A \mathbf{y} = \mathbf{y}^T A^T \hat{\mathbf{x}} = \mathbf{y}^T A \hat{\mathbf{x}}$ because A is symmetric

$$\begin{aligned} &= \frac{1}{2} \left(\hat{\mathbf{x}}^T A \hat{\mathbf{x}} + 2 \mathbf{y}^T A \hat{\mathbf{x}} + \mathbf{y}^T A \mathbf{y} \right) - \hat{\mathbf{x}}^T \mathbf{b} - \mathbf{y}^T \mathbf{b} \\ &= \frac{1}{2} \hat{\mathbf{x}}^T A \hat{\mathbf{x}} - \hat{\mathbf{x}}^T \mathbf{b} + \mathbf{y}^T A \hat{\mathbf{x}} - \mathbf{y}^T \mathbf{b} + \frac{1}{2} \mathbf{y}^T A \mathbf{y} \\ &= \frac{1}{2} \hat{\mathbf{x}}^T A \hat{\mathbf{x}} - \hat{\mathbf{x}}^T \mathbf{b} + \mathbf{y}^T (A \hat{\mathbf{x}} - \mathbf{b}) + \frac{1}{2} \mathbf{y}^T A \mathbf{y} \end{aligned}$$

We know that $\nabla \phi(\hat{\mathbf{x}}) \perp K$, therefore $\mathbf{y}^T (A \hat{\mathbf{x}} - \mathbf{b}) = 0$

$$\begin{aligned} &= \frac{1}{2} \hat{\mathbf{x}}^T A \hat{\mathbf{x}} - \hat{\mathbf{x}}^T \mathbf{b} + \frac{1}{2} \mathbf{y}^T A \mathbf{y} \\ &= \phi(\hat{\mathbf{x}}) + \frac{1}{2} \mathbf{y}^T A \mathbf{y} \end{aligned}$$

Since A is positive definite $\mathbf{y}^T A \mathbf{y} \geq 0$ and therefore

$$\phi(\mathbf{x}) \geq \phi(\hat{\mathbf{x}})$$

This $\hat{\mathbf{x}}$ minimizes $\phi(\mathbf{x})$ over K .

Now assume that $\hat{\mathbf{x}}$ minimizes $\phi(\mathbf{x})$ over K . □

3. Show that:

(a) (Forward error analysis)

$$\left| fl(\mathbf{x}^T \mathbf{a}) - \mathbf{x}^T \mathbf{a} \right| \leq n \epsilon_{machine} |\mathbf{x}|^T |\mathbf{a}| + O(\epsilon_{machine}^2)$$

where \mathbf{x} and \mathbf{a} are n -dimensional floating point vectors and $fl(\mathbf{x}^T \mathbf{a})$ represents the floating point computation of the dot product.

Proof. I will prove by induction. First consider the case when $n = 1$, then

$$|fl(xa) - xa| = |xa(1 + \epsilon) - xa|$$

Where $\epsilon = \epsilon_{machine} + O(\epsilon_{machine}^2)$

$$\begin{aligned} &= \epsilon |x| |a| \\ &= 1 \epsilon_{machine} |x| |a| + O(\epsilon_{machine}^2) \end{aligned}$$

Assume that

$$\left| fl(\mathbf{x}^T \mathbf{a}) - \mathbf{x}^T \mathbf{a} \right| \leq n \epsilon_{machine} |\mathbf{x}|^T |\mathbf{a}| + O(\epsilon_{machine}^2)$$

for $n = 1, 2, \dots, k$. Now consider the case when $n = k + 1$. In this case $\mathbf{x} = [\mathbf{x}_k, x_{k+1}]^T$ and $\mathbf{a} = [\mathbf{a}_k, a_{k+1}]^T$.

$$\begin{aligned}
|fl(\mathbf{x}^T \mathbf{a}) - \mathbf{x}^T \mathbf{a}| &= |fl(\mathbf{x}_k^T \mathbf{a}_k + v_{k+1} a_{k+1}) - \mathbf{x}_k^T \mathbf{a}_k - v_{k+1} a_{k+1}| \\
&= \left| \left(fl(\mathbf{x}_k^T \mathbf{a}_k) + fl(v_{k+1} a_{k+1}) \right) (1 + \epsilon) - \mathbf{x}_k^T \mathbf{a}_k - v_{k+1} a_{k+1} \right| \\
&= \left| fl(\mathbf{x}_k^T \mathbf{a}_k)(1 + \epsilon) - \mathbf{x}_k^T \mathbf{a}_k + fl(v_{k+1} a_{k+1})(1 + \epsilon) - v_{k+1} a_{k+1} \right| \\
&\leq \left| fl(\mathbf{x}_k^T \mathbf{a}_k)(1 + \epsilon) - \mathbf{x}_k^T \mathbf{a}_k \right| + \left| fl(v_{k+1} a_{k+1})(1 + \epsilon) - v_{k+1} a_{k+1} \right| \\
&\leq k \epsilon_{machine} |\mathbf{x}_k|^T |\mathbf{a}_k| + O(\epsilon_{machine}^2) + \left| fl(v_{k+1} a_{k+1})(1 + \epsilon) - v_{k+1} a_{k+1} \right| \\
&\leq k \epsilon_{machine} |\mathbf{x}_k|^T |\mathbf{a}_k| + O(\epsilon_{machine}^2) + (\epsilon_{machine}) |v_{k+1}| |a_{k+1}| + O(\epsilon_{machine}^2) \\
&\leq (k + 1) \epsilon_{machine} |\mathbf{x}|^T |\mathbf{a}| + O(\epsilon_{machine}^2)
\end{aligned}$$

Thus

$$|fl(\mathbf{x}^T \mathbf{a}) - \mathbf{x}^T \mathbf{a}| \leq n \epsilon_{machine} |\mathbf{x}|^T |\mathbf{a}| + O(\epsilon_{machine}^2)$$

for all n . □

(b)

(c)

4. Let $A \in \mathbb{R}^{n \times n}$ be a symmetric positive definite matrix. Let Gaussian elimination be carried out on A without pivoting. After k steps, A will be reduced to the form

$$A^{(k)} = \begin{pmatrix} A_{11}^{(k)} & A_{12}^{(k)} \\ 0 & A_{22}^{(k)} \end{pmatrix}$$

where $A_{22}^{(k)}$ is an $(n - k) \times (n - k)$ matrix. Show by induction

- (a) $A_{22}^{(k)}$ is symmetric positive definite.

Proof. □

- (b) $a_{ii}^{(k)} \leq a_{ii}^{(k-1)}$ for all $k \leq i \leq n$, $k = 1, \dots, n - 1$.

Proof. □

5. Let $A \in \mathbb{R}^{m \times n}$ with $m > n$ and

$$A = \begin{pmatrix} A_1 \\ A_2 \end{pmatrix}$$

where A_1 is a nonsingular $n \times n$ matrix, and A_2 is an $(m - n) \times n$ arbitrary matrix.

- (a) What is the pseudo-inverse A^+ of A such that $A^+A = I_n$? Express it explicitly in terms of A_1 and A_2 .

The pseudo-inverse of A is defined as

$$A^+ = (A^T A)^{-1} A^T$$

Writing this in terms of A_1 and A_2 results in

$$\begin{aligned} A^+ &= \left(\begin{pmatrix} A_1^T & A_2^T \end{pmatrix} \begin{pmatrix} A_1 \\ A_2 \end{pmatrix} \right)^{-1} \begin{pmatrix} A_1^T & A_2^T \end{pmatrix} \\ &= \left(A_1^T A_1 + A_2^T A_2 \right)^{-1} \begin{pmatrix} A_1^T & A_2^T \end{pmatrix} \end{aligned}$$

- (b) Prove that $\|A^+\|_2 \leq \|A_1^{-1}\|_2$.

Proof.

$$\|A^+\|_2 = \sup_{\mathbf{b}} \frac{\|A^+ \mathbf{b}\|_2}{\|\mathbf{b}\|_2}$$

Let P be the orthogonal projector onto the range of A , then because P is unitary and the 2-norm is unitarily invariant, $\|\mathbf{b}\|_2 = \|P\mathbf{b}\|_2$. Therefore

$$\|A^+\|_2 = \sup_{\mathbf{b}} \frac{\|A^+ \mathbf{b}\|_2}{\|P\mathbf{b}\|_2}.$$

Now considering the least squares problem $\min_{\mathbf{x}} \|A\mathbf{x} - b\|_2$, we know that $A^+ \mathbf{b} = \mathbf{x}$ and $P\mathbf{b} = A\mathbf{x}$. Therefore

$$\|A^+\|_2 = \sup_{\mathbf{x}} \frac{\|\mathbf{x}\|_2}{\|A\mathbf{x}\|_2}.$$

Note that

$$A\mathbf{x} = \begin{pmatrix} A_1 \mathbf{x} \\ A_2 \mathbf{x} \end{pmatrix}$$

Therefore $\|A\mathbf{x}\|_2 \geq \|A_1 \mathbf{x}\|_2$, so

$$\|A^+\|_2 \leq \sup_{\mathbf{x}} \frac{\|\mathbf{x}\|_2}{\|A_1 \mathbf{x}\|_2}.$$

If we let $\mathbf{y} = A_1 \mathbf{x}$, then $\mathbf{x} = A_1^{-1} \mathbf{y}$ and

$$\begin{aligned} \|A^+\|_2 &\leq \sup_{\mathbf{y}} \frac{\|A_1^{-1} \mathbf{y}\|_2}{\|\mathbf{y}\|_2} \\ &= \|A_1^{-1}\|_2 \end{aligned}$$

Thus $\|A^+\|_2 \leq \|A_1^{-1}\|_2$. □

6. Let $A \in \mathbb{C}^{m \times m}$ with $\text{rank}(A) = r$. Suppose an SVD of A is given by $A = U\Sigma V^*$, where $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_m$ denote the columns of U and $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_m$ denote the columns of V . Prove that $\langle \mathbf{v}_{r+1}, \dots, \mathbf{v}_m \rangle = (A)^\perp$.

Proof. First let $\mathbf{x} \in \langle \mathbf{v}_{r+1}, \dots, \mathbf{v}_m \rangle$, then $\mathbf{x} = \sum_{i=r+1}^m (b_i \mathbf{v}_i)$. If we let $b_i = 0$ for $i = 1, 2, \dots, r$ and $\mathbf{b} = [b_i]$, then $\mathbf{x} = V\mathbf{b}$. Now consider $A\mathbf{x}$.

$$\begin{aligned} A\mathbf{x} &= U\Sigma V^* V\mathbf{b} \\ &= U\Sigma\mathbf{b} \end{aligned}$$

However Σ is a diagonal matrix with σ_i along the diagonal, so $(\Sigma\mathbf{b})_i = \sigma_i b_i$. Since A has $\text{rank}(A) = r$, we know that $\sigma_i = 0$ for $i \geq r+1$. Therefore if $1 \leq i \leq r$, then $\sigma_i b_i = 0$ because $b_i = 0$. If $r+1 \leq i \leq m$, then $\sigma_i b_i = 0$ because $\sigma_i = 0$. Therefore we can conclude that $\Sigma\mathbf{b} = \mathbf{0}$. Thus $A\mathbf{x} = \mathbf{0}$ and $\mathbf{x} \in (A)^\perp$.

Now assume that $\mathbf{x} \in (A)^\perp$, that is $A\mathbf{x} = \mathbf{0}$.

$$\begin{aligned} A\mathbf{x} &= \mathbf{0} \\ U\Sigma V^* \mathbf{x} &= \mathbf{0} \\ U^* U \Sigma V^* \mathbf{x} &= U^* \mathbf{0} \\ \Sigma V^* \mathbf{x} &= \mathbf{0} \\ \begin{bmatrix} \sigma_1 \mathbf{v}_1^* \mathbf{x} \\ \dots \\ \sigma_r \mathbf{v}_r^* \mathbf{x} \\ \sigma_{r+1} \mathbf{v}_{r+1}^* \mathbf{x} \\ \dots \\ \sigma_m \mathbf{v}_m^* \mathbf{x} \end{bmatrix} &= \mathbf{0} \end{aligned}$$

Since $\sigma_i = 0$ for $r+1 \leq i \leq m$

$$\begin{bmatrix} \sigma_1 \mathbf{v}_1^* \mathbf{x} \\ \dots \\ \sigma_r \mathbf{v}_r^* \mathbf{x} \\ 0 \dots \\ 0 \end{bmatrix} = \mathbf{0}$$

This implies that $\mathbf{v}_i^* \mathbf{x} = 0$ for $1 \leq i \leq r$ since $\sigma_i > 0$ for $1 \leq i \leq r$. This is equivalent to $\mathbf{x} \perp \mathbf{v}_i$ for $1 \leq i \leq r$. Hence $\mathbf{x} \in \langle \mathbf{v}_1, \dots, \mathbf{v}_r \rangle^\perp$. Since V is unitary

$$\langle \mathbf{v}_1, \dots, \mathbf{v}_r \rangle^\perp = \langle \mathbf{v}_{r+1}, \dots, \mathbf{v}_m \rangle$$

Thus $\mathbf{x} \in \langle \mathbf{v}_{r+1}, \dots, \mathbf{v}_m \rangle$. □

7. Problem 33.2 (Page 255) Suppose algorithm 33.1 is executed for a particular A and \mathbf{b} until at some step n , an entry $h_{n+1,n} = 0$ is encountered.

- (a) Show how (33.13) can be simplified in this case. What does this imply about the structure of a full $m \times m$ Hessenberg reduction $A = QHQ^*$ of A ?
- (b) Show that K_n is an invariant subspace of A , i.e., $AK_n \subseteq K_n$.
- (c) Show that if the Krylov subspaces of A generated by \mathbf{b} are defined by $K_k = \langle \mathbf{b}, A\mathbf{b}, \dots, A^{k-1}\mathbf{b} \rangle$, then $K_n = K_{n+1} = K_{n+2}$.
- (d) Show that each eigenvalue of H_n is an eigenvalue of A .
- (e) Show that if A is nonsingular, then the solution \mathbf{x} to the system of equations $A\mathbf{x} = \mathbf{b}$ lies in K_n .

8. Problem 36.1 (Page 283)

9. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be twice continuously differentiable for all x in the neighborhood $\{x \in \mathbb{R} \mid |x - \xi| < r\}$ of a simple zero ξ of f such that $f(\xi) = 0$. Consider the two-step Newton method:

$$y_k = x_k - f(x_k)/f'(x_k), \quad x_{k+1} = y_k - f(y_k)/f'(y_k).$$

- (a) Show that if the method converges, then

$$\lim_{k \rightarrow \infty} \left(\frac{x_{k+1} - \xi}{(y_k - \xi)(x_k - \xi)} \right) = \frac{f''(\xi)}{f'(\xi)}$$

Proof. First we assume that the method converges, this implies that

$$\lim_{k \rightarrow \infty} (x_k - \xi) = 0$$

and

$$\lim_{k \rightarrow \infty} (y_k - \xi) = 0$$

Also the problem will use the Taylor expansion of $f(y_k)$ and $f'(x_k)$ about ξ , so these will be stated here for later use.

$$\begin{aligned} f(y_k) &= f(\xi) + (y_k - \xi)f'(\xi) + O((y_k - \xi)^2) \\ &= (y_k - \xi)(f'(\xi) + O(y_k - \xi)) \\ f'(x_k) &= f'(\xi) + (x_k - \xi)f''(\xi) + O((x_k - \xi)^2) \end{aligned}$$

Now I will consider the limit

$$\begin{aligned}
& \lim_{k \rightarrow \infty} \left(\frac{1}{(y_k - \xi)(x_k - \xi)} (x_{k+1} - \xi) \right) \\
&= \lim_{k \rightarrow \infty} \left(\frac{1}{(y_k - \xi)(x_k - \xi)} \left(y_k - \xi - \frac{f(y_k)}{f'(x_k)} \right) \right) \\
&= \lim_{k \rightarrow \infty} \left(\frac{1}{(y_k - \xi)(x_k - \xi)} \left(y_k - \xi - \frac{(y_k - \xi)(f'(\xi) + O(y_k - \xi))}{f'(x_k)} \right) \right) \\
&= \lim_{k \rightarrow \infty} \left(\frac{1}{(x_k - \xi)} \left(1 - \frac{f'(\xi) + O(y_k - \xi)}{f'(x_k)} \right) \right) \\
&= \lim_{k \rightarrow \infty} \left(\frac{1}{(x_k - \xi)} \left(1 - \frac{f'(\xi) + O(y_k - \xi)}{f'(\xi) + (x_k - \xi)f''(\xi) + O((x_k - \xi)^2)} \right) \right) \\
&= \lim_{k \rightarrow \infty} \left(\frac{1}{(x_k - \xi)} \left(\frac{f'(\xi) + (x_k - \xi)f''(\xi) + O((x_k - \xi)^2) - f'(\xi) - O(y_k - \xi)}{f'(\xi) + (x_k - \xi)f''(\xi) + O((x_k - \xi)^2)} \right) \right) \\
&= \lim_{k \rightarrow \infty} \left(\frac{1}{(x_k - \xi)} \left(\frac{(x_k - \xi)f''(\xi) + O((x_k - \xi)^2) - O(y_k - \xi)}{f'(\xi) + (x_k - \xi)f''(\xi) + O((x_k - \xi)^2)} \right) \right) \\
&= \lim_{k \rightarrow \infty} \left(\frac{f''(\xi) + O((x_k - \xi)) - O((y_k - \xi)/(x_k - \xi))}{f'(\xi) + (x_k - \xi)f''(\xi) + O((x_k - \xi)^2)} \right)
\end{aligned}$$

Taking the limit results in

$$= \frac{f''(\xi)}{f'(\xi)}$$

□

(b) The convergence is cubic:

$$\lim_{k \rightarrow \infty} \left(\frac{x_{k+1} - \xi}{(x_k - \xi)^3} \right) = \frac{1}{2} \left(\frac{f''(\xi)}{f'(\xi)} \right)^2$$

Proof. In this proof I will use the Taylor expansions of $f(x_k)$ and $f'(x_k)$ around ξ , so I will state them here

$$\begin{aligned}
f(x_k) &= f(\xi) + (x_k - \xi)f'(\xi) + O((x_k - \xi)^2) \\
&= (x_k - \xi)(f'(\xi) + O(x_k - \xi)) \\
f'(x_k) &= f'(\xi) + (x_k - \xi)f''(\xi) + O((x_k - \xi)^2)
\end{aligned}$$

Now consider the limit

$$\lim_{k \rightarrow \infty} \left(\frac{x_{k+1} - \xi}{(x_k - \xi)^3} \right) = \lim_{k \rightarrow \infty} \left(\frac{x_{k+1} - \xi}{(y_k - \xi)(x_k - \xi)} \frac{y_k - \xi}{(x_k - \xi)^2} \right)$$

From part (a)

$$= \frac{f''(\xi)}{f'(\xi)} \lim_{k \rightarrow \infty} \left(\frac{y_k - \xi}{(x_k - \xi)^2} \right)$$

Now consider this limit individually

$$\begin{aligned} & \lim_{k \rightarrow \infty} \left(\frac{y_k - \xi}{(x_k - \xi)^2} \right) \\ &= \lim_{k \rightarrow \infty} \left(\frac{1}{(x_k - \xi)^2} \left(x_k - \xi - \frac{f(x_k)}{f'(x_k)} \right) \right) \\ &= \lim_{k \rightarrow \infty} \left(\frac{1}{(x_k - \xi)^2} \left(x_k - \xi - \frac{(x_k - \xi) \left(f'(\xi) + \frac{1}{2}(x_k - \xi)f''(\xi) + O((x_k - \xi)^2) \right)}{f'(x_k)} \right) \right) \\ &= \lim_{k \rightarrow \infty} \left(\frac{1}{x_k - \xi} \left(1 - \frac{f'(\xi) + \frac{1}{2}(x_k - \xi)f''(\xi) + O((x_k - \xi)^2)}{f'(x_k)} \right) \right) \\ &= \lim_{k \rightarrow \infty} \left(\frac{1}{x_k - \xi} \left(1 - \frac{f'(\xi) + \frac{1}{2}(x_k - \xi)f''(\xi) + O((x_k - \xi)^2)}{f'(\xi) + (x_k - \xi)f''(\xi) + O((x_k - \xi)^2)} \right) \right) \end{aligned}$$

By finding a common denominator and adding the fractions

$$\begin{aligned} &= \lim_{k \rightarrow \infty} \left(\frac{1}{x_k - \xi} \left(\frac{\frac{1}{2}(x_k - \xi)f''(\xi) + O((x_k - \xi)^2)}{f'(\xi) + (x_k - \xi)f''(\xi) + O((x_k - \xi)^2)} \right) \right) \\ &= \lim_{k \rightarrow \infty} \left(\frac{\frac{1}{2}f''(\xi) + O(x_k - \xi)}{f'(\xi) + (x_k - \xi)f''(\xi) + O((x_k - \xi)^2)} \right) \end{aligned}$$

Taking the limit

$$= \frac{1}{2} \frac{f''(\xi)}{f'(\xi)}$$

Now going back to the original limit

$$\begin{aligned} \lim_{k \rightarrow \infty} \left(\frac{x_{k+1} - \xi}{(x_k - \xi)^3} \right) &= \frac{f''(\xi)}{f'(\xi)} \lim_{k \rightarrow \infty} \left(\frac{y_k - \xi}{(x_k - \xi)^2} \right) \\ \lim_{k \rightarrow \infty} \left(\frac{x_{k+1} - \xi}{(x_k - \xi)^3} \right) &= \frac{1}{2} \left(\frac{f''(\xi)}{f'(\xi)} \right)^2 \end{aligned}$$

□

10. MATLAB project

Below are the function for performing the Jacobi method, the Gauss-Seidel Method, and the Conjugate Gradient method.

```

function [x0, k, r] = Jacobi(A, b, tol, maxIter)
    M = diag(diag(A));
    N = M - A;

    x0 = zeros(size(b));
    k = 0;
    r = norm(b - A*x0, inf);
    while(r(end) > tol && k < maxIter)
        k = k + 1;
        x = M\ (N*x0) + M\b;
        x0 = x;
        r = [r, norm(b - A*x0, inf)];
    end
end

```

```

function [x0, k, r] = GaussSeidel(A, b, tol, maxIter)
    M = tril(A);
    N = M - A;

    x0 = zeros(size(b));
    k = 0;
    r = norm(b - A*x0, inf);
    while(r(end) > tol && k < maxIter)
        k = k + 1;
        x = M\ (N*x0) + M\b;
        x0 = x;
        r = [r, norm(b - A*x0, inf)];
    end
end

```

```

function [x0, k, r] = ConjugateGradient(A, b, tol, maxIter)
    x0 = zeros(size(b));
    k = 0;
    r0 = b;
    p0 = r0;
    r = norm(r0, inf);
    while(r(end) > tol && k < maxIter)
        a = (r0'*r0)/(r0'*A*p0);
        x = x0 + a*p0;
        r1 = r0 - a*A*p0;
        bn = (r1'*r1)/(r0'*r0);
        p0 = r1 + bn*p0;

        % move to next step
        k = k+1;
        x0 = x;
    end
end

```

```

        r0 = r1;
        r = [r, norm(r0,inf)];
    end
end


```

The following script uses these three methods to solve the diffusion equation $-u_{xx} = 1$ on $x \in (0,1)$. It also plots the residual against the number of iterations.

```

%% Problem 10
% Initial matrix
tol = 1e-10;
maxIter = 2e6;
m = 500;
h = 1/(m+1);
e = ones(m, 1);
A = (1/h^2)*spdiags([-e, 2*e, -e], -1:1, m,m);
[xGS, kGS, rGS] = GaussSeidel(A, e, tol, maxIter);
[xJ, kJ, rJ] = Jacobi(A, e, tol, maxIter);
[xCG, kCG, rCG] = ConjugateGradient(A, e, tol, maxIter);
figure;
semilogy(0:kGS, rGS, 'g', 0:kJ, rJ, 'b', 0:kCG, rCG, 'r');
xlabel('Number of Iterations');
ylabel('Infinity Norm of Residual');
legend('Gauss-Seidel', 'Jacobi', 'Conjugate Gradient');

```



We see in this plot that the Gauss-Seidel method converges much faster than the Jacobi method. Both of these experience linear convergence, but Gauss-Seidel is a faster linear convergence. The Conjugate Gradient method converges much faster than either of the stationary methods. The Conjugate Gradient method converges in only 250 steps which is less than $m = 500$, the size of the matrix.