

# Lecture 02

## Orthogonal Vectors and Matrices; Vector Norms

Songting Luo

Department of Mathematics  
Iowa State University

MATH 562 Numerical Analysis II

# Outline

① Orthogonal Vectors and Matrices

② Vector Norms

# Outline

## ① Orthogonal Vectors and Matrices

## ② Vector Norms

# Inner Product

- Inner product (dot product) of two column vectors  $\mathbf{u}, \mathbf{v} \in \mathbb{C}^m$  is  $\mathbf{u}^* \mathbf{v} = \sum_{i=1}^m \bar{u}_i v_i$
- In contrast, outer product of  $\mathbf{u}, \mathbf{v}$  is  $\mathbf{u} \mathbf{v}^*$ .
- Note that cross product is different

# Inner Product

- Inner product (dot product) of two column vectors  $\mathbf{u}, \mathbf{v} \in \mathbb{C}^m$  is  $\mathbf{u}^* \mathbf{v} = \sum_{i=1}^m \bar{u}_i v_i$
- In contrast, outer product of  $\mathbf{u}, \mathbf{v}$  is  $\mathbf{u} \mathbf{v}^*$ .
- Note that cross product is different
- Euclidean length of  $\mathbf{u}$  is the square root of the inner product of  $\mathbf{u}$  with itself, i.e.,  $\sqrt{\mathbf{u}^* \mathbf{u}}$

# Inner Product

- Inner product (dot product) of two column vectors  $\mathbf{u}, \mathbf{v} \in \mathbb{C}^m$  is  $\mathbf{u}^* \mathbf{v} = \sum_{i=1}^m \bar{u}_i v_i$
- In contrast, outer product of  $\mathbf{u}, \mathbf{v}$  is  $\mathbf{u} \mathbf{v}^*$ .
- Note that cross product is different
- Euclidean length of  $\mathbf{u}$  is the square root of the inner product of  $\mathbf{u}$  with itself, i.e.,  $\sqrt{\mathbf{u}^* \mathbf{u}}$
- Inner product of two unit vectors  $\mathbf{u}$  and  $\mathbf{v}$  is the cosine of the angle  $\alpha$  between  $\mathbf{u}$  and  $\mathbf{v}$ , i.e.,  $\cos \alpha = \frac{\mathbf{u}^* \mathbf{v}}{\|\mathbf{u}\| \|\mathbf{v}\|}$

# Inner Product

- Inner product (dot product) of two column vectors  $\mathbf{u}, \mathbf{v} \in \mathbb{C}^m$  is  $\mathbf{u}^* \mathbf{v} = \sum_{i=1}^m \bar{u}_i v_i$
- In contrast, outer product of  $\mathbf{u}, \mathbf{v}$  is  $\mathbf{u} \mathbf{v}^*$ .
- Note that cross product is different
- Euclidean length of  $\mathbf{u}$  is the square root of the inner product of  $\mathbf{u}$  with itself, i.e.,  $\sqrt{\mathbf{u}^* \mathbf{u}}$
- Inner product of two unit vectors  $\mathbf{u}$  and  $\mathbf{v}$  is the cosine of the angle  $\alpha$  between  $\mathbf{u}$  and  $\mathbf{v}$ , i.e.,  $\cos \alpha = \frac{\mathbf{u}^* \mathbf{v}}{\|\mathbf{u}\| \|\mathbf{v}\|}$
- Inner product is bilinear, in the sense that it is linear in each vertex separately:

$$(\mathbf{u}_1 + \mathbf{u}_2)^* \mathbf{v} = \mathbf{u}_1^* \mathbf{v} + \mathbf{u}_2^* \mathbf{v}$$

$$\mathbf{u}^* (\mathbf{v}_1 + \mathbf{v}_2) = \mathbf{u}^* \mathbf{v}_1 + \mathbf{u}^* \mathbf{v}_2$$

$$(\alpha \mathbf{u})^* (\beta \mathbf{v}) = \bar{\alpha} \beta \mathbf{u}^* \mathbf{v}$$

# Orthogonal Vectors

## Definition

A pair of vectors  $\mathbf{x}, \mathbf{y}$  are orthogonal if  $\mathbf{x}^* \mathbf{y} = 0$

In other words, the angle between them is 90 degrees



# Orthogonal Vectors

## Definition

A pair of vectors  $\mathbf{x}, \mathbf{y}$  are orthogonal if  $\mathbf{x}^* \mathbf{y} = 0$

In other words, the angle between them is 90 degrees

## Definition

Two sets of vectors  $X$  and  $Y$  are orthogonal if every  $\mathbf{x} \in X$  is orthogonal to every  $\mathbf{y} \in Y$ .

# Orthogonal Vectors

## Definition

A pair of vectors  $\mathbf{x}, \mathbf{y}$  are orthogonal if  $\mathbf{x}^* \mathbf{y} = 0$

In other words, the angle between them is 90 degrees

## Definition

Two sets of vectors  $X$  and  $Y$  are orthogonal if every  $\mathbf{x} \in X$  is orthogonal to every  $\mathbf{y} \in Y$ .

## Definition

A set of nonzero vectors  $S$  is orthogonal if they are pairwise orthogonal. They are orthonormal if it is orthogonal and in addition each vector has unit Euclidean length.

# Orthogonal Vectors

## Theorem

*The vectors in an orthogonal set  $S$  are linearly independent.*

# Orthogonal Vectors

## Theorem

*The vectors in an orthogonal set  $S$  are linearly independent.*

## Proof?

# Orthogonal Vectors

## Theorem

*The vectors in an orthogonal set  $S$  are linearly independent.*

## Proof?

**Question:** If the column vectors of an  $m \times n$  matrix  $\mathbf{A}$  are orthogonal, what is the rank of  $\mathbf{A}$ ?

# Orthogonal Vectors

## Theorem

*The vectors in an orthogonal set  $S$  are linearly independent.*

## Proof?

**Question:** If the column vectors of an  $m \times n$  matrix  $\mathbf{A}$  are orthogonal, what is the rank of  $\mathbf{A}$ ?

**Answer:**  $\text{rank}(\mathbf{A}) = \min\{m, n\}$ , i.e.,  $\mathbf{A}$  has full rank.

# Components of Vector

- Given an orthonormal set  $\{\mathbf{q}_1, \mathbf{q}_2, \dots, \mathbf{q}_m\}$  forming a basis of  $\mathbb{C}^m$ , vector  $\mathbf{v}$  can be decomposed into orthogonal components as

$$\mathbf{v} = \sum_{i=1}^m (\mathbf{q}_i^* \mathbf{v}) \mathbf{q}_i$$

# Components of Vector

- Given an orthonormal set  $\{\mathbf{q}_1, \mathbf{q}_2, \dots, \mathbf{q}_m\}$  forming a basis of  $\mathbb{C}^m$ , vector  $\mathbf{v}$  can be decomposed into orthogonal components as
$$\mathbf{v} = \sum_{i=1}^m (\mathbf{q}_i^* \mathbf{v}) \mathbf{q}_i$$
- Another way to express the condition is  $\mathbf{v} = \sum_{i=1}^m (\mathbf{q}_i \mathbf{q}_i^*) \mathbf{v}$
- $\mathbf{q}_i \mathbf{q}_i^*$  is an orthogonal projection matrix. Note that it is NOT an orthogonal matrix.



# Components of Vector

- Given an orthonormal set  $\{\mathbf{q}_1, \mathbf{q}_2, \dots, \mathbf{q}_m\}$  forming a basis of  $\mathbb{C}^m$ , vector  $\mathbf{v}$  can be decomposed into orthogonal components as
$$\mathbf{v} = \sum_{i=1}^m (\mathbf{q}_i^* \mathbf{v}) \mathbf{q}_i$$
- Another way to express the condition is  $\mathbf{v} = \sum_{i=1}^m (\mathbf{q}_i \mathbf{q}_i^*) \mathbf{v}$
- $\mathbf{q}_i \mathbf{q}_i^*$  is an orthogonal projection matrix. Note that it is NOT an orthogonal matrix.
- More generally, given an orthonormal set  $\{\mathbf{q}_1, \mathbf{q}_2, \dots, \mathbf{q}_n\}$  with  $n \leq m$ , we have

$$\mathbf{v} = \mathbf{r} + \sum_{i=1}^n (\mathbf{q}_i^* \mathbf{v}) \mathbf{q}_i = \sum_{i=1}^n (\mathbf{q}_i \mathbf{q}_i^*) \mathbf{v}, \text{ and } \mathbf{r}^* \mathbf{q}_i = 0, 1 \leq i \leq n$$

- Let  $\mathbf{Q}$  be composed of column vectors  $\{\mathbf{q}_1, \mathbf{q}_2, \dots, \mathbf{q}_n\}$ ,  $\mathbf{Q}\mathbf{Q}^* = \sum_{i=1}^n (\mathbf{q}_i \mathbf{q}_i^*)$  is an orthogonal projection matrix.

# Unitary Matrices

## Definition

A matrix is unitary if  $\mathbf{Q}^* = \mathbf{Q}^{-1}$ , i.e., if  $\mathbf{Q}^* \mathbf{Q} = \mathbf{Q} \mathbf{Q}^* = \mathbf{I}$ .

- In the real case, we say the matrix is orthogonal. Its column vectors are orthonormal.
- In other words,  $\mathbf{q}_i^* \mathbf{q}_j = \delta_{ij}$ , the Kronecker delta.

# Unitary Matrices

## Definition

A matrix is unitary if  $\mathbf{Q}^* = \mathbf{Q}^{-1}$ , i.e., if  $\mathbf{Q}^* \mathbf{Q} = \mathbf{Q} \mathbf{Q}^* = \mathbf{I}$ .

- In the real case, we say the matrix is orthogonal. Its column vectors are orthonormal.
- In other words,  $\mathbf{q}_i^* \mathbf{q}_j = \delta_{ij}$ , the Kronecker delta.

**Question:** What is the geometric meaning of multiplication by a unitary matrix?

# Unitary Matrices

## Definition

A matrix is unitary if  $\mathbf{Q}^* = \mathbf{Q}^{-1}$ , i.e., if  $\mathbf{Q}^* \mathbf{Q} = \mathbf{Q} \mathbf{Q}^* = \mathbf{I}$ .

- In the real case, we say the matrix is orthogonal. Its column vectors are orthonormal.
- In other words,  $\mathbf{q}_i^* \mathbf{q}_j = \delta_{ij}$ , the Kronecker delta.

**Question:** What is the geometric meaning of multiplication by a unitary matrix?

**Answer:** It preserves angles and Euclidean length. In the real case, multiplication by an orthogonal matrix  $\mathbf{Q}$  is a rotation (if  $\det(\mathbf{Q}) = 1$ ) or reflection (if  $\det(\mathbf{Q}) = -1$ )

# Outline

## ① Orthogonal Vectors and Matrices

## ② Vector Norms

# Definition of Norms

- Norm captures “size” of vector or “distance” between vectors
- There are many different measures for “sizes” but a norm must satisfy some requirements:

# Definition of Norms

- Norm captures “size” of vector or “distance” between vectors
- There are many different measures for “sizes” but a norm must satisfy some requirements:

## Definition

A norm is a function:  $\| \cdot \| : \mathbb{C}^m \rightarrow \mathbb{R}$  that assigns a real-valued length to each vector. It must satisfy the following conditions:

- $\|\mathbf{x}\| \geq 0$ , and  $\|\mathbf{x}\| = 0$  only if  $\mathbf{x} = \mathbf{0}$
- $\|\mathbf{x} + \mathbf{y}\| \leq \|\mathbf{x}\| + \|\mathbf{y}\|$
- $\|\alpha\mathbf{x}\| = |\alpha|\|\mathbf{x}\|$

# Definition of Norms

- Norm captures “size” of vector or “distance” between vectors
- There are many different measures for “sizes” but a norm must satisfy some requirements:

## Definition

A norm is a function:  $\|\cdot\| : \mathbb{C}^m \rightarrow \mathbb{R}$  that assigns a real-valued length to each vector. It must satisfy the following conditions:

- $\|\mathbf{x}\| \geq 0$ , and  $\|\mathbf{x}\| = 0$  only if  $\mathbf{x} = \mathbf{0}$
- $\|\mathbf{x} + \mathbf{y}\| \leq \|\mathbf{x}\| + \|\mathbf{y}\|$
- $\|\alpha\mathbf{x}\| = |\alpha|\|\mathbf{x}\|$
- An example is Euclidean length (i.e.,  $\|\mathbf{x}\| = \sqrt{\sum_{i=1}^m |x_i|^2}$ )



## p-norms

- Euclidean length is a special case of  $p$ -norms, defined as

$$\|\mathbf{x}\|_p = \left( \sum_{i=1}^m |x_i|^p \right)^{1/p}$$

for  $1 \leq p \leq \infty$ .

## p-norms

- Euclidean length is a special case of  $p$ -norms, defined as

$$\|\mathbf{x}\|_p = \left( \sum_{i=1}^m |x_i|^p \right)^{1/p}$$

for  $1 \leq p \leq \infty$ .

- Euclidean norm is 2-norm (i.e.,  $p = 2$ ).

## p-norms

- Euclidean length is a special case of  $p$ -norms, defined as

$$\|\mathbf{x}\|_p = \left( \sum_{i=1}^m |x_i|^p \right)^{1/p}$$

for  $1 \leq p \leq \infty$ .

- Euclidean norm is 2-norm (i.e.,  $p = 2$ ).
- 1-norm:  $\|\mathbf{x}\|_1 = \sum_{i=1}^m |x_i|$

## p-norms

- Euclidean length is a special case of  $p$ -norms, defined as

$$\|\mathbf{x}\|_p = \left( \sum_{i=1}^m |x_i|^p \right)^{1/p}$$

for  $1 \leq p \leq \infty$ .

- Euclidean norm is 2-norm (i.e.,  $p = 2$ ).
- 1-norm:  $\|\mathbf{x}\|_1 = \sum_{i=1}^m |x_i|$
- $\infty$ -norm:  $\|\mathbf{x}\|_\infty$ . What is its value?

## p-norms

- Euclidean length is a special case of  $p$ -norms, defined as

$$\|\mathbf{x}\|_p = \left( \sum_{i=1}^m |x_i|^p \right)^{1/p}$$

for  $1 \leq p \leq \infty$ .

- Euclidean norm is 2-norm (i.e.,  $p = 2$ ).
- 1-norm:  $\|\mathbf{x}\|_1 = \sum_{i=1}^m |x_i|$
- $\infty$ -norm:  $\|\mathbf{x}\|_\infty$ . What is its value?
  - Answer:  $\|\mathbf{x}\|_\infty = \max_{1 \leq i \leq m} |x_i|$  (Proof ?)

## p-norms

- Euclidean length is a special case of  $p$ -norms, defined as

$$\|\mathbf{x}\|_p = \left( \sum_{i=1}^m |x_i|^p \right)^{1/p}$$

for  $1 \leq p \leq \infty$ .

- Euclidean norm is 2-norm (i.e.,  $p = 2$ ).
- 1-norm:  $\|\mathbf{x}\|_1 = \sum_{i=1}^m |x_i|$
- $\infty$ -norm:  $\|\mathbf{x}\|_\infty$ . What is its value?
  - Answer:  $\|\mathbf{x}\|_\infty = \max_{1 \leq i \leq m} |x_i|$  (Proof ?)
- Why we require  $p \geq 1$ ? What happens if  $0 < p < 1$ ?

## Weighted $p$ -norms

- A generalization of  $p$ -norm is weighted  $p$ -norm, which assigns different weights (priorities) to different components. (It is anisotropic instead of isotropic)
- Algebraically,  $\|\mathbf{x}\|_W = \|\mathbf{W}\mathbf{x}\|$ , where  $\mathbf{W}$  is diagonal matrix with  $i$ -th diagonal entry  $w_i \neq 0$  being weight for  $i$ -th component
- In other words,

$$\|\mathbf{x}\|_W = \left( \sum_{i=1}^m |w_i x_i|^p \right)^{1/p}$$

- What happens if we allow  $w_i = 0$ ?

## Weighted $p$ -norms

- A generalization of  $p$ -norm is weighted  $p$ -norm, which assigns different weights (priorities) to different components. (It is anisotropic instead of isotropic)
- Algebraically,  $\|\mathbf{x}\|_W = \|\mathbf{W}\mathbf{x}\|$ , where  $\mathbf{W}$  is diagonal matrix with  $i$ -th diagonal entry  $w_i \neq 0$  being weight for  $i$ -th component
- In other words,

$$\|\mathbf{x}\|_W = \left( \sum_{i=1}^m |w_i x_i|^p \right)^{1/p}$$

- What happens if we allow  $w_i = 0$ ?
- Can we further generalize it to allow  $\mathbf{W}$  being arbitrary matrix?



## Weighted $p$ -norms

- A generalization of  $p$ -norm is weighted  $p$ -norm, which assigns different weights (priorities) to different components. (It is anisotropic instead of isotropic)
- Algebraically,  $\|\mathbf{x}\|_W = \|\mathbf{W}\mathbf{x}\|$ , where  $\mathbf{W}$  is diagonal matrix with  $i$ -th diagonal entry  $w_i \neq 0$  being weight for  $i$ -th component
- In other words,

$$\|\mathbf{x}\|_W = \left( \sum_{i=1}^m |w_i x_i|^p \right)^{1/p}$$

- What happens if we allow  $w_i = 0$ ?
- Can we further generalize it to allow  $\mathbf{W}$  being arbitrary matrix?
- No. But we can allow  $\mathbf{W}$  to be arbitrary nonsingular matrix.