

Network flows - First Algorithm


Ford-Fulkerson Algorithm

Input: Network (G, u, s, t) .

Output: and s - t -flow f of maximum value

1. $f(e) = 0$ for all $e \in E(G)$
2. while f -augmenting path P exists:
3. compute $\gamma := \min_{e \in E(P)} u_f(e)$
4. augment f along P by γ (as much as possible)

1: How many iterations (at most) the algorithm needs at the following network:



(N is a big integer. Try to trick the algorithm to do many steps by picking unlucky P .)

Solution: Always use the edge uv with capacity 1. The augmentation will be always by one. So $2N$ iterations. The running time depends on u .

2: Show that s - t -flow f is maximum if and only if there is no f -augmenting path. (That is, Ford-Fulkerson algorithm is correct.)

Solution: If there is an augmenting path, then the flow can increase. If there is no augmenting path, then from s reachable vertices in reduced graph from a cut. Recall

$$\text{value}(f) = \sum_{e \in \delta^+(A)} f(e) - \sum_{e \in \delta^-(A)} f(e).$$

Since there is no augmenting path,

$$\sum_{e \in \delta^-(A)} f(e) = 0 \qquad \sum_{e \in \delta^+(A)} f(e) = \sum_{e \in \delta^-(A)} f(e).$$

This proves the maximality of the flow.

The question gives proof to

Theorem (Ford Fulkerson) Maximum value of an s - t -flow equals minimum capacity of an s - t -cut.

3: If $c : E \rightarrow \mathbb{Z}$, is it true that the flow produced from Ford-Fulekrson is integral and that the algorithm finishes in a finite time?

Solution: Yes, the augmenting is always by an integer. Every augmentation raises the value of the flow by at least one. This gives finiteness.

This proves

Theorem Dantzig Fulkerson: If the capacities are integral, then there exists an integral maximum flow.

4: If capacities are integral, is it true that every maximum flow is integral?

Solution: No - build network that has small cut and 2 paths to/or from it.

Theorem Gallai; Ford and Fulerson

Every flow f can be decomposed into s - t -paths \mathcal{P} and circuits \mathcal{C} with weight function $w : \mathcal{P} \cup \mathcal{C} \rightarrow \mathbb{R}_+$ such that

- $f(e) = \sum_{P \in \mathcal{P}} w(P) + \sum_{C \in \mathcal{C}} w(C)$
- $value(f) = \sum_{P \in \mathcal{P}} w(P)$.
- $|\mathcal{P} + \mathcal{C}| \leq |E(G)|$.

5: Prove the theorem. Hint, use induction on number of edges e , where $f(e) > 0$.

Solution: Take the longest circuit or s - t -path W . Assign weight w to W such that $f(h) = w(h)$ for some edge h and for all other edges $f(h) \geq w(h)$. Put W to the collection and apply induction.


Network flows - Menger's theorem

Theorem Menger: Let G be a graph (directed or undirected), let $s, t \in V(G)$, and $k \in \mathbb{N}$. Then there are k edge-disjoint s - t -paths iff after deleting any $k - 1$ edges t is still reachable from s .


6: Use flows to prove the theorem in directed case.

Solution: Assign capacities one to all edges, find maximum flow and use the previous decomposition theorem. Also notice that the maximum flow is integral.

7: Use the directed case to prove undirected case. Replacing an edge by a pair of opposite directed edges does not work. (Why?) Replace edge uv by a suitable orientation of the following gadget



Solution:



Paths P_1 and P_2 are called *internally disjoint* if they do not share more than the endvertices.

Theorem Menger: Let G be a graph (directed or undirected), let s and t be two non-adjacent vertices, and $k \in \mathbb{N}$. Then there are k pairwise internally disjoint s - t -paths iff after deleting any $k - 1$ vertices (distinct from s and t) t is still reachable from s .

8: Prove the directed case. Use the edge case.