

Caleb Logemann
MATH667 Hyperbolic Partial Differential Equations
Homework 7

1. Derive the following 3rd order accuracy reconstruction for finite volume methods.

$$u_{j+1/2}^- = -\frac{1}{6}\bar{u}_{j-1} + \frac{5}{6}\bar{u}_j + \frac{1}{3}\bar{u}_{j+1}$$

$$u_{j+1/2}^+ = \frac{1}{3}\bar{u}_j + \frac{5}{6}\bar{u}_{j+1} - \frac{1}{6}\bar{u}_{j+2}$$

For this problem we would like to construct a quadratic polynomial $p(x)$ such that p matches the cell average on intervals I_{j-1} , I_j , and I_{j+1} . This means that we need to set $p(x) = ax^2 + bx + c$ and solve the following set of equations for a , b , and c .

$$\frac{1}{h} \int_{x_{j-3/2}}^{x_{j-1/2}} p(x) dx = \bar{u}_{j-1}$$

$$\frac{1}{h} \int_{x_{j-1/2}}^{x_{j+1/2}} p(x) dx = \bar{u}_j$$

$$\frac{1}{h} \int_{x_{j+1/2}}^{x_{j+3/2}} p(x) dx = \bar{u}_{j+1}$$

These equation can be simplified as follows.

$$\frac{1}{3h}a(x_{j-1/2}^3 - x_{j-3/2}^3) + \frac{1}{2h}b(x_{j-1/2}^2 - x_{j-3/2}^2) + c = \bar{u}_{j-1}$$

$$\frac{1}{3h}a(x_{j+1/2}^3 - x_{j-1/2}^3) + \frac{1}{2h}b(x_{j+1/2}^2 - x_{j-1/2}^2) + c = \bar{u}_j$$

$$\frac{1}{3h}a(x_{j+3/2}^3 - x_{j+1/2}^3) + \frac{1}{2h}b(x_{j+3/2}^2 - x_{j+1/2}^2) + c = \bar{u}_{j+1}$$

Factoring the difference of cubes and difference of squares and cancelling h gives

$$\frac{1}{3}a(x_{j-1/2}^2 + x_{j-1/2}x_{j-3/2} + x_{j-3/2}^2) + \frac{1}{2}b(x_{j-1/2} + x_{j-3/2}) + c = \bar{u}_{j-1}$$

$$\frac{1}{3}a(x_{j+1/2}^2 + x_{j+1/2}x_{j-1/2} + x_{j-1/2}^2) + \frac{1}{2}b(x_{j+1/2} + x_{j-1/2}) + c = \bar{u}_j$$

$$\frac{1}{3}a(x_{j+3/2}^2 + x_{j+3/2}x_{j+1/2} + x_{j+1/2}^2) + \frac{1}{2}b(x_{j+3/2} + x_{j+1/2}) + c = \bar{u}_{j+1}.$$

Now I will substitute in so that all the x values are in terms of $x_{j+1/2}$ and h , i.e. $x_{j-3/2} = x_{j+1/2} - 2h$, $x_{j-1/2} = x_{j+1/2} - h$, etc.

$$\frac{1}{3}a\left((x_{j+1/2} - h)^2 + (x_{j+1/2} - h)(x_{j+1/2} - 2h) + (x_{j+1/2} - 2h)^2\right) + b\left(x_{j+1/2} - \frac{3}{2}h\right) + c = \bar{u}_{j-1}$$

$$\frac{1}{3}a\left(x_{j+1/2}^2 + x_{j+1/2}(x_{j+1/2} - h) + (x_{j+1/2} - h)^2\right) + b\left(x_{j+1/2} - \frac{1}{2}h\right) + c = \bar{u}_j$$

$$\frac{1}{3}a\left((x_{j+1/2} + h)^2 + (x_{j+1/2} + h)x_{j+1/2} + x_{j+1/2}^2\right) + b\left(x_{j+1/2} + \frac{1}{2}h\right) + c = \bar{u}_{j+1}.$$

Simplifying gives

$$\frac{1}{3}a(3x_{j+1/2}^2 - 9hx_{j+1/2} + 7h^2) + b\left(x_{j+1/2} - \frac{3}{2}h\right) + c = \bar{u}_{j-1}$$

$$\frac{1}{3}a(3x_{j+1/2}^2 - 3hx_{j+1/2} + 1h^2) + b\left(x_{j+1/2} - \frac{1}{2}h\right) + c = \bar{u}_j$$

$$\frac{1}{3}a(3x_{j+1/2}^2 + 3hx_{j+1/2} + 1h^2) + b\left(x_{j+1/2} + \frac{1}{2}h\right) + c = \bar{u}_{j+1}.$$

Subtracting the last two equations gives

$$\frac{1}{3}a(-6hx_{j+1/2}) - hb = \bar{u}_j - \bar{u}_{j+1}$$

2.

3.