

Caleb Logemann,
James
Rossmanith

Introduction

Method

Convection

Diffusion

Numerical Results

Conclusion

References

Discontinuous Galerkin Method for Solving Thin Film Equations

Caleb Logemann James Rossmanith

Mathematics Department,
Iowa State University

logemann@iastate.edu

April 20, 2019

Overview

Caleb Logemann,
James
Rossmanith

Introduction

Method

Convection

Diffusion

Numerical Results

Conclusion

References

1 Introduction

2 Method

- Convection
- Diffusion

3 Numerical Results

4 Conclusion

Motivation

Caleb Logemann,
James
Rossmanith

Introduction

Method

Convection


Diffusion

Numerical Results

Conclusion

References

- Aircraft Icing
- Runback



- Industrial Coating

Model Equations

Caleb Logemann,
James
Rossmanith

Introduction

Method

Convection

Diffusion

Numerical Results

Conclusion

References

■ Navier-Stokes Equation

$$\nabla \cdot \mathbf{u} = 0$$

$$\partial_t \mathbf{u} + \nabla \cdot (\mathbf{u}\mathbf{u}) = -\frac{1}{\rho} \nabla p + \frac{1}{\rho} \nabla \cdot \sigma + \mathbf{g}$$

$$\partial_t h_s + (u, v)^T \cdot \nabla h_s = w$$

$$\partial_t h_b + (u, v)^T \cdot \nabla h_b = w$$

- Lubrication or reduced Reynolds number approximation
- Thin-Film Equation - 1D with q as fluid height.

$$q_t + (f(x, t)q^2 - g(x, t)q^3)_x = -(h(x, t)q^3 q_{xxx})_x$$

Operator Splitting

Caleb Logemann,
James
Rossmanith

Introduction

Method

Convection

Diffusion

Numerical Results

Conclusion

References

■ Simplified Model

$$q_t + (q^2 - q^3)_x = -(q^3 q_{xxx})_x \quad (0, T) \times \Omega$$

■ Operator Splitting

$$q_t + (q^2 - q^3)_x = 0$$

$$q_t + (q^3 u_{xxx})_x = 0$$

■ Strang Splitting

$\frac{1}{2}\Delta t$ step of Convection

$$q_t + (q^2 - q^3)_x = 0$$

Δt step of Diffusion

$$q_t + (q^3 u_{xxx})_x = 0$$

$\frac{1}{2}\Delta t$ step of Convection

$$q_t + (q^2 - q^3)_x = 0$$

Convection

Caleb Logemann,
James
Rossmanith

Introduction

Method

Convection

Diffusion

Numerical Results

Conclusion

References

■ Convection Equation

$$\begin{aligned}q_t + f(q)_x &= 0 & (0, T) \times \Omega \\f(q) &= q^2 - q^3\end{aligned}$$

■ Weak Form

Find q such that

$$\int_{\Omega} (q_t v - f(q) v_x) dx + \hat{f} v \Big|_{\partial\Omega} = 0$$

for all test functions v

Notation

Caleb Logemann,
James
Rossmanith

Introduction

Method

Convection

Diffusion

Numerical Results

Conclusion


References

- Partition the domain, $[a, b]$ as

$$a = x_{1/2} < \cdots < x_{j-1/2} < x_{j+1/2} < \cdots < x_{N+1/2} = b$$

- $I_j = [x_{j-1/2}, x_{j+1/2}]$

- $x_j = \frac{x_{j+1/2} + x_{j-1/2}}{2}$.



Runge Kutta Discontinuous Galerkin

Caleb Logemann,
James
Rossmanith

Introduction

Method

Convection

Diffusion

Numerical Results

Conclusion

References

- Find $Q(t, x)$ such that for each time $t \in (0, T)$,
 $Q(t, \cdot) \in V_h = \left\{ v \in L^1(\Omega) : v|_{I_j} \in P^k(I_j) \right\}$

$$\begin{aligned} \int_{I_j} Q_t v \, dx &= \int_{I_j} f(Q) v_x \, dx \\ &\quad - \left(\mathcal{F}_{j+1/2} v^-(x_{j+1/2}) - \mathcal{F}_{j-1/2} v^+(x_{j-1/2}) \right) \end{aligned}$$

for all $v \in V_h$

- Rusanov/Local Lax-Friedrichs Numerical Flux

$$\mathcal{F}_{j+1/2} = \frac{1}{2} \left(f(Q_{j+1/2}^-) + f(Q_{j+1/2}^+) \right) + \frac{1}{2} \max_q \{ |f'(q)| \} (Q_{j+1/2}^- - Q_{j+1/2}^+)$$

- Solve this system of ODEs with any Explicit Strong Stability Preserving (SSP) Runge-Kutta Method.

Explicit SSP Runge Kutta Methods

Caleb Logemann,
James
Rossmanith

Introduction

Method

Convection

Diffusion

Numerical Results

Conclusion

References

■ Forward Euler

$$q^{n+1} = q^n + \Delta t L(q^n)$$

■ Second Order

$$q^* = q^n + \Delta t L(q^n)$$

$$q^{n+1} = \frac{1}{2}(q^n + q^*) + \frac{1}{2}\Delta t L(q^*)$$

Diffusion

Caleb Logemann,
James
Rossmanith

Introduction

Method

Convection

Diffusion

Numerical Results

Conclusion

References

■ Diffusion Equation

$$q_t = -(q^3 q_{xxx})_x \quad (0, T) \times \Omega$$

■ Linearize operator at $t = t^n$, let $f(x) = q^3(t = t^n, x)$

$$q_t = -(f(x) q_{xxx})_x \quad (0, T) \times \Omega$$

Local Discontinuous Galerkin

Caleb Logemann,
James
Rossmanith

Introduction

Method

Convection

Diffusion

Numerical Results

Conclusion

References

Find $Q(t, x), R(x), S(x), U(x)$ such that for all $t \in (0, T)$

$$Q(t, \cdot), R, S, U \in V_h = \left\{ v \in L^1(\Omega) : v|_{I_j} \in P^k(I_j) \right\}$$

$$\int_{I_j} Rv \, dx = - \int_{I_j} Qv_x \, dx + \left(\hat{Q}_{j+1/2} v_{j+1/2}^- - \hat{Q}_{j-1/2} v_{j-1/2}^+ \right)$$

$$\int_{I_j} Sw \, dx = - \int_{I_j} R w_x \, dx + \left(\hat{R}_{j+1/2} w_{j+1/2}^- - \hat{R}_{j-1/2} w_{j-1/2}^+ \right)$$

$$\begin{aligned} \int_{I_j} Uy \, dx = & \int_{I_j} S_x f y \, dx - \left(S_{j+1/2}^- f_{j+1/2}^- y_{j+1/2}^- - S_{j-1/2}^+ f_{j-1/2}^+ y_{j-1/2}^+ \right) \\ & + \left(\hat{S}_{j+1/2} \hat{f}_{j+1/2} y_{j+1/2}^- - \hat{S}_{j-1/2} \hat{f}_{j-1/2} y_{j-1/2}^+ \right) \end{aligned}$$

$$\int_{I_j} Q_t z \, dx = - \int_{I_j} U z_x \, dx + \left(\hat{U}_{j+1/2} z_{j+1/2}^- - \hat{U}_{j-1/2} z_{j-1/2}^+ \right)$$

for all $I_j \in \Omega$ and all $v, w, y, z \in V_h$.

Numerical Fluxes

Caleb Logemann,
James
Rossmanith

Introduction

Method

Convection

Diffusion

Numerical Results

Conclusion

References


$$\hat{f}_{j+1/2} = \frac{1}{2} \left(f_{j+1/2}^+ + f_{j+1/2}^- \right)$$

$$\hat{Q}_{j+1/2} = Q_{j+1/2}^+$$

$$\hat{R}_{j+1/2} = R_{j+1/2}^-$$

$$\hat{S}_{j+1/2} = S_{j+1/2}^+$$

$$\hat{U}_{j+1/2} = U_{j+1/2}^-$$



LDG Complications

Caleb Logemann,
James
Rossmanith

Introduction

Method

Convection

Diffusion

Numerical Results

Conclusion

References

- Explicit time step scales with h^4
- Implicit System is difficult to solve efficiently
 - GMRES iterations scale with size of system
 - Preconditioned GMRES

$$P = A_0^{-1}$$

$$PAx = Pb$$

- Geometric Multigrid fails to converge

Finite Difference Approach

Caleb Logemann,
James
Rossmanith

Introduction

Method

Convection

Diffusion

Numerical Results

Conclusion

References

- Let cell centers, x_i , form finite difference grid.
- Finite difference space, \mathbb{R}^N .
- $Q_{DG} \in V_h \rightarrow Q_{FD} \in \mathbb{R}^N$

$$(Q_{FD})_i = \frac{1}{h} \int_{K_i} Q_{DG} \, dx$$

- $Q_{FD} \in \mathbb{R}^N \rightarrow Q_{DG} \in V_h$

$$Q_{DG}|_K \in P^1(K)$$

$$\frac{1}{h} \int_{K_i} Q_{DG} \, dx = (Q_{FD})_i$$

$$\partial_x Q_{DG}|_{K_i} = \frac{(Q_{FD})_{i+1} - (Q_{FD})_{i-1}}{2h}$$

Finite Difference Approximation

Caleb Logemann,
James
Rossmanith

Introduction

Method

Convection

Diffusion

Numerical Results

Conclusion

References

■ First derivative approximation

$$(-(f(x)q_{xxx})_x)_i \approx -\frac{q_{i+1/2}^3(q_{xxx})_{i+1/2} - q_{i-1/2}^3(q_{xxx})_{i-1/2}}{h}$$

■ Third derivative approximation

$$(q_{xxx})_{i+1/2} \approx \frac{-Q_{i-1} + 3Q_i - 3Q_{i+1} + Q_{i+2}}{h^3}$$

■ Value of Q^3 at boundary

$$q_{i+1/2}^3 = \left(\frac{Q_i + Q_{i+1}}{2} \right)^3$$

Implicit L-Stable Runge Kutta

Caleb Logemann,
James
Rossmanith

Introduction

Method

Convection

Diffusion

Numerical Results

Conclusion

References

■ Backward Euler

$$q^{n+1} = q^n + \Delta t L(q^{n+1})$$

■ 2nd Order

$$\begin{aligned} q^* &= q^n + \frac{1}{4} \Delta t (L(q^n) + L(q^*)) \\ 3q^{n+1} &= 4q^* - q^n + \Delta t L(q^{n+1}) \end{aligned}$$

Nonlinear Solvers

Caleb Logemann,
James
Rossmanith

Introduction

Method

Convection

Diffusion

Numerical Results

Conclusion

References

■ Picard Iteration

$$L(q) = A(f \approx q^3)q$$

$$q_0^{n+1} = q^n$$

$$q_{m+1}^{n+1} = q^n + \Delta t A(q_m^{n+1}) q_{m+1}^{n+1}$$

$$q_{m+1}^* = q^n + \frac{1}{4} \Delta t (L(q^n) + A(q_m^*) q_{m+1}^*)$$

$$3q_{m+1}^{n+1} = 4q^* - q^n + \Delta t A(q_m^{n+1}) q_{m+1}^{n+1}$$

■ Newton's Method

$$q_{m+1}^{n+1} = q_m^{n+1} - J(q_m^{n+1})^{-1} F(q_m^{n+1})$$

$$F(q) = q - q^n - \Delta t L(q)$$

$$J(q) = I - \Delta t L'(q)$$

Manufactured Solution

Caleb Logemann,
James
Rossmanith

Introduction

Method

Convection

Diffusion

Numerical Results

Conclusion

References

$$q_t = -(q^3 q_{xxx})_x + s(x, t)$$
$$q(x, t) = 0.1 * \sin(2\pi(x - t)) + 0.15$$

Backward Euler				
1 Iteration			2 Iterations	
N	error	order	error	order
100	.0131	-	.0053	-
200	.0064	1.0264	.0026	1.0466
400	.0033	0.96	.0013	0.9704
800	.0016	1.0069	.0007	1.0134

Manufactured Solution

Caleb Logemann,
James
Rossmanith

Introduction

Method

Convection

Diffusion

Numerical Results

Conclusion

References

$$q_t = -(q^3 q_{xxx})_x + s(x, t)$$
$$q(x, t) = 0.1 * \sin(2\pi(x - t)) + 0.15$$

2nd Order IRK						
1 Iteration			2 Iterations		3 Iterations	
N	error	order	error	order	error	order
50	.0075	-	.00047	-	.0004901	-
100	.0041	0.8601	.00012	1.9844	.0001209	2.0194
200	.0020	1.0391	.0000312	1.9451	.0000305	1.9887
400	.0010	0.9652	.0000082	1.9244	.0000078	1.9641

Manufactured Solution

Caleb Logemann,
James
Rossmanith

Introduction

Method

Convection

Diffusion

Numerical Results

Conclusion

References

$$q_t = -(q^3 q_{xxx})_x + s(x, t)$$
$$q(x, t) = \frac{2}{10} e^{-10t} e^{-300(x-\frac{1}{2})^2} + \frac{1}{10}$$

Backward Euler				
N	1 Iteration		2 Iterations	
	error	order	error	order
100	.0097	-	.0933	-
200	.0050	0.95	.0421	1.1494
400	.0027	0.87	3.756	-6.48
800	33.21	-13.5	16.51	-2.14

Manufactured Solution - Newton's Method

Caleb Logemann,
James
Rossmanith

Introduction

Method

Convection

Diffusion

Numerical Results

Conclusion

References

$$q_t = -(q^3 q_{xxx})_x + s(x, t)$$
$$q(x, t) = \frac{2}{10} e^{-10t} e^{-300(x-\frac{1}{2})^2} + \frac{1}{10}$$

Backward Euler

N	error	order
50	0.0280	-
100	0.0153	0.8765
200	0.0080	0.9249
400	5.5e75	-258

Conclusion

Caleb Logemann,
James
Rossmanith

Introduction

Method

Convection

Diffusion

Numerical Results

Conclusion

References

Observations

- Expensive computations
- Nonlinear Hyper Diffusion has subtle instabilities

Future Work

- Hybridized Discontinuous Galerkin Method
- Higher Order Convergence
 - Higher order finite difference approximations
 - More accurate transition from finite difference to discontinuous Galerkin
 - Runge Kutta IMEX
- Space and time dependent coefficients

Bibliography

Caleb Logemann,
James
Rossmanith

Introduction

Method

Convection

Diffusion

Numerical Results

Conclusion

References

- [1] Y. Ha, Y.-J. Kim, and T.G. Myers. “On the numerical solution of a driven thin film equation”. In: *J. Comp. Phys.* 227.15 (2008), pp. 7246–7263.
- [2] T.G. Myers and J.P.F. Charpin. “A mathematical model for atmospheric ice accretion and water flow on a cold surface”. In: *Int. J. Heat and Mass Transfer* 47.25 (2004), pp. 5483–5500.
- [3] Tim G Myers. “Thin films with high surface tension”. In: *SIAM review* 40.3 (1998), pp. 441–462.
- [4] NASA. URL: http://icebox.grc.nasa.gov/gallery/images/C95_03918.html.
- [5] J.A. Rossmanith. DOGPack. Available from <http://www.dogpack-code.org/>.