

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

Discontinuous Galerkin Method for Solving Thin Film Equations

Caleb Logemann James Rossmanith

Mathematics Department,
Iowa State University

logemann@iastate.edu

May 13, 2019

Overview

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

1 Thin Film Equation

- Model
- Numerical Methods
- Results

2 Generalized Shallow Water Equations

- Model
- Numerical Methods
- Results
- Future Work

Model Equations

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods
Results


Generalized
Shallow Water
Equations

Model

Numerical Methods
Results

Future Work

References



■ Incompressible Navier-Stokes Equation

$$u_x + w_z = 0$$


$$\rho(u_t + uu_x + wu_z) = -p_x + \mu\Delta u - \phi_x$$

$$\rho(w_t + uw_x + ww_z) = -p_z + \mu\Delta w - \phi_z$$

$$w = 0, u = 0 \quad \text{at } z = 0$$

$$w = h_t + uh_x \quad \text{at } z = h$$

$$\mathbf{T} \cdot \mathbf{n} = (-\kappa\sigma + \Pi)\mathbf{n} + \left(\frac{\partial\sigma}{\partial s} + \tau\right)\mathbf{t} \quad \text{at } z = h$$



Nondimensionalize, integrate over Z , and simplify, gives

$$H_T + \left(\frac{1}{2}(\tau + \Sigma_x)H^2 - \frac{1}{3}(\Phi|_{Z=H} - \Pi)_x H^3 \right)_x = -\frac{1}{3}\bar{C}^{-1}(H^3 H_{xxx})_x$$

$$q_t + (q^2 - q^3)_x = -(q^3 q_{xxx})_x$$

Method Overview

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

■ Simplified Model

$$q_t + (q^2 - q^3)_x = -(q^3 q_{xxx})_x \quad (0, T) \times \Omega$$

■ Runge Kutta Implicit Explicit (IMEX)

$$q_t = F(q) + G(q)$$

■ F evaluated explicitly

■ G solved implicitly

$$F(q) = -(q^2 - q^3)_x$$

$$G(q) = (q^3 q_{xxx})_x$$

Convection

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

■ Convection Equation

$$\begin{aligned} F(q) &= f(q)_x = 0 & (0, T) \times \Omega \\ f(q) &= q^2 - q^3 \end{aligned}$$

■ Weak Form

Find q such that

$$\int_{\Omega} (F(q)v - f(q)v_x) \, dx + \hat{f}v \Big|_{\partial\Omega} = 0$$

for all test functions v

Notation

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work


References

- Partition the domain, $[a, b]$ as

$$a = x_{1/2} < \cdots < x_{j-1/2} < x_{j+1/2} < \cdots < x_{N+1/2} = b$$

- $I_j = [x_{j-1/2}, x_{j+1/2}]$

- $x_j = \frac{x_{j+1/2} + x_{j-1/2}}{2}$.



Runge Kutta Discontinuous Galerkin

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

- Find $Q(t, x)$ such that for each time $t \in (0, T)$,
 $Q(t, \cdot) \in V_h = \left\{ v \in L^1(\Omega) : v|_{I_j} \in P^k(I_j) \right\}$

$$\begin{aligned} \int_{I_j} F(Q) v \, dx &= \int_{I_j} f(Q) v_x \, dx \\ &\quad - \left(\mathcal{F}_{j+1/2} v^-(x_{j+1/2}) - \mathcal{F}_{j-1/2} v^+(x_{j-1/2}) \right) \end{aligned}$$

for all $v \in V_h$

- Rusanov/Local Lax-Friedrichs Numerical Flux

$$\mathcal{F}_{j+1/2} = \frac{1}{2} \left(f(Q_{j+1/2}^-) + f(Q_{j+1/2}^+) \right) + \frac{1}{2} \max_q \{ |f'(q)| \} (Q_{j+1/2}^- - Q_{j+1/2}^+)$$

Diffusion

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

■ Diffusion Equation

$$G(q) = -(q^3 q_{xxx})_x \quad (0, T) \times \Omega$$

■ Local Discontinuous Galerkin

$$r = q_x$$

$$s = r_x$$

$$u = s_x$$

$$G(q) = (q^3 u)_x$$

Local Discontinuous Galerkin

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

Find $Q(t, x), R(x), S(x), U(x)$ such that for all $t \in (0, T)$

$$Q(t, \cdot), R, S, U \in V_h = \left\{ v \in L^1(\Omega) : v|_{I_j} \in P^k(I_j) \right\}$$

$$\int_{I_j} Rv \, dx = - \int_{I_j} Qv_x \, dx + \left(\hat{Q}_{j+1/2} v_{j+1/2}^- - \hat{Q}_{j-1/2} v_{j-1/2}^+ \right)$$

$$\int_{I_j} Sw \, dx = - \int_{I_j} R w_x \, dx + \left(\hat{R}_{j+1/2} w_{j+1/2}^- - \hat{R}_{j-1/2} w_{j-1/2}^+ \right)$$

$$\int_{I_j} Uy \, dx = - \int_{I_j} S y_x \, dx + \left(\hat{S}_{j+1/2} y_{j+1/2}^- - \hat{S}_{j-1/2} y_{j-1/2}^+ \right)$$

$$\int_{I_j} G(Q)z \, dx = - \int_{I_j} Q^3 U z_x \, dx + \left(\hat{U}_{j+1/2} z_{j+1/2}^- - \hat{U}_{j-1/2} z_{j-1/2}^+ \right)$$

for all $I_j \in \Omega$ and all $v, w, y, z \in V_h$.

Numerical Fluxes

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work


References

$$\hat{Q}_{j+1/2} = Q_{j+1/2}^+$$

$$\hat{R}_{j+1/2} = R_{j+1/2}^-$$

$$\hat{S}_{j+1/2} = S_{j+1/2}^+$$

$$\hat{U}_{j+1/2} = (Q^3 U)_{j+1/2}^-$$



IMEX Runge Kutta

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

■ IMEX scheme

$$\begin{aligned}q^{n+1} &= q^n + \Delta t \sum_{i=1}^s (b'_i F(t_i, u_i)) + \Delta t \sum_{i=1}^s (b_i G(t_i, u_i)) \\u_i &= q^n + \Delta t \sum_{j=1}^{i-1} (a'_{ij} F(t_j, u_j)) + \Delta t \sum_{j=1}^i (a_{ij} G(t_j, u_j)) \\t_i &= t^n + c_i \Delta t\end{aligned}$$

■ Double Butcher Tableaus

$$\begin{array}{c|c} c' & a' \\ \hline & b'^T \end{array} \quad \begin{array}{c|c} c & a \\ \hline & b^T \end{array}$$

■ 1st Order — L-Stable SSP

$$\begin{array}{c|c} 0 & 0 \\ \hline & 1 \end{array} \quad \begin{array}{c|c} 1 & 1 \\ \hline & 1 \end{array}$$

■ 2nd Order — SSP

$$\begin{array}{c|ccc} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ \hline & 0 & \frac{1}{2} & \frac{1}{2} \end{array} \quad \begin{array}{c|cc} \frac{1}{2} & \frac{1}{2} & 0 \\ 0 & -\frac{1}{2} & 0 \\ 1 & 0 & \frac{1}{2} \\ \hline & 0 & \frac{1}{2} \end{array}$$

■ 3rd Order — L-Stable SSP

0	0	0	0	0	α	α	0	0	0
0	0	0	0	0	0	$-\alpha$	α	0	0
1	0	1	0	0	1	0	$1 - \alpha$	α	0
$\frac{1}{2}$	0	$\frac{1}{4}$	$\frac{1}{4}$	0	$\frac{1}{2}$	β	η	ζ	α
<hr/>					<hr/>				
	0	$\frac{1}{6}$	$\frac{1}{6}$	$\frac{2}{3}$		0	$\frac{1}{6}$	$\frac{1}{6}$	$\frac{2}{3}$

$$\alpha = 0.24169426078821$$

$$\beta = 0.06042356519705$$

$$\eta = 0.1291528696059$$

$$\zeta = \frac{1}{2} - \beta - \eta - \alpha$$

Nonlinear Solvers

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

■ Nonlinear System

$$u_i - a_{ij} \Delta t G(u_i) = b$$

■ Picard Iteration

$$\tilde{G}(q, u) = (q^3 u_{xxx})_x$$

$$u_0 = q^n \quad u_i^0 = u_{i-1}$$
$$u_i^j - a_{ij} \Delta t \tilde{G}(u_i^{j-1}, u_i^j) = b$$

- Number of picard iterations equals order in time and space

Manufactured Solution

$$q_t + (q^2 - q^3)_x = -(q^3 q_{xxx})_x + s$$

$$s = \hat{q}_t + (\hat{q}^2 - \hat{q}^3)_x + (\hat{q}^3 \hat{q}_{xxx})_x$$

$$\hat{q} = 0.1 \times \sin(2\pi/20.0 \times (x - t)) + 0.15 \quad \text{for } (x, t) \in [0, 40] \times [0, 5.0]$$

	1st Order		2nd Order		3rd Order	
n	error	order	error	order	error	order
20	0.136	—	7.33×10^{-3}	—	5.29×10^{-4}	—
40	0.0719	0.92	1.99×10^{-3}	1.88	5.38×10^{-5}	3.30
80	0.0378	0.93	5.60×10^{-4}	1.83	7.47×10^{-6}	2.85
160	0.0191	0.99	1.56×10^{-4}	1.85	9.97×10^{-7}	2.91
320	0.00961	0.99	3.98×10^{-5}	1.97	1.26×10^{-7}	2.98
640	0.00483	0.99	1.00×10^{-5}	1.99	1.58×10^{-8}	3.00
1280	0.00242	1.00	2.50×10^{-6}	2.00	1.98×10^{-9}	3.00

Table: Convergence table with a constant, linear, quadratic polynomial bases. CFL = 0.9, 0.2, 0.1 respectively.

Wave Structure with Nonlinear Hyper Diffusion

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model


Numerical Methods

Results

Future Work

References

$$q_t + (q^2 - q^3)_x = -(q^3 q_{xxx})_x$$
$$q_r = 0.1 \quad q_l = 0.3$$



Wave Structure with Nonlinear Hyper Diffusion

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods


Results

Future Work

References

$$q_r = 0.1 \quad q_l = 0.3323$$

$$q(x, 0) = (-\tanh(x) + 1) \frac{q_l - q_r}{2} + q_r$$



Wave Structure with Nonlinear Hyper Diffusion

Caleb Logemann,
James
Rossmanith


Thin Film
Equation
Model
Numerical Methods
Results

Generalized
Shallow Water
Equations

Model
Numerical Methods
Results
Future Work
References

$$q_r = 0.1 \quad q_l = 0.3323 \quad q_m = 0.6$$

$$q(x, 0) = \begin{cases} \frac{q_m - q_l}{2} \tanh(x) + \frac{q_m + q_l}{2} & x < 5 \\ -\frac{q_m - q_r}{2} \tanh(x - 10) + \frac{q_m + q_r}{2} + q_r & x > 5 \end{cases}$$



Wave Structure with Nonlinear Hyper Diffusion

Caleb Logemann,
James
Rossmanith


Thin Film
Equation
Model
Numerical Methods
Results

Generalized
Shallow Water
Equations

Model
Numerical Methods
Results
Future Work
References

$$q_r = 0.1 \quad q_l = 0.3323 \quad q_m = 0.6$$

$$q(x, 0) = \begin{cases} \frac{q_m - q_l}{2} \tanh(x) + \frac{q_m + q_l}{2} & x < 10 \\ -\frac{q_m - q_r}{2} \tanh(x - 20) + \frac{q_m + q_r}{2} + q_r & x > 10 \end{cases}$$



Wave Structure with Nonlinear Hyper Diffusion

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods


Results

Future Work

References

$$q_r = 0.1 \quad q_l = 0.4$$

$$q(x, 0) = (-\tanh(x - 100) + 1) \frac{q_l - q_r}{2} + q_r$$



Wave Structure with Nonlinear Hyper Diffusion

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods


Results

Future Work

References

$$q_r = 0.1 \quad q_l = 0.8$$

$$q(x, 0) = (-\tanh(x - 10) + 1) \frac{q_l - q_r}{2} + q_r$$



Bibliography I

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods
Results

Generalized
Shallow Water
Equations

Model

Numerical Methods
Results

Future Work

References

- [1] Andrea L Bertozzi, Andreas Münch, and Michael Shearer. “Undercompressive shocks in thin film flows”. In: *Physica D: Nonlinear Phenomena* 134.4 (1999), pp. 431–464.
- [2] Y. Ha, Y.-J. Kim, and T.G. Myers. “On the numerical solution of a driven thin film equation”. In: *J. Comp. Phys.* 227.15 (2008), pp. 7246–7263.
- [3] T.G. Myers and J.P.F. Charpin. “A mathematical model for atmospheric ice accretion and water flow on a cold surface”. In: *Int. J. Heat and Mass Transfer* 47.25 (2004), pp. 5483–5500.
- [4] Tim G Myers. “Thin films with high surface tension”. In: *SIAM review* 40.3 (1998), pp. 441–462.
- [5] NASA. URL: http://icebox.grc.nasa.gov/gallery/images/C95_03918.html.

Bibliography II

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods
Results

Generalized
Shallow Water
Equations

Model

Numerical Methods
Results

Future Work

References

- [6] Alexander Oron, Stephen H Davis, and S George Bankoff. “Long-scale evolution of thin liquid films”. In: *Reviews of modern physics* 69.3 (1997), p. 931.
- [7] J.A. Rossmanith. DOGPACK. Available from <http://www.dogpack-code.org/>.