

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

Discontinuous Galerkin Method for Solving Thin Film and Shallow Water Equations

Caleb Logemann James Rossmanith

Mathematics Department,
Iowa State University

logemann@iastate.edu

May 13, 2019

Overview

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References


1 Thin Film Equation

- Model
- Numerical Methods
- Results

2 Generalized Shallow Water Equations

- Model
- Numerical Methods
- Results
- Future Work

Model Equations



■ Incompressible Navier-Stokes Equation

$$u_x + w_z = 0$$


$$\rho(u_t + uu_x + ww_z) = -p_x + \mu\Delta u - \phi_x$$

$$\rho(w_t + uw_x + ww_z) = -p_z + \mu\Delta w - \phi_z$$

$$w = 0, u = 0 \quad \text{at } z = 0$$

$$w = h_t + uh_x \quad \text{at } z = h$$

$$\mathbf{T} \cdot \mathbf{n} = (-\kappa\sigma + \Pi)\mathbf{n} + \left(\frac{\partial\sigma}{\partial s} + \tau\right)\mathbf{t} \quad \text{at } z = h$$



Nondimensionalize, integrate over Z , and simplify, gives

$$H_T + \left(\frac{1}{2}(\tau + \Sigma_x)H^2 - \frac{1}{3}(\Phi|_{Z=H} - \Pi)_x H^3 \right)_x = -\frac{1}{3}\bar{C}^{-1}(H^3 H_{xxx})_x$$

$$q_t + (q^2 - q^3)_x = -(q^3 q_{xxx})_x$$

Method Overview

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

■ Simplified Model

$$q_t + (q^2 - q^3)_x = -(q^3 q_{xxx})_x \quad (0, T) \times \Omega$$

■ Runge Kutta Implicit Explicit (IMEX)

$$q_t = F(q) + G(q)$$

■ F evaluated explicitly

■ G solved implicitly

$$F(q) = -(q^2 - q^3)_x$$

$$G(q) = (q^3 q_{xxx})_x$$

Convection

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

■ Convection Equation

$$\begin{aligned} F(q) = f(q)_x &= 0 & (0, T) \times \Omega \\ f(q) &= q^2 - q^3 \end{aligned}$$

■ Weak Form

Find q such that

$$\int_{\Omega} (F(q)v - f(q)v_x) \, dx + \hat{f}v \Big|_{\partial\Omega} = 0$$

for all test functions v

Notation

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work


References

- Partition the domain, $[a, b]$ as

$$a = x_{1/2} < \cdots < x_{j-1/2} < x_{j+1/2} < \cdots < x_{N+1/2} = b$$

- $I_j = [x_{j-1/2}, x_{j+1/2}]$

- $x_j = \frac{x_{j+1/2} + x_{j-1/2}}{2}$.



Runge Kutta Discontinuous Galerkin

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

- Find $Q(t, x)$ such that for each time $t \in (0, T)$,
 $Q(t, \cdot) \in V_h = \left\{ v \in L^1(\Omega) : v|_{I_j} \in P^k(I_j) \right\}$

$$\begin{aligned} \int_{I_j} F(Q) v \, dx &= \int_{I_j} f(Q) v_x \, dx \\ &\quad - \left(\mathcal{F}_{j+1/2} v^-(x_{j+1/2}) - \mathcal{F}_{j-1/2} v^+(x_{j-1/2}) \right) \end{aligned}$$

for all $v \in V_h$

- Rusanov/Local Lax-Friedrichs Numerical Flux

$$\mathcal{F}_{j+1/2} = \frac{1}{2} (f(Q_{j+1/2}^-) + f(Q_{j+1/2}^+)) + \frac{1}{2} \max_q \{ |f'(q)| \} (Q_{j+1/2}^- - Q_{j+1/2}^+)$$

Diffusion

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

■ Diffusion Equation

$$G(q) = -(q^3 q_{xxx})_x \quad (0, T) \times \Omega$$

■ Local Discontinuous Galerkin

$$r = q_x$$

$$s = r_x$$

$$u = s_x$$

$$G(q) = (q^3 u)_x$$

Local Discontinuous Galerkin

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

Find $Q(t, x), R(x), S(x), U(x)$ such that for all $t \in (0, T)$

$$Q(t, \cdot), R, S, U \in V_h = \left\{ v \in L^1(\Omega) : v|_{I_j} \in P^k(I_j) \right\}$$

$$\int_{I_j} Rv \, dx = - \int_{I_j} Qv_x \, dx + \left(\hat{Q}_{j+1/2} v_{j+1/2}^- - \hat{Q}_{j-1/2} v_{j-1/2}^+ \right)$$

$$\int_{I_j} Sw \, dx = - \int_{I_j} R w_x \, dx + \left(\hat{R}_{j+1/2} w_{j+1/2}^- - \hat{R}_{j-1/2} w_{j-1/2}^+ \right)$$

$$\int_{I_j} Uy \, dx = - \int_{I_j} S y_x \, dx + \left(\hat{S}_{j+1/2} y_{j+1/2}^- - \hat{S}_{j-1/2} y_{j-1/2}^+ \right)$$

$$\int_{I_j} G(Q)z \, dx = - \int_{I_j} Q^3 U z_x \, dx + \left(\hat{U}_{j+1/2} z_{j+1/2}^- - \hat{U}_{j-1/2} z_{j-1/2}^+ \right)$$

for all $I_j \in \Omega$ and all $v, w, y, z \in V_h$.

Numerical Fluxes

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work


References

$$\hat{Q}_{j+1/2} = Q_{j+1/2}^+$$

$$\hat{R}_{j+1/2} = R_{j+1/2}^-$$

$$\hat{S}_{j+1/2} = S_{j+1/2}^+$$

$$\hat{U}_{j+1/2} = (Q^3 U)_{j+1/2}^-$$



IMEX Runge Kutta

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

■ IMEX scheme

$$\begin{aligned}q^{n+1} &= q^n + \Delta t \sum_{i=1}^s (b'_i F(t_i, u_i)) + \Delta t \sum_{i=1}^s (b_i G(t_i, u_i)) \\u_i &= q^n + \Delta t \sum_{j=1}^{i-1} (a'_{ij} F(t_j, u_j)) + \Delta t \sum_{j=1}^i (a_{ij} G(t_j, u_j)) \\t_i &= t^n + c_i \Delta t\end{aligned}$$

■ Double Butcher Tableaus

$$\begin{array}{c|c} c' & a' \\ \hline & b'^T \end{array} \quad \begin{array}{c|c} c & a \\ \hline & b^T \end{array}$$

■ 1st Order — L-Stable SSP

$$\begin{array}{c|c} 0 & 0 \\ \hline & 1 \end{array} \quad \begin{array}{c|c} 1 & 1 \\ \hline & 1 \end{array}$$

■ 2nd Order — SSP

$$\begin{array}{c|ccc} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ \hline & 0 & \frac{1}{2} & \frac{1}{2} \end{array} \quad \begin{array}{c|ccc} \frac{1}{2} & \frac{1}{2} & 0 & 0 \\ 0 & -\frac{1}{2} & \frac{1}{2} & 0 \\ 1 & 0 & \frac{1}{2} & \frac{1}{2} \\ \hline & 0 & \frac{1}{2} & \frac{1}{2} \end{array}$$

■ 3rd Order — L-Stable SSP

0	0	0	0	0	α	α	0	0	0
0	0	0	0	0	0	$-\alpha$	α	0	0
1	0	1	0	0	1	0	$1 - \alpha$	α	0
$\frac{1}{2}$	0	$\frac{1}{4}$	$\frac{1}{4}$	0	$\frac{1}{2}$	β	η	ζ	α
<hr/>					<hr/>				
	0	$\frac{1}{6}$	$\frac{1}{6}$	$\frac{2}{3}$		0	$\frac{1}{6}$	$\frac{1}{6}$	$\frac{2}{3}$

$$\alpha = 0.24169426078821$$

$$\beta = 0.06042356519705$$

$$\eta = 0.1291528696059$$

$$\zeta = \frac{1}{2} - \beta - \eta - \alpha$$

Nonlinear Solvers

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

■ Nonlinear System

$$u_i - a_{ij} \Delta t G(u_i) = b$$

■ Picard Iteration

$$\tilde{G}(q, u) = (q^3 u_{xxx})_x$$

$$u_0 = q^n \quad u_i^0 = u_{i-1}$$
$$u_i^j - a_{ij} \Delta t \tilde{G}(u_i^{j-1}, u_i^j) = b$$

■ Number of picard iterations equals order in time and space

Manufactured Solution

$$q_t + (q^2 - q^3)_x = -(q^3 q_{xxx})_x + s$$

$$s = \hat{q}_t + (\hat{q}^2 - \hat{q}^3)_x + (\hat{q}^3 \hat{q}_{xxx})_x$$

$$\hat{q} = 0.1 \times \sin(2\pi/20.0 \times (x - t)) + 0.15 \quad \text{for } (x, t) \in [0, 40] \times [0, 5.0]$$

1st Order			2nd Order		3rd Order	
n	error	order	error	order	error	order
20	0.136	—	7.33×10^{-3}	—	5.29×10^{-4}	—
40	0.0719	0.92	1.99×10^{-3}	1.88	5.38×10^{-5}	3.30
80	0.0378	0.93	5.60×10^{-4}	1.83	7.47×10^{-6}	2.85
160	0.0191	0.99	1.56×10^{-4}	1.85	9.97×10^{-7}	2.91
320	0.00961	0.99	3.98×10^{-5}	1.97	1.26×10^{-7}	2.98
640	0.00483	0.99	1.00×10^{-5}	1.99	1.58×10^{-8}	3.00
1280	0.00242	1.00	2.50×10^{-6}	2.00	1.98×10^{-9}	3.00

Table: Convergence table with a constant, linear, quadratic polynomial bases. CFL = 0.9, 0.2, 0.1 respectively.

Wave Structure with Nonlinear Hyper Diffusion

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model


Numerical Methods

Results

Future Work

References

$$q_t + (q^2 - q^3)_x = -(q^3 q_{xxx})_x$$
$$q_r = 0.1 \quad q_l = 0.3$$



Wave Structure with Nonlinear Hyper Diffusion

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods


Results

Future Work

References

$$q_r = 0.1 \quad q_l = 0.3323$$

$$q(x, 0) = (-\tanh(x) + 1) \frac{q_l - q_r}{2} + q_r$$



Wave Structure with Nonlinear Hyper Diffusion

Caleb Logemann,
James
Rossmanith


Thin Film
Equation
Model
Numerical Methods
Results

Generalized
Shallow Water
Equations

Model
Numerical Methods
Results
Future Work
References

$$q_r = 0.1 \quad q_l = 0.3323 \quad q_m = 0.6$$

$$q(x, 0) = \begin{cases} \frac{q_m - q_l}{2} \tanh(x) + \frac{q_m + q_l}{2} & x < 5 \\ -\frac{q_m - q_r}{2} \tanh(x - 10) + \frac{q_m + q_r}{2} + q_r & x > 5 \end{cases}$$



Wave Structure with Nonlinear Hyper Diffusion

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods


Results

Future Work

References

$$q_r = 0.1 \quad q_l = 0.3323 \quad q_m = 0.6$$

$$q(x, 0) = \begin{cases} \frac{q_m - q_l}{2} \tanh(x) + \frac{q_m + q_l}{2} & x < 10 \\ -\frac{q_m - q_r}{2} \tanh(x - 20) + \frac{q_m + q_r}{2} + q_r & x > 10 \end{cases}$$



Wave Structure with Nonlinear Hyper Diffusion

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods


Results

Future Work

References

$$q_r = 0.1 \quad q_l = 0.4$$

$$q(x, 0) = (-\tanh(x - 100) + 1) \frac{q_l - q_r}{2} + q_r$$



Wave Structure with Nonlinear Hyper Diffusion

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods


Results

Future Work

References

$$q_r = 0.1 \quad q_l = 0.8$$

$$q(x, 0) = (-\tanh(x - 10) + 1) \frac{q_l - q_r}{2} + q_r$$



Generalized Shallow Water

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods
Results


Generalized
Shallow Water
Equations

Model

Numerical Methods
Results

Future Work

References



$$\nabla \cdot \mathbf{u} = 0$$

$$\mathbf{u}_t + \nabla \cdot (\mathbf{u} \mathbf{u}) = -\frac{1}{\rho} \nabla p + \frac{1}{\rho} \nabla \cdot \boldsymbol{\sigma} + \mathbf{g}$$

$$(h_s)_t + [u(t, x, y, h_s), v(t, x, y, h_s)]^T \cdot \nabla h_s = w(t, x, y, h_s)$$

$$(h_b)_t + [u(t, x, y, h_b), v(t, x, y, h_b)]^T \cdot \nabla h_b = w(t, x, y, h_b)$$

Nondimensionalization

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

Characteristic Lengths

$$\varepsilon = \frac{H}{L}, \quad x = L\hat{x}, \quad y = L\hat{y}, \quad z = H\hat{z}$$

Characteristic Velocities

$$u = U\hat{u}, \quad v = U\hat{v}, \quad w = \varepsilon U\hat{w}$$

Characteristic Time

$$t = \frac{L}{U}\hat{t}$$

Characteristic Stresses

$$p = \rho g H \hat{p}, \quad \sigma_{xz/yz} = S \hat{\sigma}_{xz/yz}, \quad \sigma_{xx/xy/yy/zz} = \varepsilon S \hat{\sigma}_{xx/xy/yy/zz}$$

Nondimensionalization

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

$$\hat{u}_{\hat{x}} + \hat{v}_{\hat{y}} + \hat{w}_{\hat{z}} = 0$$

$$\begin{aligned} \varepsilon F^2 \left(\hat{u}_{\hat{t}} + (\hat{u}^2)_{\hat{x}} + (\hat{u}\hat{v})_{\hat{y}} + (\hat{u}\hat{w})_{\hat{z}} \right) &= -\varepsilon \hat{p}_{\hat{x}} \\ &+ G \left(\varepsilon^2 (\hat{\sigma}_{xx})_{\hat{x}} + \varepsilon^2 (\hat{\sigma}_{xy})_{\hat{y}} + (\hat{\sigma}_{xz})_{\hat{z}} \right) + e_x \end{aligned}$$

$$\begin{aligned} \varepsilon F^2 \left(\hat{v}_{\hat{t}} + (\hat{u}\hat{v})_{\hat{x}} + (\hat{v}^2)_{\hat{y}} + (\hat{v}\hat{w})_{\hat{z}} \right) &= -\varepsilon \hat{p}_{\hat{y}} \\ &+ G \left(\varepsilon^2 (\hat{\sigma}_{xy})_{\hat{x}} + \varepsilon^2 (\hat{\sigma}_{yy})_{\hat{y}} + (\hat{\sigma}_{yz})_{\hat{z}} \right) + e_y \end{aligned}$$

$$\begin{aligned} \varepsilon^2 F^2 \left(\hat{w}_{\hat{t}} + (\hat{u}\hat{w})_{\hat{x}} + (\hat{v}\hat{w})_{\hat{y}} + (\hat{w}^2)_{\hat{z}} \right) &= -\hat{p}_{\hat{z}} \\ &+ \varepsilon G \left((\hat{\sigma}_{xz})_{\hat{x}} + (\hat{\sigma}_{yz})_{\hat{y}} + (\hat{\sigma}_{zz})_{\hat{z}} \right) + e_z \end{aligned}$$

$$F = \frac{U}{\sqrt{gH}} \approx 1, \quad G = \frac{S}{\rho g H} < 1$$

Nondimensionalization

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

Drop ε^2 and εG terms

$$\hat{u}_{\hat{x}} + \hat{v}_{\hat{y}} + \hat{w}_{\hat{z}} = 0$$

$$\varepsilon F^2 \left(\hat{u}_{\hat{t}} + (\hat{u}^2)_{\hat{x}} + (\hat{u}\hat{v})_{\hat{y}} + (\hat{u}\hat{w})_{\hat{z}} \right) = -\varepsilon \hat{p}_{\hat{x}} + G(\hat{\sigma}_{xz})_{\hat{z}} + e_x$$

$$\varepsilon F^2 \left(\hat{v}_{\hat{t}} + (\hat{u}\hat{v})_{\hat{x}} + (\hat{v}^2)_{\hat{y}} + (\hat{v}\hat{w})_{\hat{z}} \right) = -\varepsilon \hat{p}_{\hat{y}} + G(\hat{\sigma}_{yz})_{\hat{z}} + e_y$$

$$\hat{p}_{\hat{z}} = e_z$$

Solving for the hydrostatic pressure

$$\hat{p}(\hat{t}, \hat{x}, \hat{y}) = \left(\hat{h}_s(\hat{t}, \hat{x}, \hat{y}) - \hat{z} \right) e_z$$

Nondimensionalization

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

Dimensional Variables

$$u_x + v_y + w_z = 0$$

$$u_t + (u^2)_x + (uv)_y + (uw)_z = -\frac{1}{\rho}p_x + \frac{1}{\rho}(\sigma_{xz})_z + ge_x$$

$$v_t + (uv)_x + (v^2)_y + (vw)_z = -\frac{1}{\rho}p_y + \frac{1}{\rho}(\sigma_{yz})_z + ge_y$$

$$p(t, x, y, z) = (h_s(t, x, y) - z)\rho ge_z$$

Kinematic Boundary Conditions

$$(h_s)_t + [u(t, x, y, h_s), v(t, x, y, h_s)]^T \cdot \nabla h_s = w(t, x, y, h_s)$$

$$(h_b)_t + [u(t, x, y, h_b), v(t, x, y, h_b)]^T \cdot \nabla h_b = w(t, x, y, h_b)$$

Mapping

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations



Model

Numerical Methods

Results

Future Work

References



Transform from $z \rightarrow \zeta$, by

$$\zeta = \frac{z - h_b(t, x, y)}{h(t, x, y)},$$

or equivalently

$$z = h(t, x, y)\zeta + h_b(t, x, y)$$

where $h(t, x, y) = h_s(t, x, y) - h_b(t, x, y)$.

$$\tilde{\Psi}(t, x, y, \zeta) = \Psi(t, x, y, h(t, x, y)\zeta + h_b(t, x, y))$$

Mapping Continuity Equation

$$u_x + v_y + w_z = 0$$

Map to new space

$$(h\tilde{u})_x - ((\zeta h + h_b)_x \tilde{u})_\zeta + (h\tilde{v})_y - ((\zeta h + h_b)_y \tilde{v})_\zeta + (\tilde{w})_\zeta = 0$$

Solve for vertical velocity, w ,

$$\begin{aligned} \tilde{w}(t, x, y, \zeta') = & - \left(h \int_0^{\zeta'} \tilde{u} d\zeta \right)_x - \left(h \int_0^{\zeta'} \tilde{v} d\zeta \right)_y \\ & + (\zeta' h + h_b)_x \tilde{u}(t, x, y, \zeta') + (\zeta' h + h_b)_y \tilde{v}(t, x, y, \zeta') \end{aligned}$$

Depth averaged equation

$$h_t + \left(h \int_0^1 \tilde{u} d\zeta \right)_x + \left(h \int_0^1 \tilde{v} d\zeta \right)_y = 0$$

Let u_m and v_m denote the mean velocity

$$h_t + (hu_m)_x + (hv_m)_y = 0$$

Mapping Momentum Equations

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

$$u_t + (u^2)_x + (uv)_y + (uw)_z = -\frac{1}{\rho}p_x + \frac{1}{\rho}(\sigma_{xz})_z + ge_x$$

$$v_t + (uv)_x + (v^2)_y + (vw)_z = -\frac{1}{\rho}p_y + \frac{1}{\rho}(\sigma_{yz})_z + ge_y$$

$$(h\tilde{u})_t + \left(h\tilde{u}^2 + \frac{1}{2}ge_z h^2\right)_x + (h\tilde{u}\tilde{v})_y + \left(h\tilde{u}\omega - \frac{1}{\rho}\tilde{\sigma}_{xz}\right)_\zeta = gh(e_x - e_z(h_b)_x)$$

$$(h\tilde{v})_t + (h\tilde{u}\tilde{v})_x + \left(h\tilde{v}^2 + \frac{1}{2}ge_z h^2\right)_y + \left(h\tilde{v}\omega - \frac{1}{\rho}\tilde{\sigma}_{yz}\right)_\zeta = gh(e_x - e_z(h_b)_y)$$

where

$$\omega = \frac{1}{h} \left(- \left(h \int_0^\zeta \tilde{u} - u_m d\zeta' \right)_x - \left(h \int_0^\zeta \tilde{v} - v_m d\zeta' \right)_y \right)$$

Mapped Reference System

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

$$h_t + (hu_m)_x + (hv_m)_y = 0$$

$$(h\tilde{u})_t + \left(h\tilde{u}^2 + \frac{1}{2}ge_z h^2\right)_x + (h\tilde{u}\tilde{v})_y + \left(h\tilde{u}\omega - \frac{1}{\rho}\tilde{\sigma}_{xz}\right)_\zeta = gh(e_x - e_z(h_b)_x)$$

$$(h\tilde{v})_t + (h\tilde{u}\tilde{v})_x + \left(h\tilde{v}^2 + \frac{1}{2}ge_z h^2\right)_y + \left(h\tilde{v}\omega - \frac{1}{\rho}\tilde{\sigma}_{yz}\right)_\zeta = gh(e_x - e_z(h_b)_y)$$

$$\omega = \frac{1}{h} \left(- \left(h \int_0^\zeta \tilde{u}_d \, d\zeta' \right)_x - \left(h \int_0^\zeta \tilde{v}_d \, d\zeta' \right)_y \right)$$

with

$$\tilde{u}_d = \tilde{u} - u_m \quad \tilde{v}_d = \tilde{v} - v_m$$

Newtonian Flow

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

Newtonian Stress Tensor

$$\sigma_{xz} = \mu u_z \quad \sigma_{yz} = \mu v_z$$

Kinematic Viscosity

$$\nu = \frac{\mu}{\rho}$$

Mapped stress tensor

$$\frac{1}{\rho} \tilde{\sigma}_{xz} = \frac{\nu}{h} \tilde{u}_\zeta \quad \frac{1}{\rho} \tilde{\sigma}_{yz} = \frac{\nu}{h} \tilde{v}_\zeta$$

Boundary Conditions

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

Stress Free Condition at surface

$$u_z|_{z=h_s} = v_z|_{z=h_s} = 0$$

Mixed Slip Condition at bottom topography

$$u - \frac{\lambda}{\mu} \sigma_{xz} \Big|_{z=h_b} = v - \frac{\lambda}{\mu} \sigma_{yz} \Big|_{z=h_b} = 0$$

Mapped with Newtonian Stress

$$\tilde{u}_\zeta|_{\zeta=1} = \tilde{v}_\zeta|_{\zeta=1} = 0$$

and

$$\tilde{u} - \frac{\lambda}{h} \tilde{u}_\zeta \Big|_{\zeta=0} = \tilde{v} - \frac{\lambda}{h} \tilde{v}_\zeta \Big|_{\zeta=0} = 0$$

Moment Closure

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

Depth Averaged Momentum Equations

$$\begin{aligned} (hu_m)_t + \left(h \int_0^1 \tilde{u}^2 d\zeta + \frac{1}{2} g e_z h^2 \right)_x + \left(h \int_0^1 \tilde{u} \tilde{v} d\zeta \right)_y \\ + \frac{\nu}{\lambda} u|_{\zeta=0} = hg(e_x - e_z(h_b)_x) \\ (hv_m)_t + \left(h \int_0^1 \tilde{u} \tilde{v} d\zeta \right)_y + \left(h \int_0^1 \tilde{v}^2 d\zeta + \frac{1}{2} g e_z h^2 \right)_x \\ + \frac{\nu}{\lambda} v|_{\zeta=0} = hg(e_y - e_z(h_b)_y) \end{aligned}$$

Polynomial Ansatz

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

$$\tilde{u}(t, x, y, \zeta) = u_m(t, x, y) + u_d(t, x, y, \zeta)$$

$$= u_m(t, x, y) + \sum_{j=1}^N (\alpha_j(t, x, y) \phi_j(\zeta))$$

$$\tilde{v}(t, x, y, \zeta) = v_m(t, x, y) + v_d(t, x, y, \zeta)$$

$$= v_m(t, x, y) + \sum_{j=1}^N (\beta_j(t, x, y) \phi_j(\zeta))$$

Orthogonality Condition

$$\int_0^1 \phi_j(\zeta) \phi_i(\zeta) d\zeta = 0 \quad \text{for } j \neq i$$

$$\phi_0(\zeta) = 1, \quad \phi_1(\zeta) = 1 - 2\zeta, \quad \phi_2(\zeta) = 1 - 6\zeta + 6\zeta^2$$

Constant Moments

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

$$\begin{aligned}
 & (hu_m)_t + \left(h \left(u_m^2 + \sum_{j=1}^N \frac{\alpha_j^2}{2j+1} \right) + \frac{1}{2} g e_z h^2 \right)_x \\
 & + \left(h \left(u_m v_m + \sum_{j=1}^N \frac{\alpha_j \beta_j}{2j+1} \right) \right)_y = -\frac{\nu}{\lambda} \left(u_m + \sum_{j=1}^N \alpha_j \right) + hg (e_x - e_z (h_b)_x) \\
 & (hv_m)_t + \left(h \left(v_m^2 + \sum_{j=1}^N \frac{\alpha_j \beta_j}{2j+1} \right) + \frac{1}{2} g e_z h^2 \right)_y \\
 & + \left(h \left(u_m v_m + \sum_{j=1}^N \frac{\alpha_j \beta_j}{2j+1} \right) \right)_x = -\frac{\nu}{\lambda} \left(v_m + \sum_{j=1}^N \beta_j \right) + hg (e_y - e_z (h_b)_y)
 \end{aligned}$$

Higher Order Moments

Moment Equation

$$\int_0^1 \phi_i \left((h\tilde{u})_t + \left(h\tilde{u}^2 + \frac{1}{2} g e_z h^2 \right)_x + (h\tilde{u}\tilde{v})_y + \left(h\tilde{u}\omega - \frac{1}{\rho} \tilde{\sigma}_{xz} \right)_z \right) d\zeta$$

$$= \int_0^1 \phi_i (gh(e_x - e_z(h_b)_x)) d\zeta$$

Simplified gives

$$(h\alpha_i)_t + \left(2hu_m\alpha_i + h \sum_{j,k=1}^N A_{ijk}\alpha_j\alpha_k \right)_x$$

$$+ \left(hu_m\beta_i + hv_m\alpha_i + h \sum_{j,k=1}^N A_{ijk}\alpha_j\beta_k \right)_y$$

$$= u_m D_i - \sum_{j,k=1}^N B_{ijk} D_j \alpha_k - (2i+1) \frac{\nu}{\lambda} \left(u_m + \sum_{j=1}^N \left(1 + \frac{\lambda}{h} C_{ij} \right) \alpha_j \right)$$

Higher Order Moments

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

$$\begin{aligned}
 (h\beta_i)_t + \left(hu_m\beta_i + hv_m\alpha_i + h \sum_{j,k=1}^N A_{ijk}\alpha_j\beta_k \right)_x + \left(2hv_m\beta_i + h \sum_{j,k=1}^N A_{ijk}\beta_j\beta_k \right)_y \\
 = v_m D_i - \sum_{j,k=1}^N B_{ijk} D_j \beta_k - (2i+1) \frac{\nu}{\lambda} \left(v_m + \sum_{j=1}^N \left(1 + \frac{\lambda}{h} C_{ij} \right) \beta_j \right)
 \end{aligned}$$

$$A_{ijk} = (2i+1) \int_0^1 \phi_i \phi_j \phi_k \, d\zeta$$

$$B_{ijk} = (2i+1) \int_0^1 \phi'_i \left(\int_0^\zeta \phi_j \, d\hat{\zeta} \right) \phi_k \, d\zeta$$

$$C_{ij} = \int_0^1 \phi'_i \phi'_j \, d\zeta$$

$$D_i = (h\alpha_i)_x + (h\beta_i)_y$$

Example Systems

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

1D model with h_b constant, $e_x = e_y = 0$, and $e_z = 1$
Constant System

$$\begin{bmatrix} h \\ hu_m \end{bmatrix}_t + \begin{bmatrix} hu_m \\ hu_m^2 + \frac{1}{2}gh^2 \end{bmatrix} = -\frac{\nu}{\lambda} \begin{bmatrix} 0 \\ u_m \end{bmatrix}$$

Linear System, $s = \alpha_1$

$$\begin{bmatrix} h \\ hu_m \\ hs \end{bmatrix}_t + \begin{bmatrix} hu_m \\ hu_m^2 + \frac{1}{2}gh^2 + \frac{1}{3}hs^2 \\ 2hu_ms \end{bmatrix} = Q \begin{bmatrix} h \\ hu_m \\ hs \end{bmatrix}_x - P$$

$$Q = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & u_m \end{bmatrix} \quad P = \frac{\nu}{\lambda} \begin{bmatrix} 0 \\ u_m + s \\ 3(u_m + s + 4\frac{\lambda}{h}s) \end{bmatrix}$$

Nonconservative Flux

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

$$\int Q \mathbf{q}_x \phi \, dx$$

$$\begin{bmatrix} Q_x^1 \\ Q_x^2 \\ Q_x^3 \\ Q_x^4 \\ Q_x^5 \end{bmatrix} = \frac{1}{2\Delta x} \begin{bmatrix} \Delta Q^1 - 2\sqrt{5}\Delta Q^3 + 78\Delta Q^5 \\ \Delta Q^2 - \frac{10}{3}\sqrt{3}\sqrt{7}\Delta Q^4 \\ \Delta Q^3 - 14\sqrt{5}\Delta Q^5 \\ \Delta Q^4 \\ \Delta Q^5 \end{bmatrix}$$

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

Higer Momemt Equations

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

Quadratic Vertical Profile

$$\begin{bmatrix} h \\ hu \\ hs \\ h\kappa \end{bmatrix}_t + \begin{bmatrix} hu \\ hu^2 + \frac{1}{2}gh^2 + \frac{1}{3}hs^2 + \frac{1}{5}h\kappa^2 \\ 2hus + \frac{4}{5}hs\kappa \\ 2hu\kappa + \frac{2}{3}hs^2 + \frac{2}{7}h\kappa^2 \end{bmatrix}_x = Q \begin{bmatrix} h \\ hu \\ hs \\ h\kappa \end{bmatrix}_x - P$$

Bibliography I

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods
Results

Generalized
Shallow Water
Equations

Model

Numerical Methods
Results

Future Work

References

- [1] Andrea L Bertozzi, Andreas Münch, and Michael Shearer. “Undercompressive shocks in thin film flows”. In: *Physica D: Nonlinear Phenomena* 134.4 (1999), pp. 431–464.
- [2] Y. Ha, Y.-J. Kim, and T.G. Myers. “On the numerical solution of a driven thin film equation”. In: *J. Comp. Phys.* 227.15 (2008), pp. 7246–7263.
- [3] T.G. Myers and J.P.F. Charpin. “A mathematical model for atmospheric ice accretion and water flow on a cold surface”. In: *Int. J. Heat and Mass Transfer* 47.25 (2004), pp. 5483–5500.
- [4] Tim G Myers. “Thin films with high surface tension”. In: *SIAM review* 40.3 (1998), pp. 441–462.
- [5] NASA. URL: http://icebox.grc.nasa.gov/gallery/images/C95_03918.html.

Bibliography II

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

- [6] Alexander Oron, Stephen H Davis, and S George Bankoff. “Long-scale evolution of thin liquid films”. In: *Reviews of modern physics* 69.3 (1997), p. 931.
- [7] J.A. Rossmanith. DOGPack. Available from <http://www.dogpack-code.org/>.