

Caleb Logemann,  
James  
Rossmanith

Thin Film  
Equation

Model

Numerical Methods

Results

Generalized  
Shallow Water  
Equations

Model

Numerical Methods

Results

Future Work

References

# Discontinuous Galerkin Method for Solving Thin Film and Shallow Water Equations

Caleb Logemann    James Rossmanith

Mathematics Department,  
Iowa State University

*logemann@iastate.edu*

May 13, 2019

# Overview

Caleb Logemann,  
James  
Rossmanith

Thin Film  
Equation

Model

Numerical Methods

Results

Generalized  
Shallow Water  
Equations

Model

Numerical Methods

Results

Future Work

References

## 1 Thin Film Equation

- Model
- Numerical Methods
- Results

## 2 Generalized Shallow Water Equations

- Model
- Numerical Methods
- Results
- Future Work

# Model Equations

Caleb Logemann,  
James  
Rossmanith

Thin Film  
Equation

Model

Numerical Methods  
Results


Generalized  
Shallow Water  
Equations

Model

Numerical Methods  
Results

Future Work

References



## ■ Incompressible Navier-Stokes Equation

$$u_x + w_z = 0$$


$$\rho(u_t + uu_x + ww_z) = -p_x + \mu\Delta u - \phi_x$$

$$\rho(w_t + uw_x + ww_z) = -p_z + \mu\Delta w - \phi_z$$

$$w = 0, u = 0 \quad \text{at } z = 0$$

$$w = h_t + uh_x \quad \text{at } z = h$$

$$\mathbf{T} \cdot \mathbf{n} = (-\kappa\sigma + \Pi)\mathbf{n} + \left(\frac{\partial\sigma}{\partial s} + \tau\right)\mathbf{t} \quad \text{at } z = h$$



Nondimensionalize, integrate over  $Z$ , and simplify, gives

$$H_T + \left( \frac{1}{2}(\tau + \Sigma_x)H^2 - \frac{1}{3}(\Phi|_{Z=H} - \Pi)_x H^3 \right)_x = -\frac{1}{3}\bar{C}^{-1}(H^3 H_{xxx})_x$$

$$q_t + (q^2 - q^3)_x = -(q^3 q_{xxx})_x$$

# Method Overview

Caleb Logemann,  
James  
Rossmanith

Thin Film  
Equation

Model

Numerical Methods

Results

Generalized  
Shallow Water  
Equations

Model

Numerical Methods

Results

Future Work

References

## ■ Simplified Model

$$q_t + (q^2 - q^3)_x = -(q^3 q_{xxx})_x \quad (0, T) \times \Omega$$

## ■ Runge Kutta Implicit Explicit (IMEX)

$$q_t = F(q) + G(q)$$

■ F evaluated explicitly

■ G solved implicitly

$$F(q) = -(q^2 - q^3)_x$$

$$G(q) = (q^3 q_{xxx})_x$$

# Convection

Caleb Logemann,  
James  
Rossmanith

Thin Film  
Equation

Model

Numerical Methods

Results

Generalized  
Shallow Water  
Equations

Model

Numerical Methods

Results

Future Work

References

## ■ Convection Equation

$$\begin{aligned} F(q) &= f(q)_x = 0 & (0, T) \times \Omega \\ f(q) &= q^2 - q^3 \end{aligned}$$

## ■ Weak Form

Find  $q$  such that

$$\int_{\Omega} (F(q)v - f(q)v_x) \, dx + \hat{f}v \Big|_{\partial\Omega} = 0$$

for all test functions  $v$

# Notation

Caleb Logemann,  
James  
Rossmanith

Thin Film  
Equation

Model

Numerical Methods

Results

Generalized  
Shallow Water  
Equations

Model

Numerical Methods

Results

Future Work


References

- Partition the domain,  $[a, b]$  as

$$a = x_{1/2} < \cdots < x_{j-1/2} < x_{j+1/2} < \cdots < x_{N+1/2} = b$$

- $I_j = [x_{j-1/2}, x_{j+1/2}]$

- $x_j = \frac{x_{j+1/2} + x_{j-1/2}}{2}$ .



# Runge Kutta Discontinuous Galerkin

Caleb Logemann,  
James  
Rossmanith

Thin Film  
Equation

Model

Numerical Methods

Results

Generalized  
Shallow Water  
Equations

Model

Numerical Methods

Results

Future Work

References

- Find  $Q(t, x)$  such that for each time  $t \in (0, T)$ ,  
 $Q(t, \cdot) \in V_h = \left\{ v \in L^1(\Omega) : v|_{I_j} \in P^k(I_j) \right\}$

$$\begin{aligned} \int_{I_j} F(Q) v \, dx &= \int_{I_j} f(Q) v_x \, dx \\ &\quad - \left( \mathcal{F}_{j+1/2} v^-(x_{j+1/2}) - \mathcal{F}_{j-1/2} v^+(x_{j-1/2}) \right) \end{aligned}$$

for all  $v \in V_h$

- Rusanov/Local Lax-Friedrichs Numerical Flux

$$\mathcal{F}_{j+1/2} = \frac{1}{2} \left( f(Q_{j+1/2}^-) + f(Q_{j+1/2}^+) \right) + \frac{1}{2} \max_q \{ |f'(q)| \} (Q_{j+1/2}^- - Q_{j+1/2}^+)$$



# Diffusion

Caleb Logemann,  
James  
Rossmanith

Thin Film  
Equation

Model

Numerical Methods

Results

Generalized  
Shallow Water  
Equations

Model

Numerical Methods

Results

Future Work

References

## ■ Diffusion Equation

$$G(q) = -(q^3 q_{xxx})_x \quad (0, T) \times \Omega$$

## ■ Local Discontinuous Galerkin

$$r = q_x$$

$$s = r_x$$

$$u = s_x$$

$$G(q) = (q^3 u)_x$$

# Local Discontinuous Galerkin

Caleb Logemann,  
James  
Rossmanith

Thin Film  
Equation

Model

Numerical Methods

Results

Generalized  
Shallow Water  
Equations

Model

Numerical Methods

Results

Future Work

References

Find  $Q(t, x), R(x), S(x), U(x)$  such that for all  $t \in (0, T)$

$$Q(t, \cdot), R, S, U \in V_h = \left\{ v \in L^1(\Omega) : v|_{I_j} \in P^k(I_j) \right\}$$

$$\int_{I_j} Rv \, dx = - \int_{I_j} Qv_x \, dx + \left( \hat{Q}_{j+1/2} v_{j+1/2}^- - \hat{Q}_{j-1/2} v_{j-1/2}^+ \right)$$

$$\int_{I_j} Sw \, dx = - \int_{I_j} R w_x \, dx + \left( \hat{R}_{j+1/2} w_{j+1/2}^- - \hat{R}_{j-1/2} w_{j-1/2}^+ \right)$$

$$\int_{I_j} Uy \, dx = - \int_{I_j} S y_x \, dx + \left( \hat{S}_{j+1/2} y_{j+1/2}^- - \hat{S}_{j-1/2} y_{j-1/2}^+ \right)$$

$$\int_{I_j} G(Q)z \, dx = - \int_{I_j} Q^3 U z_x \, dx + \left( \hat{U}_{j+1/2} z_{j+1/2}^- - \hat{U}_{j-1/2} z_{j-1/2}^+ \right)$$

for all  $I_j \in \Omega$  and all  $v, w, y, z \in V_h$ .

# Numerical Fluxes

Caleb Logemann,  
James  
Rossmanith

Thin Film  
Equation

Model

Numerical Methods

Results

Generalized  
Shallow Water  
Equations

Model

Numerical Methods

Results

Future Work


References

$$\hat{Q}_{j+1/2} = Q_{j+1/2}^+$$

$$\hat{R}_{j+1/2} = R_{j+1/2}^-$$

$$\hat{S}_{j+1/2} = S_{j+1/2}^+$$

$$\hat{U}_{j+1/2} = (Q^3 U)_{j+1/2}^-$$



# IMEX Runge Kutta

Caleb Logemann,  
James  
Rossmanith

Thin Film  
Equation

Model

Numerical Methods

Results

Generalized  
Shallow Water  
Equations

Model

Numerical Methods

Results

Future Work

References

## ■ IMEX scheme

$$\begin{aligned}q^{n+1} &= q^n + \Delta t \sum_{i=1}^s (b'_i F(t_i, u_i)) + \Delta t \sum_{i=1}^s (b_i G(t_i, u_i)) \\u_i &= q^n + \Delta t \sum_{j=1}^{i-1} (a'_{ij} F(t_j, u_j)) + \Delta t \sum_{j=1}^i (a_{ij} G(t_j, u_j)) \\t_i &= t^n + c_i \Delta t\end{aligned}$$

## ■ Double Butcher Tableaus

$$\begin{array}{c|c} c' & a' \\ \hline & b'^T \end{array} \quad \begin{array}{c|c} c & a \\ \hline & b^T \end{array}$$

## ■ 1st Order — L-Stable SSP

$$\begin{array}{c|c} 0 & 0 \\ \hline & 1 \end{array} \quad \begin{array}{c|c} 1 & 1 \\ \hline & 1 \end{array}$$

## ■ 2nd Order — SSP

$$\begin{array}{c|ccc} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ \hline & 0 & \frac{1}{2} & \frac{1}{2} \end{array} \quad \begin{array}{c|cc} \frac{1}{2} & \frac{1}{2} & 0 \\ 0 & -\frac{1}{2} & 0 \\ 1 & 0 & \frac{1}{2} \\ \hline & 0 & \frac{1}{2} \end{array}$$

## ■ 3rd Order — L-Stable SSP

0	0	0	0	0	$\alpha$	$\alpha$	0	0	0
0	0	0	0	0	0	$-\alpha$	$\alpha$	0	0
1	0	1	0	0	1	0	$1 - \alpha$	$\alpha$	0
$\frac{1}{2}$	0	$\frac{1}{4}$	$\frac{1}{4}$	0	$\frac{1}{2}$	$\beta$	$\eta$	$\zeta$	$\alpha$
<hr/>					<hr/>				
	0	$\frac{1}{6}$	$\frac{1}{6}$	$\frac{2}{3}$		0	$\frac{1}{6}$	$\frac{1}{6}$	$\frac{2}{3}$

$$\alpha = 0.24169426078821$$

$$\beta = 0.06042356519705$$

$$\eta = 0.1291528696059$$

$$\zeta = \frac{1}{2} - \beta - \eta - \alpha$$

# Nonlinear Solvers

Caleb Logemann,  
James  
Rossmanith

Thin Film  
Equation

Model

Numerical Methods

Results

Generalized  
Shallow Water  
Equations

Model

Numerical Methods

Results

Future Work

References

## ■ Nonlinear System

$$u_i - a_{ij} \Delta t G(u_i) = b$$

## ■ Picard Iteration

$$\tilde{G}(q, u) = (q^3 u_{xxx})_x$$

$$u_0 = q^n \quad u_i^0 = u_{i-1}$$
$$u_i^j - a_{ij} \Delta t \tilde{G}(u_i^{j-1}, u_i^j) = b$$

- Number of picard iterations equals order in time and space

# Manufactured Solution

$$q_t + (q^2 - q^3)_x = -(q^3 q_{xxx})_x + s$$

$$s = \hat{q}_t + (\hat{q}^2 - \hat{q}^3)_x + (\hat{q}^3 \hat{q}_{xxx})_x$$

$$\hat{q} = 0.1 \times \sin(2\pi/20.0 \times (x - t)) + 0.15 \quad \text{for } (x, t) \in [0, 40] \times [0, 5.0]$$

	1st Order		2nd Order		3rd Order	
$n$	error	order	error	order	error	order
20	0.136	—	$7.33 \times 10^{-3}$	—	$5.29 \times 10^{-4}$	—
40	0.0719	0.92	$1.99 \times 10^{-3}$	1.88	$5.38 \times 10^{-5}$	3.30
80	0.0378	0.93	$5.60 \times 10^{-4}$	1.83	$7.47 \times 10^{-6}$	2.85
160	0.0191	0.99	$1.56 \times 10^{-4}$	1.85	$9.97 \times 10^{-7}$	2.91
320	0.00961	0.99	$3.98 \times 10^{-5}$	1.97	$1.26 \times 10^{-7}$	2.98
640	0.00483	0.99	$1.00 \times 10^{-5}$	1.99	$1.58 \times 10^{-8}$	3.00
1280	0.00242	1.00	$2.50 \times 10^{-6}$	2.00	$1.98 \times 10^{-9}$	3.00

**Table:** Convergence table with a constant, linear, quadratic polynomial bases. CFL = 0.9, 0.2, 0.1 respectively.



# Wave Structure with Nonlinear Hyper Diffusion

Caleb Logemann,  
James  
Rossmanith

Thin Film  
Equation

Model

Numerical Methods

Results

Generalized  
Shallow Water  
Equations

Model


Numerical Methods

Results

Future Work

References

$$q_t + (q^2 - q^3)_x = -(q^3 q_{xxx})_x$$
$$q_r = 0.1 \quad q_l = 0.3$$



# Wave Structure with Nonlinear Hyper Diffusion

Caleb Logemann,  
James  
Rossmanith

Thin Film  
Equation

Model

Numerical Methods

Results

Generalized  
Shallow Water  
Equations

Model

Numerical Methods


Results

Future Work

References

$$q_r = 0.1 \quad q_l = 0.3323$$

$$q(x, 0) = (-\tanh(x) + 1) \frac{q_l - q_r}{2} + q_r$$



# Wave Structure with Nonlinear Hyper Diffusion

Caleb Logemann,  
James  
Rossmanith


Thin Film  
Equation  
Model  
Numerical Methods  
Results

Generalized  
Shallow Water  
Equations

Model  
Numerical Methods  
Results  
Future Work  
References

$$q_r = 0.1 \quad q_l = 0.3323 \quad q_m = 0.6$$

$$q(x, 0) = \begin{cases} \frac{q_m - q_l}{2} \tanh(x) + \frac{q_m + q_l}{2} & x < 5 \\ -\frac{q_m - q_r}{2} \tanh(x - 10) + \frac{q_m + q_r}{2} + q_r & x > 5 \end{cases}$$



# Wave Structure with Nonlinear Hyper Diffusion

Caleb Logemann,  
James  
Rossmanith


Thin Film  
Equation  
Model  
Numerical Methods  
Results

Generalized  
Shallow Water  
Equations

Model  
Numerical Methods  
Results  
Future Work  
References

$$q_r = 0.1 \quad q_l = 0.3323 \quad q_m = 0.6$$

$$q(x, 0) = \begin{cases} \frac{q_m - q_l}{2} \tanh(x) + \frac{q_m + q_l}{2} & x < 10 \\ -\frac{q_m - q_r}{2} \tanh(x - 20) + \frac{q_m + q_r}{2} + q_r & x > 10 \end{cases}$$



# Wave Structure with Nonlinear Hyper Diffusion

Caleb Logemann,  
James  
Rossmanith

Thin Film  
Equation

Model

Numerical Methods

Results

Generalized  
Shallow Water  
Equations

Model

Numerical Methods


Results

Future Work

References

$$q_r = 0.1 \quad q_l = 0.4$$

$$q(x, 0) = (-\tanh(x - 100) + 1) \frac{q_l - q_r}{2} + q_r$$



# Wave Structure with Nonlinear Hyper Diffusion

Caleb Logemann,  
James  
Rossmanith

Thin Film  
Equation

Model

Numerical Methods

Results

Generalized  
Shallow Water  
Equations

Model

Numerical Methods


Results

Future Work

References

$$q_r = 0.1 \quad q_l = 0.8$$

$$q(x, 0) = (-\tanh(x - 10) + 1) \frac{q_l - q_r}{2} + q_r$$



# Generalized Shallow Water

Caleb Logemann,  
James  
Rossmanith

Thin Film  
Equation

Model

Numerical Methods

Results

Generalized  
Shallow Water  
Equations


Model

Numerical Methods

Results

Future Work

References



$$\nabla \cdot \mathbf{u} = 0$$

$$\mathbf{u}_t + \nabla \cdot (\mathbf{u}\mathbf{u}) = -\frac{1}{\rho}\nabla p + \frac{1}{\rho}\nabla \cdot \boldsymbol{\sigma} + \mathbf{g}$$

$$(h_s)_t + [u(t, x, y, h_s), v(t, x, y, h_s)]^T \cdot \nabla h_s = w(t, x, y, h_s)$$

$$(h_b)_t + [u(t, x, y, h_b), v(t, x, y, h_b)]^T \cdot \nabla h_b = w(t, x, y, h_b)$$

# Nondimensionalization

Caleb Logemann,  
James  
Rossmanith

Thin Film  
Equation

Model

Numerical Methods

Results

Generalized  
Shallow Water  
Equations

Model

Numerical Methods

Results

Future Work

References

## Characteristic Lengths

$$\varepsilon = \frac{H}{L}, \quad x = L\hat{x}, \quad y = L\hat{y}, \quad z = H\hat{z}$$

## Characteristic Velocities

$$u = U\hat{u}, \quad v = U\hat{v}, \quad w = \varepsilon U\hat{w}$$

## Characteristic Time

$$t = \frac{L}{U}\hat{t}$$

## Characteristic Stresses

$$p = \rho g H \hat{p}, \quad \sigma_{xz/yz} = S \hat{\sigma}_{xz/yz}, \quad \sigma_{xx/xy/yy/zz} = \varepsilon S \hat{\sigma}_{xx/xy/yy/zz}$$



# Nondimensionalization

Caleb Logemann,  
James  
Rossmanith

Thin Film  
Equation

Model

Numerical Methods

Results

Generalized  
Shallow Water  
Equations

Model

Numerical Methods

Results

Future Work

References

$$\hat{u}_{\hat{x}} + \hat{v}_{\hat{y}} + \hat{w}_{\hat{z}} = 0$$

$$\begin{aligned} \varepsilon F^2 \left( \hat{u}_{\hat{t}} + (\hat{u}^2)_{\hat{x}} + (\hat{u}\hat{v})_{\hat{y}} + (\hat{u}\hat{w})_{\hat{z}} \right) &= -\varepsilon \hat{p}_{\hat{x}} \\ &+ G \left( \varepsilon^2 (\hat{\sigma}_{xx})_{\hat{x}} + \varepsilon^2 (\hat{\sigma}_{xy})_{\hat{y}} + (\hat{\sigma}_{xz})_{\hat{z}} \right) + e_x \end{aligned}$$

$$\begin{aligned} \varepsilon F^2 \left( \hat{v}_{\hat{t}} + (\hat{u}\hat{v})_{\hat{x}} + (\hat{v}^2)_{\hat{y}} + (\hat{v}\hat{w})_{\hat{z}} \right) &= -\varepsilon \hat{p}_{\hat{y}} \\ &+ G \left( \varepsilon^2 (\hat{\sigma}_{xy})_{\hat{x}} + \varepsilon^2 (\hat{\sigma}_{yy})_{\hat{y}} + (\hat{\sigma}_{yz})_{\hat{z}} \right) + e_y \end{aligned}$$

$$\begin{aligned} \varepsilon^2 F^2 \left( \hat{w}_{\hat{t}} + (\hat{u}\hat{w})_{\hat{x}} + (\hat{v}\hat{w})_{\hat{y}} + (\hat{w}^2)_{\hat{z}} \right) &= -\hat{p}_{\hat{z}} \\ &+ \varepsilon G \left( (\hat{\sigma}_{xz})_{\hat{x}} + (\hat{\sigma}_{yz})_{\hat{y}} + (\hat{\sigma}_{zz})_{\hat{z}} \right) + e_z \end{aligned}$$

$$F = \frac{U}{\sqrt{gH}} \approx 1, \quad G = \frac{S}{\rho g H} < 1$$

# Nondimensionalization

Caleb Logemann,  
James  
Rossmanith

Thin Film  
Equation

Model

Numerical Methods

Results

Generalized  
Shallow Water  
Equations

Model

Numerical Methods

Results

Future Work

References

Drop  $\varepsilon^2$  and  $\varepsilon G$  terms

$$\hat{u}_{\hat{x}} + \hat{v}_{\hat{y}} + \hat{w}_{\hat{z}} = 0$$

$$\varepsilon F^2 \left( \hat{u}_{\hat{t}} + (\hat{u}^2)_{\hat{x}} + (\hat{u}\hat{v})_{\hat{y}} + (\hat{u}\hat{w})_{\hat{z}} \right) = -\varepsilon \hat{p}_{\hat{x}} + G(\hat{\sigma}_{xz})_{\hat{z}} + e_x$$

$$\varepsilon F^2 \left( \hat{v}_{\hat{t}} + (\hat{u}\hat{v})_{\hat{x}} + (\hat{v}^2)_{\hat{y}} + (\hat{v}\hat{w})_{\hat{z}} \right) = -\varepsilon \hat{p}_{\hat{y}} + G(\hat{\sigma}_{yz})_{\hat{z}} + e_y$$

$$\hat{p}_{\hat{z}} = e_z$$

Solving for the hydrostatic pressure

$$\hat{p}(\hat{t}, \hat{x}, \hat{y}) = \left( \hat{h}_s(\hat{t}, \hat{x}, \hat{y}) - \hat{z} \right) e_z$$

# Nondimensionalization

Caleb Logemann,  
James  
Rossmanith

Thin Film  
Equation

Model

Numerical Methods

Results

Generalized  
Shallow Water  
Equations

Model

Numerical Methods

Results

Future Work

References

## Dimensional Variables

$$u_x + v_y + w_z = 0$$

$$u_t + (u^2)_x + (uv)_y + (uw)_z = -\frac{1}{\rho}p_x + \frac{1}{\rho}(\sigma_{xz})_z + ge_x$$

$$v_t + (uv)_x + (v^2)_y + (vw)_z = -\frac{1}{\rho}p_y + \frac{1}{\rho}(\sigma_{yz})_z + ge_y$$

$$p(t, x, y, z) = (h_s(t, x, y) - z)\rho ge_z$$

## Kinematic Boundary Conditions

$$(h_s)_t + [u(t, x, y, h_s), v(t, x, y, h_s)]^T \cdot \nabla h_s = w(t, x, y, h_s)$$

$$(h_b)_t + [u(t, x, y, h_b), v(t, x, y, h_b)]^T \cdot \nabla h_b = w(t, x, y, h_b)$$

# Mapping

Caleb Logemann,  
James  
Rossmanith

Thin Film  
Equation

Model

Numerical Methods

Results

Generalized  
Shallow Water  
Equations



Model

Numerical Methods

Results

Future Work

References



Transform from  $z \rightarrow \zeta$ , by

$$\zeta = \frac{z - h_b(t, x, y)}{h(t, x, y)},$$

or equivalently

$$z = h(t, x, y)\zeta + h_b(t, x, y)$$

where  $h(t, x, y) = h_s(t, x, y) - h_b(t, x, y)$ .

$$\tilde{\Psi}(t, x, y, \zeta) = \Psi(t, x, y, h(t, x, y)\zeta + h_b(t, x, y))$$

# Mapping Continuity Equation

Caleb Logemann,  
James  
Rossmanith

Thin Film  
Equation

Model

Numerical Methods

Results

Generalized  
Shallow Water  
Equations

Model

Numerical Methods

Results

Future Work

References

$$u_x + v_y + w_z = 0$$

Map to new space

$$(h\tilde{u})_x - ((\zeta h + h_b)_x \tilde{u})_\zeta + (h\tilde{v})_y - ((\zeta h + h_b)_y \tilde{v})_\zeta + (\tilde{w})_\zeta = 0$$

Solve for vertical velocity,  $w$ ,

$$\begin{aligned} \tilde{w}(t, x, y, \zeta') = & - \left( h \int_0^{\zeta'} \tilde{u} d\zeta \right)_x - \left( h \int_0^{\zeta'} \tilde{v} d\zeta \right)_y \\ & + (\zeta' h + h_b)_x \tilde{u}(t, x, y, \zeta') + (\zeta' h + h_b)_y \tilde{v}(t, x, y, \zeta') \end{aligned}$$

Depth averaged equation

$$h_t + \left( h \int_0^1 \tilde{u} d\zeta \right)_x + \left( h \int_0^1 \tilde{v} d\zeta \right)_y = 0$$

Let  $u_m$  and  $v_m$  denote the mean velocity

$$h_t + (hu_m)_x + (hv_m)_y = 0$$

# Mapping Momentum Equations

Caleb Logemann,  
James  
Rossmanith

Thin Film  
Equation

Model

Numerical Methods

Results

Generalized  
Shallow Water  
Equations

Model

Numerical Methods

Results

Future Work

References

$$u_t + (u^2)_x + (uv)_y + (uw)_z = -\frac{1}{\rho}p_x + \frac{1}{\rho}(\sigma_{xz})_z + ge_x$$

$$v_t + (uv)_x + (v^2)_y + (vw)_z = -\frac{1}{\rho}p_y + \frac{1}{\rho}(\sigma_{yz})_z + ge_y$$

$$(h\tilde{u})_t + \left(h\tilde{u}^2 + \frac{1}{2}ge_z h^2\right)_x + (h\tilde{u}\tilde{v})_y + \left(h\tilde{u}\omega - \frac{1}{\rho}\tilde{\sigma}_{xz}\right)_\zeta = gh(e_x - e_z(h_b)_x)$$

$$(h\tilde{v})_t + (h\tilde{u}\tilde{v})_x + \left(h\tilde{v}^2 + \frac{1}{2}ge_z h^2\right)_y + \left(h\tilde{v}\omega - \frac{1}{\rho}\tilde{\sigma}_{yz}\right)_\zeta = gh(e_x - e_z(h_b)_y)$$

where

$$\omega = \frac{1}{h} \left( - \left( h \int_0^\zeta \tilde{u} - u_m d\zeta' \right)_x - \left( h \int_0^\zeta \tilde{v} - v_m d\zeta' \right)_y \right)$$

# Mapped Reference System

Caleb Logemann,  
James  
Rossmanith

Thin Film  
Equation

Model

Numerical Methods

Results

Generalized  
Shallow Water  
Equations

Model

Numerical Methods

Results

Future Work

References

$$h_t + (hu_m)_x + (hv_m)_y = 0$$

$$(h\tilde{u})_t + \left(h\tilde{u}^2 + \frac{1}{2}ge_z h^2\right)_x + (h\tilde{u}\tilde{v})_y + \left(h\tilde{u}\omega - \frac{1}{\rho}\tilde{\sigma}_{xz}\right)_\zeta = gh(e_x - e_z(h_b)_x)$$

$$(h\tilde{v})_t + (h\tilde{u}\tilde{v})_x + \left(h\tilde{v}^2 + \frac{1}{2}ge_z h^2\right)_y + \left(h\tilde{v}\omega - \frac{1}{\rho}\tilde{\sigma}_{yz}\right)_\zeta = gh(e_x - e_z(h_b)_y)$$

$$\omega = \frac{1}{h} \left( - \left( h \int_0^\zeta \tilde{u}_d \, d\zeta' \right)_x - \left( h \int_0^\zeta \tilde{v}_d \, d\zeta' \right)_y \right)$$

with

$$\tilde{u}_d = \tilde{u} - u_m \quad \tilde{v}_d = \tilde{v} - v_m$$

# Newtonian Flow

Caleb Logemann,  
James  
Rossmanith

Thin Film  
Equation

Model

Numerical Methods

Results

Generalized  
Shallow Water  
Equations

Model

Numerical Methods

Results

Future Work

References

## Newtonian Stress Tensor

$$\sigma_{xz} = \mu u_z \quad \sigma_{yz} = \mu v_z$$

## Kinematic Viscosity

$$\nu = \frac{\mu}{\rho}$$

## Mapped stress tensor

$$\frac{1}{\rho} \tilde{\sigma}_{xz} = \frac{\nu}{h} \tilde{u}_\zeta \quad \frac{1}{\rho} \tilde{\sigma}_{yz} = \frac{\nu}{h} \tilde{v}_\zeta$$



# Boundary Conditions

Caleb Logemann,  
James  
Rossmanith

Thin Film  
Equation

Model

Numerical Methods

Results

Generalized  
Shallow Water  
Equations

Model

Numerical Methods

Results

Future Work

References

Stress Free Condition at surface

$$u_z|_{z=h_s} = v_z|_{z=h_s} = 0$$

Mixed Slip Condition at bottom topography

$$u - \frac{\lambda}{\mu} \sigma_{xz} \Big|_{z=h_b} = v - \frac{\lambda}{\mu} \sigma_{yz} \Big|_{z=h_b} = 0$$

Mapped with Newtonian Stress

$$\tilde{u}_\zeta|_{\zeta=1} = \tilde{v}_\zeta|_{\zeta=1} = 0$$

and

$$\tilde{u} - \frac{\lambda}{h} \tilde{u}_\zeta \Big|_{\zeta=0} = \tilde{v} - \frac{\lambda}{h} \tilde{v}_\zeta \Big|_{\zeta=0} = 0$$

# Moment Closure

Caleb Logemann,  
James  
Rossmanith

Thin Film  
Equation

Model

Numerical Methods

Results

Generalized  
Shallow Water  
Equations

Model

Numerical Methods

Results

Future Work

References

## Depth Average Momentum Equations

$$\begin{aligned} (hu_m)_t + \left( h \int_0^1 \tilde{u}^2 d\zeta + \frac{1}{2} g e_z h^2 \right)_x + \left( h \int_0^1 \tilde{u} \tilde{v} d\zeta \right)_y \\ + \frac{\nu}{\lambda} u|_{\zeta=0} = hg(e_x - e_z(h_b)_x) \\ (hv_m)_t + \left( h \int_0^1 \tilde{u} \tilde{v} d\zeta \right)_y + \left( h \int_0^1 \tilde{v}^2 d\zeta + \frac{1}{2} g e_z h^2 \right)_x \\ + \frac{\nu}{\lambda} v|_{\zeta=0} = hg(e_y - e_z(h_b)_y) \end{aligned}$$

# Polynomial Ansatz

Caleb Logemann,  
James  
Rossmanith

Thin Film  
Equation

Model

Numerical Methods

Results

Generalized  
Shallow Water  
Equations

Model

Numerical Methods

Results

Future Work

References

$$\tilde{u}(t, x, y, \zeta) = u_m(t, x, y) + u_d(t, x, y, \zeta)$$

$$= u_m(t, x, y) + \sum_{j=1}^N (\alpha_j(t, x, y) \phi_j(\zeta))$$

$$\tilde{v}(t, x, y, \zeta) = v_m(t, x, y) + v_d(t, x, y, \zeta)$$

$$= v_m(t, x, y) + \sum_{j=1}^N (\beta_j(t, x, y) \phi_j(\zeta))$$

Orthogonality Condition

$$\int_0^1 \phi_j(\zeta) \phi_i(\zeta) d\zeta = 0 \quad \text{for } j \neq i$$

$$\phi_0(\zeta) = 1, \quad \phi_1(\zeta) = 1 - 2\zeta, \quad \phi_2(\zeta) = 1 - 6\zeta + 6\zeta^2$$

Caleb Logemann,  
James  
Rossmanith

Thin Film  
Equation

Model

Numerical Methods

Results

Generalized  
Shallow Water  
Equations

**Model**

Numerical Methods

Results

Future Work

References

$(h)$

# Example Systems

Caleb Logemann,  
James  
Rossmanith

Thin Film  
Equation

Model

Numerical Methods

Results

Generalized  
Shallow Water  
Equations

Model

Numerical Methods

Results

Future Work

References

1D model with  $h_b$  constant,  $e_x = e_y = 0$ , and  $e_z = 1$   
Constant System

$$\begin{bmatrix} h \\ hu_m \end{bmatrix}_t + \begin{bmatrix} hu_m \\ hu_m^2 + \frac{1}{2}gh^2 \end{bmatrix} = -\frac{\nu}{\lambda} \begin{bmatrix} 0 \\ u_m \end{bmatrix}$$

Linear System,  $s = \alpha_1$

$$\begin{bmatrix} h \\ hu_m \\ hs \end{bmatrix}_t + \begin{bmatrix} hu_m \\ hu_m^2 + \frac{1}{2}gh^2 + \frac{1}{3}hs^2 \\ 2hu_ms \end{bmatrix} = Q \begin{bmatrix} h \\ hu_m \\ hs \end{bmatrix}_x - P$$

$$Q = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & u_m \end{bmatrix} \quad P = \frac{\nu}{\lambda} \begin{bmatrix} 0 \\ u_m + s \\ 3(u_m + s + 4\frac{\lambda}{h}s) \end{bmatrix}$$

# Nonconservative Flux

Caleb Logemann,  
James  
Rossmanith

Thin Film  
Equation

Model

Numerical Methods

Results

Generalized  
Shallow Water  
Equations

Model

Numerical Methods

Results

Future Work

References

$$\int Q \mathbf{q}_x \phi \, dx$$

$$\begin{bmatrix} Q_x^1 \\ Q_x^2 \\ Q_x^3 \\ Q_x^4 \\ Q_x^5 \end{bmatrix} = \frac{1}{2\Delta x} \begin{bmatrix} \Delta Q^1 - 2\sqrt{5}\Delta Q^3 + 78\Delta Q^5 \\ \Delta Q^2 - \frac{10}{3}\sqrt{3}\sqrt{7}\Delta Q^4 \\ \Delta Q^3 - 14\sqrt{5}\Delta Q^5 \\ \Delta Q^4 \\ \Delta Q^5 \end{bmatrix}$$

# Higer Momemt Equations

Caleb Logemann,  
James  
Rossmanith

Thin Film  
Equation

Model

Numerical Methods

Results

Generalized  
Shallow Water  
Equations

Model

Numerical Methods

Results

Future Work

References

## Quadratic Vertical Profile

$$\begin{bmatrix} h \\ hu \\ hs \\ h\kappa \end{bmatrix}_t + \begin{bmatrix} hu \\ hu^2 + \frac{1}{2}gh^2 + \frac{1}{3}hs^2 + \frac{1}{5}h\kappa^2 \\ 2hus + \frac{4}{5}hs\kappa \\ 2hu\kappa + \frac{2}{3}hs^2 + \frac{2}{7}h\kappa^2 \end{bmatrix}_x = Q \begin{bmatrix} h \\ hu \\ hs \\ h\kappa \end{bmatrix}_x - P$$

# Bibliography I

Caleb Logemann,  
James  
Rossmanith

Thin Film  
Equation

Model

Numerical Methods  
Results

Generalized  
Shallow Water  
Equations

Model

Numerical Methods  
Results

Future Work

References

- [1] Andrea L Bertozzi, Andreas Münch, and Michael Shearer. “Undercompressive shocks in thin film flows”. In: *Physica D: Nonlinear Phenomena* 134.4 (1999), pp. 431–464.
- [2] Y. Ha, Y.-J. Kim, and T.G. Myers. “On the numerical solution of a driven thin film equation”. In: *J. Comp. Phys.* 227.15 (2008), pp. 7246–7263.
- [3] T.G. Myers and J.P.F. Charpin. “A mathematical model for atmospheric ice accretion and water flow on a cold surface”. In: *Int. J. Heat and Mass Transfer* 47.25 (2004), pp. 5483–5500.
- [4] Tim G Myers. “Thin films with high surface tension”. In: *SIAM review* 40.3 (1998), pp. 441–462.
- [5] NASA. URL: [http://icebox.grc.nasa.gov/gallery/images/C95\\_03918.html](http://icebox.grc.nasa.gov/gallery/images/C95_03918.html).



# Bibliography II

Caleb Logemann,  
James  
Rossmanith

Thin Film  
Equation

Model

Numerical Methods  
Results

Generalized  
Shallow Water  
Equations

Model

Numerical Methods  
Results

Future Work

References

- [6] Alexander Oron, Stephen H Davis, and S George Bankoff. “Long-scale evolution of thin liquid films”. In: *Reviews of modern physics* 69.3 (1997), p. 931.
- [7] J.A. Rossmanith. DOGPACK. Available from <http://www.dogpack-code.org/>.