

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

Discontinuous Galerkin Method for Solving Thin Film and Shallow Water Equations

Caleb Logemann James Rossmanith

Mathematics Department,
Iowa State University

logemann@iastate.edu

January 14, 2019

Overview

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

1 Thin Film Equation

- Model
- Numerical Methods
- Results

2 Generalized Shallow Water Equations

- Model
- Numerical Methods
- Results
- Future Work

Model Equations

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods
Results


Generalized
Shallow Water
Equations

Model

Numerical Methods
Results

Future Work

References



■ Incompressible Navier-Stokes Equation

$$u_x + w_z = 0$$


$$\rho(u_t + uu_x + wu_z) = -p_x + \mu\Delta u - \phi_x$$

$$\rho(w_t + uw_x + ww_z) = -p_z + \mu\Delta w - \phi_z$$

$$w = 0, u = 0 \quad \text{at } z = 0$$

$$w = h_t + uh_x \quad \text{at } z = h$$

$$\mathbf{T} \cdot \mathbf{n} = (-\kappa\sigma + \Pi)\mathbf{n} + \left(\frac{\partial\sigma}{\partial s} + \tau\right)\mathbf{t} \quad \text{at } z = h$$



Nondimensionalize, integrate over Z , and simplify, gives

$$H_T + \left(\frac{1}{2}(\tau + \Sigma_x)H^2 - \frac{1}{3}(\Phi|_{Z=H} - \Pi)_x H^3 \right)_x = -\frac{1}{3}\bar{C}^{-1}(H^3 H_{xxx})_x$$

$$q_t + (q^2 - q^3)_x = -(q^3 q_{xxx})_x$$

Method Overview

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

■ Simplified Model

$$q_t + (q^2 - q^3)_x = -(q^3 q_{xxx})_x \quad (0, T) \times \Omega$$

■ Runge Kutta Implicit Explicit (IMEX)

$$q_t = F(q) + G(q)$$

■ F evaluated explicitly

■ G solved implicitly

$$F(q) = -(q^2 - q^3)_x$$


$$G(q) = (q^3 q_{xxx})_x$$

Notation

- Partition the domain, $[a, b]$ as

$$a = x_{1/2} < \cdots < x_{j-1/2} < x_{j+1/2} < \cdots < x_{N+1/2} = b$$

- $I_j = [x_{j-1/2}, x_{j+1/2}]$
- $x_j = \frac{x_{j+1/2} + x_{j-1/2}}{2}$
- $\Delta x_j = x_{j+1/2} - x_{j-1/2}$
- $\Delta x_j = \Delta x$ for all j .



Convection

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

■ Convection Equation

$$\begin{aligned} F(q) &= f(q)_x = 0 & (0, T) \times \Omega \\ f(q) &= q^2 - q^3 \end{aligned}$$

■ Weak Form

Find q such that

$$\int_{\Omega} (F(q)v - f(q)v_x) \, dx + \hat{f}v \Big|_{\partial\Omega} = 0$$

for all test functions v

Runge Kutta Discontinuous Galerkin

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

- Find $Q(t, x)$ such that for each time $t \in (0, T)$,
 $Q(t, \cdot) \in V_h = \left\{ v \in L^1(\Omega) : v|_{I_j} \in P^k(I_j) \right\}$

$$\begin{aligned} \int_{I_j} F(Q) v \, dx &= \int_{I_j} f(Q) v_x \, dx \\ &\quad - \left(\mathcal{F}_{j+1/2} v^-(x_{j+1/2}) - \mathcal{F}_{j-1/2} v^+(x_{j-1/2}) \right) \end{aligned}$$

for all $v \in V_h$

- Rusanov/Local Lax-Friedrichs Numerical Flux

$$\mathcal{F}_{j+1/2} = \frac{1}{2} \left(f(Q_{j+1/2}^-) + f(Q_{j+1/2}^+) \right) + \frac{1}{2} \max_q \{ |f'(q)| \} (Q_{j+1/2}^- - Q_{j+1/2}^+)$$

Diffusion

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

■ Diffusion Equation

$$G(q) = -(q^3 q_{xxx})_x \quad (0, T) \times \Omega$$

■ Local Discontinuous Galerkin

$$r = q_x$$

$$s = r_x$$

$$u = s_x$$

$$G(q) = (q^3 u)_x$$

Local Discontinuous Galerkin

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

Find $Q(t, x), R(x), S(x), U(x)$ such that for all $t \in (0, T)$

$$Q(t, \cdot), R, S, U \in V_h = \left\{ v \in L^1(\Omega) : v|_{I_j} \in P^k(I_j) \right\}$$

$$\int_{I_j} Rv \, dx = - \int_{I_j} Qv_x \, dx + \left(\hat{Q}_{j+1/2} v_{j+1/2}^- - \hat{Q}_{j-1/2} v_{j-1/2}^+ \right)$$

$$\int_{I_j} Sw \, dx = - \int_{I_j} R w_x \, dx + \left(\hat{R}_{j+1/2} w_{j+1/2}^- - \hat{R}_{j-1/2} w_{j-1/2}^+ \right)$$

$$\int_{I_j} Uy \, dx = - \int_{I_j} S y_x \, dx + \left(\hat{S}_{j+1/2} y_{j+1/2}^- - \hat{S}_{j-1/2} y_{j-1/2}^+ \right)$$

$$\int_{I_j} G(Q)z \, dx = - \int_{I_j} Q^3 U z_x \, dx + \left(\hat{U}_{j+1/2} z_{j+1/2}^- - \hat{U}_{j-1/2} z_{j-1/2}^+ \right)$$

for all $I_j \in \Omega$ and all $v, w, y, z \in V_h$.

Numerical Fluxes

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work


References

$$\hat{Q}_{j+1/2} = Q_{j+1/2}^+$$

$$\hat{R}_{j+1/2} = R_{j+1/2}^-$$

$$\hat{S}_{j+1/2} = S_{j+1/2}^+$$

$$\hat{U}_{j+1/2} = (Q^3 U)_{j+1/2}^-$$



IMEX Runge Kutta

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

■ IMEX scheme

$$\begin{aligned}q^{n+1} &= q^n + \Delta t \sum_{i=1}^s (b'_i F(t_i, u_i)) + \Delta t \sum_{i=1}^s (b_i G(t_i, u_i)) \\u_i &= q^n + \Delta t \sum_{j=1}^{i-1} (a'_{ij} F(t_j, u_j)) + \Delta t \sum_{j=1}^i (a_{ij} G(t_j, u_j)) \\t_i &= t^n + c_i \Delta t\end{aligned}$$

■ Double Butcher Tableaus

$$\begin{array}{c|c} c' & a' \\ \hline & b'^T \end{array} \quad \begin{array}{c|c} c & a \\ \hline & b^T \end{array}$$

■ 1st Order — L-Stable SSP

$$\begin{array}{c|c} 0 & 0 \\ \hline & 1 \end{array} \quad \begin{array}{c|c} 1 & 1 \\ \hline & 1 \end{array}$$

■ 2nd Order — SSP

$$\begin{array}{c|ccc} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ \hline & 0 & \frac{1}{2} & \frac{1}{2} \end{array} \quad \begin{array}{c|cc} \frac{1}{2} & \frac{1}{2} & 0 \\ 0 & -\frac{1}{2} & 0 \\ 1 & 0 & \frac{1}{2} \\ \hline & 0 & \frac{1}{2} \end{array}$$

■ 3rd Order — L-Stable SSP

| | | | | | | | | | |
|---------------|---|---------------|---------------|---------------|---------------|-----------|---------------|---------------|---------------|
| 0 | 0 | 0 | 0 | 0 | α | α | 0 | 0 | 0 |
| 0 | 0 | 0 | 0 | 0 | 0 | $-\alpha$ | α | 0 | 0 |
| 1 | 0 | 1 | 0 | 0 | 1 | 0 | $1 - \alpha$ | α | 0 |
| $\frac{1}{2}$ | 0 | $\frac{1}{4}$ | $\frac{1}{4}$ | 0 | $\frac{1}{2}$ | β | η | ζ | α |
| <hr/> | | | | | <hr/> | | | | |
| | 0 | $\frac{1}{6}$ | $\frac{1}{6}$ | $\frac{2}{3}$ | | 0 | $\frac{1}{6}$ | $\frac{1}{6}$ | $\frac{2}{3}$ |

$$\alpha = 0.24169426078821$$

$$\beta = 0.06042356519705$$

$$\eta = 0.1291528696059$$

$$\zeta = \frac{1}{2} - \beta - \eta - \alpha$$

Nonlinear Solvers

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

■ Nonlinear System

$$u_i - a_{ij}\Delta t G(u_i) = b$$

■ Picard Iteration

$$\tilde{G}(q, u) = (q^3 u_{xxx})_x$$

$$\begin{aligned} u_0 &= q^n & u_i^0 &= u_{i-1} \\ u_i^j - a_{ij}\Delta t \tilde{G}(u_i^{j-1}, u_i^j) &= b \end{aligned}$$

Manufactured Solution

$$q_t + (q^2 - q^3)_x = -(q^3 q_{xxx})_x + s$$

$$s = \hat{q}_t + (\hat{q}^2 - \hat{q}^3)_x + (\hat{q}^3 \hat{q}_{xxx})_x$$

$$\hat{q} = 0.1 \times \sin(2\pi/20.0 \times (x - t)) + 0.15 \quad \text{for } (x, t) \in [0, 40] \times [0, 5.0]$$

| | 1st Order | | 2nd Order | | 3rd Order | |
|------|-----------|-------|-----------------------|-------|-----------------------|-------|
| n | error | order | error | order | error | order |
| 20 | 0.136 | — | 7.33×10^{-3} | — | 5.29×10^{-4} | — |
| 40 | 0.0719 | 0.92 | 1.99×10^{-3} | 1.88 | 5.38×10^{-5} | 3.30 |
| 80 | 0.0378 | 0.93 | 5.60×10^{-4} | 1.83 | 7.47×10^{-6} | 2.85 |
| 160 | 0.0191 | 0.99 | 1.56×10^{-4} | 1.85 | 9.97×10^{-7} | 2.91 |
| 320 | 0.00961 | 0.99 | 3.98×10^{-5} | 1.97 | 1.26×10^{-7} | 2.98 |
| 640 | 0.00483 | 0.99 | 1.00×10^{-5} | 1.99 | 1.58×10^{-8} | 3.00 |
| 1280 | 0.00242 | 1.00 | 2.50×10^{-6} | 2.00 | 1.98×10^{-9} | 3.00 |

Table: Convergence table with a constant, linear, quadratic polynomial bases. CFL = 0.9, 0.2, 0.1 respectively.

Wave Structure with Nonlinear Hyper Diffusion

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model


Numerical Methods

Results

Future Work

References

$$q_t + (q^2 - q^3)_x = -(q^3 q_{xxx})_x$$
$$q_r = 0.1 \quad q_l = 0.3$$



Wave Structure with Nonlinear Hyper Diffusion

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods


Results

Future Work

References

$$q_r = 0.1 \quad q_l = 0.3323$$

$$q(x, 0) = (-\tanh(x) + 1) \frac{q_l - q_r}{2} + q_r$$



Wave Structure with Nonlinear Hyper Diffusion

Caleb Logemann,
James
Rossmanith


Thin Film
Equation
Model
Numerical Methods
Results

Generalized
Shallow Water
Equations

Model
Numerical Methods
Results
Future Work
References

$$q_r = 0.1 \quad q_l = 0.3323 \quad q_m = 0.6$$

$$q(x, 0) = \begin{cases} \frac{q_m - q_l}{2} \tanh(x) + \frac{q_m + q_l}{2} & x < 5 \\ -\frac{q_m - q_r}{2} \tanh(x - 10) + \frac{q_m + q_r}{2} + q_r & x > 5 \end{cases}$$



Wave Structure with Nonlinear Hyper Diffusion

Caleb Logemann,
James
Rossmanith


Thin Film
Equation
Model
Numerical Methods
Results

Generalized
Shallow Water
Equations

Model
Numerical Methods
Results
Future Work
References

$$q_r = 0.1 \quad q_l = 0.3323 \quad q_m = 0.6$$

$$q(x, 0) = \begin{cases} \frac{q_m - q_l}{2} \tanh(x) + \frac{q_m + q_l}{2} & x < 10 \\ -\frac{q_m - q_r}{2} \tanh(x - 20) + \frac{q_m + q_r}{2} + q_r & x > 10 \end{cases}$$



Wave Structure with Nonlinear Hyper Diffusion

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods


Results

Future Work

References

$$q_r = 0.1 \quad q_l = 0.4$$

$$q(x, 0) = (-\tanh(x - 100) + 1) \frac{q_l - q_r}{2} + q_r$$



Wave Structure with Nonlinear Hyper Diffusion

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods


Results

Future Work

References

$$q_r = 0.1 \quad q_l = 0.8$$

$$q(x, 0) = (-\tanh(x - 10) + 1) \frac{q_l - q_r}{2} + q_r$$



Generalized Shallow Water

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods
Results


Generalized
Shallow Water
Equations

Model

Numerical Methods
Results

Future Work

References



$$\nabla \cdot \mathbf{u} = 0$$

$$\mathbf{u}_t + \nabla \cdot (\mathbf{u}\mathbf{u}) = -\frac{1}{\rho} \nabla p + \frac{1}{\rho} \nabla \cdot \sigma + \mathbf{g}$$

$$(h_s)_t + [u(t, x, y, h_s), v(t, x, y, h_s)]^T \cdot \nabla h_s = w(t, x, y, h_s)$$

$$(h_b)_t + [u(t, x, y, h_b), v(t, x, y, h_b)]^T \cdot \nabla h_b = w(t, x, y, h_b)$$

Nondimensionalization

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

Characteristic Lengths

$$\varepsilon = \frac{H}{L}, \quad x = L\hat{x}, \quad y = L\hat{y}, \quad z = H\hat{z}$$

Characteristic Velocities

$$u = U\hat{u}, \quad v = U\hat{v}, \quad w = \varepsilon U\hat{w}$$

Characteristic Time

$$t = \frac{L}{U}\hat{t}$$

Characteristic Stresses

$$p = \rho g H \hat{p}, \quad \sigma_{xz/yz} = S \hat{\sigma}_{xz/yz}, \quad \sigma_{xx/xy/yy/zz} = \varepsilon S \hat{\sigma}_{xx/xy/yy/zz}$$

Nondimensionalization

Nondimensional Equations

$$\hat{u}_{\hat{x}} + \hat{v}_{\hat{y}} + \hat{w}_{\hat{z}} = 0$$

$$\begin{aligned} \varepsilon F^2 \left(\hat{u}_{\hat{t}} + (\hat{u}^2)_{\hat{x}} + (\hat{u}\hat{v})_{\hat{y}} + (\hat{u}\hat{w})_{\hat{z}} \right) &= -\varepsilon \hat{p}_{\hat{x}} \\ &+ G \left(\varepsilon^2 (\hat{\sigma}_{xx})_{\hat{x}} + \varepsilon^2 (\hat{\sigma}_{xy})_{\hat{y}} + (\hat{\sigma}_{xz})_{\hat{z}} \right) + e_x \end{aligned}$$

$$\begin{aligned} \varepsilon F^2 \left(\hat{v}_{\hat{t}} + (\hat{u}\hat{v})_{\hat{x}} + (\hat{v}^2)_{\hat{y}} + (\hat{v}\hat{w})_{\hat{z}} \right) &= -\varepsilon \hat{p}_{\hat{y}} \\ &+ G \left(\varepsilon^2 (\hat{\sigma}_{xy})_{\hat{x}} + \varepsilon^2 (\hat{\sigma}_{yy})_{\hat{y}} + (\hat{\sigma}_{yz})_{\hat{z}} \right) + e_y \end{aligned}$$

$$\begin{aligned} \varepsilon^2 F^2 \left(\hat{w}_{\hat{t}} + (\hat{u}\hat{w})_{\hat{x}} + (\hat{v}\hat{w})_{\hat{y}} + (\hat{w}^2)_{\hat{z}} \right) &= -\hat{p}_{\hat{z}} \\ &+ \varepsilon G \left((\hat{\sigma}_{xz})_{\hat{x}} + (\hat{\sigma}_{yz})_{\hat{y}} + (\hat{\sigma}_{zz})_{\hat{z}} \right) + e_z \end{aligned}$$

$$F = \frac{U}{\sqrt{gH}} \approx 1, \quad G = \frac{S}{\rho g H} < 1$$

Nondimensionalization

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

Drop ε^2 and εG terms

$$\hat{u}_{\hat{x}} + \hat{v}_{\hat{y}} + \hat{w}_{\hat{z}} = 0$$

$$\varepsilon F^2 \left(\hat{u}_{\hat{t}} + (\hat{u}^2)_{\hat{x}} + (\hat{u}\hat{v})_{\hat{y}} + (\hat{u}\hat{w})_{\hat{z}} \right) = -\varepsilon \hat{p}_{\hat{x}} + G(\hat{\sigma}_{xz})_{\hat{z}} + e_x$$

$$\varepsilon F^2 \left(\hat{v}_{\hat{t}} + (\hat{u}\hat{v})_{\hat{x}} + (\hat{v}^2)_{\hat{y}} + (\hat{v}\hat{w})_{\hat{z}} \right) = -\varepsilon \hat{p}_{\hat{y}} + G(\hat{\sigma}_{yz})_{\hat{z}} + e_y$$

$$\hat{p}_{\hat{z}} = e_z$$

Solving for the hydrostatic pressure

$$\hat{p}(\hat{t}, \hat{x}, \hat{y}) = \left(\hat{h}_s(\hat{t}, \hat{x}, \hat{y}) - \hat{z} \right) e_z$$

Nondimensionalization

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

Dimensional Variables

$$u_x + v_y + w_z = 0$$

$$u_t + (u^2)_x + (uv)_y + (uw)_z = -\frac{1}{\rho}p_x + \frac{1}{\rho}(\sigma_{xz})_z + ge_x$$

$$v_t + (uv)_x + (v^2)_y + (vw)_z = -\frac{1}{\rho}p_y + \frac{1}{\rho}(\sigma_{yz})_z + ge_y$$



$$p(t, x, y, z) = (h_s(t, x, y) - z)\rho ge_z$$

Kinematic Boundary Conditions

$$(h_s)_t + [u(t, x, y, h_s), v(t, x, y, h_s)]^T \cdot \nabla h_s = w(t, x, y, h_s)$$

$$(h_b)_t + [u(t, x, y, h_b), v(t, x, y, h_b)]^T \cdot \nabla h_b = w(t, x, y, h_b)$$

Mapping



Transform from $z \rightarrow \zeta$, by

$$\zeta = \frac{z - h_b(t, x, y)}{h(t, x, y)},$$

or equivalently

$$z = h(t, x, y)\zeta + h_b(t, x, y)$$

where $h(t, x, y) = h_s(t, x, y) - h_b(t, x, y)$.

$$\tilde{\Psi}(t, x, y, \zeta) = \Psi(t, x, y, h(t, x, y)\zeta + h_b(t, x, y))$$

Mapping Continuity Equation

$$u_x + v_y + w_z = 0$$

Map to new space

$$(h\tilde{u})_x - ((\zeta h + h_b)_x \tilde{u})_\zeta + (h\tilde{v})_y - ((\zeta h + h_b)_y \tilde{v})_\zeta + (\tilde{w})_\zeta = 0$$

Solve for vertical velocity, w ,

$$\begin{aligned} \tilde{w}(t, x, y, \zeta) = & - \left(h \int_0^\zeta \tilde{u} d\zeta' \right)_x - \left(h \int_0^\zeta \tilde{v} d\zeta' \right)_y \\ & + (\zeta h + h_b)_x \tilde{u}(t, x, y, \zeta) + (\zeta h + h_b)_y \tilde{v}(t, x, y, \zeta) \end{aligned}$$

Depth averaged equation

$$h_t + \left(h \int_0^1 \tilde{u} d\zeta \right)_x + \left(h \int_0^1 \tilde{v} d\zeta \right)_y = 0$$

Let u_m and v_m denote the mean velocity

$$h_t + (hu_m)_x + (hv_m)_y = 0$$

Mapping Momentum Equations

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

$$u_t + (u^2)_x + (uv)_y + (uw)_z = -\frac{1}{\rho}p_x + \frac{1}{\rho}(\sigma_{xz})_z + ge_x$$

$$v_t + (uv)_x + (v^2)_y + (vw)_z = -\frac{1}{\rho}p_y + \frac{1}{\rho}(\sigma_{yz})_z + ge_y$$

$$(h\tilde{u})_t + \left(h\tilde{u}^2 + \frac{1}{2}ge_z h^2\right)_x + (h\tilde{u}\tilde{v})_y + \left(h\tilde{u}\omega - \frac{1}{\rho}\tilde{\sigma}_{xz}\right)_\zeta = gh(e_x - e_z(h_b)_x)$$

$$(h\tilde{v})_t + (h\tilde{u}\tilde{v})_x + \left(h\tilde{v}^2 + \frac{1}{2}ge_z h^2\right)_y + \left(h\tilde{v}\omega - \frac{1}{\rho}\tilde{\sigma}_{yz}\right)_\zeta = gh(e_x - e_z(h_b)_y)$$

where

$$\omega = \frac{1}{h} \left(- \left(h \int_0^\zeta \tilde{u} - u_m d\zeta' \right)_x - \left(h \int_0^\zeta \tilde{v} - v_m d\zeta' \right)_y \right)$$

Mapped Reference System

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

$$h_t + (hu_m)_x + (hv_m)_y = 0$$

$$(h\tilde{u})_t + \left(h\tilde{u}^2 + \frac{1}{2}ge_z h^2\right)_x + (h\tilde{u}\tilde{v})_y + \left(h\tilde{u}\omega - \frac{1}{\rho}\tilde{\sigma}_{xz}\right)_\zeta = gh(e_x - e_z(h_b)_x)$$

$$(h\tilde{v})_t + (h\tilde{u}\tilde{v})_x + \left(h\tilde{v}^2 + \frac{1}{2}ge_z h^2\right)_y + \left(h\tilde{v}\omega - \frac{1}{\rho}\tilde{\sigma}_{yz}\right)_\zeta = gh(e_x - e_z(h_b)_y)$$

$$\omega = \frac{1}{h} \left(- \left(h \int_0^\zeta \tilde{u}_d \, d\zeta' \right)_x - \left(h \int_0^\zeta \tilde{v}_d \, d\zeta' \right)_y \right)$$

with

$$\tilde{u}_d = \tilde{u} - u_m \quad \tilde{v}_d = \tilde{v} - v_m$$

Newtonian Flow

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

Newtonian Stress Tensor

$$\sigma_{xz} = \mu u_z \quad \sigma_{yz} = \mu v_z$$

Kinematic Viscosity

$$\nu = \frac{\mu}{\rho}$$

Mapped stress tensor

$$\frac{1}{\rho} \tilde{\sigma}_{xz} = \frac{\nu}{h} \tilde{u}_\zeta \quad \frac{1}{\rho} \tilde{\sigma}_{yz} = \frac{\nu}{h} \tilde{v}_\zeta$$

Boundary Conditions

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

Stress Free Condition at surface

$$u_z|_{z=h_s} = v_z|_{z=h_s} = 0$$

Mixed Slip Condition at bottom topography

$$u - \frac{\lambda}{\mu} \sigma_{xz} \Big|_{z=h_b} = v - \frac{\lambda}{\mu} \sigma_{yz} \Big|_{z=h_b} = 0$$

Mapped with Newtonian Stress

$$\tilde{u}_\zeta|_{\zeta=1} = \tilde{v}_\zeta|_{\zeta=1} = 0$$

and

$$\tilde{u} - \frac{\lambda}{h} \tilde{u}_\zeta \Big|_{\zeta=0} = \tilde{v} - \frac{\lambda}{h} \tilde{v}_\zeta \Big|_{\zeta=0} = 0$$

Mapped Reference System

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

$$h_t + (hu_m)_x + (hv_m)_y = 0$$

$$(h\tilde{u})_t + \left(h\tilde{u}^2 + \frac{1}{2}ge_z h^2\right)_x + (h\tilde{u}\tilde{v})_y + \left(h\tilde{u}\omega - \frac{\nu}{h}\tilde{u}_\zeta\right)_\zeta = gh(e_x - e_z(h_b)_x)$$

$$(h\tilde{v})_t + (h\tilde{u}\tilde{v})_x + \left(h\tilde{v}^2 + \frac{1}{2}ge_z h^2\right)_y + \left(h\tilde{v}\omega - \frac{\nu}{h}\tilde{v}_\zeta\right)_\zeta = gh(e_x - e_z(h_b)_y)$$

Boundary Conditions

$$\tilde{u}_\zeta|_{\zeta=1} = \tilde{v}_\zeta|_{\zeta=1} = 0$$

and

$$\tilde{u} - \frac{\lambda}{h}\tilde{u}_\zeta\Big|_{\zeta=0} = \tilde{v} - \frac{\lambda}{h}\tilde{v}_\zeta\Big|_{\zeta=0} = 0$$

Moment Closure

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

Depth Averaged Momentum Equations

$$\begin{aligned} (hu_m)_t + \left(h \int_0^1 \tilde{u}^2 d\zeta + \frac{1}{2} g e_z h^2 \right)_x + \left(h \int_0^1 \tilde{u} \tilde{v} d\zeta \right)_y \\ + \frac{\nu}{\lambda} u|_{\zeta=0} = hg(e_x - e_z(h_b)_x) \\ (hv_m)_t + \left(h \int_0^1 \tilde{u} \tilde{v} d\zeta \right)_y + \left(h \int_0^1 \tilde{v}^2 d\zeta + \frac{1}{2} g e_z h^2 \right)_x \\ + \frac{\nu}{\lambda} v|_{\zeta=0} = hg(e_y - e_z(h_b)_y) \end{aligned}$$

Polynomial Ansatz

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

$$\tilde{u}(t, x, y, \zeta) = u_m(t, x, y) + u_d(t, x, y, \zeta)$$

$$= u_m(t, x, y) + \sum_{j=1}^N (\alpha_j(t, x, y) \phi_j(\zeta))$$

$$\tilde{v}(t, x, y, \zeta) = v_m(t, x, y) + v_d(t, x, y, \zeta)$$

$$= v_m(t, x, y) + \sum_{j=1}^N (\beta_j(t, x, y) \phi_j(\zeta))$$

Orthogonality Condition

$$\int_0^1 \phi_j(\zeta) \phi_i(\zeta) d\zeta = 0 \quad \text{for } j \neq i$$

$$\phi_0(\zeta) = 1, \quad \phi_1(\zeta) = 1 - 2\zeta, \quad \phi_2(\zeta) = 1 - 6\zeta + 6\zeta^2$$

Constant Moments

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

$$\begin{aligned}
 & (hu_m)_t + \left(h \left(u_m^2 + \sum_{j=1}^N \frac{\alpha_j^2}{2j+1} \right) + \frac{1}{2} g e_z h^2 \right)_x \\
 & + \left(h \left(u_m v_m + \sum_{j=1}^N \frac{\alpha_j \beta_j}{2j+1} \right) \right)_y = -\frac{\nu}{\lambda} \left(u_m + \sum_{j=1}^N \alpha_j \right) + hg (e_x - e_z (h_b)_x) \\
 & (hv_m)_t + \left(h \left(v_m^2 + \sum_{j=1}^N \frac{\alpha_j \beta_j}{2j+1} \right) + \frac{1}{2} g e_z h^2 \right)_y \\
 & + \left(h \left(u_m v_m + \sum_{j=1}^N \frac{\alpha_j \beta_j}{2j+1} \right) \right)_x = -\frac{\nu}{\lambda} \left(v_m + \sum_{j=1}^N \beta_j \right) + hg (e_y - e_z (h_b)_y)
 \end{aligned}$$

Higher Order Moments

Moment Equation

$$\int_0^1 \phi_i \left((h\tilde{u})_t + \left(h\tilde{u}^2 + \frac{1}{2} g e_z h^2 \right)_x + (h\tilde{u}\tilde{v})_y + \left(h\tilde{u}\omega - \frac{1}{\rho} \tilde{\sigma}_{xz} \right)_z \right) d\zeta$$

$$= \int_0^1 \phi_i (gh(e_x - e_z(h_b)_x)) d\zeta$$

Simplified gives

$$(h\alpha_i)_t + \left(2hu_m\alpha_i + h \sum_{j,k=1}^N A_{ijk}\alpha_j\alpha_k \right)_x$$

$$+ \left(hu_m\beta_i + hv_m\alpha_i + h \sum_{j,k=1}^N A_{ijk}\alpha_j\beta_k \right)_y$$

$$= u_mD_i - \sum_{j,k=1}^N B_{ijk}D_j\alpha_k - (2i+1)\frac{\nu}{\lambda} \left(u_m + \sum_{j=1}^N \left(1 + \frac{\lambda}{h} C_{ij} \right) \alpha_j \right)$$

Higher Order Moments

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

$$\begin{aligned}
 (h\beta_i)_t + \left(hu_m\beta_i + hv_m\alpha_i + h \sum_{j,k=1}^N A_{ijk}\alpha_j\beta_k \right)_x + \left(2hv_m\beta_i + h \sum_{j,k=1}^N A_{ijk}\beta_j\beta_k \right)_y \\
 = v_m D_i - \sum_{j,k=1}^N B_{ijk} D_j \beta_k - (2i+1) \frac{\nu}{\lambda} \left(v_m + \sum_{j=1}^N \left(1 + \frac{\lambda}{h} C_{ij} \right) \beta_j \right)
 \end{aligned}$$

$$A_{ijk} = (2i+1) \int_0^1 \phi_i \phi_j \phi_k \, d\zeta$$

$$B_{ijk} = (2i+1) \int_0^1 \phi'_i \left(\int_0^\zeta \phi_j \, d\hat{\zeta} \right) \phi_k \, d\zeta$$

$$C_{ij} = \int_0^1 \phi'_i \phi'_j \, d\zeta$$

$$D_i = (h\alpha_i)_x + (h\beta_i)_y$$

Example Systems

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

1D model with h_b constant, $e_x = e_y = 0$, and $e_z = 1$
Constant System

$$\begin{bmatrix} h \\ hu_m \end{bmatrix}_t + \begin{bmatrix} hu_m \\ hu_m^2 + \frac{1}{2}gh^2 \end{bmatrix}_x = -\frac{\nu}{\lambda} \begin{bmatrix} 0 \\ u_m \end{bmatrix}$$

Linear System, $\tilde{u} = u_m + s\phi_1$, $s = \alpha_1$

$$\begin{bmatrix} h \\ hu_m \\ hs \end{bmatrix}_t + \begin{bmatrix} hu_m \\ hu_m^2 + \frac{1}{2}gh^2 + \frac{1}{3}hs^2 \\ 2hu_ms \end{bmatrix}_x = Q \begin{bmatrix} h \\ hu_m \\ hs \end{bmatrix}_x - P$$

$$Q = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & u_m \end{bmatrix} \quad P = \frac{\nu}{\lambda} \begin{bmatrix} 0 \\ u_m + s \\ 3(u_m + s + 4\frac{\lambda}{h}s) \end{bmatrix}$$

Numerical Methods

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

Model Equation

$$\mathbf{q}_t + \mathbf{f}(\mathbf{q})_x = Q(\mathbf{q})\mathbf{q}_x - \mathbf{P}(\mathbf{q}) \quad \text{for } (x, t) \in [a, b] \times [0, T]$$

Weak Form, find \mathbf{q} such that

$$\int_a^b \mathbf{q}_t v \, dx + \int_a^b \mathbf{f}(\mathbf{q})_x v \, dx = \int_a^b Q(\mathbf{q})\mathbf{q}_x v \, dx - \int_a^b \mathbf{P}(\mathbf{q}) v \, dx$$

for all $v \in L^2([a, b] \times [0, T])$

Notation

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results


Future Work

References

- Partition the domain, $[a, b]$ as

$$a = x_{1/2} < \cdots < x_{j-1/2} < x_{j+1/2} < \cdots < x_{N+1/2} = b$$

- $I_j = [x_{j-1/2}, x_{j+1/2}]$
- $x_j = \frac{x_{j+1/2} + x_{j-1/2}}{2}$
- $\Delta x_j = x_{j+1/2} - x_{j-1/2}$
- $\Delta x_j = \Delta x$ for all j .



Discontinuous Galerkin Space

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

Finite Dimensional DG Space

$$V^k = \left\{ v \in L^2([a, b]) \mid v|_{I_j} \in P^k(I_j) \right\}$$

Basis for V^k

$$\{\phi_j^\ell\} \text{ where } \phi_j^\ell(x)|_{I_j} = \phi^\ell(\xi_j(x)) \text{ and } \phi_j^\ell(x)|_{\bar{I}_j} = 0$$

for $j = 1, \dots, N$ and $\ell = 1, \dots, k$.

Legendre Polynomials

$$\phi^k \in P^k([-1, 1]) \text{ with } \frac{1}{2} \int_{-1}^1 \phi^k(\xi) \phi^\ell(\xi) d\xi = \delta_{k\ell}$$

and

$$\xi_j(x) = \frac{2}{\Delta x_j}(x - x_j)$$

Numerical Methods

Find $\mathbf{q}_h \in V^k$ such that

$$\begin{aligned} \int_{I_j} (\mathbf{q}_h)_t \phi_j^\ell(x) \, dx &= \int_{I_j} \mathbf{f}(\mathbf{q}_h)_x \phi_j^\ell \, dx \\ &\quad - F_{j+1/2} \phi_j^\ell(x_{j+1/2}) + F_{j-1/2} \phi_j^\ell(x_{j-1/2}) \\ &\quad + \int_{I_j} Q(\mathbf{q}_h)(\mathbf{q}_h)_x \phi_j^\ell \, dx - \int_{I_j} \mathbf{P}(\mathbf{q}_h) \phi_j^\ell \, dx \end{aligned}$$

for all ϕ_j^l .

Local Lax-Friedrichs Flux

$$\mathbf{q}_h^+ = \lim_{x \rightarrow x_{j+1/2}^+} (\mathbf{q}_h(x))$$

$$\mathbf{q}_h^- = \lim_{x \rightarrow x_{j+1/2}^-} (\mathbf{q}_h(x))$$

$$\lambda = \max_{\mathbf{q} \in [\mathbf{q}_h^-, \mathbf{q}_h^+]} \{ \rho(\mathbf{f}'(\mathbf{q}) - Q(\mathbf{q})) \}$$

$$F_{j+1/2} = \frac{1}{2} (\mathbf{f}(\mathbf{q}_h^+) + \mathbf{f}(\mathbf{q}_h^-)) - \frac{1}{2} \lambda (\mathbf{q}_h^+ - \mathbf{q}_h^-)$$

Nonconservative Flux

Need to evaluate

$$\int^{I_j} Q \mathbf{q}_x \phi_j^\ell dx$$

$$\mathbf{q}|_{I_j} = \sum_{\ell=1}^k (Q_j^\ell \phi_j^\ell(x)), \quad \mathbf{q}_x|_{I_j} = \sum_{\ell=1}^k (Q_x^\ell \phi_j^\ell(x))$$

where

$$\begin{bmatrix} Q_x^1 \\ Q_x^2 \\ Q_x^3 \\ Q_x^4 \\ Q_x^5 \end{bmatrix} = \frac{1}{2\Delta x} \begin{bmatrix} \Delta Q^1 - 2\sqrt{5}\Delta Q^3 + 78\Delta Q^5 \\ \Delta Q^2 - \frac{10}{3}\sqrt{3}\sqrt{7}\Delta Q^4 \\ \Delta Q^3 - 14\sqrt{5}\Delta Q^5 \\ \Delta Q^4 \\ \Delta Q^5 \end{bmatrix}$$

$$\Delta Q^\ell = Q_{i+1}^\ell - Q_{i-1}^\ell$$

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

Inviscid Example

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References




$$x \in [-1, 1] \quad t \in [0, 2.0]$$

$$h(t = 0, x) = 1 + e^{3 \cos(\pi(x+0.5))} - 4$$

$$\tilde{u}(t = 0, x) = \begin{cases} 0.25 & \text{constant} \\ 0.5\zeta & \text{linear} \end{cases}$$

$$u_m = 0.25$$

$$s = -0.25$$



Inviscid Example

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations






Model

Numerical Methods

Results

Future Work

References



Higher Moment Equations

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations

Model

Numerical Methods

Results

Future Work

References

1 dimensional with h_b constant, $e_x = e_y = 0$, and $e_z = 1$ Quadratic Vertical Profile, $\tilde{u} = u_m + s\phi_1 + \kappa\phi_2$

$$\begin{bmatrix} h \\ hu \\ hs \\ h\kappa \end{bmatrix}_t + \begin{bmatrix} hu \\ hu^2 + \frac{1}{2}gh^2 + \frac{1}{3}hs^2 + \frac{1}{5}h\kappa^2 \\ 2hus + \frac{4}{5}hs\kappa \\ 2hu\kappa + \frac{2}{3}hs^2 + \frac{2}{7}h\kappa^2 \end{bmatrix}_x = Q \begin{bmatrix} h \\ hu \\ hs \\ h\kappa \end{bmatrix}_x - P$$

$$Q = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & u_m - \frac{\kappa}{5} & \frac{s}{5} \\ 0 & 0 & s & u_m + \frac{\kappa}{7} \end{bmatrix} \quad P = \frac{\nu}{\lambda} \begin{bmatrix} 0 \\ u_m + s + \kappa \\ 3(u_m + s + \kappa + 4\frac{\lambda}{h}s) \\ 5(u_m + s + \kappa + 12\frac{\lambda}{h}\kappa) \end{bmatrix}$$

Future Work

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods
Results

Generalized
Shallow Water
Equations

Model

Numerical Methods
Results

Future Work

References

- Higher Order Numerical Methods
- Slope Limiters
- Two Dimensional Meshes
- Icosahedral Spherical Mesh
- Positivity Preserving Limiters

Icosahedral Mesh

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations


Model

Numerical Methods

Results

Future Work

References



Spherical Test Cases

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations



Model

Numerical Methods

Results

Future Work

References



Spherical Test Cases

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods
Results


Generalized
Shallow Water
Equations

Model

Numerical Methods
Results

Future Work

References



Spherical Test Cases

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods

Results

Generalized
Shallow Water
Equations


Model

Numerical Methods

Results

Future Work

References



Bibliography I

Caleb Logemann,
James
Rossmanith

Thin Film
Equation

Model

Numerical Methods
Results

Generalized
Shallow Water
Equations

Model

Numerical Methods
Results

Future Work

References

- [1] Andrea L Bertozzi, Andreas Münch, and Michael Shearer. “Undercompressive shocks in thin film flows”. In: *Physica D: Nonlinear Phenomena* 134.4 (1999), pp. 431–464.
- [2] Bernardo Cockburn and Chi-Wang Shu. “The local discontinuous Galerkin method for time-dependent convection-diffusion systems”. In: *SIAM Journal on Numerical Analysis* 35.6 (1998), pp. 2440–2463.
- [3] Bernardo Cockburn and Chi-Wang Shu. “The Runge–Kutta discontinuous Galerkin method for conservation laws V: multidimensional systems”. In: *Journal of Computational Physics* 141.2 (1998), pp. 199–224.
- [4] Y. Ha, Y.-J. Kim, and T.G. Myers. “On the numerical solution of a driven thin film equation”. In: *J. Comp. Phys.* 227.15 (2008), pp. 7246–7263.

Bibliography II

- [5] Julia Kowalski and Manuel Torrilhon. “Moment Approximations and Model Cascades for Shallow Flow”. In: *arXiv preprint arXiv:1801.00046* (2017).
- [6] Randall J LeVeque et al. *Finite volume methods for hyperbolic problems*. Vol. 31. Cambridge university press, 2002.
- [7] T.G. Myers and J.P.F. Charpin. “A mathematical model for atmospheric ice accretion and water flow on a cold surface”. In: *Int. J. Heat and Mass Transfer* 47.25 (2004), pp. 5483–5500.
- [8] Tim G Myers. “Thin films with high surface tension”. In: *SIAM review* 40.3 (1998), pp. 441–462.
- [9] NASA. URL: http://icebox.grc.nasa.gov/gallery/images/C95_03918.html.
- [10] Alexander Oron, Stephen H Davis, and S George Bankoff. “Long-scale evolution of thin liquid films”. In: *Reviews of modern physics* 69.3 (1997), p. 931.

Caleb Logemann,
James
Rossmanith

Thin Film
Equation
Model
Numerical Methods
Results

Generalized
Shallow Water
Equations
Model
Numerical Methods
Results
Future Work

References

Bibliography III

Caleb Logemann,
James
Rossmanith

Thin Film
Equation
Model
Numerical Methods
Results

Generalized
Shallow Water
Equations
Model
Numerical Methods
Results
Future Work

References

- [11] J.A. Rossmanith. DOGPack. Available from <http://www.dogpack-code.org/>.
- [12] James A Rossmanith. "A wave propagation method for hyperbolic systems on the sphere". In: *Journal of Computational Physics* 213.2 (2006), pp. 629–658.
- [13] David L Williamson et al. "A standard test set for numerical approximations to the shallow water equations in spherical geometry". In: *Journal of Computational Physics* 102.1 (1992), pp. 211–224.